

Please cite the Published Version

Brearley, FQ (2011) Below-ground secondary succession in tropical forests of Borneo. *Journal of Tropical Ecology*, 27 (4). pp. 413-420. ISSN 0266-4674

DOI: <https://doi.org/10.1017/S0266467411000149>

Publisher: Cambridge University Press

Version: Accepted Version

Downloaded from: <https://e-space.mmu.ac.uk/475/>

Usage rights: © In Copyright

Additional Information: This is an author accepted manuscript of a paper accepted for publication in *Journal of Tropical Ecology*, published by and copyright Cambridge University Press.

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

Below-ground secondary succession in tropical forests of Borneo

Francis Q. Brearley^{1,2}

¹ Department of Botany, Trinity College, University of Dublin, Dublin 2, Ireland

² School of Science and the Environment, Manchester Metropolitan University, Chester Street, Manchester, M1 5GD, UK, +44 (0)161 2471717, f.q.brearley@mmu.ac.uk

ABSTRACT

As the destruction and severe disturbance of primary tropical forest continues, it is important to understand how these forests may recover from perturbations. Considerable work has been done on above-ground recovery but below-ground processes are less well understood. To determine changes in root mass during tropical secondary succession in lowland forests of Central Borneo, samples were taken from stands of increasing ages since abandonment of agriculture (1, 3, 14 & 31 y) with a primary forest control (six plots from 1-y-old stands and three from all other ages). Root mass and elemental concentrations were determined and soils were chemically analysed. There was no increase in root mass with stand age for fine-root (<2 mm diameter) or small-root (<5 mm diameter) mass but there was a trend for coarse-root mass (5-10 mm diameter) to increase with stand age. Negative correlations were shown between root mass and soil nutrient status. Fine-root C concentrations increased with stand age but there was no clear effect of stand age on fine-root N or P. Fine-root mass did not increase significantly with stand age suggesting a rapid recovery; instead, soil nutrient status appeared to be the most important factor controlling root mass. Of the soil nutrients measured in this study, N had a stronger control over root mass than P suggesting that this element may be limiting during secondary succession in tropical lowland forests of Borneo.

Keywords: fine-root biomass, Indonesia, nitrogen, shifting cultivation, soils, tropical lowland evergreen rain forest

Running head: Roots in tropical secondary forests

Text pages: **15**; Tables: **3**; Figures: **2**

INTRODUCTION

With the continued disturbance and destruction of tropical forests, secondary forests now comprise an expanding proportion of all tropical forest land and understanding how their ecological processes recover from disturbances (e.g. logging, shifting cultivation or burning) is therefore becoming increasingly important (Brown & Lugo 1990, Chazdon *et al.* 2007, Corlett 1995, Finegan 1996, Guariguata & Ostertag 2001). In Kalimantan (Indonesian Borneo) shifting cultivation is still an important form of agriculture (de Jong 1997, Lawrence *et al.* 1998, Nagy & Proctor 1999) and creates a mosaic of agricultural fields and secondary forest stands of various ages. A reasonable body of work has accrued describing the recovery of tropical forests following shifting cultivation but this has focussed mainly on above-ground processes (reviews by Brown & Lugo 1990, Chazdon *et al.* 2007, Guariguata & Ostertag 2001). Generally, there is an increase in above-ground biomass and species diversity as the stands age. However, much less work has been done on examining below-ground processes during tropical secondary succession or on the accumulation of carbon (C) and nutrients in roots, soils or mycorrhizal fungi.

Roots comprise up to 25% of total plant biomass in lowland tropical forests (Cairns *et al.* 1997, Sanford & Cuevas 1996) and are essential in structural support and water and nutrient uptake, transport and storage. They are therefore equally as important as above-ground

components for the functioning of the ecosystem. During root decomposition, C and nutrients will be returned to the soil; therefore, to improve our knowledge of elemental cycling in tropical secondary forests we need to understand root biomass and distribution in secondary forest soils. Studies examining the effects of secondary succession on root biomass in tropical forests have mainly been conducted in montane forests. For example, Berish (1982) and Hertel *et al.* (2003) showed large increases in root mass during secondary succession in forests of Costa Rica but, in contrast, Cavelier *et al.* (1996) found no change in root mass in Colombian successional forests. Indeed, it has been suggested by Raich (1980) that ‘fine-roots regrow rapidly after forest felling’ as he found that fine-root mass of a 1-y-old successional forest in lowland Costa Rica was not different from that in an undisturbed primary forest.

Negative correlations between soil nutrient status and fine-root mass have been found across a range of Australian and Neotropical forests (Maycock & Congdon 2000, Powers *et al.* 2005) but there are few comparable smaller-scale studies which have determined the relative importance of soil nutrient status on root mass at the landscape scale or in secondary successional forests (but see Ostertag 1998, 2001). Soil nutrient status is also likely to have an effect on root nutrient status and, given that foliar nutrient concentrations have been reported to decrease during secondary succession (Bonafant *et al.* 2007, Ellsworth & Reich 1996, Reich *et al.* 1995), it is of interest to determine if similar changes in nutrient concentrations occur in roots during secondary succession. Therefore, in terms of understanding C and nutrient return to the soil, it is important to understand the factors controlling root nutrient concentrations in more detail (Gordon & Jackson 2000).

In this study, I examine changes in root mass and element concentrations along a successional chronosequence of secondary forest stands (1-31 y, plus primary forest) and test the following

hypotheses: (1) root mass increases with stand age, (2) root element concentrations increase with stand age, and (3) there is a negative relationship between soil nutrient status and root mass.

METHODS

Study area

The Project Barito Ulu (PBU) research area is situated in Central Kalimantan, Indonesia, at 114°0'E, 0°06'S, in the centre of the island of Borneo and is within the Heart of Borneo protected area. The research area contains a range of forest types including tropical lowland evergreen rain forest and heath forest (kerangas) as well as areas of shifting cultivation fallows of various ages (Brearley *et al.* 2004, Mirmanto *et al.* 1999, Nagy & Proctor 1999, Prajadinata 1996). The geology is based on a Tertiary sedimentary formation which has given rise to sandy ultisols that are acidic and low in nutrients (Brearley *et al.* 2004, Mirmanto *et al.* 1999). The base-camp is around 150 m asl. Mean maximum and mean minimum temperatures are 34.0°C and 22.6°C with mean annual rainfall around 3800 mm with no month having a mean of less than 200 mm of rain; the climate is therefore considered to be perhumid aseasonal. However, there are consistent annual fluctuations in rainfall with the wettest months being November to April (with the exception of February) and the driest months being June to September (Brearley *et al.* 2007).

Forest plots

The plots were situated in stands of various ages along the Busang and Joloi rivers; all were within 10 km of the PBU base-camp. In each stand (with the exception of the 1-y-old stands) three plots of the same age were set up in representative areas of vegetation. For the 1-y-old stands, two plots were set up at each of three separate stands (Table 1). The experimental

design was therefore somewhat pseudo-replicated. The primary forest plots are those of Brearley *et al.* (2004), the 31-y-old and 14-y-old plots were set up by Prajadinata (1996) at an earlier stage of succession, and the 3-y-old and 1-y-old plots were set up specifically for this study. Previous crops would have included rice (*Oryza* sp.), maize (*Zea* sp.), cassava (*Manihot* sp.), and a small range of vegetables. Sites were burnt before cultivation, the length of which is normally 2 y (Brearley *et al.* 2004). Unfortunately, the number of times the sites were cultivated prior to secondary succession is not known in most cases.

The diameter at breast height (dbh; 1.3 m) of all trees and lianas greater than a given dbh within each plot was recorded using standard methodologies (Brearley *et al.* 2004). Differing number and sizes of plots and minimum tree dbh for enumeration depended upon the age, and therefore heterogeneity, of the vegetation (Table 1). There was a general increase in basal area with stand age (Table 1) and also in mean tree height (pers. obs.); hence stand biomass increased with age since abandonment of agriculture.

Roots

Sampling and mass Locations for root coring were selected in a stratified random fashion within each plot. Increasing numbers of cores were taken from plots of older age due to increasing heterogeneity of the vegetation (Table 1). The unconsolidated surface litter layer was gently removed and a corer (internal diameter 4.2 cm) was inserted into the soil to a given depth. The core containing the soil and roots was then extracted. There were two sets of cores: one set was taken to a depth of 10 cm and another set was taken to a depth of 20 cm (in separate randomisations). Cores were soaked in stream water overnight (there is no indication that this process affects root nitrogen or phosphorus; Green 1992) and soil was then washed from the samples through a 2-mm sieve over a 0.5-mm sieve in order to extract the roots

which were retained on the sieves. Roots were then picked off the sieves using forceps with no attempt made to separate live and dead roots. All samples were processed within 24 h. Roots from 20-cm depth were dried in the sun in the field, returned to Ireland, and then dried at 75°C for 96 h. Roots from 10-cm depth were stored in 70% ethanol, returned to Ireland, and then dried as above. The roots were divided into three size classes: <2 mm (fine roots), 2-5 mm (small roots), and 5-10 mm (coarse roots) and each size class weighed separately.

Carbon and nutrient concentrations Root C and nutrient concentrations were determined on the roots to 20-cm depth. Each sample was ground in liquid nitrogen (N) and homogenised prior to analysis. In some cases, for roots <2 mm diameter, there was insufficient material for analysis so 50 out of 61 samples were analysed for C and N and 59 out of 61 samples were analysed for phosphorus (P). Carbon and N concentrations were analysed on c. 0.15-g subsamples using a LECO CNS-1000 elemental analyser. For P, c. 0.5-g subsamples (plus two or three anti-bumping granules) were refluxed in 10 ml concentrated nitric acid at 190°C for about 5 h. They were then made up to 50 ml with deionised water and P was determined colorimetrically using molybdenum blue methodology on a Shimadzu UV-1601 spectrophotometer. For roots 2-5 mm in diameter, samples were composited to provide sufficient material (usually two original samples from the same plot were bulked to give one new sample) for a total of 28 samples. These were then prepared and analysed for C, N, and P as above.

Soils

Soil samples were taken from the faces of the holes used to take the roots to 10 cm depth. They were air-dried in the sun and packed in plastic bags in the field for return to Ireland where they were sieved to pass a 2-mm mesh. Sample pH was measured by adding 10 g of soil to 25

ml of distilled water. It was stirred and left to equilibrate for 1 h before measurement with a pH meter (pH 510, Eutech Instruments). Total C and N were analysed on *c.* 0.2-g subsamples using a LECO CNS-1000 elemental analyser. Phosphorus was extracted from *c.* 5-g subsamples with 20 ml Modified Kelowna reagent (Ashworth & Mrazek 1995) by shaking them on a rotary shaker for 30 min. Phosphorus was then determined colorimetrically using molybdenum blue methodology on a Hitachi U-1100 spectrophotometer. Moisture content of the air-dried soil was determined by heating *c.* 2.5-g subsamples to 105°C for 24 h and all results are expressed on a soil oven-dry basis.

Statistics

Changes in root mass, and C and nutrient concentrations with stand age (primary forest was set at an arbitrary 500 y old) were analysed using linear mixed-effects models with the stand considered as a random effect using R 2.4.1; Box-Cox transformations were carried out as appropriate. Pearson's correlation coefficients were calculated between root and soil characteristics using Minitab 15.1; for roots to 10-cm depth this was done on an individual core basis, for roots to 20-cm depth it was done on a plot basis as soil samples were taken to a depth of 10-cm only.

RESULTS

Soils

There was a trend of increasing soil acidity as succession proceeded (Table 2, $P < 0.10$). The 31-y-old and 3-y-old stands had the highest concentrations of soil C and N, with the lowest concentrations found in the 14-y-old and 1-y-old stands. Primary forest generally had intermediate values for both of these elements (Table 2). There were no significant changes in extractable soil P during succession (Table 2).

Root mass

For roots to a depth of 10 cm, there was no significant increase with stand age for those <2 mm diameter, those <5 mm diameter, or the total root mass (Figure 1). For roots 5-10 mm diameter, the primary forest had a six-fold increase in mass relative to the 1-y-old stand although this was not significant due to large variation (Figure 1).

For roots to a depth of 20 cm, again, there was no significant increase with stand age for those <2 mm diameter, those <5 mm diameter, or the total root mass (Figure 1). For roots 5-10 mm diameter, the primary forest had an eight-fold increase in mass relative to the 1-y-old stand although, again, this was not significant due to large variation (Figure 1).

Root carbon and nutrient concentrations

For fine roots <2 mm diameter, C concentrations showed no increase with increasing stand age ($P < 0.10$), and small-roots 2-5 mm diameter also showed no increase in C concentration with stand age (Figure 2). For fine roots, N concentrations were greatest in the 31-y-old stand; for small-roots, N concentrations were greatest in the 1-y-old stand (Figure 2). There was no effect of stand age on root P concentrations (Figure 2). Root N and P concentrations were greater in fine roots <2 mm diameter when compared with small roots 2-5 mm diameter but the opposite pattern was seen for root C concentrations which were greater in the 2-5-mm-diameter roots (t -test, $P < 0.001$ in all three cases).

Correlations between root mass and soil characteristics

The only significant negative correlation between root mass and soil elemental concentration was for fine roots < 2 mm diameter to 20-cm depth and soil N ($P < 0.05$; Table 3).

DISCUSSION

Root mass

Fine- and small-root mass was not significantly different between stands suggesting that these roots recover rapidly after forest disturbance (within 1 y). Following this initial recovery, root mass appears to be controlled more by soil fertility than stand age. However, an assertion of rapid regrowth assumes that root mass in agricultural fields is lower than in secondary forests which may not always be the case (Powers 2004), and was not measured in this study. Above-ground biomass increased by a factor of at least thirty during secondary succession (F. Q. Brearley, unpubl. data) whereas below-ground root mass did not increase, indicating that trees are allocating proportionately more resources below-ground (to roots) than above-ground during the early stages of secondary succession.

Raich (1980) found rapid recovery of fine-root mass within 1 y and Cavelier *et al.* (1996) found no effect of forest stand age on root mass. In contrast, Berish (1982), Hertel *et al.* (2003) and Muthukumar *et al.* (2003) all showed root mass to increase with stand age. Whilst Hertel *et al.* (2006) found a significant positive correlation ($r^2 = 0.32$) between stand age and total fine-root biomass across a number of studies, they also found no significant differences in total root mass between secondary forest stands of various ages and primary forest (D. Hertel, pers. comm.). The above contrasts show that the relationship between successional stand age and root mass is not always clear. It should be noted that some of the stand ages in this study were pseudoreplicated which limits the generalisation of the results as differences detected in root characteristics may be controlled not only by the stand age but also differences between stands in terms of soils, hydrology or site history. Nevertheless, this was

the only design possible to include older stands of a known age as secondary forests are usually re-cut within 15 y in the Barito Ulu area (pers. obs.).

When compared with other tropical forest data sets, the Barito Ulu mean root mass (658 g m^{-2} for roots $<2 \text{ mm}$ diameter to 20 cm depth) was well above the mean values presented elsewhere. For example, Jackson *et al.* (1997) found a mean value of 570 g m^{-2} although the majority of the studies in that paper examined roots to a greater depth than I did. The Barito Ulu values are also at least 2 SE greater than the mean presented by Hertel & Leuschner (2010) of $451 \pm 45 \text{ g m}^{-2}$ (live root mass only) but their studies were generally to $40\text{--}50 \text{ cm}$ depth. Of course, differences in sampling depth, inclusion of live/dead roots, and differing root diameters make comparisons between studies difficult but, where strict comparisons were made, the Barito Ulu values were still higher than values from many other forests (Cavelier *et al.* 1999, Muthukumar *et al.* 2003, Powers *et al.* 2005, Raich 1980, Yamashita *et al.* 2003).

Seasonal variation in root mass and nutrient concentration has been found in a similar site at Danum Valley in Sabah by Green *et al.* (2005) who showed root mass to be lower in the drier season. I examined root mass in the drier season at Barito Ulu and my results may therefore underestimate the maximum root mass to a certain degree. I also did not separate live and dead root mass in this study (estimates of dead root mass in similar forests are $<5\%$: Green *et al.* 2005; $<15\%$: Powers *et al.* 2005). I hypothesise that root turnover is more rapid in the younger secondary stands (as trees grow faster and die younger) and therefore a larger proportion of roots in the younger stands may have been dead. There may also have been roots of some crop species remaining in the soil which had not fully decomposed over the intervening 1-y period although this is unlikely. Finally, as samples were only taken to 20 cm

depth this may have underestimated root mass in the older stands in relation to the younger stands as I hypothesise that the maximum rooting depth would increase with stand age.

Relationship between root mass and soil fertility

When nutrients are limiting to growth, trees may be expected to allocate resources preferentially below-ground in order to increase their growth and/or reproductive rates. Lack of a particular nutrient in a soil may therefore lead to increased root biomass if that nutrient is considered limiting to growth (Bloom *et al.* 1985). Previous studies found negative correlations between root mass and soil N (Maycock & Congdon 2000, Powers *et al.* 2005), P (Gower 1987, Powers *et al.* 2005) and Ca (Gower 1987). It was interesting to note that soil fertility had a greater effect on root mass than did stand age and this was also shown by Powers (pers. comm.) for tropical dry forests in Costa Rica. Negative correlations between root mass and soil fertility suggests nutrient limitations to tree growth in these successional forests. The only significant correlation was with soil N, this is likely due to the volatilisation of N during biomass burning whereas P and cations are returned to the soil in ash (Kleinman *et al.* 1995) which will also raise soil pH. Davidson *et al.* (2007) have recently shown that secondary forest stands in the Amazon cycle N more conservatively than mature forests lending weight to the hypothesis of N limitation in these successional forests.

Root carbon and nutrients

The most notable change in root element concentration during succession was an increase in root C for roots <2 mm diameter with primary forest roots having about 40% C compared to about 28% C in 1-y-old-stand roots. Jaramillo *et al.* (2003) found a similar, but less pronounced, pattern for roots in Mexican forests and pastures. These differences in C concentration are probably due to differences in root age and/or morphological differences

between early- and late-successional species, with older and late-successional species (and larger diameter roots) containing a higher concentration of C due to a higher proportion of woody tissue (with a greater proportion of structural components) and fewer younger cells with higher nutrient concentrations.

There was no clear effect of succession on root N concentrations with differing root diameters having greater concentrations at differing ages. If it were possible to separate out the roots of certain common species it may have been easier to detect any potential trends by avoiding the confounding factor of changing tree species composition during succession. There was no difference in root P concentration with different stand ages. Arunachalam *et al.* (1997) also found no notable effects of secondary forest stand age on root nutrient concentrations.

Root nutrient concentrations, especially P, were low when compared with data presented by Gordon & Jackson (2000) and Jackson *et al.* (1997). The value for root P is likely to be a slight underestimate as the recovery of P by nitric acid digestion in our Dublin laboratory is around 87% (M. Kavanagh & F. Q. Brearley unpubl. data). Seasonal variation in root nutrient concentration has also been found at Danum Valley in Sabah by Green *et al.* (2005) who showed that roots extracted in the drier season had higher N concentrations (1.66% in the drier season vs. 1.41% yearly mean), but lower P concentrations (0.028% in the drier season vs. 0.049% yearly mean). Concentrations of N and P in this study decreased with increasing root diameter in agreement with other studies (Arunachalam *et al.* 1997, Gordon & Jackson 2000, Soethe *et al.* 2007).

Conclusions

Fine roots regrew rapidly during secondary succession in this ecosystem indicating that above-ground disturbance does not always have a long lasting effect on fine-root biomass. In these successional forest stands, soil N was the factor with the strongest control over fine-root mass suggesting that this element may be limiting in secondary forests of central Borneo.

ACKNOWLEDGEMENTS

This research results from work undertaken by the joint cooperative project (Project Barito Ulu), established between the Ministry of Forestry, Indonesia and the University of Cambridge, UK. I thank the Carnegie Trust for the Universities of Scotland and the Trinity College Dublin Association and Trust for their financial support; Kursani and Afri Mardinata for field assistance; Rupert Ridgeway for logistical support in Indonesia; Lembaga Ilmu Pengetahuan Indonesia for research permission; Muhammad Mansur (Herbarium Bogoriense) for acting as my research counterpart; Mark Kavanagh, Fran O'Donovan and Jacqueline Stone for technical assistance; Robert Bagchi for statistical advice; and Dietrich Hertel, Jennifer Powers and journal reviewers for their helpful comments on the manuscript.

LITERATURE CITED

- ARUNACHALAM, A., PANDEY, H. N., MAITHANI, K. & TRIPATHI, R. S. 1997. Organic matter and nutrient dynamics of fine and coarse roots during recovery of a felled subtropical humid forest. *Russian Journal of Ecology* 28:380-385.
- ASHWORTH, J. & MRAZEK, K. 1995. "Modified Kelowna" test for available phosphorus and potassium in soil. *Communications in Soil Science and Plant Analysis* 26:731-739.
- BERISH, C. W. 1982. Root biomass and surface area in three successional tropical forests. *Canadian Journal of Forest Research* 12:699-704.

- BLOOM, A. J., CHAPIN, F. S. & MOONEY, H. A. 1985. Resource limitation in plants: an economic analogy. *Annual Review of Ecology and Systematics* 16:363-392.
- BONAL, D., BORN, C., BRECHET, C., COSTE, S., MARCON, E., ROGGY, J.-C. & GUEHL, J.-M. 2007. The successional status of tropical rain forest tree species is associated with differences in leaf carbon isotope discrimination and functional traits. *Annals of Forest Science* 64:169-176.
- BREARLEY, F. Q., PRAJADINATA, S., KIDD, P. S., PROCTOR, J. & SURIANTATA 2004. Structure and floristics of an old secondary rain forest in Central Kalimantan, Indonesia, and a comparison with adjacent primary forest. *Forest Ecology and Management* 195:385-397.
- BREARLEY, F. Q., PROCTOR, J., SURIANTATA, NAGY, L., DALRYMPLE, G. & VOYSEY, B. C. 2007. Reproductive phenology over a 10-year period in a lowland evergreen rain forest of central Borneo. *Journal of Ecology* 95:828-839.
- BROWN, S. & LUGO, A. E. 1990. Tropical secondary forests. *Journal of Tropical Ecology* 6:1-32.
- CAIRNS, M. A., BROWN, S., HELMER, E. H. & BAUMGARDNER, G. A. 1997. Root biomass allocation in the world's upland forests. *Oecologia* 111:1-11.
- CAVELIER, J., ESTEVEZ, J. & ARJONA, B. 1996. Fine-root biomass in three successional stages of an Andean cloud forest in Colombia. *Biotropica* 28:728-736.
- CAVELIER, J., WRIGHT, S. J. & SANTAMARÍA, J. 1999. Effects of irrigation on litterfall, fine root biomass and production in a semideciduous lowland forest in Panama. *Plant and Soil* 211:207-213.
- CHAZDON, R. L., LETCHER, S. G., VAN BRUEGEL, M., MARTÍNEZ-RAMOS, M., BONGERS, F. & FINEGAN, B. 2007. Rates of change in tree communities of

- secondary Neotropical forests following major disturbances. *Philosophical Transactions of the Royal Society Series B - Biological Sciences* 362:273-289.
- CORLETT, R. T. 1995. Tropical secondary forests. *Progress in Physical Geography* 19:159-172.
- DAVIDSON, E. A., REIS DE CARVALHO, C. J., FIGUEIRA, A. M., ISHIDA, F. Y., OMETTO, J. P. H. B., NARDOTO, G. B., SABÁ, R. T., HAYASHI, S. N., LEAL, E. C., VIEIRA, I. C. G. & MARTINELLI, L. A. 2007. Recuperation of nitrogen cycling in Amazonian forests following agricultural abandonment. *Nature* 447:995-998.
- DE JONG, W. 1997. Developing swidden agriculture and the threat of biodiversity loss. *Agriculture Ecosystems and Environment* 62:187-197.
- ELLSWORTH, D. S. & REICH, P. B. 1996. Photosynthesis and leaf nitrogen in five Amazonian tree species during early secondary succession. *Ecology* 77:581-594.
- FINEGAN, B. 1996. Patterns and process in Neotropical secondary rain forests: the first 100 years of succession. *Trends in Ecology and Evolution* 11:119-124.
- GORDON, W. S. & JACKSON, R. B. 2000 Nutrient concentrations in fine roots. *Ecology* 81:275-280.
- GOWER, S. T. 1987 Relations between mineral nutrient availability and fine root biomass in two Costa Rican tropical wet forests: a hypothesis. *Biotropica* 19:171-175.
- GREEN, J. J. 1992. *Fine root dynamics in a Bornean rain forest*. Ph.D. thesis, University of Stirling, UK.
- GREEN, J. J., DAWSON, L. A., PROCTOR, J., DUFF, E. I. & ELSTON, D. A. 2005. Fine root dynamics in a tropical rain forest is influenced by rainfall. *Plant and Soil* 276:23-32.

- GUARIGUATA, M. R. & OSTERTAG, R. 2001. Neotropical secondary succession: changes in structural and functional characteristics. *Forest Ecology and Management* 148:185–206.
- HERTEL, D. & LEUSCHNER, C. 2010. Fine root mass and fine root production in tropical moist forests as dependent on soil, climate and elevation. Pp. 428-444 in Bruijnzeel, L. A., Scatena, F. N. & Hamilton, L. S. (eds.). *Tropical montane cloud forest: science for conservation and management*. Cambridge University Press, Cambridge.
- HERTEL, D., LEUSCHNER, C. & HÖLSCHER, D. 2003. Size and structure of fine root systems in old-growth and secondary tropical montane forests (Costa Rica). *Biotropica* 35:143-153.
- HERTEL, D., HÖLSCHER, D., KOHLER, L. & LEUSCHNER, C. 2006. Changes in fine root system size and structure during secondary succession in a Costa Rican montane oak forest. Pp. 283-297 in Kapelle, M. (ed.). *Ecology and conservation of neotropical montane oak forests*. Ecological Studies Volume 185. Springer-Verlag, Berlin.
- JACKSON, R. B., MOONEY, H. A. & SCHULZE, E.-D. 1997. A global budget for fine root biomass, surface area, and nutrient contents. *Proceedings of the National Academy of Science of the USA* 94:7362-7366.
- JARAMILLO, V. J., AHEDO-HERNÁNDEZ, R. & KAUFFMAN, J. B. 2003. Root biomass and carbon in a tropical evergreen forest of Mexico: changes with secondary succession and conversion to pasture. *Journal of Tropical Ecology* 19:457-464.
- KLEINMAN, P. J. A., PIMENTEL, D. & BRYANT, R. B. 1995. The ecological sustainability of slash-and-burn agriculture. *Agriculture Ecosystems and Environment* 52:235-249.

- LAWRENCE, D. C., PEART, D. R. & LEIGHTON, M. 1998. The impact of shifting cultivation on a rainforest landscape in West Kalimantan: spatial and temporal dynamics. *Landscape Ecology* 13:135-148.
- MAYCOCK, C. R. & CONGDON, R. A. 2000. Fine root biomass and soil N and P in north Queensland rain forests. *Biotropica* 32:185-190.
- MIRMANTO, E., PROCTOR, J., GREEN, J. J., NAGY, L. & SURIANTATA 1999. Effects of nitrogen and phosphorus fertilisation in a lowland evergreen rain forest. *Philosophical Transactions of the Royal Society Series B - Biological Sciences* 354:1825-1829.
- MUTHUKUMAR, T., SHA, L., YANG, X., CAO, M., TANG, J. & ZHENG, Z. 2003. Distribution of roots and arbuscular mycorrhizal associations in tropical forest types of Xishuangbanna, southwest China. *Applied Soil Ecology* 22:241-253.
- NAGY, L. & PROCTOR, J. 1999. Early secondary forest growth after shifting cultivation. Pp. 1-12 in Sist, P., Sabogal, C. & Byron, Y. (eds.). *Management of secondary and logged-over forest in Indonesia*. Centre for International Forestry Research, Bogor.
- OSTERTAG, R. 1998. Belowground effects of canopy gaps in a lowland tropical wet forest. *Ecology* 79:1294-1304.
- OSTERTAG, R. 2001. The effects of nitrogen and phosphorus availability on fine root dynamics in Hawaiian montane forests. *Ecology* 82:485-499.
- POWERS, J. S. 2004. Changes in soil carbon and nitrogen after contrasting land-use transitions in northeastern Costa Rica. *Ecosystems* 7:134-146.
- POWERS, J. S., TRESEDER, K. K. & LERDAU, M. T. 2005. Fine roots, arbuscular mycorrhizal hyphae and soil nutrients in four Neotropical rain forests: patterns across large geographical distances. *New Phytologist* 165:913-921.

- PRAJADINATA, S. 1996. *Studies on tree regrowth on shifting cultivation sites near Muara Joloi, Central Kalimantan, Indonesia*. M.Sc. thesis, University of Stirling, UK.
- RAICH, J. W. 1980. Fine roots regrow rapidly after forest felling. *Biotropica* 12:230-231.
- REICH, P. B., ELLSWORTH, D. S. & UHL, C. 1995. Leaf carbon and nutrient assimilation and conservation in species of differing successional status in an oligotrophic Amazonian forest. *Functional Ecology* 9:65-76.
- SANFORD, R. L. & CUEVAS, E. 1996. Root growth and rhizosphere interactions in tropical forests. Pp. 268-300 in Mulkey, S. S., Chazdon, R. L. & Smith, A. P. (eds.). *Tropical forest plant ecophysiology*. Chapman & Hall, London.
- SOETHE, N., LEHMANN, J. & ENGELS, C. 2007. Carbon and nutrient stocks in roots of forests at different altitudes in the Ecuadorian Andes. *Journal of Tropical Ecology* 23:319-328.
- YAMASHITA, T., KASUYA, N., KADIR, W. R., CHIK, S. W., QUAH, E. S. & OKUDA, T. 2003. Soil and belowground characteristics of Pasoh Forest Reserve. Pp. 89-109 in Okuda, T., Manokaran, N., Matsumoto, Y., Niiyama, K., Thomas, S. C. & Ashton, P. S. (eds.). *Pasoh: ecology of a lowland rain forest in Southeast Asia*. Springer-Verlag, Tokyo.

FIGURE LEGENDS

Figure 1 Changes in root mass during secondary succession in tropical forests in the Barito Ulu area (Central Indonesian Borneo) assessed by soil coring for fine roots <2 mm diameter (a), small roots <5 mm diameter (b), coarse roots 5-10 mm diameter (c), and total (<10 mm) root mass (d). All values are mean \pm SE.

Figure 2 Changes in root element concentrations during secondary succession in tropical forests in the Barito Ulu area (Central Indonesian Borneo) assessed by soil coring. The figure shows carbon (a), nitrogen (b), and phosphorus (c) concentrations in roots to 20 cm depth . All values are mean \pm SE.

Table 1 Forest plot characteristics in the Barito Ulu area (Central Indonesian Borneo) used for the study of below-ground secondary succession. All values are mean \pm SE. In each stand (with the exception of the 1-y-old stands) three plots of the same age were set up in representative areas of vegetation. For the 1-y-old forest, two plots were set up in each of three separate stands.

| Age (y) | No. of plots | Plot size (ha) | Cores per plot | Min. tree dbh (cm) | No. stems ha ⁻¹ | Basal area (m ² ha ⁻¹) |
|---------|--------------|--|----------------|--------------------|----------------------------|---|
| 1 | 6 | 0.01 | 1 | 1 | 5570 \pm 900 | 3.28 \pm 0.60 |
| 3 | 3 | 1 \times 0.03125 & 2 \times 0.0625 | 2 & 4 | 5 | 1800 \pm 35 | 7.06 \pm 0.37 |
| 14 | 3 | 0.0625 | 4 | 5 | 1690 \pm 47 | 25.6 \pm 0.37 |
| 31 | 3 | 0.25 | 5 | 10 | 581 \pm 32 | 24.0 \pm 1.81 |
| Primary | 3 | 0.25 | 6 | 10 | 632 \pm 13 | 32.5 \pm 2.27 |

Table 2 Changes in soil characteristics during secondary succession in tropical forests in the Barito Ulu area (Central Indonesian Borneo). All values are mean \pm SE and are expressed on an oven-dry (105°C) basis.

| Age (y) | 1 | 3 | 14 | 31 | Primary |
|--|-----------------|-----------------|-----------------|-----------------|-----------------|
| n | 6 | 10 | 12 | 12 | 18 |
| pH | 4.66 \pm 0.09 | 4.74 \pm 0.06 | 4.52 \pm 0.04 | 4.32 \pm 0.04 | 4.34 \pm 0.03 |
| Total N (%) | 0.14 \pm 0.02 | 0.30 \pm 0.03 | 0.15 \pm 0.02 | 0.31 \pm 0.02 | 0.23 \pm 0.02 |
| Total C (%) | 1.93 \pm 0.23 | 3.84 \pm 0.47 | 2.11 \pm 0.31 | 4.50 \pm 0.51 | 2.71 \pm 0.37 |
| Extractable P ($\mu\text{g g}^{-1}$) | 0.45 \pm 0.06 | 0.41 \pm 0.04 | 0.34 \pm 0.03 | 0.30 \pm 0.05 | 0.44 \pm 0.06 |

Table 3 Correlations between root mass assessed by soil coring and soil characteristics for roots < 10 cm depth (assessed on an individual core basis) and < 20 cm depth (assessed on a plot basis) in the Barito Ulu area (Central Indonesian Borneo) (* = $P < 0.05$).

| | Soil characteristic | | | |
|---------------|---------------------|---------|----------|---------------|
| | pH | Total C | Total N | Extractable P |
| < 10 cm depth | | | | |
| <2 mm | -0.143 | -0.065 | -0.222 | -0.171 |
| <5 mm | -0.146 | -0.085 | -0.216 | -0.185 |
| 5-10 mm | 0.010 | -0.190 | -0.114 | -0.158 |
| Total mass | -0.119 | -0.177 | -0.215 | -0.247 |
| < 20 cm depth | | | | |
| < 2 mm | 0.077 | -0.296 | -0.528 * | -0.409 |
| < 5 mm | -0.304 | 0.036 | -0.273 | -0.444 |
| 5-10 mm | -0.161 | -0.305 | 0.351 | 0.185 |
| Total mass | -0.229 | 0.015 | -0.250 | -0.386 |

Figure 1 Changes in root mass during secondary succession in tropical forests in the Barito Ulu area (Central Indonesian Borneo) assessed by soil coring to two depths for fine roots <2 mm diameter (a), small roots <5 mm diameter (b), coarse roots 5-10 mm diameter (c), and total (<10 mm) root mass (d). All values are mean \pm SE.




Figure 2 Changes in root element concentrations in two diameter classes during secondary succession in tropical forests in the Barito Ulu area (Central Indonesian Borneo) assessed by soil coring. The figure shows carbon (a), nitrogen (b), and phosphorus (c) concentrations in roots to 20 cm depth . All values are mean \pm SE.

