

Please cite the Published Version

Olivero, J, Fa, JE, Real, R, Farfan, MA, Marquez, AL, Vargas, JM, Gonzalez, JP, Cunningham, AA and Nasi, R (2017) Mammalian biogeography and the Ebola virus in Africa. *Mammal Review*, 47 (1). pp. 24-37. ISSN 0305-1838

DOI: <https://doi.org/10.1111/mam.12074>

Publisher: Wiley

Version: Accepted Version

Downloaded from: <https://e-space.mmu.ac.uk/611996/>

Usage rights: © In Copyright

Additional Information: This is the peer reviewed version of the article which has been published in final form at 10.1111/mam.12074. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving. Copyright The Mammal Society and John Wiley & Sons Ltd.

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

Mammalian biogeography and the Ebola virus in Africa

Jesús OLIVERO *Universidad de Málaga, Grupo de Biogeografía, Diversidad y Conservación, Dept. Biología Animal, Campus de Teatinos s/n, 29071 Málaga, Spain.*

Email: jesusolivero@uma.es

John E. FA* *Division of Biology and Conservation Ecology, School of Science and the Environment, Manchester Metropolitan University, Manchester M1 5GD, UK and Center for International Forestry Research, CIFOR Headquarters, Bogor 16115, Indonesia.*

Email: jfa949@gmail.com

Raimundo REAL *Universidad de Málaga, Grupo de Biogeografía, Diversidad y Conservación, Dept. Biología Animal, Campus de Teatinos s/n, 29071 Málaga, Spain.*

Email: rrgimenez@uma.es

Miguel Ángel FARFÁN *Universidad de Málaga, Grupo de Biogeografía, Diversidad y Conservación, Dept. Biología Animal, Campus de Teatinos s/n, 29071 Málaga, Spain.*

Email: mafarfanaquilar@hotmail.com

Ana Luz MÁRQUEZ *Universidad de Málaga, Grupo de Biogeografía, Diversidad y Conservación, Dept. Biología Animal, Campus de Teatinos s/n, 29071 Málaga, Spain.*

Email: almarquez@uma.es

J. Mario VARGAS *Universidad de Málaga, Grupo de Biogeografía, Diversidad y Conservación, Dept. Biología Animal, Campus de Teatinos s/n, 29071 Málaga, Spain.*

Email: jmvy@uma.es

J. Paul GONZALEZ *Metabiota, Inc., 8757 Georgia Ave. Suite 420, Silver Spring, MD 20910, USA.*

Email: jpgonzalez@metabiota.com

Andrew A. CUNNINGHAM *Institute of Zoology, Zoological Society of London, Regent's Park, London NW1 4RY, UK.*

Email: a.cunningham@ioz.ac.uk

Robert NASI *Center for International Forestry Research, CIFOR Headquarters, Bogor 16115, Indonesia.*

Email: R.NASI@cgiar.org

ABSTRACT

1. Ebola virus is responsible for the fatal Ebola virus disease (EVD).
2. Identifying the distribution area of the Ebola virus is crucial for understanding risk factors conditioning the emergence of new EVD cases. Existing distribution models have underrepresented the potential contribution that reservoir species and vulnerable species make in sustaining the presence of the virus.
3. In this paper, we map favourable areas for Ebola virus in Africa according to environmental and zoogeographic descriptors, independent of human-to-human transmissions. We combine two different biogeographic approaches: analysis of mammalian distribution types (chorotypes), and distribution modelling of the Ebola virus.
4. We first obtain a model defining the distribution of environmentally favourable areas for the presence of Ebola virus. Based on a review of mammal taxa affected by or suspected of exposure to the Ebola virus, we model favourable areas again, this time according to mammalian chorotypes. We then build a combined model in which both the environment and mammalian distributions explain the favourable areas for Ebola virus in the wild.
5. We demonstrate that mammalian biogeography contributes to explaining the distribution of Ebola virus in Africa, although vegetation may also underscore clear limits to the presence of the virus. Our model suggests that the Ebola virus may be even more widespread than previously suspected, given that additional favourable areas are found throughout the coastal areas of West and Central Africa, stretching from Cameroon to Guinea, and extend further East into the East African Lakes region.

6. Our findings show that the most favourable area for the Ebola virus is significantly associated with the presence of the virus in non-human mammals. Core areas are surrounded by regions of intermediate favourability, in which human infections of unknown source were found. The difference in association between humans and other mammals and the virus may offer further insights on how EVD can spread.

Running head: Mammalian distributions and the Ebola virus

Keywords: Chorotypes, distribution modelling, favourability, fuzzy logic, reservoirs

*Correspondence author

Submitted: 15 October 2015

Returned for revision: 23 November 2015

Revision accepted: 29 March 2016

Editor: KH/DR

INTRODUCTION

EVD is a zoonosis caused by filoviruses of the genus *Ebolavirus* (hereafter Ebola virus), of which 4 species (*Zaire ebolavirus*, *Sudan ebolavirus*, *Tai Forest ebolavirus* and *Budibugyo ebolavirus*) are known in Africa (Kuhn et al. 2010). These viruses cause often-fatal haemorrhagic fever upon infection in humans (Kuhn et al. 2011). Ebola virus transmission from wildlife (here defined as wild mammals) has mostly been linked to people handling and butchering wild animals for bushmeat (Leroy et al. 2004a). Because bushmeat is important for human nutrition in Sub-Saharan Africa (Fa et al. 2015), it is fundamentally important to understand how reservoir factors, together with ecological conditions and human behaviour, contribute to Ebola virus outbreaks (Groseth et al. 2007). Recent biogeographical analyses have highlighted the importance of potential reservoirs (animals that can harbour the

pathogen indefinitely with no ill effects) in explaining the spatial assemblage of human infectious diseases worldwide (Murray et al. 2015). Biogeography has contributed broadly to questions of infectious disease ecology, management and surveillance (e.g. Cliff & Hagget 1995, Guernier et al. 2004, Smith & Guégan 2010). In this study, we build on this and propose a way to integrate virological, zoogeographical and environmental information through a combination of biogeographical approaches. Using these approaches we define the areas where Ebola virus may find suitable conditions to occur in the wild.

Despite methodological advances and increasing data availability, a limited understanding of the animals potentially implicated in the zoonosis has hampered mapping the extent of Ebola virus. Although likely reservoir species for the Ebola virus have been highlighted by some authors (Peterson et al. 2007), existing models describing the distribution of the virus have either not considered the contribution of reservoirs in sustaining its presence (Peterson et al. 2004), or have assumed that only a small number of species, suspected to be the reservoirs for the virus, are meaningful in the biogeography of the virus (Pigott et al. 2014). However, the ecology of the Ebola virus is complex and widely unresolved (Groseth et al. 2007). Thus, imposing restrictions to the selection of animal species considered in a distribution model might underrepresent the zoological substrate that could be determining the distribution of the virus. In fact, the role of particular bat species as true reservoirs of Ebola virus is still under discussion, and it is almost certain that there is a significant virus spillover among mammal species not suspected to be the reservoirs (Leroy et al. 2004a, 2004b, Groseth et al. 2007, Lahm et al. 2007, Olival et al. 2014).

Murray et al. (2015) suggested that mammalian biodiversity, as a whole, could be the strongest predictor explaining similarities between pathogeographic regions of the world. In

this study, we analysed the spatial distribution of Ebola virus in Africa, independent of human-to-human transmissions, under the hypothesis that it is influenced by how mammal species are distributed throughout the region. Thus, instead of taking sides with an uncertain selection of probable natural reservoirs and victims, we tested the explanatory potential of the biogeographic patterns of mammals (involving species co-occurrence and thus potential interactions) in Africa. Our prediction was that a distribution model of Ebola virus, based on variables defining the existing types of mammalian distributions in Africa, should better describe the virus occurrences recorded in wildlife than a model based on environmental descriptors alone. We examined all known literature regarding events of Ebola virus emergence, either EVD outbreaks or recorded presence of the virus in non-human mammals (Appendix S1). We then combined two different biogeographic approaches: the analysis of distribution types for mammals (chorotypes), and distribution modelling of Ebola virus, to develop a map of favourable areas for the virus according to both environmental conditions and mammal distributions in Africa.

METHODS

Identifying spatial links between Ebola virus and wildlife

Our model to define the favourable areas for Ebola virus was derived from occurrence data of any recorded presence of the virus in wildlife, disregarding whether it was detected in victims (i.e. organisms experiencing EVD symptoms, including human index cases) or in reservoirs. From the literature, we found a total of 91 geo-referenced events (Appendix S1) in which the Ebola virus was transmitted to humans from wildlife or was present in other non-human mammals. These events were recognised either via the: 1) detection of viral antibodies or viral nucleic acid ($n = 58$ events, including 22 in non-human mammal carcasses), 2) observed abnormal increase of mortality in non-human mammal populations, associated with

EVD outbreaks ($n = 51$ events), or 3) identification of index cases of EVD in humans ($n = 40$ events; the total number of occurrence records is 91 because recognition methods overlap).

We found published evidence from cases of serological and/or polymerase chain reaction (PCR) positivity of EVD in non-human mammal, or of EVD-linked mortality, in 28 mammal species: 10 primates, 3 rodents, 1 shrew, 8 bats, 1 carnivore and 5 ungulates (Fig 1; Appendix S1).




Fig 1. Non-human mammal species for which there is a record of naturally occurring Ebola virus infection either from serological or PCR positivity, or from increased mortality attributed to EVD. Top map: localities with a record of Ebola virus presence in wildlife. Bottom map: 1° x 1° squares with a record of Ebola virus presence (outlined in white); spatial model of Ebola virus (spatial favourability increases from white to black). Maps were generated using ArcGIS.

In 45 of the 58 laboratory-confirmed events of Ebola virus presence in wildlife, the virus was either isolated or was detected by PCR. In 11 out of the other 12 events, the virus was detected using enzyme-linked immunosorbent assay serology. The high sensitivity of this method, recommended for laboratory diagnosis of EVD by the World Health Organization (Anonymous 2014), was demonstrated by Ksiazek et al. (1992, 1999). The remaining case was serologically identified by both immunofluorescence antibody test and western blot analysis (Hayman et al. 2010).

A large number of incidents of abnormally high non-human mammal mortality attributed to Ebola virus were reported since November 1994 (Formenty et al. 1999). These data are of high significance for the development of a model of Ebola virus distribution in wildlife; the detection of unusual wild animal mortality can uncover the propagation of Ebola virus (Anonymous 2003a, 2003b, Rouquet et al. 2005). Indeed, increased animal mortality preceded the first human cases in most EVD outbreaks that occurred in Gabon and the Republic of Congo between 2001 and 2003 (Leroy et al. 2004a). Of the 51 mortality events considered here, 19 cases were temporally and spatially linked to recorded EVD outbreaks in humans, and/or were confirmed by PCR. The remaining 32 cases represented extensions of disease from confirmed events (Leroy et al. 2004a, Rouquet et al. 2005, Caillud et al. 2006, Lahm et al. 2007).

Only one of the 40 index cases of haemorrhagic fever syndrome considered in humans was not confirmed to be EVD through laboratory testing of clinical samples. This episode took place in Olloba (Republic of the Congo) in May 2002, when sample collection was not possible (Anonymous 2002; Rouquet et al. 2005). However, a previous EVD index case in Olloba had been confirmed just five months before (Leroy et al. 2004a).

Environmental model of Ebola virus

Individual data points were represented as 5-km buffers around the original virus occurrence locations, and we overlapped these buffers to a 1°×1°-resolution grid covering the whole of Africa (n = 2547 grid cells). With this buffer approach, we aimed to provide the occurrence-allocation in grid cells with some flexibility. In central Africa, 5 km is less than 0.05 times the side-length of a 1°×1° grid cell; given that the Ebola virus is hosted by non-human mammals, being less than 5 km far from a cell border is virtually equivalent to being located at the limits of two cells. The spatial resolution employed prevented autocorrelation that could exist as a consequence of spatial dependence among very close (< 1°) observations (Legendre & Legendre 1998). The final number of grid cells for detected presence of Ebola virus was 40, of which 33 corresponded to *Zaire ebolavirus* (Fig. 1). From these, an environmental model of Ebola virus revealing the distribution of environmentally favourable areas for its presence was made by employing the Favourability Function (Real et al. 2006, Acevedo & Real 2012):

$$F = \frac{P}{1-P} / \left(\frac{n_1}{n_0} + \frac{P}{1-P} \right) \quad (1),$$

where F is environmental favourability (0-1), P is probability of occurrence, and n_1 and n_0 are presence and absence numbers, respectively (absences were considered to be those squares not included in the presences subset). P was calculated for the entire African continent through a forward-stepwise logistic regression, according to 24 predictor variables describing types of ecosystems, abiotic factors and anthropogenic pressures on wildlife (see variable descriptions and sources in Appendix S2). Land-cover variables were computed as cover percentages in every grid, and the rest of the variables were estimated by averaged grid-

values. The logistic regression was based on the 40 Ebola virus occurrence squares ($= n_I$), considering the four Ebola virus species together. The cases of serological or PCR positivity of EVD, of non-human mammal mortality related to EVD, and of index cases of EVD in humans were not differentiated in the occurrence data employed for model building. Events of abnormal non-human mammal mortality not directly confirmed in the laboratory only contributed towards 5 presences, all of which were adjacent to the main core of EVD outbreaks in Gabon and the Republic of Congo. Models based on the Favourability Function distinguish those localities with environmental conditions that favour the species' existence from those with detrimental characteristics for its presence, irrespective of the species' proportion of occurrences within the study area. This property is essential for our objectives, as it enables direct comparison between models when several species are involved in the analytical design, and allows for model combinations through fuzzy logic (Barbosa & Real 2012). Fuzzy logic provides metrics for fuzzy sets, i.e. classes of objects with a continuum of membership degrees; these sets are characterised by a membership function that assigns to each object a real number in the interval [0, 1] (Zadeh 1965). In biogeography, environmental favourability, chorotypes, biotic boundaries and species richness can be identified as fuzzy sets (Estrada et al. 2008, Olivero et al. 2011, 2013, Romero et al. in press), thus fuzzy logic can be easily applied to these concepts (see below).

Adding a spatial descriptor to this variable set allowed us to consider autocorrelation resulting from the spatial structure of the distribution of the virus (Sokal & Oden 1978), and so take into account the impact of dispersal barriers, geological history and biotic interactions as potential predictors (Fa et al. 2014). To this end, we followed the 'trend surface approach' (Legendre & Legendre 1998). Thus, a series of spatial variables resulting from average X and average Y combinations for every square of the grid were examined through a backward

stepwise logistic regression. Then we used as spatial descriptor the lineal combination (logit) of spatial variables resulting from the logistic regression.

In order to control Type 1 errors caused by the large number of variables employed in the process, we used Benjamini and Hochberg's (1995) False Discovery Rate. This control was performed before the stepwise variable selection, so that only significant variables with a False Discovery Rate of $q < 0.05$ were accepted in a multivariate environmental model. To minimise multicollinearity, we also avoided Pearson correlation values higher than 0.8 between predictor variables.

Zoogeographic model of Ebola virus

We defined a zoogeographic model of Ebola virus as the distribution of favourable areas for the presence of Ebola virus according to mammalian chorotypes. A chorotype is a distribution pattern followed by one or several species, which can be recognised operatively within an area (Baroni-Urbani et al. 1978, Real et al. 2008); chorotypes, or types of distribution, represent the shared geographical, ecological and evolutionary context of several species (Real et al. 2008). The zoogeographic model of Ebola virus was produced in three steps: (1) we generated a list of mammal species to be considered; (2) we then defined the mammalian chorotypes with potential for explaining the geographical range of the Ebola virus in Africa; and (3) we finally built a model sustained on a set of predictor variables based on the chorotypes.

SELECTION OF MAMMAL SPECIES TO BE CONSIDERED

Two different criteria were used to select the list of mammal species based on the *a priori* analysis of probable (*sensu lato*) links between mammals and Ebola virus, via (1)

taxonomic proximity; and (2) biogeographic coincidence. Our aim here was to describe mammalian distribution patterns with enough potential to explain the distribution of Ebola virus; this does not mean that the species selected are proposed to be Ebola virus reservoirs or susceptible taxa.

Taxonomically related species might be susceptible to similar pathogens (Plyusnin & Morzunov 2001). Therefore, using the 28 mammal species recorded to be linked to the Ebola virus (see Appendix S1), we generated a preliminary list of 216 species (Appendix S3), by considering congeners: *Mus*, *Praomys*, *Sylvisorex*, *Tadarida*; and whole suprageneric groups if they were represented by more than 5% of their species (a lower proportion was considered to be anecdotic) among the probable reservoirs and susceptible species:

Pteropodidae, Hystricidae, Hominidae, Cercopithecidae, Viverridae and Suidae families, and forest tribes of the family Bovidae. We also included pouched rats *Cricetomys* and grasscutters *Thryonomys* in our preliminary list, though there is no current evidence of their relationship with the Ebola virus. We considered that, because members of these rodent genera are so numerically important in the bushmeat trade in central and West Africa (Alexander et al. 2014), their inclusion in the analysis was warranted. Thus, our taxonomic criterion was defined so that it discarded groups very far from any suspect reservoirs (e.g. proboscideans, equids or felines), whilst it considered all those taxa including species whose membership of the Ebola virus reservoir system cannot be categorically denied.

We then preselected, from our preliminary list of 216 species, those whose biogeography was potentially able to explain the distribution of Ebola virus. To this end, those species geographically coinciding with or nested within the range of the Ebola virus were chosen. Given that the area covered by the Ebola virus is poorly known, we estimated the

geographical coincidence between the virus and mammalian distributions according to models defining the spatial structure of their respective ranges. Thus, for every mammal species, and for Ebola virus, we built spatial models that described their distributions according to purely spatial variables (i.e. combinations of average X and average Y). These models were generated by employing the Favourability Function (see above). For mammals, presences/absences were derived from polygon shapefiles for each species available from the International Union for Conservation of Nature's website (IUCN Red List of Threatened Species. Version 2012.1. <http://www.iucnredlist.org>.) projected to a $1^{\circ} \times 1^{\circ}$ -grid system, which is the maximum spatial resolution at which 'extent-of-occurrence' range maps (such as those provided by the IUCN) are suitable for analysis (Hurlbert & Jetz 2007). The spatial model of Ebola virus was based on the same presence/absence data set considered for the environmental model. The degree of geographic coincidence was quantified through the fuzzy overlap (or fuzzy similarity) index, whilst the degree of nesting was measured using the fuzzy inclusion index (Dubois & Prade 1980, Olivero et al. 2011); both measures ranged from 0 to 1. The mathematic formulation of the fuzzy overlap (i.e. intersection divided by union) is equivalent to Jaccard's similarity index (Olivero et al. 2011), the theoretical distribution of which is randomly distributed around 0.33 (Baroni-Urbani 1980). Thus, we considered a geographic coincidence to be significant when the fuzzy overlap was > 0.33 (i.e. when both Ebola virus and the mammal had more than one third of their ranges in common). There is no theoretical reference for estimating significance of fuzzy inclusion values; hence, we considered that a mammal species' geographical range was nested within that of Ebola virus when at least two thirds of the mammal's range was included within the Ebola virus's range (i.e. fuzzy inclusion > 0.66). A final list of 96 mammal species was ultimately considered for entry in the zoogeographic model of Ebola virus (Appendix S3), 89 of which

fulfilled the two selection criteria. Another 7 species were considered because there is published record of their probable infection by Ebola virus.

DEFINING AND MAPPING MAMMALIAN CHOROTYPES

The geographical ranges of the 96 mammal species were classified hierarchically according to the Baroni-Urbani and Buser (1976) similarity index, using the Unweighted Pair Group Method with Arithmetic Mean agglomerative algorithm (Sneath & Sokal 1973). All clusters in the resulting classification dendrogram were assessed for statistical significance with the method given by Olivero et al. (2011), using RMacoqui 1.0 software (<http://rmacoqui.r-forge.r-project.org/>). The number of resulting chorotypes was not predefined; all groups of distributions that were significantly clustered were considered chorotypes.

The resulting chorotypes were mapped following the accumulated favourability approach as in Fa et al. (2014). The accumulated favourability is considered a fuzzy-logic method for estimating species richness (Estrada et al. 2008). For all species forming part of the same chorotype, we built an environmental model using the method shown above for Ebola virus; then, in every grid cell of the study area, we added the favourability values defined by these environmental models. So, a cell showing high accumulated favourability for the species of a chorotype is defined to have favourable conditions for the presence of a large number of these species. Mapping chorotypes this way allowed for further downscaling to a higher spatial resolution (see below).

BUILDING THE ZOOGEOGRAPHIC MODEL OF EBOLA VIRUS

To build the zoogeographic model of Ebola virus, we followed the same procedure as for the environmental model. In this case, the model was built using every chorotype as a predictor variable. The accumulated favourability for the species of a chorotype was employed as a variable.

Although every chorotype was based on a finite cluster of species (i.e. the chorotypical cluster), the distribution of every species has a certain degree of membership in all detected chorotypes (Olivero et al. 2011). We proffered a biogeographically-justified list of mammal species whose link with Ebola virus is worth investigating, based on a species membership degree > 0.5 in at least one of the chorotypes entered in the zoogeographic model of Ebola virus. The degree of membership of every species in each chorotype was calculated as the average of the Baroni-Urbani and Buser similarities between a species and all distributions in the chorotypical cluster. The theoretical distribution of this similarity index is randomly distributed around 0.5 (Baroni-Urbani & Buser 1976); we considered membership to be significant above this value.

Environmental/zoogeographic model assessment and combination

Both the environmental and zoogeographic models were assessed by using calibration (Hosmer & Lemeshow 2000), i.e. testing whether favourability values reflected the existing observations of Ebola virus presence in wildlife. We also compared the performance of both models in relation to goodness-of-fit, by using $-2 \times \log$ -likelihood; discrimination capacity using the Area Under the receiver-operating-characteristic Curve (AUC; Lobo et al. 2008); and classification capacity using sensitivity, specificity, correct classification rate, Cohen's Kappa (Fielding & Bell 1997), and under-prediction and over-prediction rates (Barbosa et al.

2013). Classification measures were based on the 0.5 favourability threshold because probability is equal to the overall prevalence at this level (Acevedo & Real 2012). In a calibrated model, the Hosmer-Lemeshow index should be non-significant; for higher goodness of fit, $-2 \times \log$ -likelihood should be lower; for better discrimination, AUC should be higher; for better classification, sensitivity, specificity, correct classification rate and Kappa should be higher, whereas under-prediction and over-prediction should be lower.

The environmental and the zoogeographic models of Ebola virus were combined so that the final model gave favourability values based on the degree to which conditions are both environmentally and zoogeographically favourable for Ebola virus presence. To this end, we used the fuzzy intersection between the environmental model and the zoogeographic model of Ebola virus (Romero et al. in press), by measuring the minimum favourability value for either of the two models in each grid cell.

We analysed the relative contribution of environment and zoogeography to the combined model. To do this, we mapped where favourability values were derived from either the environmental or the zoogeographic model. We, thus, identified where, and to what extent, one of these models acted as a limiting factor whilst the other model showed a higher favourability for presence of Ebola virus. Finally, we employed the sensitivity index to assess the capacity of the combined model to classify recorded presences of Ebola virus and of the four African Ebola virus species separately.

We compared the distribution of known EVD outbreaks in humans with our resulting favourability maps for the Ebola virus. For this purpose, we overlapped the locations of index cases in humans with a favourability map divided into three regions depending on their

favourability values, according to the thresholds proposed by Muñoz & Real (2006). If the predicted favourability was higher than 0.8, which means that the odds are more than 4:1 favourable to Ebola virus, the square was considered as highly favourable. Those areas with a favourability value lower than 0.2 (odds less than 1:4) were considered of low favourability for Ebola virus. The remaining squares were regarded as intermediate favourability areas.

The potential for contacts between human populations and wildlife could have influenced detection of Ebola virus infections beyond environmental and zoogeographic factors that could favour the presence of Ebola virus. Therefore, we tested whether Ebola virus occurrences poorly explained by the combined model of Ebola virus (i.e. not explained by environmental conditions or by mammalian biogeography) could be accounted for by the presence of human populations. To do this, we performed a logistic regression of the 40 Ebola virus occurrence squares, using favourability values for Ebola virus as the independent variable. The residuals of this regression were then related to rural population density and with distance to roads by using linear regression (for variable sources, see Appendix S2).

To increase the potential uses of our output for management, surveillance or analyses requiring an ecological context for Ebola virus, the combined model was downscaled to $0.1^{\circ} \times 0.1^{\circ}$ resolution squares, by employing the direct downscaling approach (Bombi & d'Amen 2012). This method was classified by Bierkens et al. (2000) as “downscaling based on mechanistic models through a deterministic [favourability] function”. To do this, we applied the favourability equations involved in the combined model to predictor variables at this resolution. A 10-fold shortening of the grain size (referring to pixel side length) does not severely affect predictions of species distributions (Bombi & d'Amen 2012).

RESULTS

The environmental model of Ebola virus

Three variables (*terra firme* rain forests, natural vegetation/cropland mosaics and small annual temperature range) were significantly associated with the areas of high environmental favourability for Ebola virus presence ($P < 0.01$; Figs 1 and 2). Pearson correlation values between these variables were lower than 0.51. Vegetation/cropland mosaics and relatively constant temperatures complemented the predictive power of forests, especially within deforested areas. Crucially, the large swamp forest areas along the lower course of the Congo River (“cuvette congolaise”, where EVD outbreaks have so far not been recorded) did not appear to be as favourable as other *terra firme* central and West African forest areas (Fig 2).




Fig 2. Modelling of the environmental/zoogeographic favourability for the presence of Ebola virus in wildlife. The environmental model is based on *terra firme* Rain Forests (TFRF), Natural Vegetation/Cropland Mosaics (NVCM) and Annual Temperature Range (ATR, with increasing values from yellow to red). The zoogeographic model is composed from four types of mammalian distributions or chorotypes (see Fig 3). The two models are combined according to fuzzy logic, requiring both environmentally and zoogeographically favourable conditions. Maps were generated using ArcGIS.

The zoogeographic model of Ebola virus

Our analyses revealed the presence of 16 significantly different types of mammalian distributions, or chorotypes (Fig 3). Four of these chorotypes significantly supported the zoogeographic model of Ebola virus ($P < 0.001$): Rainforest chorotype, West-African Forest chorotype, North-Western Congolian Forest chorotype, and the single-species chorotype of *Mus gundae* (Fig 3). The zoogeographic model of Ebola virus, based on these chorotypes, showed high favourability values within the rain forests of Central and West Africa, with a decline in favourability towards the East, dropping even more dramatically south of the Congo River.




Fig 3. Classification of 96 mammal species according to distributional similarities ($S =$ Baroni-Urbani & Buser similarity index). *: Branches defining significant chorotypes ($P < 0.05$). For every chorotype, accumulated favourability values (increasing from white to black) of all species are mapped. Chorotypes are abbreviated as: NWCF: North-Western Congolian Forest; SECF: South-Eastern Congolian Forest; NCF: Northern Congolian Forest; SCF: Southern Congolian Forest; CSBF: Cross-Sanaga-Bioko Forest; RF: Rain Forest; WAF: West-African Forest; SS: Sub-Saharan; RA: *Rousettus aegyptiacus*; TB: *Tadarida bemmeleni*; MO: *Mus oubanguii*; AS: *Allochrocebus solatus*; CD: *Cercopithecus dryas*; MG:

Mus goundae; PM: *Praomys mutoni*; SK: *Sylvisorex konganensis*. Solid circles indicate the 64 mammals that are significant members (degree > 0.5) of chorotypes explaining the zoogeographic model of Ebola virus (RF, WAF, NWCF and MG). Red circles indicate published records of probable contact with the Ebola virus. Maps were generated using ArcGIS.

Having identified the four chorotypes that, according to the zoogeographic model, favour Ebola virus presence, we propose a list of 64 mammal species as a guide for future investigations in the search for Ebola virus reservoirs and potentially susceptible species (see Fig 3; Appendix S3).

Environmental/zoogeographic model assessment and combination

Both the environmental and zoogeographic models appear to be significantly well calibrated (Table 1). The zoogeographic model shows a better goodness of fit, higher discrimination and greater classification power than the environmental model. However, both provide significant complementary information about the virus's distribution. Thus, our environmental model suggests that waterlogged areas (swamps and swamp forests) could limit the presence of Ebola virus (Fig 2). Vegetation/cropland mosaics away from the rain forest block appeared unfavourable in the zoogeographic model, the only exception being a small area of the northern Central African Republic savannas; this area corresponds with the discovery of Ebola virus genetic sequences in rodents in gallery forests around Bohou River, 490 km North of Bangui (Morvan et al. 1999).

Table 1. Model assessment and comparison based on calibration, goodness of fit, and discrimination and classification capacities. In a calibrated model, H-L should be non-significant; for higher goodness of fit, -2logL should be lower; for better discrimination, AUC should be higher; for better classification, sensitivity, specificity, correct classification rate and Kappa should be higher, whereas under-prediction and over-prediction should be lower.

Indices	Model	
	Environmental	Zoogeographic
H-L*	$\chi^2_{\tau}=13.02, P>0.05$	$\chi^2_{\tau}=7.03, P>0.05$
-2logL**	226.26	225.48
AUC***	0.943	0.965
Sensitivity	0.925	0.975
Specificity	0.885	0.889
Kappa	0.180	0.196
Correct classification rate	0.886	0.890
Under-prediction	0.00135	0.00045
Over-prediction	0.886	0.877

* Hosmer-Lemeshow calibration index

** -2 x ln(Likelihood)

***Area Under the receiver-operating-characteristic Curve

Zoogeography was the most extended limiting factor for the presence of Ebola virus, covering 76% of the African continent; instead, environment was the limiting factor in the remaining 34%. These percentages turned into 55% and 45%, respectively, if completely unfavourable areas for Ebola virus (favourability < 0.05) were excluded; and they became 48% and 52% within the intermediate or high favourability areas (favourability > 0.2; see Appendix S4). The combined model inherited, from the zoogeographic model, the eastward decline of favourability values (Fig 2), where zoogeographic favourability was lower than environmental favourability (Appendix S4). Instead, the combined model excluded swamp forests as Ebola virus favourable areas as a legacy of the environmental model.

Due to the sensitivity of the combined model, more than 92% of the 1° x 1° squares with records of the Ebola virus ($n = 40$) were classified correctly in areas with favourability values higher than 0.5. The combined model explained 94% of the *Zaire ebolavirus* presences ($n = 33$), 100% of the *Budibugyo ebolavirus* presences ($n = 2$), 75% of the *Sudan ebolavirus*

presences ($n = 4$), and 100% of the *Tai Forest ebolavirus* presences ($n = 1$). Up to 63% of Ebola virus presences appeared within the highly favourable areas (i.e. value > 0.8 ; Fig 4a), largely coinciding with non-flooded rain forests. The 0.8-favourability threshold significantly separated virus occurrences in humans from presences in other mammals (Fig 4a); the highly favourable region included a significantly higher proportion of presences in non-human mammals ($\chi^2_1 = 6.22$, $P < 0.05$), as well as in both humans and non-human mammals ($\chi^2_1 = 8.00$, $P < 0.01$). In contrast, presences recorded only in humans were significantly located within the intermediate favourability areas (i.e. value between 0.2 and 0.8; $\chi^2_1 = 19.16$, $P < 0.001$). This regionalisation reveals an emergent property that arises from the combination of the environmental and zoogeographic models. Residual analyses (to explore the differences between predicted and observed probabilities of presence) indicate that there is a significant positive correlation between Ebola virus points recorded outside the highly favourable region and both rural population density ($r = 0.589$, $P = 0.001$) and distance to roads ($r = -0.466$, $P = 0.014$). A downscaled geographical representation of the combined model, at a $0.1^\circ \times 0.1^\circ$ spatial resolution, is shown in Fig 4b.




Fig 4. Combination of the environmental and the zoogeographic models of Ebola virus at the original $1^{\circ} \times 1^{\circ}$ resolution (A) and downscaled to a $0.1^{\circ} \times 0.1^{\circ}$ resolution (B). Red tones distinguish regions with high (value > 0.8), intermediate ($0.2-0.8$), and low favourability (< 0.2) for the presence of Ebola virus. White areas are completely unfavourable for Ebola virus (< 0.05). Circles represent squares with cases of serological or PCR positivity of Ebola virus infection in non-human mammals, animal mortality attributed to EVD, and zoonotic transmission to humans. G: Guinea; SL: Sierra Leone; L: Liberia; CI: Cote d'Ivoire; GH: Ghana; T: Togo; Be: Benin; N: Nigeria; CA: Cameroon; CAR: Central African Republic; S: South Sudan; U: Uganda; R: Rwanda; B: Burundi; DRC: Democratic Republic of Congo; A: Angola; CO: Congo; GA: Gabon; EG: Equatorial Guinea. Maps were generated using ArcGIS.

A total of 17 countries contained high favourability areas (> 0.8) for Ebola virus (Fig 4b). In decreasing order, according to the extent of highly favourable areas in each, these were: Democratic Republic of the Congo, Gabon, Cameroon, Republic of Congo, Côte

d'Ivoire, Liberia, Ghana, Central African Republic, Nigeria, Equatorial Guinea, Sierra Leone, Uganda, Benin, Angola, Tanzania, Guinea and Togo. Another four countries comprised areas of intermediate favourability (0.2 – 0.8): Rwanda, Burundi, South Sudan and Kenya.

DISCUSSION

Spatial modelling, applied to pathogen systems, can generate statistically robust predictions of the geographic distributions of the organism causing a disease, and its maintenance reservoir(s) (Peterson 2006, Purse & Golding 2015). However, species distribution modelling applied to EVD transmission in Africa has so far underestimated the contribution reservoirs and vulnerable species may make in sustaining the virus presence (Peterson et al. 2004, Pigott et al. 2014; see above). Our models are the first to analyse the contribution of mammalian distribution patterns to the biogeography of Ebola virus in Africa. In our study, we considered the four African Ebola virus species together because of the scarcity of information available on the presence of species other than *Zaire ebolavirus*. The latter is the most lethal (Kuhn et al. 2011), was responsible for the EVD outbreaks in central Africa (Bausch & Schward 2014), and has also been associated with the current epidemic in West Africa (Dudas & Rambaut 2014). Our models have very high sensitivity (0.94) and specificity (0.90) to the presence of *Zaire ebolavirus*, but high favourability areas also adequately predicted the presence of *Budibugyo ebolavirus*, and of *Tai Forest ebolavirus* (sensitivity = 1). However, our models poorly predicted *Sudan ebolavirus*. With better knowledge of the distribution of all Ebola virus species, more robust models could have been developed. In particular, models focused on the different Ebola species should be developed promptly, since some mammalian chorotypes that were not entered in the model could explain the presence of these Ebola virus species better, e.g. chorotype NECF in the case of *Budibugyo ebolavirus* (compare Figs. 1 and 3).

Peterson et al. (2007) identified 55 groups of mammalian taxa of interest as potential reservoirs, based on a detailed inspection of distributional overlap patterns with known filovirus disease outbreaks. Our list of 68 species biogeographically linked to Ebola virus differs from the list of likely reservoir species presented by Peterson et al. (2007). These authors generated an inventory of candidate reservoir species, assuming that the reservoir supports persistent, largely asymptomatic Ebola virus infections, which *de facto* eliminates most primates and ungulates. Our list, unlike Peterson et al.'s, is 33% primates (families Hominidae and Cercopithecidae) and 20% ungulates (families Suidae and Bovidae). Members of the fruit bat family (Pteropodidae), Molossid bats, viverrid carnivores, and the rodent *Praomys jacksoni* are included in both lists. In our list of species, however, we also included other fruit bats (*Casinycteris argynnis*, *Scotonycteris ophiodon* and *Scotonycteris zenkeri*) as members of the chorotypes significantly linked to Ebola virus distribution. Of the two rodent genera *Cricetomys* and *Thryonomys* included in our analyses given their prominence in the bushmeat trade, *Cricetomys emini* emerged as worth investigating. Thus, unlike Peterson et al.'s (2007), our list is not a proposal of potential reservoir species; instead, we present a list of mammals biogeographically associated with Ebola virus, and thus with the geographic potential of being involved in the virus cycle as reservoirs or susceptible species. This is not a closed list, even though we have restricted the analysis to a limited number of species. This is because any species (i.e. also those not considered in the analysis) are members of all chorotypes to a certain degree (Olivero et al. 2011). Thus, it is possible to evaluate whether unassessed taxa can be considered part of the zoogeographic factor as defined in the Ebola virus model.

Of the 28 mammal species with published record of probable contact with Ebola virus, only 8 were not included in our 64-species list: *Rousettus aegyptiacus*, *Eidolon helvum*, *Epomophorus gambianus*, *Micropteropus pusillus*, *Tadarida condylura*, *Civettictis civetta*, *Papio anubis* and *Mandrillus leucophaeus* (Appendix S3). What these species have in common is that a large proportion of their populations are found distant from where Ebola virus has been detected. We do not challenge the evidenced relationship of these species with Ebola virus, but note that they do not show strong overlap with it (even though there is overlap in some parts of the range). It could be argued that subspecific taxa of the same species may differ in their ability to serve as Ebola virus reservoirs. In the case of these 8 mammal species, however, all subspecies currently recognised (Kingdon et al. 2013) are partially distributed outside the Ebola virus's range, with the exception of *Tadarida condylura osborni*. These species might be more interesting from a phylogeographic perspective, but their distributions cannot explain the geographic distribution of the African Ebola viruses in wildlife; the virus might potentially be hosted by individuals of these species, but factors other than the distribution of these reservoirs could explain the observed geographic limits of the Ebola virus in Africa (e.g. other reservoirs or environmental conditions).

We show, as Pigott et al. (2014) clearly indicated, that the high relative contribution of vegetation in the model might underscore clear limits to the presence of Ebola virus. As in the Pigott et al. (2014) and Murray et al. (2015) maps (and less patently in Peterson et al.'s 2004), we confirm West Africa as a highly favourable region for Ebola virus. Our map also suggests that the virus distribution may be even more widespread than previously suspected, given that additional favourable areas for the virus are found throughout the coastal areas of West and Central Africa, stretching from Cameroon to Guinea, and extending further East

into the East African Lakes region (i.e. Uganda). Of the 17 countries with highly favourable areas (> 0.8) for Ebola virus occurrence in wildlife in our model, 16 are among the top 17 considered as high-risk from Ebola by both Pigott et al. (2014) and Murray et al. (2016), and 10 of these are common to both sources. The Democratic Republic of the Congo, Gabon, Republic of Congo, Côte d'Ivoire, Uganda and Guinea have records of EVD index cases in humans, whereas Cameroon, Liberia, Central African Republic, Nigeria, Angola, Ghana, Sierra Leone, Benin, Tanzania and Togo have not. Identifying 'at risk' countries that are currently Ebola virus-free, according to various different evidence types, could be fundamental in achieving adequate levels of preparedness. For example, countries such as Equatorial Guinea, with 94% of its territory highly favourable for Ebola virus in our model, though it was ranked number 20 by Pigott et al. (2014), should not escape attention. Similarly, Burundi, Rwanda and Kenya, containing (like South Sudan) areas of intermediate favourability —covering all of Burundi and Rwanda— but so far free from EVD cases, should be targeted for further investigation.

In our map, the favourability for the disease in areas south of the Congo River is remarkably lower than in other rainforest areas; a pattern not apparent in Pigott et al.'s (2014) map. A possible explanation may be a lack of adequate zoogeographic conditions for the virus to flourish linked to a lower diversity of potential reservoir species in this area. There is evidence that the richness and population structure of mammals found in the central part of the Congo basin is considerably lower than in areas along the western, northern and eastern side of the Congo River (Fa et al. 2014). The limiting role of the zoogeographic factor south to the Congo River (Appendix S4) supports this possibility, but we cannot discard the possibility of observation bias, since the central Congo is not an easily accessible area.

Serological surveys among human populations support a rainforest origin of the potential reservoir of Ebola virus (Gonzalez et al. 1989, 2000). Living or spending significant time deep in the forest is positively correlated with seropositivity in Pygmies and in other human groups (Gonzalez et al. 2000, Becquart et al. 2010, Schoepp et al. 2014). This is consistent with our finding of the highest favourability for Ebola virus in the rainforest, and with the recorded concurrence of cases of spillover from other animals to humans there, even though this information was not used to build the model. However, epidemic outbreaks also originated outside of the rain forest block. This is the case for the first known outbreaks in humans, recorded by the World Health Organization in Zaire and Sudan in 1976 (Anonymous 1978a,b), for the last and most important episode in West Africa (Bausch & Schward 2014), and for some other cases in Uganda (Albariño et al. 2013), Congo (Pourrut et al. 2005), the Democratic Republic of the Congo (Anonymous 2009) and Sudan (Onyango et al. 2007). All of them have in common that the human index cases were not documented to be related to cases of EVD in non-human mammals, and that they occurred in areas of intermediate favourability for Ebola virus according to our model. Thus, a main contribution of our model is the finding of a geographical context for exploring which mechanisms differentiate Ebola virus transmission to humans in highly favourable areas, compared to in areas of intermediate favourability for Ebola virus; this is of high epidemiological relevance. In areas of intermediate favourability, gallery forests may play a significant role in harbouring and spreading the Ebola virus.

EVD epidemics seem to be related more to human behaviour, which increases the risk of contact with the reservoir, than to the emergence of a highly pathogenic viral strain (Gonzalez et al. 2000). The significant relationship between human presence (i.e. population density and roads) and EVD cases in areas only moderately favourable to Ebola virus

suggests that EVD transmission to humans in these areas could be influenced by anthropogenic factors. More frequent contact between humans and forest fauna would amplify the chances of Ebola virus transmission to humans, even where environmental or zoogeographic conditions are not the most favourable for the virus. About half of the EVD index cases reported in the region of intermediate favourability happened at the limits of highly favourable areas (Fig 4), which could have facilitated contacts. Even so, some index cases have been located away from the forest domain. One possible explanation for the arrival of *Zaire Ebolavirus* in West Africa, during the 2013 Guinean outbreak, far from its usual haunts in central Africa, is that the virus was introduced by travelling bats (Bausch & Schward 2014). Sáez et al. (2015) also suggested that the index case in Guinea may have been infected by a colony of insectivorous free-tailed bats *Mops condylurus*. Do Chiropterans have a more significant role in zoonotic spillover in the intermediate-favourability areas than in the deep rainforest haunts of Ebola virus? In the first documented case of EVD in Sudan in 1976, the index case was located (by the World Health Organization) in a cotton factory far from the forest block, where the only wild significantly abundant species was an insectivorous bat species (Anonymous 1978b). Additional analysis and study of the human-wildlife interface are still required to delineate areas in which human populations are at risk of zoonotic transmission of Ebola virus. In this task, our model could be used to delineate the geographic context of the analysis. As risk factors for zoonotic transmission of Ebola virus are better understood, it will be possible to incorporate this information into future risk mapping assessments and to develop mitigation or prevention measures.

ACKNOWLEDGEMENTS

This study was supported by USAID as part of the Bushmeat Research Initiative of the CGIAR research program on Forests, Trees and Agroforestry. We thank Dr Kris Murray for his valuable help in improving the manuscript.

REFERENCES

- Acevedo P, Real R (2012) Favourability: concept, distinctive characteristics and potential usefulness. *Naturwissenschaften* 99: 515-522.
- Albariño CG, Shoemaker T, Khristova ML, Wamala JF, Muyembe JJ, Balinandi S et al. (2013) Genomic analysis of filoviruses associated with four viral hemorrhagic fever outbreaks in Uganda and the Democratic Republic of the Congo in 2012. *Virology* 442: 97-100.
- Alexander JS, McNamara J, Rowcliffe JM, Oppong J, Milner-Gulland EJ (2014) The role of bushmeat in a West African agricultural landscape. *Oryx* 49: 643-651.
- Anonymous (1978a) Ebola haemorrhagic fever in Sudan, 1976. *Bulletin of the World Health Organization* 56: 247-270.
- Anonymous (1978b) Ebola haemorrhagic fever in Zaire, 1976. *Bulletin of the World Health Organization* 56: 271-293.
- Anonymous (2002) Suspected acute haemorrhagic fever syndrome, Gabon. World Health Organization. *Weekly Epidemiological Record* 77: 213.
- Anonymous (2003a) Outbreak(s) of Ebola haemorrhagic fever, Congo and Gabon, October 2001 - July 2002. World Health Organization. *Weekly Epidemiological Record* 26: 223-228.

- Anonymous (2003b) Outbreak(s) of Ebola haemorrhagic fever in the Republic of the Congo, January-April 2003. World Health Organization. *Weekly Epidemiological Record* 33: 285-296.
- Anonymous (2009) *Ebola Hemorrhagic Fever, Information Packet*. Special Pathogens Branch Division of High-Consequence Pathogens and Pathology. National Center for Emerging Zoonotic Infectious Diseases, Center for Disease Control and Prevention, U.S. Department of Health and Human Services, Atlanta, USA.
- Anonymous (2014) Laboratory diagnosis of Ebola virus disease. World Health Organization Interim Guideline reference number WHO/EVD/GUIDANCE/LAB/14.1.
- Barbosa AM, Real R, Muñoz AR, Brown JA (2013) New measures for assessing model equilibrium and prediction mismatch in species distribution models. *Diversity and Distributions* 29: 1333-1338.
- Barbosa AM, Real R (2012) Applying fuzzy logic to comparative distribution modelling: a case study with two sympatric amphibians. *The Scientific World Journal* 2012; ID 428206. doi: 10.1100/2012/428206.
- Baroni-Urbani C (1980) A statistical table for the degree of coexistence between two species. *Oecologia* 44: 287-289.
- Baroni-Urbani C, Buser MW (1976) Similarity of binary data. *Systematic Zoology* 25: 251-259.
- Baroni-Urbani C, Rufo S, Vigna-Taglianti A (1978) Materiali per una biogeografia italiana fondata su alcuni generi di Coleotteri, Cicindelidi, Carabidi e Crisomelidi. *Estratto della Memorie della Societa Entomologica Italiana* 56: 35–92.
- Bausch DG, Schward L (2014) Outbreak of Ebola Virus Disease in Guinea: where ecology meets economy. *PLOS Neglected Tropical Diseases* 2014; 8:e3056. doi: 10.1371/journal.pntd.0003056.

- Becquart P, Wauquier N, Mahlakõiv T, Nkoghe D, Padilla C, Souris M et al. (2010) High prevalence of both humoral and cellular immunity to Zaire ebolavirus among rural populations in Gabon. *PLOS ONE* 2010; 5: e9126. doi: 10.1371/journal.pone.0009126.
- Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B* 57: 289-300.
- Bierkens MFP, Finke PA, de Willigen P (2000) *Upscaling and Downscaling Methods for Environmental Research*. Kluwer Academic Publishers, Dordrecht, the Netherlands.
- Bombi P, d'Amen M (2012) Scaling down distribution maps from atlas data: a test of different approaches with virtual species. *Journal of Biogeography* 39: 640-651.
- Caillud D, Levréro F, Cristescu R, Gatti S, Dewas M, Douadl M, Gutier-Hion A, Raymond M, Ménard N (2006) *Gorilla* susceptibility to Ebola virus: the cost of sociality. *Current Biology* 16: 489-491.
- Cliff AD, Haggett P (1995) The epidemiological significance of islands. *Health & Place* 1: 199-209.
- Dubois D, Prade H (1980) *Fuzzy Sets and Systems: Theory and Applications*. Academic Press, New York, USA.
- Dudas G, Rambaut A (2014) Phylogenetic analysis of Guinea 2014 EBOV Ebolavirus outbreak. *PLOS Currents Outbreaks* 6: ecurrents.outbreaks.84eefe5ce43ec9dc0bf0670f7b8b417d.
- Estrada A, Real R, Vargas JM (2008) Using crisp and fuzzy modelling to identify favourability hotspots useful to perform gap analysis. *Biodiversity and Conservation* 17: 857–871.

- Fa JE, Olivero J, Farfán MA, Márquez AL, Vargas JM, Real R, Nasi R (2014) Integrating sustainable hunting in biodiversity protection in Central Africa: hot spots, weak spots, and strong spots. *PLOS ONE* 2014; 9: e112367. doi: 10.1371/journal.pone.0112367.
- Fa JE, Olivero J, Real R, Farfán MA, Márquez AL, Vargas JM et al. (2015) Disentangling the relative effects of bushmeat availability on human nutrition in central Africa. *Scientific Reports* 5: 8168. doi: 10.1038/srep08168.
- Fielding AH, Bell JF (1997) A review of methods for the assessment of prediction errors in conservation presence-absence models. *Environmental Conservation* 24: 38-49.
- Formenty P, Boesch C, Wyers M, Steiner C, Donati F, Dind F, Walker F, Le Guenno B (1999) Ebola virus outbreak among wild chimpanzees living in a rain forest of Côte d'Ivoire. *Journal of Infectious Diseases* 179: 120-126.
- Gonzalez J-P, Josse R, Johnson ED, Merlin M, Georges AJ, Abandja J et al. (1989) Antibody prevalence against haemorrhagic fever viruses in randomized representative Central African populations. *Research Virology* 140: 319-331.
- Gonzalez J-P, Nakoune E, Slenczka W, Vidal P, Morvan JM (2000) Ebola and Marburg virus antibody prevalence in selected populations of the Central African Republic. *Microbes and Infection* 2: 39-44.
- Groseth A, Feldmann H, Strong JE (2007) The ecology of Ebola virus. *Trends in Microbiology* 15: 408-416.
- Guernier V, Hochberg ME, Guégan J-F (2004) Ecology drives the worldwide distribution of human diseases. *PLoS Biology* 2: e141.
- Hayman DTS, Emmerich P, Yu M, Wang L-F, Suu-Ire R, Fooks AR, Cunningham AA, Wood JLN (2010) Long-term survival of an urban fruit bat seropositive for Ebola and Lagos bat viruses. *PLoS ONE* 5: e11978.

- Hosmer DW, Lemeshow S. (2000) *Applied Logistic Regression*. Wiley-Interscience, New York, USA.
- Hurlbert AH, Jetz W (2007) Species richness, hotspots, and the scale dependence of range maps in ecology and conservation. *PNAS* 104: 13384–13389.
- Kingdon J, Happold D, Butynski T, Hoffman M, Happold M, Kalina J (2013) *Mammals of Africa: 6 Vols*. Bloomsbury Publishing, London, UK.
- Ksiazek TG, Rollin PE, Jahrling PB, Johnson E, Dalgard DW, Peters CJ (1992) Enzyme immunosorbent assay for Ebola virus antigens in tissues of infected primates. *Journal of Clinical Microbiology* 30: 947-950.
- Ksiazek TG, West CP, Rollin PE, Jahrling PB, Peters CJ (1999) ELISA for the detection of antibodies to Ebola viruses. *Journal of Infectious Diseases* 179: S192-S198.
- Kuhn JH, Becker S, Ebihara H, Geisbert TW, Johnson KM, Kawaoka Y et al. (2010) Proposal for a revised taxonomy of the family Filoviridae: classification, names of taxa and viruses, and virus abbreviations. *Archives of Virology* 155: 2083-2103.
- Kuhn JH, Dodd LE, Wahl-Jensen V, Radoshitzky SR, Bavari S, Jahrling PB (2011) Evaluation of perceived threat differences posed by filovirus variants. *Biosecurity and Bioterrorism* 9: 361-371.
- Lahm SA, Kombila M, Swanepoel R, Barnes RFW (2007) Morbidity and mortality of wild animals in relation to outbreaks of Ebola haemorrhagic fever in Gabon, 1994-2003. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 101: 64-78.
- Legendre P, Legendre L (1998) *Numerical Ecology*. Second English edition. Elsevier Science, Amsterdam, the Netherlands.

- Leroy EM, Rouquet P, Formenty P, Souquière S, Kilbourne A, Froment J-M et al. (2004a) Multiple Ebola virus transmission events and rapid decline of Central African wildlife. *Science* 303: 387-390.
- Leroy EM, Telfer P, Kumulungui B, Yaba P, Rouquet P, Roques P, Gonzalez J-P, Jsiasek TG, Rollin PE, Nerrienet E (2004b) A serological survey of Ebola virus infection in Central African nonhuman primates. *Journal of Infectious Diseases* 90: 1895-1899.
- Lobo JM, Jiménez-Valverde A, Real R (2008) AUC: a misleading measure of the performance of predictive distribution models. *Global Ecology and Biogeography* 17: 145-151.
- Morvan JM, Deubel V, Gounon P, Nakouné E, Barrière P, Murri S et al. (1999) Identification of Ebola virus sequences present as RNA or DNA in organs of terrestrial small mammals of the Central African Republic. *Microbes and Infection* 1: 1193-1201.
- Muñoz A-R, Real R (2006) Assessing the potential range expansion of the exotic monk parakeet in Spain. *Diversity and Distributions* 12: 656-665.
- Murray KA, Preston N, Allen T, Zambrana-Torrel C, Hosseini PR, Daszak P (2015) Global biogeography of human infectious diseases. *PNAS* 112: 12746-12751.
- Olival KJ, Hayman DT (2014) Filoviruses in bats: current knowledge and future directions. *Viruses* 6: 1759-1788.
- Olivero J, Real R, Márquez AL (2011) Fuzzy chorotypes as a conceptual tool to improve insight into biogeographic patterns. *Systematic Biology* 60: 645-660.
- Olivero J, Márquez AL, Real R (2013) Integrating fuzzy logic and statistics to improve the reliable delimitation of biogeographic regions and transition zones. *Systematic Biology* 62: 1-21.

- Onyango CO, Opoka ML, Ksiazek TG, Formenty P, Ahmed A, Tukei PM et al. (2007) Laboratory diagnosis of Ebola hemorrhagic fever during an outbreak in Yambio, Sudan, 2004. *Journal of Infectious Diseases* 196: 193-198.
- Peterson AT, Bauer JT, Mills JN (2004) Ecologic and geographic distribution of Filovirus disease. *Emerging Infectious Diseases* 10: 40-47.
- Peterson AT, Papeş M, Carroll DS, Leirs H, Johnson KM (2007) Mammal taxa constituting potential coevolved reservoirs of filoviruses. *Journal of Mammalogy* 88: 1544-1554.
- Peterson AT (2006) Ecological niche modeling and spatial patterns of disease transmission. *Emerging Infectious Diseases* 12: 1822–1826.
- Pigott DM, Golding N, Mylne A, Huang Z, Henry AJ, Weiss DJ et al. (2014) Mapping the zoonotic niche of Ebola virus disease in Africa. *eLife* 3: e04395.
- Plyusnin A, Morzunov S (2001) Virus evolution and genetic diversity of hantaviruses and their rodent hosts. *Current Topics in Microbiology and Immunology* 256: 47-75.
- Pourrut X, Kumulungui B, Wittmann T, Moussavou G, Délicat A, Yaba P, Nkoghe D, Gonzalez J-P, Leroy EM (2005) The natural history of Ebola virus in Africa. *Microbes and Infection* 7: 1005-1014.
- Purse BV, Golding N (2015) Tracking the distribution and impacts of diseases with biological records and distribution modelling. *Biological Journal of the Linnean Society* 115: 664-677.
- Real R, Barbosa AM, Vargas JM (2006) Obtaining environmental favourability functions from logistic regression. *Environmental Ecological Statistics* 13: 237-245.
- Real R, Olivero J, Vargas JM (2008) Using chorotypes to deconstruct biogeographical and biodiversity patterns: the case of breeding waterbirds in Europe. *Global Ecology and Biogeography* 17: 735-746.

- Romero D, Olivero J, Brito JC, Real R (in press) Comparison of approaches to combine species distribution models based on different sets of predictors. *Ecography* In press; doi: 10.1111/ecog.01477.
- Rouquet P, Froment J-M, Bermejo M, Kilbourn A, Karesh W, Reed P et al. (2005) Wild animal mortality monitoring and human Ebola outbreaks, Gabon and Republic of Congo, 2001-2003. *Emerging Infectious Diseases* 11: 283-290.
- Sáez AM, Weiss S, Nowak K, Lapeyre V, Zimmermann F, Dux A et al. (2015) Investigating the zoonotic origin of the West African Ebola epidemic. *EMBO Molecular Medicine* 7: 17-23.
- Schoepp RJ, Rossi CA, Khan SH, Goba A, Fair JN (2014) Undiagnosed acute viral febrile illnesses, Sierra Leone. *Emerging Infectious Diseases* 20: 1176-1182.
- Smith KF, Guégan J-F (2010) Changing geographic distributions of human pathogens. *Annual Review of Ecology, Evolution and Systematics* 41: 231-250.
- Sneath PHA, Sokal RR (1973) *Numerical Taxonomy. The Principles and Practices of Numerical Classification*. Freeman, San Francisco, USA.
- Sokal RR, Oden NL (1978) Spatial autocorrelation in biology. 1. Methodology. *Biological Journal of the Linnean Society* 10: 199-228.
- Zadeh LA (1965) Fuzzy sets. *Information and Control* 8:338-353.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of this article at the publisher's web-site.

Appendix S1. Referenced locations of Ebola virus presence in wildlife.

Appendix S2. Predictor-variable description and sources.

Appendix S3. List and features of the 216 mammal species considered in this study.

Appendix S4. Contribution of environment and zoogeography to the Ebola virus distribution model based on combination of explanatory factors.