



**Manchester
Metropolitan
University**

Rojer, AGM, Kruizenga, HM, Trappenburg, MC, Reijnierse, EM, Sipilä, S, Narici, MV, Hogrel, JY, Butler-Browne, G, McPhee, JS, Pääsuke, M, Meskers, CGM, Maier, AB and de van der Schueren, MAE (2016) The prevalence of malnutrition according to the new ESPEN definition in four diverse populations. *Clinical Nutrition*, 35 (3). pp. 758-762. ISSN 0261-5614

Downloaded from: <https://e-space.mmu.ac.uk/617227/>

Version: Accepted Version

Publisher: Elsevier

DOI: <https://doi.org/10.1016/j.clnu.2015.06.005>

Usage rights: Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Please cite the published version

<https://e-space.mmu.ac.uk>

1 **Short communication**

2 **The prevalence of malnutrition according to the new ESPEN definition in four clinically**
3 **relevant populations**

4 A.G.M. Rojer

5 Department of Internal Medicine, Section of Gerontology and Geriatrics, VU University Medical
6 Center, Amsterdam, The Netherlands

7 H.M. Kruizenga PhD, RD

8 Department of Nutrition and Dietetics, Internal Medicine, VU University Medical Center, Amsterdam,
9 The Netherlands

10 Dutch Malnutrition Steering Group, Amsterdam, The Netherlands

11 M.C. Trappenburg, PhD, MD

12 Department of Internal Medicine, Section of Gerontology and Geriatrics, VU University Medical
13 Center, Amsterdam, The Netherlands

14 Department of Internal Medicine, Amstelland Hospital, Amstelveen, The Netherlands

15 E.M. Reijnierse, MSc

16 Department of Internal Medicine, Section of Gerontology and Geriatrics, VU University Medical
17 Center, Amsterdam, The Netherlands

18 S. Sipilä, PhD Gerontology Research Centre and Department of Health Sciences, University of
19 Jyväskylä, Jyväskylä, Finland

20 M.V. Narici, PhD Division of Medical Sciences & Graduate Entry Medicine, MRC-ARUK Centre of
21 Excellence for Musculoskeletal Ageing Research, University of Nottingham, Royal Derby Hospital
22 Centre.

23 J.Y. Hogrel, PhD UPMC UM 76, INSERM U 974, CNRS 7215, Institut de Myologie, Paris, France

24 G. Butler-Browne UPMC UM 76, INSERM U 974, CNRS 7215, Institut de Myologie, Paris, France

25 J.S. McPhee, PhD School of Healthcare Science, John Dalton Building, Manchester Metropolitan
26 University, Manchester M1 5GD, UK

27 H. Gapeyeva, PhD Institute of Exercise Biology and Physiotherapy, University of Tartu, Centre of
28 Behavioral and Health Sciences, University of Tartu, Tartu, Estonia

29 C.G.M. Meskers, PhD, MD
30 Department of Rehabilitation Medicine, VU University Medical Center, Amsterdam, The Netherlands

31 A.B. Maier, PhD, MD
32 Department of Internal Medicine, Section of Gerontology and Geriatrics, VU University Medical
33 Center, Amsterdam, The Netherlands

34 M.A.E. de van der Schueren, PhD, RD
35 Department of Nutrition and Dietetics, Internal Medicine, VU University Medical Center, Amsterdam,
36 The Netherlands

37 Department of Nutrition, Sports and Health, Faculty of Health and Social Studies, HAN University of
38 Applied Sciences, Nijmegen, The Netherlands

39

40 Corresponding Author: Dr. M.A.E. de van der Schueren
41 E-mail: m.devanderschueren@vumc.nl

42 Tel: +31 204443410

43 Text word count: 2201

44 Tables and figures: 4

45 Abstract word count: 403

46 References: 13

47 **Abstract**

48 **Background & Aims:** International consensus on the definition of malnutrition has not yet been
49 reached. Recently, the European Society for Clinical Nutrition and Metabolism (ESPEN) proposed a
50 new consensus definition of malnutrition. The aim of the present study was to describe the prevalence
51 of malnutrition according to the new ESPEN consensus definition in four clinically relevant
52 populations: acutely ill middle-aged patients, geriatric outpatients, healthy old individuals and healthy
53 young individuals.

54 **Methods:** The recently released ESPEN consensus definition of malnutrition was applied to the four
55 different populations. This definition consists of two different options: option one requires body mass
56 index (BMI, kg/m^2) $<18.5 \text{ kg}/\text{m}^2$ to define malnutrition. Option two requires the combined finding of
57 unintentional weight loss (mandatory) and at least one of either reduced BMI or low fat free mass
58 index (FFMI, kg/m^2). Unintentional weight loss could be either $>10\%$ of habitual weight independent
59 of time, or $>5\%$ over the previous 3 months. Reduced BMI is defined as $<20 \text{ kg}/\text{m}^2$ or $<22 \text{ kg}/\text{m}^2$ in
60 subjects younger and older than 70 years, respectively. Low FFMI is $<15 \text{ kg}/\text{m}^2$ and $<17 \text{ kg}/\text{m}^2$ in
61 females and males, respectively. Only individuals for whom all data on diagnostic options were
62 complete were included in the present analysis: acutely ill middle-aged patients ($n=349$), geriatric
63 outpatients ($n=135$), healthy old individuals ($n=306$) and healthy young individuals ($n=179$).

64 **Results:** According to the new ESPEN consensus definition of malnutrition, the prevalence of
65 malnutrition ranged from 1% in healthy old individuals to 15% in the acutely ill middle-aged patients.
66 The different options that compose the new ESPEN consensus definition of malnutrition were
67 represented in the four populations in various ways, i.e., high prevalence rates of low FFMI in all four
68 populations, a relatively high prevalence of BMI $<18.5 \text{ kg}/\text{m}^2$ in healthy young individuals but low
69 prevalence of BMI $<18.5 \text{ kg}/\text{m}^2$ in all other populations and relatively low prevalence rates of the
70 combination of weight loss with either low BMI or low FFMI.

71 **Conclusions:** Combining the diagnostic options that compose the new ESPEN consensus definition of
72 malnutrition results in prevalence rates lower than expected in acutely ill middle-aged patients and
73 geriatric outpatients. In contrast, healthy young individuals are (most likely falsely) defined
74 malnourished based on a low BMI $<18.5 \text{ kg}/\text{m}^2$. Future studies should further determine the cut-off

75 points for FFMI and BMI in older persons. In addition, the association of the new ESPEN consensus
76 definition of malnutrition with clinically relevant outcomes needs further study.

77

78 **Keywords:** Malnutrition, definition, prevalence

79 **Introduction**

80 Malnutrition is an increasingly recognized problem that is associated with morbidity, mortality, and
81 increased costs of care. To enhance early recognition and treatment of malnutrition, an easy and
82 widely accepted definition of malnutrition is necessary. Such a definition should be easy applied for
83 all health care professionals and in all health care settings. Furthermore, the definition of malnutrition
84 should be widely accepted to be able to compare prevalence rates among health care settings and
85 countries, and to improve communication among health care providers and politicians worldwide.

86 International consensus on the definition of malnutrition has not yet been reached. Recently, the
87 European Society for Clinical Nutrition and Metabolism (ESPEN) proposed a new consensus
88 definition including two options for the diagnosis of malnutrition(1). The first diagnostic option
89 requires a low body mass index (BMI), following the recommendation by the World Health
90 Organization: subjects are defined as malnourished if they have a BMI <18.5 kg/m²(2). The second
91 diagnostic option encompasses unintentional weight loss (>10% independent of time or >5% in the
92 last three months), always combined with either a low BMI (<20 kg/m² if <70 years old or <22 kg/m²
93 if ≥70 years old) or a low Fat Free Mass Index (FFMI). Given the increasingly recognized importance
94 of body protein reserves, the preferred diagnostic trajectory involves the assessment of the FFMI, with
95 cut-off points of 15 kg/m² for women and 17 kg/m² for men.

96 As the new ESPEN consensus definition of malnutrition has been released only recently, validation
97 studies have not yet been published. The aim of the present study was to describe the prevalence rates
98 of malnutrition according to the newly proposed ESPEN consensus definition of malnutrition in four
99 clinically relevant populations including acutely ill middle-aged patients, geriatric outpatients, healthy
100 old individuals and healthy young individuals. This study will provide a first overview of the
101 applicability of the newly proposed consensus definitions of malnutrition in various target populations.

102 **Methods**

103 The recently released ESPEN consensus definition of malnutrition (see Fact box 1), was applied to
104 four different populations. Only individuals for whom all data on diagnostic options were complete
105 were included for the present analysis.

106

107 *Population 1: acutely ill, middle-aged patients*

108 This population consisted of 349 patients (57.6 years, SD 17.7) who were admitted to a general
109 internal ward (general internal medicine, gastroenterology, dermatology, rheumatology, nephrology)
110 or a general surgical ward (general surgery and surgical oncology) of the VU University Medical
111 Center (Amsterdam, the Netherlands) in two periods respectively from April 2002 until October 2002
112 and from February until June 2003 (3).

113

114 *Population 2: geriatric outpatients*

115 This population consisted of 135 geriatric outpatients (80.8 years, SD 7.3) who were referred to the
116 geriatric outpatient clinic of the Bronovo Hospital (The Hague, the Netherlands) for a comprehensive
117 geriatric assessment due to mobility problems between March 2011 and January 2012 (4).

118

119 *Population 3 and 4: healthy old individuals and healthy young individuals*

120 The European MYOAGE study consisted of old and young healthy individuals. Individuals in the
121 MYOAGE study were recruited from five different sites across Europe, including: Manchester, UK;
122 Paris, France; Leiden, the Netherlands; Jyväskylä, Finland and Tartu, Estonia. Data was collected
123 between 2010 and 2013(5).

124 Old and young healthy individuals from the MYOAGE study were analysed separately; included were
125 306 healthy old individuals (74.4 years, SD 3.3) and 179 healthy young individuals (23.4 years SD
126 2.9).

127 Individuals in all four populations were screened with the Short Nutritional Assessment Questionnaire
128 (SNAQ), with ≥ 3 points indicating high risk of malnutrition (3). Independent of the SNAQ screening
129 results the diagnosis of malnutrition was assessed by measured weight and height, calculated BMI,
130 self-reported unintentional weight loss; FFMI was derived differently across the populations. In the
131 acutely ill middle-aged population FFMI was assessed using Xitron 4000B multiple frequency Bio-
132 electrical Impedance Spectroscopy, using its 50KHz frequency and the Geneva equations (6). In
133 geriatric outpatients, FFM was assessed using a direct segmental multi-frequency Bio-electrical
134 Impedance Analyser, which provided direct values for FFM, which were then divided by height²
135 (InBody 720, Biospace Co., Ltd, Seoul, Korea). In both the old and young healthy individuals FFMI
136 was assessed with dual-energy x-ray absorptiometry.

137

138 The prevalence of malnutrition according to the new ESPEN consensus definition, as well as to the
139 individual diagnostic options, was calculated for each population.

140

141 **Results**

142 Screening with SNAQ (≥ 3 points) identified 105 acutely ill middle-aged patients at risk of
143 malnutrition, 14 geriatric outpatients, 1 healthy old individual and none of the healthy young
144 individuals. Assessment according to the new ESPEN definition (independent of initial screening with
145 SNAQ) yielded 54 malnourished patients (15%) in the acutely ill, middle-aged patients, 10
146 malnourished geriatric outpatients (7%), 3 malnourished healthy old (1%) and 14 malnourished
147 healthy young (8%). Five malnourished patients in the acutely ill middle-aged were not identified to
148 be at risk by the initial SNAQ screening; this was 2 in the geriatric outpatients, 3 in the healthy old and
149 14 in the healthy young.

150 Table 1 depicts the prevalence data for each population. Furthermore, it shows the prevalence of the
151 individual diagnostic options of the definition. For example: in the acutely ill middle-aged population,
152 the prevalence of malnutrition was 15%. Out of the total population of 349 individuals, 116 had a
153 FFMI below the proposed cut-off points; 44 individuals (13%) out of these 116 were defined as
154 malnourished, based on the combination low FFMI and unintentional weight loss.

155 Figures 1A and 1B display the overlap of the new ESPEN consensus definition of malnutrition and its
156 individual diagnostic options in the acutely ill middle-aged population and in the geriatric outpatient
157 population. Overlap figures are not displayed for the healthy old individuals and healthy young
158 individuals due to low number of malnourished cases in the healthy old individuals ($n=3$) and
159 unilateralism in healthy young individuals ($n=14$ of which 13 were identified malnourished by having
160 only a low BMI). Furthermore, in the healthy old individuals low BMI and low FFMI were never
161 combined with unintentional weight loss.

162

163 **Discussion**

164 The description of the prevalence of malnutrition according to the recently released ESPEN consensus
165 definition showed relatively low prevalence rates of malnutrition in all four populations. A low BMI
166 and a low FFMI were observed in approximately 20% of the individuals in each population. However,
167 most individuals were eventually not identified as malnourished as low BMI/low FFMI was not
168 combined with unintentional weight loss. Thus, the criterion of unintentional weight loss has a
169 dominant influence when determining prevalence rates.

170 The first diagnostic option of the new ESPEN consensus definition of malnutrition consists of a BMI
171 $< 18.5 \text{ kg/m}^2$. A BMI $< 18.5 \text{ kg/m}^2$ was mostly observed in acutely ill middle-aged patients. However,
172 both in geriatric outpatients and in healthy old individuals, a BMI less than 18.5 kg/m^2 was rare (1% in
173 each population). Thus, a BMI $< 18.5 \text{ kg/m}^2$ is rare in older individuals, which is in line with other
174 studies that report higher BMI's in older populations (7).

175 Thirteen (7%) young healthy individuals were defined malnourished according to a low BMI. The
176 ESPEN diagnostic process suggests screening first, and further assessment only for those at risk. In the
177 cohorts described, we used the SNAQ for initial screening. This resulted in no young healthy
178 individuals at risk. i.e. no need for further assessment. However, recent research has shown that the
179 SNAQ is not a valid screening tool for outpatients as it does not comprise BMI (8). If we had used
180 MUST (9), for example, the 13 healthy individuals would have passed screening and been identified
181 malnourished in the process of diagnosis, most likely falsely, as they were all selected for their
182 excellent health. They were probably 'healthy and slim' or very athletic.

183 The second diagnostic option of the new ESPEN consensus definition of malnutrition consists of a
184 combination of unintentional weight loss and either low BMI or low FFMI. In the acutely ill middle-
185 aged population, 25% of all patients had unintentional weight loss. This is in line with expectations, as
186 unintentional weight loss is a frequently described phenomenon accompanying acute disease.
187 However, only 15% of the population was defined as malnourished according to the new ESPEN
188 consensus definition of malnutrition, indicating that in 10% of the cases unintentional weight loss did

189 not occur in combination with a low BMI or a low FFMI. We believe that the infrequent concurrence
190 of unintentional weight loss with low BMI (30 out of 54 malnourished acutely ill middle-aged
191 patients) is due to the relatively high BMI's at the population level. The combination of unintentional
192 weight loss and a low FFMI was present in 44 out of the 54 malnourished acutely ill patients.

193 In the geriatric outpatient population the combination of unintentional weight loss and low BMI (<22
194 kg/m² if ≥70 years old) (9 out of 10 malnourished outpatients) overlapped reasonably well with the
195 combination of unintentional weight loss and low FFMI (8 out of 10 malnourished outpatients).

196 As the new definition suggests that unintentional weight loss should be combined with either a low
197 BMI or a low FFMI to be defined as malnourished, this also suggests that a low BMI and a low FFMI
198 can be used interchangeably. Although in geriatric outpatients, malnutrition based on low BMI or on
199 low FFMI was equivalent, the correspondence in the acutely ill patients was lower. Larger numbers of
200 patients are required, however, to determine how well BMI and FFMI correlate in different
201 populations.

202

203 A low FFMI was highly prevalent (14-33%) in all populations, however prevalence of the
204 combination of unintentional weight loss and low FFMI showed a lower prevalence (0-13%). The high
205 prevalence of a low FFMI may be explained by the chosen cut-off points in the ESPEN consensus
206 definition of malnutrition. The cut-off point of FFMI below 15 kg/m² for women represents the 50th
207 percentile, according to Schutz's reference tables (10). For men, a cut-off point of FFMI below 17
208 kg/m² represents the 10th percentile, which is probably a much more realistic percentile to apply. This
209 raises the question of whether the cut-off point for women should be amended, for example to 14
210 kg/m², which represents the 10th percentile for women (10), and what consequences that cut-off point
211 would have for the prevalence rates. A future study should look into a possible revision of the FFMI
212 cut-off points, their overlap with unintentional weight loss and the consequences for malnutrition
213 prevalence rates.

214 Of the acutely ill middle-aged patients with a BMI $<20 \text{ kg/m}^2$ (<70 years) or $<22 \text{ kg/m}^2$ (≥ 70 years),
215 approximately half were defined as malnourished as they also met the second diagnostic option:
216 unintentional weight loss. In the geriatric outpatient population a low BMI ($<20 \text{ kg/m}^2$ (<70 years) or
217 $<22 \text{ kg/m}^2$ (≥ 70 years)) occurred in 28 (21%) outpatients. Remarkably, in only 9 (7%) geriatric
218 outpatients a low BMI was combined with unintentional weight loss; this might be one explanation for
219 the lower than expected prevalence rates in this geriatric outpatient population. Previous studies have
220 reported prevalence rates of malnutrition in approximately 50% of geriatric outpatients (11-13). Since
221 geriatric outpatients usually suffer from multiple age-related problems and many co-morbidities,
222 unintentional weight loss is most likely a problem that has occurred only slowly and thereby has not
223 reached the cut-off level of 10%, or that has gone by unnoticed. In the healthy old individuals, 39
224 (13%) had a BMI $<20 \text{ kg/m}^2$ (<70 years) or $<22 \text{ kg/m}^2$ (≥ 70 years) but none were defined as
225 malnourished based on the concurrence with unintentional weight loss; three healthy old individuals
226 were defined as malnourished based on a BMI $<18.5 \text{ kg/m}^2$. For older persons, either a BMI cut-off
227 point higher than 22 kg/m^2 or a different cut-off point for unintentional weight loss are more
228 reasonable indicators of malnutrition.

229

230

231 **Conclusion:**

232 The prevalence rates of positive scores when using the different ESPEN consensus definitions of
233 malnutrition were high. However, when combining the different diagnosis pathways, the prevalence
234 rates were lower than expected in acutely ill middle-aged patients and in geriatric outpatients. Old
235 healthy individuals were probably identified as malnourished too infrequently, due to missing
236 concurrence of low BMI/low FFMI and weight loss, whereas in contrast young healthy individuals
237 were (most likely falsely) defined malnourished based on a low BMI.

238 Some suggestions for further studies:

239 - To study the importance of the relative contribution of unintentional weight loss versus low BMI or
240 low FFMI in the new ESPEN consensus definition of malnutrition.

241 - To reconsider the proposed cut-off points for FFMI, specifically for women. Both absolute cut-off
242 points and age- and sex- specific percentiles should be studied.

243 - To study whether a low BMI and a low FFMI are interchangeable and whether this is different
244 between populations.

245 - To evaluate the proposed BMI cut-off point of $< 22 \text{ kg/m}^2$ or the degree of unintentional weight loss
246 in older adults. This descriptive study even raises the question whether BMI is a relevant parameter for
247 nutritional status in older adults at all or whether we should more strongly rely on FFMI in older
248 adults.

249

250 In future analyses, we will report on the association between the new ESPEN consensus definition of
251 malnutrition, its individual diagnostic options and clinically relevant outcome measures such as
252 functionality and survival, which will shed a further light on the chosen cut-off points for BMI and
253 FFMI.

254 **References**

- 255 (1) Cederholm T, Bosaeus I, Barazzoni R, Bauer J, Van Gossum A, Klek S, et al. Diagnostic criteria
256 for malnutrition - an ESPEN Consensus Statement. *Clinical Nutrition*, submitted for publication.
- 257 (2) WHO. Physical status: the use and interpretation of anthropometry. Report of a WHO Expert
258 Committee. 1995; World Health Organization Technical Report Series 854.
- 259 (3) Kruizenga HM, Seidell JC, de Vet HCW, Wierdsma NJ, Van Bokhorst-de Van der Schueren,
260 Development and validation of a hospital screening tool for malnutrition: the short nutritional
261 assessment questionnaire (SNAQ). *Clin Nutr* 2005 02;24(1):75-82.
- 262 (4) Bijlsma AY, Pasma JH, Lambers D, Stijntjes M, Blauw GJ, Meskers CG, et al. Muscle strength
263 rather than muscle mass is associated with standing balance in elderly outpatients. *J Am Med Dir
264 Assoc* 2013 Jul;14(7):493-498.
- 265 (5) McPhee JS, Hogrel JY, Maier AB, Seppet E, Seynnes OR, Sipila S, et al. Physiological and
266 functional evaluation of healthy young and older men and women: design of the European MyoAge
267 study. *Biogerontology* 2013 Jun;14(3):325-337.
- 268 (6) Kyle UG, Genton L, Karsegard L, Slosman DO, Pichard C. Single prediction equation for
269 bioelectrical impedance analysis in adults aged 20--94 years. *Nutrition* 2001 03;17(0899-9007; 3):248-
270 253.
- 271 (7) Winter JE, MacInnis RJ, Wattanapenpaiboon N, Nowson CA. BMI and all-cause mortality in older
272 adults: a meta-analysis. *Am J Clin Nutr* 2014 Apr;99(4):875-890.
- 273 (8) Leistra E, Langius JA, Evers AM, van Bokhorst-de van der Schueren, M.A., Visser M, de Vet HC,
274 et al. Validity of nutritional screening with MUST and SNAQ in hospital outpatients. *Eur J Clin Nutr*
275 2013 May 1.
- 276 (9) Elia M. The 'MUST' report. Nutritional screening of adults: a multidisciplinary responsibility.
277 2003.
- 278 (10) Schutz Y, Kyle UU, Pichard C. Fat-free mass index and fat mass index percentiles in Caucasians
279 aged 18-98 y. *Int J Obes Relat Metab Disord* 2002 07;26(0307-0565; 7):953-960.
- 280 (11) van Bokhorst-de van der Schueren, M.A., Lonterman-Monasch S, de Vries OJ, Danner SA,
281 Kramer MH, Muller M. Prevalence and determinants for malnutrition in geriatric outpatients. *Clin
282 Nutr* 2013 May 17;32:1007-1011.
- 283 (12) Saka B, Kaya O, Ozturk GB, Erten N, Karan MA. Malnutrition in the elderly and its relationship
284 with other geriatric syndromes. *Clin Nutr* 2010 12;29(1532-1983; 0261-5614; 6):745-748.
- 285 (13) Ulger Z, Halil M, Kalan I, Yavuz BB, Cankurtaran M, Gungor E, et al. Comprehensive
286 assessment of malnutrition risk and related factors in a large group of community-dwelling older
287 adults. *Clin Nutr* 2010 08;29(1532-1983; 0261-5614; 4):507-511.

288

289

290

291 **Attachments:**

- 292 - Fact box: Two alternative ways to diagnose malnutrition.
- 293 - Figure 1A and 1B: The overlap of the new ESPEN consensus definition of malnutrition
294 and its individual diagnostic options in acutely ill middle-aged patients and geriatric
295 outpatients.
- 296 - Table 1: Prevalence rates of malnutrition according to the new ESPEN consensus
297 definition and to its individual diagnostic options in four populations.

298

299 **Fact box: Two alternative ways to diagnose malnutrition.**

300 Before diagnosis of malnutrition is considered it is mandatory to fulfil criteria for being “at risk” of
301 malnutrition by any validated risk screening tool.

302

303 Alternative 1:

- 304 • BMI $<18.5 \text{ kg/m}^2$

305 Alternative 2:

- 306 • Weight loss (unintentional) $>10\%$ indefinite of time, or $>5\%$ over the last 3 months combined
307 with either

- 308 • BMI $<20 \text{ kg/m}^2$ if <70 years of age, or $<22 \text{ kg/m}^2$ if ≥ 70 years of age

309 or

- 310 • FFMI <15 and 17 kg/m^2 in women and men, respectively.

311

312 **Figure 1A and 1B: The overlap of the new ESPEN consensus definition of malnutrition and its**
 313 **individual diagnostic options in acutely ill middle-aged patients and geriatric outpatients.**

314

315 **1A: Acutely ill middle-aged patients N = 349**

316

Of the 54 malnourished patients (new ESPEN diagnostic options):

- 317 BMI < 18.5 kg/m² N=21
- 318 Unintentional weight loss + low BMI <20 kg/m² (<70 years) or 22 kg/m² (≥70 years) N=30
- 319 Unintentional weight loss + low FFMI <15 kg/m² (females) or 17 kg/m²(males) N=44

320

321

322

323

324

325

326

327

328


329

330

331

332

333



334 **1B: Geriatric outpatients N = 135**

335

Of the 10 malnourished outpatients (new ESPEN diagnostic options):

336 **——** BMI < 18.5 kg/m² N=2

337 **- - -** Unintentional weight loss + low BMI <20 kg/m² (<70 years) or 22 kg/m² (≥70 years) N=9

338 **.....** Unintentional weight loss + low FFMI <15 kg/m² (females) or 17 kg/m² (males) N=8

339

340

341

342

343

344

345

346

347

