

**Please cite the Published Version**

Rojer, AGM, Kruizenga, HM, Trappenburg, MC, Reijnierse, EM, Sipilä, S, Narici, MV, Hogrel, JY, Butler-Browne, G, McPhee, JS, Pääsuke, M, Meskers, CGM, Maier, AB and de van der Schueren, MAE (2016) The prevalence of malnutrition according to the new ESPEN definition in four diverse populations. *Clinical Nutrition*, 35 (3). pp. 758-762. ISSN 0261-5614

**DOI:** <https://doi.org/10.1016/j.clnu.2015.06.005>

**Publisher:** Elsevier

**Version:** Accepted Version

**Downloaded from:** <https://e-space.mmu.ac.uk/617227/>

**Usage rights:**



[Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0](#)

**Additional Information:** This is an author accepted manuscript of a paper accepted for publication in *Clinical Nutrition*, published by and copyright Elsevier.

**Enquiries:**

If you have questions about this document, contact [openresearch@mmu.ac.uk](mailto:openresearch@mmu.ac.uk). Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

1    **Short communication**

2    **The prevalence of malnutrition according to the new ESPEN definition in four clinically**  
3    **relevant populations**

4    A.G.M. Rojer

5    Department of Internal Medicine, Section of Gerontology and Geriatrics, VU University Medical  
6    Center, Amsterdam, The Netherlands

7    H.M. Kruizenga PhD, RD

8    Department of Nutrition and Dietetics, Internal Medicine, VU University Medical Center, Amsterdam,  
9    The Netherlands

10   Dutch Malnutrition Steering Group, Amsterdam, The Netherlands

11   M.C. Trappenburg, PhD, MD

12   Department of Internal Medicine, Section of Gerontology and Geriatrics, VU University Medical  
13   Center, Amsterdam, The Netherlands

14   Department of Internal Medicine, Amstelland Hospital, Amstelveen, The Netherlands

15   E.M. Reijnierse, MSc

16   Department of Internal Medicine, Section of Gerontology and Geriatrics, VU University Medical  
17   Center, Amsterdam, The Netherlands

18   S. Sipilä, PhD Gerontology Research Centre and Department of Health Sciences, University of  
19   Jyväskylä, Jyväskylä, Finland

20   M.V. Narici, PhD Division of Medical Sciences & Graduate Entry Medicine, MRC-ARUK Centre of  
21   Excellence for Musculoskeletal Ageing Research, University of Nottingham, Royal Derby Hospital  
22   Centre.

23   J.Y. Hogrel, PhD UPMC UM 76, INSERM U 974, CNRS 7215, Institut de Myologie, Paris, France

24   G. Butler-Browne UPMC UM 76, INSERM U 974, CNRS 7215, Institut de Myologie, Paris, France

25 J.S. McPhee, PhD School of Healthcare Science, John Dalton Building, Manchester Metropolitan  
26 University, Manchester M1 5GD, UK

27 H. Gapeyeva, PhD Institute of Exercise Biology and Physiotherapy, University of Tartu, Centre of  
28 Behavioral and Health Sciences, University of Tartu, Tartu, Estonia

29 C.G.M. Meskers, PhD, MD  
30 Department of Rehabilitation Medicine, VU University Medical Center, Amsterdam, The Netherlands

31 A.B. Maier, PhD, MD  
32 Department of Internal Medicine, Section of Gerontology and Geriatrics, VU University Medical  
33 Center, Amsterdam, The Netherlands

34 M.A.E. de van der Schueren, PhD, RD  
35 Department of Nutrition and Dietetics, Internal Medicine, VU University Medical Center, Amsterdam,  
36 The Netherlands

37 Department of Nutrition, Sports and Health, Faculty of Health and Social Studies, HAN University of  
38 Applied Sciences, Nijmegen, The Netherlands

39

40 Corresponding Author: Dr. M.A.E. de van der Schueren  
41 E-mail: [m.devanderschueren@vumc.nl](mailto:m.devanderschueren@vumc.nl)

42 Tel: +31 204443410

43 Text word count: 2201

44 Tables and figures: 4

45 Abstract word count: 403

46 References: 13

## Abstract

**Background & Aims:** International consensus on the definition of malnutrition has not yet been reached. Recently, the European Society for Clinical Nutrition and Metabolism (ESPEN) proposed a new consensus definition of malnutrition. The aim of the present study was to describe the prevalence of malnutrition according to the new ESPEN consensus definition in four clinically relevant populations: acutely ill middle-aged patients, geriatric outpatients, healthy old individuals and healthy young individuals.

**Methods:** The recently released ESPEN consensus definition of malnutrition was applied to the four different populations. This definition consists of two different options: option one requires body mass index (BMI,  $\text{kg/m}^2$ )  $<18.5 \text{ kg/m}^2$  to define malnutrition. Option two requires the combined finding of unintentional weight loss (mandatory) and at least one of either reduced BMI or low fat free mass index (FFMI,  $\text{kg/m}^2$ ). Unintentional weight loss could be either  $>10\%$  of habitual weight independent of time, or  $>5\%$  over the previous 3 months. Reduced BMI is defined as  $<20 \text{ kg/m}^2$  or  $<22 \text{ kg/m}^2$  in subjects younger and older than 70 years, respectively. Low FFMI is  $<15 \text{ kg/m}^2$  and  $<17 \text{ kg/m}^2$  in females and males, respectively. Only individuals for whom all data on diagnostic options were complete were included in the present analysis: acutely ill middle-aged patients ( $n=349$ ), geriatric outpatients ( $n=135$ ), healthy old individuals ( $n=306$ ) and healthy young individuals ( $n=179$ ).

**Results:** According to the new ESPEN consensus definition of malnutrition, the prevalence of malnutrition ranged from 1% in healthy old individuals to 15% in the acutely ill middle-aged patients. The different options that compose the new ESPEN consensus definition of malnutrition were represented in the four populations in various ways, i.e., high prevalence rates of low FFMI in all four populations, a relatively high prevalence of BMI  $<18.5 \text{ kg/m}^2$  in healthy young individuals but low prevalence of BMI  $<18.5 \text{ kg/m}^2$  in all other populations and relatively low prevalence rates of the combination of weight loss with either low BMI or low FFMI.

**Conclusions:** Combining the diagnostic options that compose the new ESPEN consensus definition of malnutrition results in prevalence rates lower than expected in acutely ill middle-aged patients and geriatric outpatients. In contrast, healthy young individuals are (most likely falsely) defined malnourished based on a low BMI  $<18.5 \text{ kg/m}^2$ . Future studies should further determine the cut-off

75 points for FFMI and BMI in older persons. In addition, the association of the new ESPEN consensus  
76 definition of malnutrition with clinically relevant outcomes needs further study.

77

78 **Keywords:** Malnutrition, definition, prevalence

## 79 Introduction

80 Malnutrition is an increasingly recognized problem that is associated with morbidity, mortality, and  
81 increased costs of care. To enhance early recognition and treatment of malnutrition, an easy and  
82 widely accepted definition of malnutrition is necessary. Such a definition should be easily applied for  
83 all health care professionals and in all health care settings. Furthermore, the definition of malnutrition  
84 should be widely accepted to be able to compare prevalence rates among health care settings and  
85 countries, and to improve communication among health care providers and politicians worldwide.

86 International consensus on the definition of malnutrition has not yet been reached. Recently, the  
87 European Society for Clinical Nutrition and Metabolism (ESPEN) proposed a new consensus  
88 definition including two options for the diagnosis of malnutrition(1). The first diagnostic option  
89 requires a low body mass index (BMI), following the recommendation by the World Health  
90 Organization: subjects are defined as malnourished if they have a BMI  $<18.5 \text{ kg/m}^2$ (2). The second  
91 diagnostic option encompasses unintentional weight loss ( $>10\%$  independent of time or  $>5\%$  in the  
92 last three months), always combined with either a low BMI ( $<20 \text{ kg/m}^2$  if  $<70$  years old or  $<22 \text{ kg/m}^2$   
93 if  $\geq 70$  years old) or a low Fat Free Mass Index (FFMI). Given the increasingly recognized importance  
94 of body protein reserves, the preferred diagnostic trajectory involves the assessment of the FFMI, with  
95 cut-off points of  $15 \text{ kg/m}^2$  for women and  $17 \text{ kg/m}^2$  for men.

96 As the new ESPEN consensus definition of malnutrition has been released only recently, validation  
97 studies have not yet been published. The aim of the present study was to describe the prevalence rates  
98 of malnutrition according to the newly proposed ESPEN consensus definition of malnutrition in four  
99 clinically relevant populations including acutely ill middle-aged patients, geriatric outpatients, healthy  
100 old individuals and healthy young individuals. This study will provide a first overview of the  
101 applicability of the newly proposed consensus definitions of malnutrition in various target populations.

## Methods

The recently released ESPEN consensus definition of malnutrition (see Fact box 1), was applied to four different populations. Only individuals for whom all data on diagnostic options were complete were included for the present analysis.

### *Population 1: acutely ill, middle-aged patients*

This population consisted of 349 patients (57.6 years, SD 17.7) who were admitted to a general internal ward (general internal medicine, gastroenterology, dermatology, rheumatology, nephrology) or a general surgical ward (general surgery and surgical oncology) of the VU University Medical Center (Amsterdam, the Netherlands) in two periods respectively from April 2002 until October 2002 and from February until June 2003 (3).

### *Population 2: geriatric outpatients*

This population consisted of 135 geriatric outpatients (80.8 years, SD 7.3) who were referred to the geriatric outpatient clinic of the Bronovo Hospital (The Hague, the Netherlands) for a comprehensive geriatric assessment due to mobility problems between March 2011 and January 2012 (4).

### *Population 3 and 4: healthy old individuals and healthy young individuals*

The European MYOAGE study consisted of old and young healthy individuals. Individuals in the MYOAGE study were recruited from five different sites across Europe, including: Manchester, UK; Paris, France; Leiden, the Netherlands; Jyväskylä, Finland and Tartu, Estonia. Data was collected between 2010 and 2013(5).

Old and young healthy individuals from the MYOAGE study were analysed separately; included were 306 healthy old individuals (74.4 years, SD 3.3) and 179 healthy young individuals (23.4 years SD 2.9).

Individuals in all four populations were screened with the Short Nutritional Assessment Questionnaire (SNAQ), with  $\geq 3$  points indicating high risk of malnutrition (3). Independent of the SNAQ screening results the diagnosis of malnutrition was assessed by measured weight and height, calculated BMI, self-reported unintentional weight loss; FFMI was derived differently across the populations. In the acutely ill middle-aged population FFMI was assessed using Xitron 4000B multiple frequency Bio-electrical Impedance Spectroscopy, using its 50KHz frequency and the Geneva equations (6). In geriatric outpatients, FFM was assessed using a direct segmental multi-frequency Bio-electrical Impedance Analyser, which provided direct values for FFM, which were then divided by height<sup>2</sup> (InBody 720, Biospace Co., Ltd, Seoul, Korea). In both the old and young healthy individuals FFMI was assessed with dual-energy x-ray absorptiometry.

The prevalence of malnutrition according to the new ESPEN consensus definition, as well as to the individual diagnostic options, was calculated for each population.



## Results

Screening with SNAQ ( $\geq 3$  points) identified 105 acutely ill middle-aged patients at risk of malnutrition, 14 geriatric outpatients, 1 healthy old individual and none of the healthy young individuals. Assessment according to the new ESPEN definition (independent of initial screening with SNAQ) yielded 54 malnourished patients (15%) in the acutely ill, middle-aged patients, 10 malnourished geriatric outpatients (7%), 3 malnourished healthy old (1%) and 14 malnourished healthy young (8%). Five malnourished patients in the acutely ill middle-aged were not identified to be at risk by the initial SNAQ screening; this was 2 in the geriatric outpatients, 3 in the healthy old and 14 in the healthy young.

Table 1 depicts the prevalence data for each population. Furthermore, it shows the prevalence of the individual diagnostic options of the definition. For example: in the acutely ill middle-aged population, the prevalence of malnutrition was 15%. Out of the total population of 349 individuals, 116 had a FFMI below the proposed cut-off points; 44 individuals (13%) out of these 116 were defined as malnourished, based on the combination low FFMI and unintentional weight loss.

Figures 1A and 1B display the overlap of the new ESPEN consensus definition of malnutrition and its individual diagnostic options in the acutely ill middle-aged population and in the geriatric outpatient population. Overlap figures are not displayed for the healthy old individuals and healthy young individuals due to low number of malnourished cases in the healthy old individuals ( $n=3$ ) and unilateralism in healthy young individuals ( $n=14$  of which 13 were identified malnourished by having only a low BMI). Furthermore, in the healthy old individuals low BMI and low FFMI were never combined with unintentional weight loss.

## Discussion

The description of the prevalence of malnutrition according to the recently released ESPEN consensus definition showed relatively low prevalence rates of malnutrition in all four populations. A low BMI and a low FFMI were observed in approximately 20% of the individuals in each population. However, most individuals were eventually not identified as malnourished as low BMI/low FFMI was not combined with unintentional weight loss. Thus, the criterion of unintentional weight loss has a dominant influence when determining prevalence rates.

The first diagnostic option of the new ESPEN consensus definition of malnutrition consists of a BMI  $< 18.5 \text{ kg/m}^2$ . A BMI  $< 18.5 \text{ kg/m}^2$  was mostly observed in acutely ill middle-aged patients. However, both in geriatric outpatients and in healthy old individuals, a BMI less than  $18.5 \text{ kg/m}^2$  was rare (1% in each population). Thus, a BMI  $< 18.5 \text{ kg/m}^2$  is rare in older individuals, which is in line with other studies that report higher BMI's in older populations (7).

Thirteen (7%) young healthy individuals were defined malnourished according to a low BMI. The ESPEN diagnostic process suggests screening first, and further assessment only for those at risk. In the cohorts described, we used the SNAQ for initial screening. This resulted in no young healthy individuals at risk. i.e. no need for further assessment. However, recent research has shown that the SNAQ is not a valid screening tool for outpatients as it does not comprise BMI (8). If we had used MUST (9), for example, the 13 healthy individuals would have passed screening and been identified malnourished in the process of diagnosis, most likely falsely, as they were all selected for their excellent health. They were probably 'healthy and slim' or very athletic.

The second diagnostic option of the new ESPEN consensus definition of malnutrition consists of a combination of unintentional weight loss and either low BMI or low FFMI. In the acutely ill middle-aged population, 25% of all patients had unintentional weight loss. This is in line with expectations, as unintentional weight loss is a frequently described phenomenon accompanying acute disease. However, only 15% of the population was defined as malnourished according to the new ESPEN consensus definition of malnutrition, indicating that in 10% of the cases unintentional weight loss did

not occur in combination with a low BMI or a low FFMI. We believe that the infrequent concurrence of unintentional weight loss with low BMI (30 out of 54 malnourished acutely ill middle-aged patients) is due to the relatively high BMI's at the population level. The combination of unintentional weight loss and a low FFMI was present in 44 out of the 54 malnourished acutely ill patients.

In the geriatric outpatient population the combination of unintentional weight loss and low BMI ( $<22 \text{ kg/m}^2$  if  $\geq 70$  years old) (9 out of 10 malnourished outpatients) overlapped reasonably well with the combination of unintentional weight loss and low FFMI (8 out of 10 malnourished outpatients).

As the new definition suggests that unintentional weight loss should be combined with either a low BMI or a low FFMI to be defined as malnourished, this also suggests that a low BMI and a low FFMI can be used interchangeably. Although in geriatric outpatients, malnutrition based on low BMI or on low FFMI was equivalent, the correspondence in the acutely ill patients was lower. Larger numbers of patients are required, however, to determine how well BMI and FFMI correlate in different populations.

A low FFMI was highly prevalent (14-33%) in all populations, however prevalence of the combination of unintentional weight loss and low FFMI showed a lower prevalence (0-13%). The high prevalence of a low FFMI may be explained by the chosen cut-off points in the ESPEN consensus definition of malnutrition. The cut-off point of FFMI below  $15 \text{ kg/m}^2$  for women represents the 50<sup>th</sup> percentile, according to Schutz's reference tables (10). For men, a cut-off point of FFMI below  $17 \text{ kg/m}^2$  represents the 10<sup>th</sup> percentile, which is probably a much more realistic percentile to apply. This raises the question of whether the cut-off point for women should be amended, for example to  $14 \text{ kg/m}^2$ , which represents the 10<sup>th</sup> percentile for women (10), and what consequences that cut-off point would have for the prevalence rates. A future study should look into a possible revision of the FFMI cut-off points, their overlap with unintentional weight loss and the consequences for malnutrition prevalence rates.

Of the acutely ill middle-aged patients with a BMI  $<20 \text{ kg/m}^2$  ( $<70$  years) or  $<22 \text{ kg/m}^2$  ( $\geq 70$  years), approximately half were defined as malnourished as they also met the second diagnostic option: unintentional weight loss. In the geriatric outpatient population a low BMI ( $<20 \text{ kg/m}^2$  ( $<70$  years) or  $<22 \text{ kg/m}^2$  ( $\geq 70$  years)) occurred in 28 (21%) outpatients. Remarkably, in only 9 (7%) geriatric outpatients a low BMI was combined with unintentional weight loss; this might be one explanation for the lower than expected prevalence rates in this geriatric outpatient population. Previous studies have reported prevalence rates of malnutrition in approximately 50% of geriatric outpatients (11-13). Since geriatric outpatients usually suffer from multiple age-related problems and many co-morbidities, unintentional weight loss is most likely a problem that has occurred only slowly and thereby has not reached the cut-off level of 10%, or that has gone by unnoticed. In the healthy old individuals, 39 (13%) had a BMI  $<20 \text{ kg/m}^2$  ( $<70$  years) or  $<22 \text{ kg/m}^2$  ( $\geq 70$  years) but none were defined as malnourished based on the concurrence with unintentional weight loss; three healthy old individuals were defined as malnourished based on a BMI  $<18.5 \text{ kg/m}^2$ . For older persons, either a BMI cut-off point higher than  $22 \text{ kg/m}^2$  or a different cut-off point for unintentional weight loss are more reasonable indicators of malnutrition.

## Conclusion:

The prevalence rates of positive scores when using the different ESPEN consensus definitions of malnutrition were high. However, when combining the different diagnosis pathways, the prevalence rates were lower than expected in acutely ill middle-aged patients and in geriatric outpatients. Old healthy individuals were probably identified as malnourished too infrequently, due to missing concurrence of low BMI/low FFMI and weight loss, whereas in contrast young healthy individuals were (most likely falsely) defined malnourished based on a low BMI.

Some suggestions for further studies:

- To study the importance of the relative contribution of unintentional weight loss versus low BMI or low FFMI in the new ESPEN consensus definition of malnutrition.

- To reconsider the proposed cut-off points for FFMI, specifically for women. Both absolute cut-off points and age- and sex- specific percentiles should be studied.

- To study whether a low BMI and a low FFMI are interchangeable and whether this is different between populations.

- To evaluate the proposed BMI cut-off point of  $< 22 \text{ kg/m}^2$  or the degree of unintentional weight loss in older adults. This descriptive study even raises the question whether BMI is a relevant parameter for nutritional status in older adults at all or whether we should more strongly rely on FFMI in older adults.

In future analyses, we will report on the association between the new ESPEN consensus definition of malnutrition, its individual diagnostic options and clinically relevant outcome measures such as functionality and survival, which will shed a further light on the chosen cut-off points for BMI and FFMI.

## References

- (1) Cederholm T, Bosaeus I, Barazzoni R, Bauer J, Van Gossum A, Klek S, et al. Diagnostic criteria for malnutrition - an ESPEN Consensus Statement. *Clinical Nutrition*, submitted for publication.
- (2) WHO. Physical status: the use and interpretation of anthropometry. Report of a WHO Expert Committee. 1995; World Health Organization Technical Report Series 854.
- (3) Kruizenga HM, Seidell JC, de Vet HCW, Wierdsma NJ, Van Bokhorst-de Van der Schueren, Development and validation of a hospital screening tool for malnutrition: the short nutritional assessment questionnaire (SNAQ). *Clin Nutr* 2005 24;24(1):75-82.
- (4) Bijlsma AY, Pasma JH, Lambers D, Stijntjes M, Blauw GJ, Meskers CG, et al. Muscle strength rather than muscle mass is associated with standing balance in elderly outpatients. *J Am Med Dir Assoc* 2013 Jul;14(7):493-498.
- (5) McPhee JS, Hogrel JY, Maier AB, Seppet E, Seynnes OR, Sipila S, et al. Physiological and functional evaluation of healthy young and older men and women: design of the European MyoAge study. *Biogerontology* 2013 Jun;14(3):325-337.
- (6) Kyle UG, Genton L, Karsegard L, Slosman DO, Pichard C. Single prediction equation for bioelectrical impedance analysis in adults aged 20--94 years. *Nutrition* 2001 03;17(0899-9007; 3):248-253.
- (7) Winter JE, MacInnis RJ, Wattanapenpaiboon N, Nowson CA. BMI and all-cause mortality in older adults: a meta-analysis. *Am J Clin Nutr* 2014 Apr;99(4):875-890.
- (8) Leistra E, Langius JA, Evers AM, van Bokhorst-de van der Schueren, M.A., Visser M, de Vet HC, et al. Validity of nutritional screening with MUST and SNAQ in hospital outpatients. *Eur J Clin Nutr* 2013 May 1.
- (9) Elia M. The 'MUST' report. Nutritional screening of adults: a multidisciplinary responsibility. 2003.
- (10) Schutz Y, Kyle UU, Pichard C. Fat-free mass index and fat mass index percentiles in Caucasians aged 18-98 y. *Int J Obes Relat Metab Disord* 2002 07;26(0307-0565; 7):953-960.
- (11) van Bokhorst-de van der Schueren, M.A., Lonterman-Monasch S, de Vries OJ, Danner SA, Kramer MH, Muller M. Prevalence and determinants for malnutrition in geriatric outpatients. *Clin Nutr* 2013 May 17;32:1007-1011.
- (12) Saka B, Kaya O, Ozturk GB, Erten N, Karan MA. Malnutrition in the elderly and its relationship with other geriatric syndromes. *Clin Nutr* 2010 12;29(1532-1983; 0261-5614; 6):745-748.
- (13) Ulger Z, Halil M, Kalan I, Yavuz BB, Cankurtaran M, Gungor E, et al. Comprehensive assessment of malnutrition risk and related factors in a large group of community-dwelling older adults. *Clin Nutr* 2010 08;29(1532-1983; 0261-5614; 4):507-511.

290

291   **Attachments:**

- 292           -   Fact box: Two alternative ways to diagnose malnutrition.
- 293           -   Figure 1A and 1B: The overlap of the new ESPEN consensus definition of malnutrition
- 294               and its individual diagnostic options in acutely ill middle-aged patients and geriatric
- 295               outpatients.
- 296           -   Table 1: Prevalence rates of malnutrition according to the new ESPEN consensus
- 297               definition and to its individual diagnostic options in four populations.

298

299 **Fact box: Two alternative ways to diagnose malnutrition.**

300 Before diagnosis of malnutrition is considered it is mandatory to fulfil criteria for being “at risk” of  
301 malnutrition by any validated risk screening tool.

302

303 Alternative 1:

- 304 • BMI  $<18.5 \text{ kg/m}^2$

305 Alternative 2:

- 306 • Weight loss (unintentional)  $>10\%$  indefinite of time, or  $>5\%$  over the last 3 months combined  
307 with either

- 308 • BMI  $<20 \text{ kg/m}^2$  if  $<70$  years of age, or  $<22 \text{ kg/m}^2$  if  $\geq 70$  years of age

309 or

- 310 • FFMI  $<15$  and  $17 \text{ kg/m}^2$  in women and men, respectively.

311




**Figure 1A and 1B: The overlap of the new ESPEN consensus definition of malnutrition and its individual diagnostic options in acutely ill middle-aged patients and geriatric outpatients.**

**1A: Acutely ill middle-aged patients N = 349**

Of the 54 malnourished patients (new ESPEN diagnostic options):

—	BMI < 18.5 kg/m <sup>2</sup>	N=21
- - -	Unintentional weight loss + low BMI <20 kg/m <sup>2</sup> (<70 years) or 22 kg/m <sup>2</sup> (≥70 years)	N=30
.....	Unintentional weight loss + low FFMI <15 kg/m <sup>2</sup> (females) or 17 kg/m <sup>2</sup> (males)	N=44



1B: Geriatric outpatients N = 135

Of the 10 malnourished outpatients (new ESPEN diagnostic options ):

- BMI < 18.5 kg/m<sup>2</sup> N=2
- - - Unintentional weight loss + low BMI <20 kg/m<sup>2</sup> (<70 years) or 22 kg/m<sup>2</sup> (≥70 years) N=9
- ..... Unintentional weight loss + low FFMI <15 kg/m<sup>2</sup> (females) or 17 kg/m<sup>2</sup> (males) N=8

