

Please cite the Published Version

Redfern, J and Verran, J (2017) Effect of humidity and temperature on the survival of *Listeria monocytogenes* on surfaces. *Letters in Applied Microbiology*, 64 (4). pp. 276-282. ISSN 0266-8254

DOI: <https://doi.org/10.1111/lam.12714>

Publisher: Wiley/The Society for Applied Microbiology

Version: Accepted Version

Downloaded from: <https://e-space.mmu.ac.uk/617893/>

Usage rights: © In Copyright

Additional Information: This is an Author Accepted Manuscript of a paper accepted for publication in *Letters in Applied Microbiology*, published by Wiley and copyright The Society for Applied Microbiology.

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16

Title page

Effect of humidity and temperature on the survival of *Listeria monocytogenes* on surfaces

James Redfern¹ and Joanna Verran¹.

¹School of Healthcare Science, Manchester Metropolitan University, Chester Street, Manchester, M1 5GD, United Kingdom.

James Redfern – J.Redfern@mmu.ac.uk

Joanna Verran – J.Verran@mmu.ac.uk

Joanna Verran to be corresponding author once published.

Running title: *Humidity and L. monocytogenes*

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Significance and Impact

Understanding survival of potential food-borne pathogens is essential to the safe production and preparation of food. Whilst it has long been ‘common knowledge’ that relative humidity can affect the growth and survival of microorganisms, this study systematically describes the survival of *L. monocytogenes* on stainless steel under varying humidity and temperatures for the first time. The outcomes from this paper will allow those involved with food manufacture and preparation to make informed judgement on environmental conditions relating to humidity control, which is lacking in the food standards guidelines.

Abstract

Listeria monocytogenes is a pathogenic bacterium, with human disease and infection linked to dairy products, seafood, ready-to-eat meat and raw & undercooked meats. Stainless steel is the most common food preparation surface and therefore, it is important to understand how food storage conditions such as surface materials, temperature and relative humidity can affect survival of *L. monocytogenes*. In this study, survival of *L. monocytogenes* on stainless steel was investigated at three temperatures (4, 10 and 21°C), each approx. 11%, 50% and 85% humidity. Results indicate that the lower the temperature, the more cells were recovered in all three humidity environments, whilst medium humidity enhances survival, irrespective of temperature. Lower humidity decreases recovery at all temperatures. These data support the guidance noted above that humidity control is important, and that lower humidity environments are less likely to support retention of viable *L. monocytogenes* on a stainless steel surface.

Keywords

42 Food
43 Listeria
44 Food preparation
45 Humidity
46 Stainless steel

47 **Introduction**

48 *Listeria monocytogenes* is a pathogenic bacterium, with human disease and infection
49 linked to dairy products, seafood, ready-to-eat meat and raw & undercooked meats.
50 Listeriosis, encompassing bacterial meningitis, sepsis, endocarditis, neonatal abortion
51 and stillbirth in humans (Schlech et al. 1983), usually presenting in those already
52 immunosuppressed, pregnant, old or young (Scholing et al. 2007; Barocchi et al. 2015).
53 During the late 1990s there was a large outbreak of listeriosis linked primarily to
54 consumption of pâté (McLauchlin et al. 1991). Investigations resulted in the discovery of
55 *Listeria* in cheese and other cook-chill foods, subsequently leading to an increase in
56 regulation surrounding chilled food storage (ACMSF 2003).

57 Studies on the interaction between *L. monocytogenes* and stainless steel, the most
58 common surface used in food preparation, have found that the survival of the
59 microorganism on the surface alters depending on contact time, temperature, nutrients,
60 moisture and the presence of other microorganisms (Bremer et al. 2001; Poimenidou et
61 al. 2009; Skovager et al. 2013a). Additionally, survival of *L. monocytogenes* can be
62 decreased by introducing antimicrobial compounds such as Lauric Arginate into
63 stainless steel (Saini et al. 2013), or by coating a stainless steel surface with an
64 antimicrobial film, for example, TiN/Ag (Skovager et al. 2013b). However, inert stainless
65 steel is the most suitable for the food industry due to its non-toxic, easy-clean,
66 mechanically stable and corrosion-resistant properties (EHEDG 2004). In short, if
67 contaminated food product requires preparation prior to packaging/cooking, for

68 example in a food processing plant, surfaces such as stainless steel worktops or
69 conveyor belts pose cross-contamination potential. Whilst this is not the only source of
70 contamination, with factors such as hygiene and disinfection being important, the
71 environmental conditions are critical to ensure there is little opportunity for growth of
72 microorganisms on surfaces and that survival is minimal.

73 Although surface characteristics such as roughness and wettability are important
74 variables when considering survival of microorganisms on steel, other environmental
75 conditions are likely to play a key role. An increase in relative humidity (RH), a measure
76 relating to amount of water vapour in the atmosphere, has been shown to prolong
77 survival of *L. monocytogenes*, as well as encourage growth when inoculated on fresh
78 produce (Likotrafiti et al. 2013), whilst a decrease in RH has demonstrated a decreased
79 survival of *L. monocytogenes* (Zoz et al. 2016). Conversely, reduction in RH has been
80 shown to enhance transfer of *L. monocytogenes* from biofilm to meat products
81 potentially due to increased capillary action within the food (Rodríguez et al. 2007).

82 Control of relative humidity in relation to control of microbial contamination in food
83 processing environments is suggested by many governments around the world (e.g. FDA
84 2009; Abu Dhabi Food Control Authority 2010; FSA 2015), and advice is available
85 (EHEDG 2006). However these documents do not recommend specific levels of RH,
86 likely due to the complex and unique nature of each food processing environment.

87 The ability for *L. monocytogenes* not only to survive but also to grow across a relatively
88 wide temperature range, often described in the literature as between 2°C to 45°C, means
89 that refrigerated food is not necessarily protected from microbial colonisation by *L.*
90 *monocytogenes* (Gandhi and Chikindas 2007). Given the variety of surface materials,
91 temperatures and RH combinations possible in the manufacture, transport and
92 consumption of food, it is important therefore to understand the effect of temperature
93 and RH on the survival of *L. monocytogenes* on surfaces. This study will investigate the

survival of *L. monocytogenes* on stainless steel in three different humidity-controlled environments, selected as examples of the possible range of humidity in a food processing location (although not all are likely to be encountered - approx. 11%, 50% and 85%), at three different temperatures.

Results and Discussion

The aim of this study was to investigate the survival of *L. monocytogenes* on stainless steel over time with respect to temperature and humidity. The experiment used *L. monocytogenes* in its planktonic state as inoculum. Biofilm is unlikely to form in this environment because good hygiene practice should remove the possibility of *L. monocytogenes* building a biofilm on a food preparation surface. The focus was survival since growth was unlikely.

Surface profiles

The average Ra value for SS 304 was 42.65nm whilst the average Ra value for SS 316 was 41.12nm. There was no significant difference ($P>0.05$) in the Ra values between the two surface types, but surfaces were visually different, with SS 304 appearing smoother with fewer defects compared to SS 316 (figure 1).

Recovery of cells from SS 304 following incubation in controlled humidity and temperature

The viability of cells recovered from the sample in low humidity decreased as time and temperature increased (figure 2). After one hour, no cells were recovered from any surface.

At medium humidity (figure 3), as temperature increased, viability decreased, although this is less obvious than at low humidity. At 4°C there was no decrease in survival, indeed the opposite was observed, with the number of cells recovered increasing.

As time and temperature increased, viability was also reduced at high humidity (figure 4). This decrease was statistically significant ($p < 0.05$) between 5h and 7h at 4°C and 10 °C ($P > 0.05$).

Overall, it appears a medium level of humidity is optimum for survival of *L. monocytogenes* on SS 304, with the change of humidity being most important in supporting survival irrespective of temperature.

Recovery of living cells from SS 316 following incubation in controlled humidity and temperature

No cells were recovered at low humidity/21°C on SS 316 after incubation (figure 5). Cells recovered after incubation at high humidity/4°C (figure 5) reduced following a similar trend to that observed on SS 304.

Acridine orange (AO) staining of SS 304 and SS 316 to assess retention on surface after swabbing

The average percentage coverages of cells on SS 304 and SS 316 were 74.97% and 65.65% respectively, when unswabbed coupons were visualised with AO. After swabbing the coverage decreased significantly ($p < 0.05$). There was no significant difference ($P > 0.05$) in the percentage average of cells on the surfaces, with SS 304 and SS 316 presenting 2.08% and 3.59% respectively, indicating effective swabbing.

During the study it was observed that samples incubated at medium or high humidity became wet, despite being dried before incubation, likely due to the water vapour in the environment. It has been shown previously that the presence of moisture on a surface can loosen cells from a surface and increase the number of cells recovered by swabbing (Verran et al. 2010), which is a possible explanation for the varied counts recovered. It is also possible that as the inoculum is rehydrated, any cell division initiated might continue, increasing the number of recovered cells.

143 A critique of this methodology is the equal drying time and conditions each sample
144 received prior to incubation in different temperatures and humidity. Whilst it was
145 important in this study to control the drying conditions to be able to draw comparisons,
146 the authors acknowledge that within a real life scenario it is possible that contamination
147 will 'dry' dependant on the ambient humidity it is stored in, which is likely to vary the
148 survival time of the microorganism.

149 Findings show that the lower the temperature, the more cells are recovered from steel
150 when incubated in any of the three humidity environments. Not many cells are retained
151 on the surface, so essentially viability is indicated by recovery. Interestingly, studies on
152 survival of *L. monocytogenes* on biotic surfaces, for example Likotrafiti et al. (2013), have
153 shown that a reduced temperature decreases the number of recovered cells when in low
154 humidity environments.

155 Results relating to SS 316 show no significant difference between survival in relation to
156 temperature and humidity, with very few cells remaining on the surface after swabbing.
157 These data indicate that the application of a finish to steel (for example, bright
158 annealed) did not affect ease of cleanliness.

159 However, the data suggest that "medium" humidity enhances survival, irrespective of
160 temperature, presumably because of a decrease in stress to cells. Lower humidity
161 decreases recovery at all temperatures, whilst high humidity decreases recovery at high
162 temperatures, presumably due to an increase in stress.

163 It is likely *in situ* that humidity will be controlled within the food industry environments,
164 however, as discussed in the introduction, humidity control is not dictated by
165 legislation, and is therefore likely to be variable across the sector. Low and high humid
166 environments can be uncomfortable and potentially dangerous to human health (Davis
167 et al. 2016), and therefore a humidity closer to 50% is more likely. However, in a food
168 processing environment, personnel are not the focus: the results of this study suggest

this is the least favourable option for reducing viable *L. monocytogenes* on stainless steel.

It is likely that environments may where food is prepared and/or stored with no humidity control. Whilst no specific guidance could be found for humidity control in such circumstances, it is recognised as one measure for the control of bacterial contamination. Our data support the guidance referenced earlier that humidity control is important, and that lower humidity environments are less likely to support retention and survival of viable *L. monocytogenes* on a stainless steel surface. It is likely that storage will always be at a low temperature, so humidity control is critical if the low temperature itself increases survival.

Materials and Methods

Microorganisms

Listeria monocytogenes Scott A, serotype 4 (kindly donated by Professor Lone Gram (Danish Institute of Fisheries Research (DIFRES), Technical University of Denmark) (Briers et al. 2011) was maintained on Tryptone Soya Agar (TSA) (Oxoid, Basingstoke) at 5°C and inoculated into 100 ml⁻¹ Tryptone Soya Broth (TSB) (Oxoid). Cultures were grown overnight (22 ± 1h) at 30°C with agitation (225 rpm). Cells were harvested by centrifugation (3600 rpm, 10 min, room temperature) and washed once in 0.85% NaCl (Oxoid), resuspended to optical density (540nm) of 1.0. A 1 ml⁻¹ sample from the cell suspension was serially diluted, plated out onto NA and CFU counted, finding the cell concentration to be 3.18 ± 0.65 x10⁹ CFU/ml⁻¹. This was used for the initial inoculum of stainless steel coupons.

Preparation of stainless steel

193 Bright annealed 304 stainless steel (SS 304) and 2B 316 stainless steel (SS 316)
194 (Outokumpu, Sheffield, UK) were cut into coupons (2cm x 2cm x 1mm) using a
195 guillotine. The steel coupons were soaked in 96% ethanol overnight to
196 remove/inactivate microorganisms and remove grease from the surface (BSSA n.d.),
197 after which they were rinsed with distilled water and air dried for one hour in a class
198 two cabinet (BH-EN 2003, Faster, Cornaredo).

199 **White light profilometry**

200 A MicroXAM (phase shift) surface mapping microscope (ADE; Omniscan, Wrexham) with an
201 analogue to digital (AD) phase shift controller (Omniscan) was coupled with an image
202 analysis system (Mapview AE 2.17; Omniscan) to visualise the surface and provide Ra values.

203 **Humidity control**

204 Humidity was controlled using saturated salt solutions contained within a desiccator
205 chamber (250mm diameter, Fischer Scientific, Loughborough UK). Salts used were;
206 lithium chloride (Fischer Scientific) to achieve a low humidity approximately 11%RH,
207 magnesium chloride (Fischer Scientific) to achieve a medium humidity approximately
208 50%RH and potassium sulphate (Fischer Scientific) to achieve a high humidity
209 approximately 85%RH (Rockland 1960). Water was added to the salts until a slushy
210 mixture filled the bottom of the chamber. The saturated salt solution was left in the
211 chamber for 24 hours prior to the start of the experiment to allow the desired RH to be
212 attained. Relative humidity and temperature were monitored with a mobile USB data
213 logger (RHT10, Extech Instruments, Boston, USA).

214 **The effect of humidity and temperature on the survival of *Listeria monocytogenes*** 215 **on stainless steel**

216 Stainless steel coupons were inoculated with $10\mu\text{l}^{-1}$ of standardised *Listeria*
217 *monocytogenes* Scott A planktonic cell suspension, and spread across the surface using a

sterile pipette tip. Coupons were left to dry for 30 minutes in a class two cabinet at room temperature prior to being placed in the desiccator containing the appropriate saturated salt solution on a platform approximately 4cm above the salt solution. The desiccator was then placed inside an incubator at the appropriate temperature. At each sample time, each coupon was swabbed with a moist swab which was placed in 10ml⁻¹ of 0.85% saline and diluted to 10⁻⁸. Dilutions were plated out onto TSA, incubated for 24h at 30°C and colonies counted.

Variables investigated were low, medium and high humidity, each at 4°C, 10°C and 21°C on SS 304. Low humidity and 21°C and high humidity and 4°C were investigated on SS 316. All temperatures were maintained to within 1°C, except at sampling time when temperature could vary $\pm 3^\circ\text{C}$. Sampling was carried out at 0h, 1h, 5h, 7h and 24h hours. Three replicates of each surface were tested at each time point. Experiments were repeated once.

Bacterial staining to assess swabbing effectiveness adapted from Airey and Verran (2007).

Cells retained on sample coupons, pre and post swabbing, were stained with acridine orange (Sigma, Dorset) (0.03% in 2% glacial acetic acid) (VWR, Lutterworth), and the surfaces were rinsed and dried before examination with epifluorescence microscopy (x100) (Nikon Eclipse E600; Nikon UK Ltd, Surry). Ten random fields of each replicate surface were examined. The percentage of an area of each microscopic field covered by cells was calculated by using cell F software (Olympus Soft Imaging Solutions). The experiment was repeated once.

Data analysis

Data were analysed in SPSS® 21 for Windows (IBM, USA) and Excel® 2013 (Microsoft, USA). Statistically significant differences were tested for using a one-way ANOVA. Data

243 are presented as percentage changes compared to the CFU ml⁻¹ recovered from steel
244 sample before incubation. Initial recovered CFU ml⁻¹ can be found in the caption for the
245 corresponding figure.

246

247 **Conflict of Interest**

248 No conflict of interest declared

249

250

251

252 References

253

- 254 Abu Dhabi Food Control Authority (2010) Food Hygiene Throughout The Food Chain. In *REG-*
255 *006/2010* ed. Authority, A.D.F.C. Abu Dhabi: Abu Dhabi Food Control Authority.
- 256 ACMSF (2003) Recent Trends in Listeriosis in the UK ed. Agency, F.S. London: Food Standards
257 Agency.
- 258 Airey, P. and Verran, J. (2007) Potential use of copper as a hygienic surface; problems
259 associated with cumulative soiling and cleaning. *J Hosp Infect* **67**, 271-277.
- 260 Barocci, S., Mancini, A., Canovari, B., Petrelli, E., Sbriscia-Fioretti, E., Licci, A., D'Addesa, S.,
261 Petrini, G., Giacomini, M., Renzi, A., Migali, A. and Briscolini, S. (2015) Listeria
262 monocytogenes meningitis in an immunocompromised patient. *New Microbiol* **38**, 113-118.
- 263 Bremer, P.J., Monk, I. and Osborne, C.M. (2001) Survival of Listeria monocytogenes Attached
264 to Stainless Steel Surfaces in the Presence or Absence of Flavobacterium spp. *J Food Prot* **64**,
265 1369-1376.
- 266 Briers, Y., Klumpp, J., Schuppler, M. and Loessner, M.J. (2011) Genome Sequence of Listeria
267 monocytogenes Scott A, a Clinical Isolate from a Food-Borne Listeriosis Outbreak. *J Bacteriol*
268 **193**, 4284-4285.
- 269 BSSA (n.d.) Cleaning methods for stainless steel. Sheffield: BSSA.
- 270 Davis, R.E., McGregor, G.R. and Enfield, K.B. (2016) Humidity: A review and primer on
271 atmospheric moisture and human health. *Environ Res* **144**, Part A, 106-116.
- 272 EHEDG (2004) *EHEDG Guidelines: Hygienic Equipment Design Criteria*. Frankfurt, Germany:
273 EHEDG.
- 274 EHEDG (2006) Guidelines on air handling in the food industry. *Trends Food Sci Technol* **17**,
275 331-336.
- 276 FDA (2009) FDA Food Code 2009. In *Chapter 3 - Food* ed. Administration, U.S.F.a.D. MD, USA:
277 FDA.
- 278 FSA (2015) Meat Industry Guide - Chapter 10 Temperature Controls ed. Agency, F.S. London:
279 Food Standards Agency.
- 280 Gandhi, M. and Chikindas, M.L. (2007) Listeria: A foodborne pathogen that knows how to
281 survive. *Int J Food Microbiol* **113**, 1-15.
- 282 Likotrafiti, E., Smirniotis, P., Nastou, A. and Rhoades, J. (2013) Effect of Relative Humidity
283 and Storage Temperature on the Behavior of Listeria monocytogenes on Fresh Vegetables. *J*
284 *Food Saf* **33**, 545-551.
- 285 McLauchlin, J., Hall, S.M., Velani, S.K. and Gilbert, R.J. (1991) Human Listeriosis And Pate;: A
286 Possible Association. *BMJ: British Medical Journal* **303**, 773-775.
- 287 Poimenidou, S., Belessi, C.A., Giaouris, E.D., Gounadaki, A.S., Nychas, G.-J.E. and Skandamis,
288 P.N. (2009) Listeria monocytogenes Attachment to and Detachment from Stainless Steel
289 Surfaces in a Simulated Dairy Processing Environment. *Appl Environ Microbiol* **75**, 7182-
290 7188.
- 291 Rockland, L.B. (1960) Saturated Salt Solutions for Static Control of Relative Humidity
292 between 5 and 40 C. *Anal Chem* **32**, 1375-1376.
- 293 Rodríguez, A., Autio, W.R. and McLandsborough, L.A. (2007) Effect of Biofilm Dryness on the
294 Transfer of Listeria monocytogenes Biofilms Grown on Stainless Steel to Bologna and Hard
295 Salami. *J Food Prot* **70**, 2480-2484.
- 296 Saini, J.K., Barrios, M.A., Marsden, J.L., Getty, K.J.K. and Fung, D.Y.C. (2013) Efficacy of
297 Antimicrobial Lauric Arginate against Listeria monocytogenes on Stainless Steel Coupons.
298 *Adv Microbiol* **3**, 4.

Schlech, W.F., Lavigne, P.M., Bortolussi, R.A., Allen, A.C., Haldane, E.V., Wort, A.J., Hightower, A.W., Johnson, S.E., King, S.H., Nicholls, E.S. and Broome, C.V. (1983) Epidemic Listeriosis — Evidence for Transmission by Food. *N Engl J Med* **308**, 203-206.

Scholing, M., Schneeberger, P.M., van den Dries, P. and Drenth, J.P.H. (2007) Clinical Features of Liver Involvement in Adult Patients with Listeriosis. Review of the Literature. *Infection* **35**, 212-218.

Skovager, A., Larsen, M.H., Castro-Mejia, J.L., Hecker, M., Albrecht, D., Gerth, U., Arneborg, N. and Ingmer, H. (2013a) Initial adhesion of *Listeria monocytogenes* to fine polished stainless steel under flow conditions is determined by prior growth conditions. *Int J Food Microbiol* **165**, 35-42.

Skovager, A., Whitehead, K., Wickens, D., Verran, J., Ingmer, H. and Arneborg, N. (2013b) A comparative study of fine polished stainless steel, TiN and TiN/Ag surfaces: Adhesion and attachment strength of *Listeria monocytogenes* as well as anti-listerial effect. *Colloids Surf B Biointerfaces* **109**, 190-196.

Verran, J., J.Redfern, Smith, L.A. and Whitehead, K.A. (2010) A critical evaluation of sampling methods used for assessing microorganisms on surfaces. *Food and Bioprocess Processing* **88**.

Zoz, F., Iaconelli, C., Lang, E., Iddir, H., Guyot, S., Grandvalet, C., Gervais, P. and Beney, L. (2016) Control of Relative Air Humidity as a Potential Means to Improve Hygiene on Surfaces: A Preliminary Approach with *Listeria monocytogenes*. *PLoS ONE* **11**, e0148418.





Figure 1 –Example WLP images of SS 304 (left) and SS 316 (right) taken at x50 magnification.




326
327

328 Figure 2 – Percentage of viable cells recovered from SS 304 over 24 hours in a low
 329 humidity environment (approximately 11%RH) at three different temperatures (4°C,
 330 10°C and 21°C). Percentages are based on the number of recovered cells before applying
 331 treatment: 4°C = 2.4×10^4 cfu/ml, 10°C = 3.37×10^4 cfu/ml, 21°C = 7.27×10^4 cfu/ml. n=30
 332 for each time point.

333

334

335




336

337 Figure 3 - Percentage of viable cells recovered from SS 304 over 24 hours in a medium
 338 humidity environment (approximately 52%RH) at three different temperatures (4°C,
 339 10°C and 21°C). Percentages are based on the number of recovered cells before applying
 340 treatment: 4°C = 7.93×10^3 cfu/ml, 10°C = 2.01×10^4 cfu/ml, 21°C = 3.96×10^4 cfu/ml.
 341 n=30 for each time point.

342

343

344




345

346 Figure 4 - Percentage of viable cells recovered from SS 304 over 24 hours in a high
347 humidity environment (approximately 86%RH) at three different temperatures (4°C,
348 10°C and 21°C). Percentages are based on the number of recovered cells before applying
349 treatment: 4°C = 2.93×10^5 cfu/ml, 10°C = 9.09×10^5 cfu/ml, 21°C = 7.89×10^4 cfu/ml.
350 n=30 for each time point.

351

352



353

354 Figure 5 – Percentage of viable cells recovered from SS 316 over 24 hours in either a
 355 high humidity and low temperature environment or a low humidity high temperature
 356 environment. Percentages are based on the number of recovered cells before applying
 357 treatment: high humidity/4°C = 1.83×10^4 cfu/ml, low humidity/21°C = 6×10^4 cfu/ml.
 358 n=30 for each time point.

359