

1 Two centuries of mastling data for European beech and Norway spruce across the European  
2 continent

3 DAVIDE ASCOLI,<sup>1,20</sup> JANET MARINGER,<sup>2,4</sup> ANDY HACKET-PAIN<sup>3</sup>, MARCO CONEDERA<sup>4</sup>, IGOR  
4 DROBYSHEV<sup>5,6</sup>, RENZO MOTTA<sup>7</sup>, MARA CIROLI<sup>7</sup>, WŁADYSŁAW KANTOROWICZ<sup>8</sup>, CHRISTIAN ZANG<sup>9</sup>,  
5 SILVIO SCHUELER<sup>10</sup>, LUC CROISÉ<sup>11</sup>, PIETRO PIUSSI<sup>12</sup>, ROBERTA BERRETTI<sup>7</sup>, CIPRIAN PALAGHIANU<sup>13</sup>,  
6 MARJANA WESTERGREN<sup>14</sup>, JONATHAN G.A. LAGEARD<sup>15</sup>, ANTON BURKART<sup>16</sup>, REGULA GEHRIG  
7 BICHSEL<sup>17</sup>, PETER A. THOMAS<sup>18</sup>, BURKHARD BEUDERT<sup>19</sup>, ROLF ÖVERGAARD<sup>5</sup>, AND GIORGIO  
8 VACCHIANO<sup>7</sup>

9

10 <sup>1</sup> *Dipartimento di Agraria, University of Naples Federico II, via Università 100, 80055 Portici,*  
11 *Napoli, Italy*

12 <sup>2</sup> *Institute for Landscape Planning and Ecology, University of Stuttgart, Keplerstr. 11, 70174*  
13 *Stuttgart, Germany*

14 <sup>3</sup> *Fitzwilliam College, Storeys Way, Cambridge, United Kingdom and St Catherine's College,*  
15 *Manor Road, Oxford, United Kingdom*

16 <sup>4</sup> *Insubric Research Group, Swiss Federal Research Institute WSL, a Ramèl 18, 6593 Cadenazzo,*  
17 *Switzerland*

18 <sup>5</sup> *Southern Swedish Forest Research Centre, Swedish University of Agricultural Sciences, P.O. Box*  
19 *49, 230 53 Alnarp, Sweden*

20 <sup>6</sup> *Chaire industrielle CRSNG-UQAT-UQAM en aménagement forestier durable, Université du*  
21 *Québec en Abitibi-Témiscamingue (UQAT), 445 boul. de l'Université, Rouyn-Noranda, Québec,*  
22 *J9X 5E4, Canada*

23 <sup>7</sup> *Department of Agriculture, Forest and Food Sciences, University of Turin, Largo Braccini 2,*  
24 *10095 Grugliasco, Italy*

25 <sup>8</sup> *Department of Silviculture and Genetics of Forest Trees, Forest Research Institute, Sekocin Stary,*  
26 *Poland*

- 27      <sup>9</sup> TUM School of Life Sciences Weihenstephan, Technische Universität München, Freising,  
28      Germany
- 29      <sup>10</sup> Department of Forest Genetics, Federal Research and Training Centre for Forests, Natural  
30      Hazards and Landscapes (BFW), Seckendorff-Gudent-Weg 8, 1131 Vienna, Austria
- 31      <sup>11</sup> Département recherche-développement-innovation, RENECAFOR, Office national des forêts,  
32      Boulevard de Constance, 77300 Fontainebleau, France
- 33      <sup>12</sup> Former professor at the University of Firenze, Italy
- 34      <sup>13</sup> Forestry Faculty, Applied Ecology Lab, Stefan cel Mare University of Suceava, Universitatii  
35      Street 13, 720229, Romania
- 36      <sup>14</sup> Slovenian Forestry Institute, Ljubljana, Slovenia
- 37      <sup>15</sup> Division of Geography and Environmental Management, Manchester Metropolitan University,  
38      Manchester, United Kingdom
- 39      <sup>16</sup> Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Disturbance Ecology,  
40      Zürcherstrasse 111, 8903 Birmensdorf, Switzerland
- 41      <sup>17</sup> Federal Department of Home Affairs FDHA, Federal Office of Meteorology and Climatology  
42      MeteoSwiss, Zurich, Switzerland
- 43      <sup>18</sup> School of Life Sciences, Keele University, Newcastle under Lyme, United Kingdom
- 44      <sup>19</sup> Nationalparkverwaltung Bayerischer Wald, Sachgebiet Naturschutz und Forschung, Freyungen  
45      Str. 2, D-94481 Grafenau, Germany
- 46
- 47      <sup>20</sup> E-mail: davide.ascoli@unina.it
- 48
- 49
- 50
- 51

52    **Abstract**

53    Tree masting is one of the most intensively studied ecological processes. It affects nutrient fluxes of  
54    trees, regeneration dynamics in forests, animal population densities, and ultimately influences  
55    ecosystem services. Despite a large volume of research focused on masting, its evolutionary  
56    ecology, spatial and temporal variability and environmental drivers are still matter of debate.  
57    Understanding the proximate and ultimate causes of masting at broad spatial and temporal scales  
58    will enable us to predict tree reproductive strategies and their response to changing environment.  
59    Here we provide broad spatial (distribution range-wide) and temporal (century) masting data for the  
60    two main masting tree species in Europe, European beech (*Fagus sylvatica* L.) and Norway spruce  
61    (*Picea abies* (L.) H. Karst.). We collected masting data from a total of 359 sources through an  
62    extensive literature review and from unpublished surveys. The dataset has a total of 1747 series and  
63    18348 yearly observations from 28 countries and covering a time span of years 1677-2016 and  
64    1791-2016 for beech and spruce, respectively. For each record, the following information is  
65    available: identification code; species; year of observation; proxy of masting (flower, pollen, fruit,  
66    seed, dendrochronological reconstructions); statistical data type (ordinal, continuous); data value;  
67    unit of measurement (only in case of continuous data); geographical location (country,  
68    Nomenclature of Units for Territorial Statistics NUTS-1 level, municipality, coordinates); first and  
69    last record year and related length; type of data source (field survey, peer reviewed scientific  
70    literature, grey literature, personal observation); source identification code; date when data were  
71    added to the database; comments. To provide a ready-to-use masting index we harmonized ordinal  
72    data into five classes. Furthermore, we computed an additional field where continuous series with  
73    length >4 years were converted into a five classes ordinal index. To our knowledge, this is the  
74    most comprehensive published database on species-specific masting behaviour. It is useful to study  
75    spatial and temporal patterns of masting and its proximate and ultimate causes, to refine studies  
76    based on tree-ring chronologies, to understand dynamics of animal species and pests vectored by

77 these animals affecting human health, and it may serve as calibration-validation data for dynamic  
78 forest models.

79

80 **Key words:** mast seeding; mast fruiting; pollen; fructification; reproduction; synchrony; tree  
81 regeneration

82

### 83 INTRODUCTION

84 Masting, i.e., the synchronous and highly variable production of large crops of flowers, fruit or  
85 seeds by a population of plants, is a widespread reproductive strategy in tree species (Crone and  
86 Rapp 2014, Pearse et al. 2016). It has immediate effects on the regeneration of forest species and  
87 cascading effects on the food web, as it provides large quantities of pollen for insects and seeds for  
88 frugivore animals (Koenig and Knops 2005). For example, mast years have frequently been linked  
89 with animal population dynamics and migrations (Perrins 1965, Boutin et al. 2006). In forestry,  
90 masting in trees is critical for scheduling silvicultural treatments (Ascoli et al. 2015). In tree-ring  
91 studies, masting usually overlaps and affects the climate signals in tree ring chronologies due to  
92 reduced growth in mast years (Mencuccini and Piussi 1995, Koenig and Knops 1998, Drobyshev et  
93 al. 2014, Hacket-Pain et al. 2015). Finally, it has important consequences on human health, because  
94 of pollen allergies and epidemic diseases vectored by frugivorous (Reil et al. 2015, Bogdziewicz  
95 and Szymkowiak 2016).

96 Despite the extensive literature on masting ecology, its evolutionary context, spatial and temporal  
97 variability, and the related proximate drivers are still a matter of debate (e.g., Kelly et al. 2013,  
98 Koenig et al. 2015, Pearse et al. 2014, Pesendorfer et al. 2016). Similarly, the effects of climate  
99 warming on masting remain to be fully tested (Schauber et al. 2002, Monks et al. 2016).  
100 Understanding proximate and ultimate causes (*sensu* Pearse et al. 2016) of masting on a broad  
101 spatial (range-wide) and temporal (century) scale could enable better prediction of these  
102 reproductive events (Koenig and Knops 2005). In the light of climate change, the calibration and

103 validation of vegetation models accounting for masting-climate interactions could improve models  
104 accuracy in predicting species range shifts (Snell et al. 2014) and support the development of  
105 adaptive management strategies.

106 To date, masting data have been largely available at site and regional level to test hypotheses and to  
107 build models, but restrictions occur because of their temporal limitation to only a few decades.

108 Several studies have collected extensive data to study masting behaviour over large geographical  
109 areas for many plant species. Valuable datasets which contributed greatly to improve masting  
110 studies include those published by Herrera et al. (1998), Koenig and Knops (2000), Kelly and Sork  
111 (2002), Schaub et al. (2002) and Kelly et al. (2013). However, these datasets consisted of data  
112 from many diverse species resulting in a reduced number of observations at single species level e.g.,  
113 mean observation number is 179 per species in Koenig and Knops (2000; Table 1). In addition, they  
114 have rarely exceeded a span of few decades (range in Herrera et al. 1998, Table A1: 4-33 years;  
115 range in Kelly and Sork 2002: 6-35 years). In contrast, long-term studies (> century) based on  
116 single species are often not continuous and limited in their geographical extent (e.g., *Fagus*  
117 *sylvatica* L. in Southern Sweden in Drobyshev et al. 2014). These shortcomings have restricted the  
118 possibility of testing hypotheses on masting and modelling at adequate spatial and temporal scale.

119 We collected extensive data on masting of two of the most important masting tree species of the  
120 European continent: European beech (*Fagus sylvatica* L.) and Norway spruce (*Picea abies* (L.) H.  
121 Karst.). The database covers the complete distribution range of European beech and a large  
122 proportion of that of Norway spruce in Europe, extending over a period of two centuries. It provides  
123 information on annual flowering, airborne pollen, fruit and seed production and consists of both  
124 ordinal and continuous data. We included also two mast year series reconstructed using  
125 dendrochronology, and a series of pollen concentration in lake sediments assessed at an annual-  
126 resolution. To provide a ready-to-use masting index we harmonized ordinal data into five classes.  
127 Furthermore, we computed an additional field where continuous series with length > 4 years where

128 converted into a five classes ordinal index. We collected data from published and unpublished  
129 studies. Data sources are fully documented.  
130 Potential uses of this database (here after MASTREE) include testing hypotheses on proximate and  
131 ultimate causes of masting, calibration and validation of tree masting models, assessing the effects  
132 of climate change on tree reproduction investment, and an enhanced understanding of the effects of  
133 masting on tree ring chronologies. Furthermore, MASTREE is a reference masting database that is  
134 not restricted to its initial component species or to geographical region.

135

136 **METADATA**

137 **CLASS I. DATA SET DESCRIPTORS**

138

139 **A. Data set identity:** The tree masting database (the MASTREE database)

140

141 **B. Data set identification code:** MASTREE\_2016.11.csv

142

143 **C. Data set description**

144 **1. Originators:** Davide Ascoli, University of Naples Federico II, via Università 100, 80055 Portici,  
145 Napoli, Italy; Janet Maringer, Institute for Landscape Planning and Ecology, Keplerstr. 11, 70174  
146 Stuttgart, Germany.

147

148

149 **2. Abstract**

150 Tree masting is one of the most intensively studied ecological processes. It affects nutrient fluxes of  
151 trees, regeneration dynamics in forests, animal population densities, and ultimately influences  
152 ecosystem services. Despite a large volume of research focused on masting, its evolutionary  
153 ecology, spatial and temporal variability and environmental drivers are still matter of debate.

154 Understanding the proximate and ultimate causes of masting at broad spatial and temporal scales  
155 will enable us to predict tree reproductive strategies and their response to changing environment.  
156 Here we provide broad spatial (distribution range-wide) and temporal (century) masting data for the  
157 two main masting tree species in Europe, European beech (*Fagus sylvatica* L.) and Norway spruce  
158 (*Picea abies* (L.) H. Karst.). We collected masting data from a total of 359 sources through an  
159 extensive literature review and from unpublished surveys. The dataset has a total of 1747 series and  
160 18348 yearly observations from 28 countries and covering a time span of years 1677-2016 and  
161 1791-2016 for beech and spruce, respectively. For each record, the following information is  
162 available: identification code; species; year of observation; proxy of masting (flower, pollen, fruit,  
163 seed, dendrochronological reconstructions); statistical data type (ordinal, continuous); data value;  
164 unit of measurement (only in case of continuous data); geographical location (country,  
165 Nomenclature of Units for Territorial Statistics NUTS-1 level, municipality, coordinates); first and  
166 last record year and related length; type of data source (field survey, peer reviewed scientific  
167 literature, grey literature, personal observation); source identification code; date when data were  
168 added to the database; comments. To provide a ready-to-use masting index we harmonized ordinal  
169 data into five classes. Furthermore, we computed an additional field where continuous series with  
170 length >4 years were converted into a five classes ordinal index. To our knowledge, this is the  
171 most comprehensive published database on species-specific masting behaviour. It is useful to study  
172 spatial and temporal patterns of masting and its proximate and ultimate causes, to refine studies  
173 based on tree-ring chronologies, to understand dynamics of animal species and pests vectored by  
174 these animals affecting human health, and it may serve as calibration-validation data for dynamic  
175 forest model.

176

177 **D. Key words:** mast seeding; mast fruiting; pollen; fructification; reproduction; synchrony; tree  
178 regeneration

179

180 CLASS II. RESEARCH ORIGIN DESCRIPTORS

181

182 A. Overall project description

183 1. Identity: The tree masting database (the MASTREE database)

184 2. Originators: Davide Ascoli, University of Naples Federico II, via Università 100, 80055 Portici,  
185 Napoli, Italy; Janet Maringer, Institute for Landscape Planning and Ecology, Keplerstr. 11, 70174  
186 Stuttgart, Germany.

187 3. Period of study: 2015–2016

188 4. Objectives:

- 189 1. To improve knowledge of masting patterns at a broad spatial and temporal scale.  
190 2. To enable hypotheses testing related to proximate and ultimate causes of masting.  
191 3. To support improvement of vegetation dynamics models.

192 5. Abstract: same as above.

193 6. Sources of funding: The paper was partly funded by the “Fondo Ricerca Locale 2015-2016” of  
194 the University of Torino and by the Stiftelsen Stina Werners fond (grant SSWF 10-1/29-3 to I.D.).

195

196 B. Specific subproject description

197 1. Site description: Data were obtained for most of the distribution range of both beech and spruce.  
198 The distribution area of beech covered by the database includes the lowland plains in southern  
199 Scandinavia (Denmark, Sweden) and northern Germany, Poland and Ukraine to United Kingdom,  
200 France and Benelux countries; the colline and the submontane elevation zone (600 – 1,100 m a.s.l.)  
201 in Central and Eastern Europe (Austria, Bosnia Herzegovina, Croatia, Czech Republic, Hungary,  
202 Romania, Slovakia, Slovenia, Switzerland); the montane-altimontane elevation zones (1,100 –  
203 1,900 m a.s.l.) of Southern Europe (Italy, Greece, Spain) (Bohn et al. 2003). For spruce, the data  
204 covers the mountainous regions in Central and Eastern Europe (Austria, Bosnia Herzegovina, Czech  
205 Republic, France, Germany, Italy, Romania, Switzerland) and northern Europe (Estonia, Finland,

206 Norway, Poland, United Kingdom, Sweden), and as far as the Russian Federation on the eastern  
207 most sites (Bohn et al. 2003).

208 **2. Experimental or sampling design:** Data were obtained from published sources, unpublished  
209 surveys, and from observations made by the authors. See research methods below.

210 **3. Research methods:** We conducted a systematic review of the published data to reconstruct beech  
211 and spruce masting. Peer-reviewed journals were searched in ISI Web of Knowledge and Google  
212 Scholar. In the case of secondary literature, the original source data was used. Mast data published  
213 in reviews were cross-checked for redundancy and the original data source was used whenever  
214 possible (e.g., Jenny 1987 in Hilton and Packam 2003). We also searched Google Scholar, Google  
215 search engine, OPACplus of the Bavarian Central Library, the global Karlsruhe Virtual Catalog and  
216 the Austrian BFW literature database for non-peer-reviewed articles and unpublished data, which  
217 were for the most part published or collected by foresters (e.g., Burkhardt 1875). Book searches  
218 were also conducted (e.g., Dengler 1944) using Google books. The search terms were beech or  
219 spruce masting in an appropriate selection of European languages: Austria, Germany and German  
220 speaking Switzerland = Samenjahr, Mastjahr, Ernteaussichten, Blühen and Fruktifizieren; Czech  
221 Republic= semenný rok; France, French speaking Switzerland, and Belgium= fainée (specific for  
222 beech); Denmark= oldenår; Hungary= bükkmakk (specific for beech); Italy and Italian speaking  
223 Switzerland = pasciona; Netherland= mastjaar; Poland= urodzaju nasion; Romania= fructificatie  
224 abundenta, an de samamta; Russian= год с обильным плодоношением; Spain= vecería; Sweden=  
225 ollonår. Additionally, we contacted experts from governmental and private forest nurseries,  
226 ministries for the environment, and research institutes. For each data record, the column *SourceType*  
227 reflects the type of source used for data collection (Field survey, Scientific literature, Grey  
228 literature, Personal observation), which can also be seen as an indicator of data accuracy (Class  
229 IV.B.9). Information on the data sources is coded in the column *SourceCode* and the reference (full  
230 reference if published, responsible agency or person if unpublished) is given below (Class IV.B.10).

231 To minimize loss of information from the original source, we have designed the database to include  
232 quantitative data on flower, pollen, fruit, seeds, and tree-ring proxies. Masting proxies such as  
233 animal population dynamics, seedling age, or disease carriers (e.g., *Hantavirus*) were not included.

234

235 **4. Project personnel:**

236 **Principal investigator:** Davide Ascoli

237 **Main associated investigator:** Janet Maringer

238 **Contributors:** Andy Hacket-Pain, Marco Conedera, Igor Drobyshev, Renzo Motta, Mara Cirolli,  
239 Władysław Kantorowicz, Christian Zang, Silvio Schueler, Luc Croisé, Pietro Piussi, Roberta  
240 Berretti, Ciprian Palaghianu, Marjana Westergren, Jonathan G.A. Lageard, Anton Burkhard, Regula  
241 Gehrig Bichsel, Peter A. Thomas, Burkhard Beudert, Rolf Övergaard, Giorgio Vacchiano

242

243 **CLASS III. DATA SET STATUS AND ACCESSIBILITY**

244 **A. Status**

245 **1. Latest Update:** January 2017

246 **2. Latest Archive data:** January 2017

247 **3. Metadata status:** The metadata are complete and up to date as January 2017.

248 **4. Data verification:** The quality of the data has been carefully reviewed by the authors. Data has  
249 undergone substantial checking throughout preliminary statistical analysis (e.g., cross-check for  
250 redundancies, spatial correlation, testing of common proximate masting cues). All records are  
251 associated to a specific source and a related reference.

252

253 **B. Accessibility**

254 **1. Storage location and medium:** Supporting Information associated with this Data Paper  
255 published in *Ecology*. An original data file exists on the server of the University of Naples and  
256 University of Turin, Italy, and University of Stuttgart, Germany.

257   **2. Contact person:** Davide Ascoli, Dipartimento di Agraria, University of Naples Federico II, via  
258   Università 100, 80055 Portici, Napoli, Italy. E-mail: [davide.ascoli@unina.it](mailto:davide.ascoli@unina.it), URL:  
259   <https://www.docenti.unina.it/davide.ascoli>

260   **3. Copyright restrictions:** None

261   **4. Proprietary restrictions:** None

262   **5. Costs:** None

263

264   **CLASS IV. DATA STRUCTURAL DESCRIPTORS**

265

266   **A. Data Set File**

267   **1. Identity:** MASTREE\_2016.11.csv

268   **2. Size:** 19 columns and 18348 records (not including header row)

269   **3. Format and storage mode:** comma-separated values (.csv). No compression scheme was used.

270   **4. Header information:** Headers describe the content of each column and are: ID, Species, Yr,

271   Proxy, VarType, Value, Unit, ORDmast, Country, NUTS1, Location, Coordinates, Start, End,

272   Length, SourceType, SourceCode, AccessionDate, Comments.

273   **5. Alphanumeric attributes:** mixed

274   **6. Special characters/fields:** in the *Location* and *SourceCode* columns we removed the following

275   special characters: å, à, á, ä, â, ã, č, è, é, ě, ì, ī, ñ, ò, ó, ö, ř, š, ù, ü, ý, ž, č, to avoid complications in

276   uploading and using the file.

277   **7. Authentication procedure:**

278   The sum of column ORDmast column is 42622. The number of characters in the whole dataset is

279   1,705,019 (excluding spaces and separations between columns and headers).

280

281   **B. Variable information**

282   1. Variable definition

<b>Variable name</b>	<b>Definition</b>	<b>Data format</b>
ID	Unique identifier (see B.2)	Alphanumeric, 9 characters
Species	Species identifier	Character string, up to 4 characters
Yr	Year of observation	Numeric, integer
Proxy	Proxy used to quantify masting (see B.3)	Character string, up to 15 characters
VarType	Variable type: O – categorical ordinal, C – continuous	Character string, up to 1 characters
Value	A number that gives the level of the masting proxy (see B.4)	Numeric, integer
Unit	The unit of measurement of the masting proxy, only if the variable is continuous, i.e., VarType= C	Character string, up to 63 characters (spaces included)
ORDmast	An ordinal index (1 to 5) of the intensity of masting (see B.5)	Numeric, integer
Country	The country where the observation was recorded	Character string, 9 characters (spaces included)
NUTS1	The Nomenclature of Units for Territorial Statistics (NUTS-1) level where the observation was recorded (see B.6)	Alphanumeric, 5 characters
Location	The municipality or specific site (e.g., Nature Reserve) where the observation was recorded (see B.7)	Character string, up to 29 characters (spaces included)

Coordinates	Geographical coordinates (UTM lat/long in degrees, minutes, seconds) of the stand where data were collected	Numeric, integer
Start	First year of a continuous segment of observations	Numeric, integer
End	Last year of a continuous segment of observations	Numeric, integer
Length	Length in years of a continuous segment of observations (see B.8)	Numeric, integer
SourceType	Field survey, Scientific literature, Compilation, Grey literature, Personal observation (see B.9)	Character string, up to 2 characters
SourceCode	Identification code for the source (published or unpublished references) from which the data have been obtained. See B.10 for the complete list.	Character string, up to 74 characters
AccessionDate	Date when the observation was uploaded in the database	Date in month-year format
Comments	Additional comments in free format	Character string, up to 171 characters

283 Only Unit, ORDmast, Location, Coordinates and Comments include empty cells; the other columns

284 have entries for all rows.

285

286 2. ID: unique identifier code

287 The unique identifier code is composed by 8 or 9 alphanumeric characters.

1 <sup>st</sup> position	Species identifier (FASY = <i>Fagus sylvatica</i> ; PIAB = <i>Picea abies</i> ).
2 <sup>nd</sup> , 3 <sup>rd</sup> positions	Country code (AT=Austria, BE=Belgium, BA=Bosnia Herzegovina, HR=Croatia, CZ=Czech Republic, DK=Denmark, EE=Estonia, FI=Finland, FR=France, DE=Germany, EL=Greece, HU=Hungary, HR=Croatia, IT=Italy, LU=Luxemburg, NL=Netherlands, NO=Norway, PL=Poland, RO=Romania, RU=Russia, RS=Serbia, SK=Slovakia, SI=Slovenia, ES=Spain, SE=Sweden, CH=Switzerland, UA=Ukraine, UK=United Kingdom).
4 <sup>th</sup> to 7 <sup>th</sup> positions	Numeric code that identifies a series collected with the same method at a single location in a given country (unique combination of the 10 <sup>th</sup> , 11 <sup>th</sup> , 12 <sup>th</sup> columns). The code starts from 0001 for each country. In few long surveys (e.g., SourceCode = UK survey..., SourceCode = AFZ...) location and methods were maintained constant at a specific location although the source publishing the survey changed through time. In these few cases we inserted the same numeric code (4 <sup>th</sup> to 7 <sup>th</sup> position of the ID) although there might correspond to more than one source.
8 <sup>th</sup> to 9 <sup>th</sup> positions	Alphabetical code that identifies a specific temporal segment of a series (i.e., A, B, C, ..., Z, AA, AB, AC, ..., AZ). To avoid existing hiatuses in the mast series we divided them into multiple segments, excluding in this way the periods with missing observations. Despite missing years, the structure of the record ID makes it possible to identify discontinuous segments collected using the same method at the same location by exhibiting the same values in the ID positions 4 <sup>th</sup> to 7 <sup>th</sup> but different letters in 8 <sup>th</sup> to 9 <sup>th</sup> positions.

290 The 4<sup>th</sup> columns reports the type of proxy used to quantify beech and spruce masting. The database  
291 is designed to collect as much information as possible; consequently, we included:

- 292 1. Flowering: mass flowering is a common and direct indicator of masting (e.g., Schaub et al.  
293 2002). However, cancelling factors of masting such as late frost may occur during or after  
294 flowering inhibiting the pollination or subsequent fruit and seed development (Kelly and  
295 Sork 2002).
- 296 2. Pollen: a strong positive relationship has been found between beech and spruce airborne  
297 pollen and seed crop (e.g., Pidek et al. 2010, Kasprzyk et al. 2014). Quantity of pollen  
298 directly affects pollination efficiency and thus the percentage of sound seeds (Nilsson and  
299 Wastljung 1987, Norton and Kelly 1988, Koenig et al. 2015).
- 300 3. Fruit/Cone: a strong linear relationship has been found between fruits of beech and spruce  
301 and their respective seeds (e.g., Ascoli et al. 2015).
- 302 4. Seed: the most common indicator to assess masting (Pearse et al. 2016).
- 303 5. Dendro: dendrochronological reconstruction of mast years, based on the split calibration-  
304 verification of the growth depressions in regional master chronologies (Drobyshev et al.  
305 2014).
- 306 6. Pollen\_sediment: Similar as point 2 (airborne pollen), but using pollen influx data from  
307 varved (laminated at annual resolution) lake sediments as masting indicator.

308

309 4. Value: value of the proxy

310 The 6<sup>th</sup> column reports the value of the masting proxy. According to the original source, the value is  
311 expressed as a continuous value (VarType=C) or as an ordinal scale ranging from 1 to 5  
312 (VarType=O).

313 Continuous data accounted for 33% of the observations in the database. If the measure is expressed  
314 as a continuous number, the original annual value as reported by the published or unpublished

315 source is reported (in the case of data published in scientific journals, any available Figure or Table  
316 number from which data were taken is indicated in the column “Comments”).  
317 Ordinal data accounted for the remaining 67% of observations. As with previous attempts at  
318 creating masting databases using ordinal data (Koenig and Knops 2000, Kelly and Sork 2002), we  
319 faced the problem of varying number of categories (range 3-9) when recording masting data from  
320 different sources. Following the approach of Koenig and Knops (2000), we harmonized the number  
321 of classes for all ordinal series, adopting a five class standard, as this had been used by several pre-  
322 existing official surveys (e.g., United Kingdom survey, European Aerobiological Network – EAN,  
323 Italian State Forest Service – CFS), in many scientific papers (e.g., Jenny 1987, Hilton and Packam  
324 2003, Watcher 1964) and in the longest recorded series (i.e., Hase 1964). The five ordinal  
325 categories expressing the annual masting level are as follow: 1 – very poor mast; 2 – poor; 3 –  
326 moderate; 4 – good; and 5 – full mast year.

327 Here after, we provide some examples of how we harmonized ordinal data to this five class system.  
328

329 Series ID: FASYDE0051A; SourceCode: Maurer 1964  
330 Maurer (1964) presented data of seed masting on a three-class scale and expressed the mast events  
331 as poor, half-mast, and full-mast. We converted this three-class scale to the five-class one assigning  
332 poor to class 1; half-mast to class 3; full-mast to class 5, as suggested by Koenig and Knops (2000).

333  
334 Series ID: from FASYDE0098A to FASYDE0182A and from PIABDE0090A to PIABDE0136A;  
335 SourceCode AFZ Year(issue n.)  
336 The German survey published by the Allgemeine Forst Zeitschrift für Wald und Forstwirtschaft  
337 (AFZ) uses a four-class scale. The annual intensity of flowering and the yield of seeds of the  
338 previous year are published in the ‘Allgemeine Forstzeitschrift’ (later ‘AFZ der Wald’). Since 1991,  
339 the intensity of flowering is systematically categorized in: class 1: no mast year (0-10% blossoms);  
340 class 2: local mast (11-30% blossoms); class 3: half mast (31-60% blossoms); class 4: full mast

341 (>60% blossoms). In some cases, flowering is reported as “half to full mast” (i.e., 3<sup>rd</sup> to 4<sup>th</sup> AFZ  
342 class). The data were transformed to the five-class system assigning “no mast year” to class 1,  
343 “local mast year” to class 2, “half mast year” to class 3, “half to full mast year” to class 4 and a “full  
344 mast year” to class 5.

345

346 Series ID: from FASYAT0051A to FASYAT0278A; SourceCode: BFW archive

347 The Austrian survey is based on a 4 class system similar to the German one. However, the  
348 categories are given in ascending order from 1: full mast to 4: no mast (Nather 1962). For the  
349 database, the Austrian classification (AS) system was transferred into the common five classes  
350 where 4-AS is 1 (very poor mast), 3-AS and 2-AS are 3 (moderate mast), and 1-AS is 5 (full mast).  
351 Based on practical difficulties of differentiating between categories 2-AS and 3-AS, these were  
352 combined in one class. If masting was reported as 1-2-AS these observations were assigned to class  
353 4.

354

355 Series ID: FASYUK0022A; SourceCode: Perrins 1965

356 Perrins (1965) presented data of seed masting on a nine-class scale and expressed the mast events as  
357 nil, nil-poor, poor, poor-moderate, moderate, moderate-good, good, good-abundant, abundant. We  
358 converted the nine-class scale to the five-class one assigning “nil” to class 1; “nil-poor, poor” to  
359 class 2; “poor-moderate, moderate” to class 3; “moderate-good, good” to class 4; “good-abundant,  
360 abundant” to class 5.

361

362 Series ID: FASYBE0008A and FASYSE0005A; SourceCode: Latte et al. 2016 and Drobyshev et  
363 al. 2014

364 In case of sources reporting single observations of full-mast years (e.g., Latte et al. 2016, page 199:  
365 *“However, for three mature beech stands located in the same locality, 1995, 2000, 2002, 2004 and*  
366 *2011 were qualified as heavy mast years”*), or dendrochronological reconstruction based on annual

367 tree ring growth depressions in regional master chronologies (Drobyshev et al. 2014), we assigned  
368 an ordinal value equal to class 5.

369

370 5. Ordinal masting index

371 The 8<sup>th</sup> column is a ready-to-use ordinal index of masting in 5 classes (class 1=very poor; class 5:  
372 very abundant) which includes all ordinal series (note: for ordinal series, ORDmast reports the same  
373 value displayed in the column Value) and all continuous series longer than four years, after being  
374 converted into the 1 to 5 ordinal scale. The procedure for data conversion from continuous to  
375 ordinal is described below.

376 i) For each ordinal series (VarType=O) with length > 4 years, we calculated the relative frequencies  
377 of the five ordinal masting classes; ii) for each species separately, we computed the mean relative  
378 frequency of each class across all series; iii) we re-classified each continuous series (VarType=C)  
379 with length > 4 years into 5 classes, using as percentile cut-offs the mean relative frequencies of the  
380 respective species.

381 Mean relative frequencies used for the conversion were:

382 Beech = class 1: 0.352; class 2: 0.279; class 3: 0.189; class 4: 0.082; class 5: 0.098

383 Spruce = class 1: 0.425; class 2: 0.237; class 3: 0.161; class 4: 0.080; class 5: 0.097

384

385 6. NUTS1: Nomenclature of Units for Territorial Statistics

386 The 10<sup>th</sup> column reports the code of the Nomenclature of Units for Territorial Statistics (NUTS-1)  
387 administrative level where data were collected. Non-EU countries where beech masting data were  
388 recorded (i.e., Russia, Ukraine, Serbia, Croatia, Bosnia and Herzegovina) were also included in the  
389 database with dummy NUTS-1 codes. When the source did not provide sufficient information to  
390 assign the observation to a specific NUTS-1, we give the country code followed by “#”, e.g., AT#,  
391 DE#, UK#. In the case of the German masting survey, we assigned each region of the survey to the  
392 most overlapping NUTS-1 level.

393

394 7. Location

395 The 11<sup>th</sup> column reports more detailed geographical information than the NUTS-1 level (e.g.,  
396 region, municipality, Nature Reserve name). In some cases, observations were made from different  
397 stands at the same general location, without further specific locational information (i.e.,  
398 coordinates). In these cases we report the name of the location followed by the stand number (e.g.,  
399 Asiago\_stand1, Asiago\_stand2, ..., Asiago\_stand5). If there was no geographic information apart  
400 from the NUTS-1 level, the location cell was left empty, i.e., the NUTS-1 level represents the only  
401 geo-referencing for the observation.

402

403 8. Length

404 The 15<sup>th</sup> column results from the difference between the last and first year of a continuous segment  
405 in a given series plus one (i.e., End – Start + 1). It refers to the length of any single segment of a  
406 series (see the ID description Class IV.B.2). It is equal to 1 in case of one or more discontinuous  
407 single year observations at a specific location, or when continuous series present missing data,  
408 resulting a segment of length 1 year (e.g., as recorded in the Hase 1964 series for the Schleswig-  
409 Holstain location in the years 1685, 1687, 1712, 1714, 1720, 1730, 1734, 1742, 1744, 1746, 1838  
410 and 1843).

411

412 9. SourceType

413 The 16<sup>th</sup> column describes the general methods of gathering the information and the related  
414 accuracy. Four possible cases are considered:  
415 FS = Field survey. Published or unpublished data obtained from an official survey. The data  
416 collection followed the same method for several years on permanent sites.  
417 SL = Scientific literature. Published data obtained from a scientific, peer-reviewed journal.

418 GL = Grey literature. Published data obtained from a research produced outside of the academic  
419 publishing (e.g., administrative reports, Masters thesis).

420 PO = Personal observation. Data from visual estimation or personal experience.

421 The first and second categories are considered the most accurate information, while the others are  
422 viewed as less accurate.

423

424 10. SourceCode

425 The 17<sup>th</sup> column provides a code that refers to the data source (SourceCode). Complete references  
426 are listed below. Note that the references include field surveys, published articles, grey literature  
427 and personal observations.

428

<b>Source Code</b>	<b>Full reference</b>
Abt. Waldbau Ib	Abteilung Waldbau I b der Forstlichen Bundesversuchsanstalt Mariabrunn in Schönbrunn. 1960. Waldsamen-Ernteaussichten für 1960/61. Fachzeitschrift für das gesamte Forstwesen; Mitteilungsbl. D. forstl. Forstvereine u. Standesorganisation Österreichs.- Wien: Österr. Agrarverlag. Band 71 (19-20): 225-226.
AFZ 1954(31/32)	Siegl, H. 1954. Prognosen der Waldsamenernte 1954. Allgemeine Forstzeitschrift. 31/32: 333.
AFZ 1955(33/34)	von Schönborn, A. 1955. Prognosen der Waldsamenernte 1955. Allgemeine Forstzeitschrift. 33/34: 381.
AFZ 1956(35/36)	von Schönborn, A. 1956. Prognosen der Waldsamenernte 1956. Allgemeine Forstzeitschrift. 35/36: 453.
AFZ 1957(39)	von Schönborn, A. 1957. Prognosen der Waldsamenernte 1957. Allgemeine Forstzeitschrift. 39: 460-462.

AFZ 1958(33)	von Schönborn, A. 1958. Prognosen der Waldsamenernte 1958. Allgemeine Forstzeitschrift. 33: 472/473.
AFZ 1959(40)	von Schönborn, A. 1959. Prognosen der Waldsamenernte 1959. Allgemeine Forstzeitschrift. 40: 703-705.
AFZ 1960(40)	von Schönborn, A. 1960. Prognosen der Waldsamenernte 1960. Allgemeine Forstzeitschrift. 40: 584-586.
AFZ 1961(35)	von Schönborn, A. 1961. Prognosen der Waldsamenernte 1961. Allgemeine Forstzeitschrift. 35: 518-520.
AFZ 1962(38)	von Schönborn, A. 1962. Prognosen der Waldsamenernte 1962. Allgemeine Forstzeitschrift. 38: 597-599.
AFZ 1963(38)	von Schönborn, A. 1963. Prognosen der Waldsamenernte 1963. Allgemeine Forstzeitschrift. 38: 586-588.
AFZ 1964(36)	von Schönborn, A. 1964. Prognosen der Waldsamenernte 1964. Allgemeine Forstzeitschrift. 36: 539-542.
AFZ 1965(45)	von Schönborn, A. 1965. Prognosen der Waldsamenernte 1965. Allgemeine Forstzeitschrift. 36: 539-542.
AFZ 1967(41)	von Schönborn, A. 1967. Prognosen der Waldsamenernte 1967. Allgemeine Forstzeitschrift. 41: 695-698.
AFZ 1968(41)	von Schönborn, A. 1968. Prognosen der Waldsamenernte 1968. Allgemeine Forstzeitschrift. 41: 719-722.
AFZ 1969(44)	von Schönborn, A. 1969. Prognosen der Waldsamenernte 1969. Allgemeine Forstzeitschrift. 44: 862-865.
AFZ 1970(39)	von Schönborn, A. 1970. Prognosen der Waldsamenernte 1970. Allgemeine Forstzeitschrift. 39: 814-818.
AFZ 1971(42)	v. Schönborn, A. 1971. Prognosen der Waldsamenernte 1971. Allgemeine

	Forstzeitschrift. 42: 877-879.
AFZ 1972(36)	Eicke, G. 1972. Prognosen der Waldsamenernte 1972. Allgemeine Forstzeitschrift. 36: 717-718.
AFZ 1973(43)	Eicke, G. 1973. Prognosen der Waldsamenernte 1973. Allgemeine Forstzeitschrift. 43: 969-972.
AFZ 1974(36)	Eicke, G. 1974. Prognosen der Waldsamenernte 1974. Allgemeine Forstzeitschrift. 36: 784-785.
AFZ 1975	Eicke, G. 1975. Prognosen der Waldsamenernte 1975. Allgemeine Forstzeitschrift. 907-908.
AFZ 1976	Eicke, G. 1976. Prognosen der Waldsamenernte 1976. Allgemeine Forstzeitung. 926-928.
AFZ 1978	Eicke, G. 1978. Prognosen der Waldsamenernte 1978. Allgemeine Forstzeitschrift. 998-999.
AFZ 1979	Eicke, G. 1979. Prognosen der Waldsamenernte 1979. Allgemeine Forstzeitschrift. 1005-1006.
AFZ 1981(37)	Eicke, G. 1981. Prognosen der Waldsamenernte 1981. Allgemeine Forstzeitschrift 37, pp. 948
AFZ 1982(37)	Eicke, G. 1982. Prognosen der Waldsamenernte 1982. Allgemeine Forstzeitschrift. 37: 1118.
AFZ 1983(37)	Eicke, G. 1983. Prognosen der Waldsamenernte 1983. 37: 950-951.
AFZ 1984(36)	Eicke, G. 1984. Das Blühen der Waldbaumarten 1984. Allgemeine Forstzeitschrift. 36: 888.
AFZ 1985(33)	Eicke, G. 1985. Das Blühen der Waldbaumarten 1985. Allgemeine Forst Zeitung. 33: 855-856.
AFZ 1986(33)	Eicke, G. 1986. Das Blühen der Waldbaumarten 1986. Allgemeine

	Forstzeitschrift 33: 812-813.
AFZ 1987(39)	Eicke, G. 1987. Das Blühen der Waldbäume 1987. Allgemeine Forstzeitschrift. 39: 1005-1007.
AFZ 1988(33)	Eicke, G. 1988. Das Blühen der Waldbäume 1988. Allgemeine Forstzeitschrift. 33: 901.
AFZ 1989(32)	Eicke, G. 1989. Das Blühen der Waldbäume 1989. Allgemeine Forstzeitschrift. 32: 833-835.
AFZ 1990(32)	Eicke, G. 1990. Das Blühen der Waldbäume 1990. Allgemeine Forst Zeitung. 32: 811-814.
AFZ 1991(17)	Eicke, G. 1991. Das Blühen der Waldbäume 1991. Allgemeine Forstzeitschrift. 17: 858-860.
AFZ 1992(17)	Eicke, G. 1992. Das Blühen der Waldbäume 1992. Allgemeine Forstzeitschrift. 17: 886-887.
AFZ 1993(18)	Eicke, G. 1993. Das Blühen der Waldbäume 1993. Allgemeine Forstzeitschrift. 18: 916-917
AFZ 1994(18)	Eicke, G. 1994. Das Blühen der Waldbäume 1994. Allgemeine Forst Zeitung. 18: 978-979.
AFZ 1995(18)	Eicke, G. 1995. Das Blühen der Waldbäume 1996. AFZ der Wald. 18: 958-959.
AFZ 1996(18)	Eicke, G. 1996. Das Blühen der Waldbäume 1996. AFZ der Wald 18, 982-983
AFZ 1997(18)	Eicke, G. 1997. Das Blühen der Waldbäume 1997. AFZ der Wald. 18: 958-956
AFZ 1998(18)	Eicke, G. 1998. das Blühen der Waldbäume 1998. AFZ der Wald. 18: 926-927.

AFZ 1999(16)	Schneck, D. 1999. Das Blühen der Waldbäume 1999. AFZ der Wald. 16: 828-829.
AFZ 2000(16)	Schneck, D. 2000. Das Blühen der Waldbäume 2000. AFZ der Wald. 16: 844-845.
AFZ 2001(16)	Schneck, D. 2001. Das Blühen der Waldbäume 2001. AFZ der Wald. 16: 812-813.
AFZ 2002(16)	Schneck, D. 2002. Das Blühen der Waldbäume 2002. AFZ der Wald. 16: 820-821.
AFZ 2003(16)	Schneck, D. 2003. Das Blühen der Waldbäume 2003. AFZ der Wald. 16: 816-817.
AFZ 2004(16)	Schneck, D. 2004. Das Blühen der Waldbäume. AFZ der Wald. 16: 848-849.
AFZ 2005(16)	Schneck, D. 2005. Das Blühen der Waldbäume. AFZ der Wald. 16: 836-837.
AFZ 2006(16)	Schneck, D. 2006. Das Blühen der Waldbäume. AFZ der Wald. 16: 852-853.
AFZ 2007(16)	Schneck, D. 2007. Das Blühen der Waldbäume. AFZ der Wald. 16: 844-845.
AFZ 2008(16)	Schneck, D. 2008. Das Blühen der Waldbäume. AFZ der Wald. 16: 844-845.
AFZ 2009(16)	Schneck, D. 2009. Das Blühen der Waldbäume 2009. AFZ der Wald. 16: 844-845.
AFZ 2010(16)	Schneck, D. 2010. Das Blühen der Waldbäume 2010. AFZ der Wald. 16: 4-5.
AFZ 2011(16)	Schneck, D. 2011. Das Blühen der Waldbäume 2011. AFZ der Wald. 16: 4-5.
AFZ 2012(16)	Schneck, D. 2012. Das Blühen der Waldbäume 2012. AFZ der Wald. 16: 4-5.
AFZ 2013(16)	Schneck, D. 2013. Das Blühen der Waldbäume 2013. AFZ der Wald. 16: 18-19.

AFZ 2014(16)	Schneck, D. 2014. Das Blühen der Waldbäume 2014. AFZ der Wald. 16: 27-28.
AFZ 2015(16)	Schneck, D. 2015. Das Blühen der Waldbäume 2015. AFZ der Wald. 16: 9-10.
AFZ 2016(16)	Schneck, D. 2016. Das Blühen der Waldbäume 2016. AFZ der Wald. 16: 16-18.
Albrecht 1977	Albrecht J. 1977. Die Keimung der Bucheckern aus der Mast 1976 im südlichen Niedersachsen. Unveröff. Diplomarbeit, Institut für Waldbau, Universität Göttingen.
Ambroz et al. 2015	Ambrož, R., Vacek, S., Vacek, Z., Král, J. and I. Štefančík. 2015. Current and simulated structure, growth parameters and regeneration of beech forests with different game management in the Lány Game Enclosure/Struktura, růstové parametry, obnova a modelový vývoj bukových porostů s odlišným způsobem mysliveckého hospodaření v podmírkách Lánské obory. Forestry Journal. 61(2): 78-88.
Andersen 1980	Andersen, S. T. 1980. Influence of climatic variation on pollen season severity in wind-pollinated trees and herbs. Grana. 19(1): 47-52.
Anderson 1949	Anderson, M. L. 1949. Some observations on Belgian forestry. Empire Forestry Review. 28(2): 117-130.
Andersson 1965	Andersson, E. 1965. Cone and seed studies in Norway spruce ( <i>Picea abies</i> (L.) Karst.). Studia Forestalia Suecica. 23: 1-278.
Angst 2004	Angst, C. 2004. Vielfältige Waldentwicklung auf Lothar-Versuchsflächen. Informationsblatt Forschungsbereich Wald. 17: 1-4.
Ardo & Lindquist 1947	Ardö, P. and B. Lindquist .1947. Om Laspeyresia grossana Haw. som skadedjur i de nordväst-europeiska bokskogarna, 36. Swedish Institute for

	Forest Research, Stockholm. Online:  <a href="http://pub.epsilon.slu.se/view/series/Meddelanden_fr=E5n_Statens_skogsfor-skningsinstitut.creators_name.html">http://pub.epsilon.slu.se/view/series/Meddelanden_fr=E5n_Statens_skogsfor-skningsinstitut.creators_name.html</a> (Access 2016-06-30)
Arpa Toscana	Agenzia regionale per la protezione ambientale della Toscana (ARPAT).  2015. Dati concentrazioni pollini e spore fungine in Toscana - anni 1996-2015. ( <a href="http://www.arpat.toscana.it/datiemappe/dati/dati-concentrazioni-pollini-e-spore-fungine-in-toscana">http://www.arpat.toscana.it/datiemappe/dati/dati-concentrazioni-pollini-e-spore-fungine-in-toscana</a> )
Ascoli et al. 2013	Ascoli, D., Castagneri, D., Valsecchi, C., Conedera, M. and G. Bovio. 2013. Post-fire restoration of beech stands in the Southern Alps by natural regeneration. Ecological Engineering. 54: 210-217.
Ascoli et al. 2015	Ascoli, D., Vacchiano, G., Maringer, J., Bovio, G. and M. Conedera. 2015. The synchronicity of masting and intermediate severity fire effects favors beech recruitment. Forest Ecology and Management. 353: 126-135.
Ascoli; Personal observation	Ascoli, D.; Personal observation
ASP Bayern	Bayerisches Amt für forstliche Saat- und Pflanzenzucht Kontinuität und Wandel (Bavarian Office for Forest Seeding and Planting Continuity and Change)
Baldwin 1942	Baldwin, H. I. 1942. Forest Tree Seed of The North Temperate Regions. Chronica Botanica Company, USA. p 1-240.
Barbeta et al. 2011	Barbeta, A., Peñuelas, J., Ogaya, R. and A. Jump. 2011. Reduced tree health and seedling production in fragmented <i>Fagus sylvatica</i> forest patches in the Montseny Mountains (NE Spain). Forest Ecology and Management. 261(11): 2029-2037.
Barnekow et al.	Barnekow, L., Loader, N. J., Hicks, S., Froyd, C. A., and T. Goslar. 2007.

2007	Strong correlation between summer temperature and pollen accumulation rates for <i>Pinus sylvestris</i> , <i>Picea abies</i> and <i>Betula</i> spp. in a high-resolution record from northern Sweden. Journal of Quaternary Science. 22(7): 653-658.
Barrios et al. 2010	Barrios, J. M., Verstraeten, W. W., Maes, P., Clement, J., Aerts, J. M., Haredasht, S. A., Wambacq, J., Lagrou, K., Ducoffre, G., Van Ranst, M., Bercjmans, D. and P. Coppin. 2010. Satellite derived forest phenology and its relation with nephropathia epidemica in Belgium. International journal of environmental research and public health. 7(6): 2486-2500.
Bastide & Vredenburch 1970	Bastide la, J. G. A. and C. L. H. van Vredenburch. 1970. The influence of weather conditions on the seed production of some forest trees in the Netherlands. Mededeling, Stichting Bosbouwproefstation 'De Dorschkamp', Wageningen.
Baumler 1981	Bäumler, W. 1981. Distribution, nutrition and population dynamics of the bank vole ( <i>Clethrionomys glareolus</i> ) and the yellow-necked field mouse ( <i>Apodemus flavicollis</i> ) in a mountain forest of the Bavarian Alps. Anzeiger für Schädlingskunde Pflanzenschutz Umweltschutz. 54: 49-53.
Beling 1877	Beling. 1877. Über die Samenjahre der Eiche, Buche und Fichte. In: Monatsschrift für das Forst- und Jagdwesen.- Stuttgart: Schweizerbart. Heft 21: 49-81.
Belmonte et al. 2008	Belmonte, J., Alarcón, M., Avila, A., Scialabba, E. and D. Pino. 2008. Long-range transport of beech ( <i>Fagus sylvatica</i> L.) pollen to Catalonia (north-eastern Spain). International journal of biometeorology. 52(7): 675-687.
Bergstedt 1965	Bergstedt, B. O. 1965. Distribution, reproduction, growth and dynamics of the rodent species <i>Clethrionomys glareolus</i> (Schreber), <i>Apodemus flavicollis</i>

	(Melchior) and <i>Apodemus sylvaticus</i> (Linne) in southern Sweden. <i>Oikos</i> . 16:132-160.
Beudert & Dieffenbach-Fries 2016	Beudert, B. and H. Dieffenbach-Fries. 2016. UNECE Integrated Monitoring programme in the Forellenbach area of the Bavarian Forest National park - litterfall data from ongoing monitoring activities. On behalf of the German Environment Agency (UBA), Grafenau (Germany).
Bezdeckova & Matejka 2015	Bezděcková, L. and K. Matějka. 2015. The quality and the depth of dormancy of beechnuts in individual stand groups with varying climatic conditions within a single unit of approval. <i>Journal of Forest Science</i> . 61(9): 382-392.
BFW archive	Department of Forest Genetics, Federal Research and Training Centre for Forests, Natural Hazards and Landscape (BFW), Vienna, Austria
BFW online	Department of Forest Genetics, Federal Research and Training Centre for Forests, Natural Hazards and Landscape (BFW). 2015. Pollen-Samenproduktion österreichischer Waldbäume. <a href="http://bfw.ac.at/rz/pollen.main?bart_in=01.0&amp;jahr_in=2009">(<a href="http://bfw.ac.at/rz/pollen.main?bart_in=01.0&amp;jahr_in=2009">http://bfw.ac.at/rz/pollen.main?bart_in=01.0&amp;jahr_in=2009</a>)</a>
Bieber & Ruf 2004	Bieber, C. and T. Ruf. 2005. Population dynamics in wild boar <i>Sus scrofa</i> : ecology, elasticity of growth rate and implications for the management of pulsed resource consumers. <i>Journal of Applied Ecology</i> . 42(6): 1203-1213.
Bieber 1998	Bieber, C. 1998. Population dynamics, sexual activity, and reproduction failure in the fat dormouse ( <i>Myoxus glis</i> ). <i>Journal of Zoology</i> . 244(02): 223-229.
Bilek et al. 2009	Bílek, L., Remeš, J. and D. Zahradník. 2009. Natural regeneration of senescent even-aged beech ( <i>Fagus sylvatica</i> L.) stands under the conditions of Central Bohemia. <i>Journal of Forest Science</i> . 55: 145-155.

Bisi et al. 2016	Bisi, F., von Hardenberg, J., Bertolino, S., Wauters, L. A., Imperio, S., Preatoni, D. G., Provenzale A., Mazzamuto M. V., and A. Martinoli. 2016. Current and future conifer seed production in the Alps: testing weather factors as cues behind masting. European Journal of Forest Research. 135(4): 743-754.
Bjedov et al. 2016	Bjedov, L., Svoboda, P., Tadin, A., Habuš, J., Štritof, Z., Labaš, N., Vučelja, M., Marcotić, A., Turk, N., and J. Margaletić. 2016. Utjecaj uroda sjemena obične bukve ( <i>Fagus sylvatica</i> l.) na populacije sitnih glodavaca i pojavnosti hantavirusa u šumama nacionalnog parka “Plitvička Jezera” i parka prirode “Medvednica”. Šumarski list. 140(9-10): 455-463.
BMEL	Bundesanstalt für Landwirtschaft und Ernährung. Referat 324- Wald und Holz, Waldklimafonds. Bonn (Germany). Archive data.
Bodyl & Sulkowska 2007	Bodyl, M. and M. Sulkowska. 2007. Ocena zroznicowania nasion buka zwyczajnego ( <i>Fagus sylvatica</i> L.) w Polsce w latach 1992-2004. Sylwan. 151(09): 12-21.
Borchers 1958	Borchers, K. 1958. Auswirkungen rezenter Klimaschwankungen auf die Häufigkeit von Buchen-Samenjahren in Niedersachsen. Forst-und Holzwirtschaft. 13: 330.
Bourne 1942	Bourne, R. 1942. A note on beech regeneration in southern England. Quarterly Journal of Forestry. 34: 42-49.
Bourne 1945	Bourne, R. .1945. Neglect of natural regeneration. Forestry. 19: 33-40.
Braun; Personal observation	Braun, H. U.; Personal observation
Brodovych; Personal	Brodovych, R. I.; Personal observation

observation	
Broome et al. 2007	Broome, A., Hendry, S. and A. Peace. 2007. Annual and spatial variation in coning shown by the Forest Condition Monitoring programme data for Norway spruce, Sitka spruce and Scots pine in Britain. <i>Forestry</i> . 80(19): 17-28.
Buhler 1918	Bühler, A. 1918. Der Waldbau nach wissenschaftlicher Forschung und praktischer Erfahrung: ein Hand-und Lehrbuch. Ulmer, Stuttgart.
Burckhardt 1875	Burckhardt, H. 1875. Das Mastjahr 1875. Von den Niedersächsischen Landesforsten: Schriftreihe "Aus dem Walde". 255-264.
Burkart; Personal observation	Burkart, A.; Personal observation
Burri et al. 2016	Burri, A., Burkart, A., Moritzi, M., Moser, B., Wasem, U. and T. Wohlgemuth. 2016. Samenproduktion bei Waldbäumen: eine neue Webseite. <i>Zürcher Wald</i> . 1: 23-27.
Burschell 1966	Burschel, P. 1966. Untersuchungen in Buchenmastjahren. <i>Forstwissenschaftliches Centralblatt</i> 85(7): 204-219.
Busgen et al. 1929	Busgen, M., Munch, E., and T. Thomson. 1929. The structure and life of forest trees. Chapman & Hall.
Cansiglio; Regione Veneto	Survey of the "Foresta Demaniale Regionale del Cansiglio", Regione Veneto
Cejkova & Kolar 2009	Čejková, A., and T. Kolář. 2009. Extreme radial growth reaction of Norway spruce along an altitudinal gradient in the Šumava Mountains. <i>Geochronometria</i> . 33(1): 41-47.
Cermak & Jezek 2005	Čermák, P. and J. Ježek. 2005. Effect of tree seed crop on small mammal populations and communities in oak and beech forests in the Drahany

	Upland (Czech Republic). Journal of Forest Science. 51(1): 6-14.
CFS_UTB_Peri	Corpo Forestale dello Stato, Ministero delle Politiche Agricole, Alimentari e Forestali, Ufficio territoriale per la Biodiversità di Verona - CNBF di Peri, Italiano
CFS_UTB_Pieve Santo Stefano	Corpo Forestale dello Stato, Ministero delle Politiche Agricole, Alimentari e Forestali, Ufficio territoriale per la Biodiversità di Pieve Santo Stefano, Italiano
Chalupka & Giertych 1973	Chalupka W. and Giertych M. 1973. Seed years in <i>Picea abies</i> (L.) Karst. Arbor. Kornickie. 18: 183-186.
Chamberlain et al. 2007	Chamberlain, D. E., Gosler, A. G. and D. E. Glue. 2007. Effects of the winter beechmast crop on bird occurrence in British gardens: Capsule Woodland birds were significantly less likely to occur in gardens in years of high beechmast crop. Bird Study. 54(1): 120-126.
Clark et al. 1923	Clark, J. E. and I. D. Margary. 1923. Report on the phenological observations in the British isles, from December 1921 to November 1922. Quarterly Journal of the Royal Meteorological Society. 49(208): 239-271.
Conrad 2005	Conrad, B. 2005. Regenerationsdynamik buchendominiert Laubwälder auf Kalkstandorten. Inaugural-Dissertation zur Erlangung der Doktorwürde der Fakultät für Forst- und Umweltwissenschaften der Albert-Ludwigs-Universität Freiburg im Breisgau.
Cornwallis & Townsend 1968	Cornwallis, R. K. and A. D. Townsend. 1968. Waxwings in Britain and Europe during 1965/66. British Birds. 61: 97-118.
Croatian forests Ltd. Zagreb	Croatian forests, Ltd., Zagreb
Cutini et al. 2009	Cutini, A., Chianucci, F., and Giannini T. 2009. Effetti del trattamento

	selvicolturale su caratteristiche della copertura, produzione di lettiera e di seme in cedui di faggio in conversione. Annals of Silvicultural Research. (36): 109-124.
Cutini et al. 2013	Cutini, A., Chianucci, F., Chirichella, R., Donaggio, E., Mattioli, L. and M. Apollonio. 2013. Mast seeding in deciduous forests of the northern Apennines (Italy) and its influence on wild boar population dynamics. Annals of Forest Science. 70(5): 493-502.
Dallimore 1935	Dallimore, W. 1935. Notes on the dry seasons of 1933 and 1934. Quarterly Journal of Forestry. Ixix: 115—17.
de Wavrin et al. 1991	de Wavrin, H., Walravens, M. and D. Rabosee. 1991. Nidifications exceptionnelles du Hibou moyen-duc ( <i>Asio otus</i> ) et du Faucon Crêcerelle ( <i>Falco tinnunculus</i> ) en 1991 en forêt de Soignes (Brabant). Aves. 28(4): 169-188.
Dengler 1944	Dengler, A. 1944. Waldbau auf ökologischer Grundlage, ed. Springer-Verlag, Berlin and Heidelberg.
Di Pierro et al. 2010	Di Pierro, E., Ghisla, A., Wauters, L. A., Molinari, A., Martinoli, A., Gurnell, J. and G. Tosi. 2011. The effects of seed availability on habitat use by a specialist seed predator. European Journal of Wildlife Research. 57(3): 585-595.
Dingemanse et al. 2004	Dingemanse, N. J., Both, C., Drent, P. J. and J.M. Tinbergen. 2004. Fitness consequences of avian personalities in a fluctuating environment. Proceedings of the Royal Society of London, Series B: Biological Sciences. 271(1541): 847-852.
Dirnberger 1959	Dirnberger, H. 1959. Waldsamenernteaussicht 1959/60. Holz-Kurier. 14(40): 246-249.

Dittmar & Elling 2006	Dittmar, C. and W. Elling. 2006. Dendroökologische Untersuchungen von Buchenbeständen in der Programm-Region des INTERREG III A-Projektes. Mitteilungen aus der Forschungsanstalt für Waldökologie und Forstwirtschaft Rheinland-Pfalz. 59: 31-40.
Dobrovolny & Tesař 2010	Dobrovolný, L. and V. Tesař. 2010. Growth and characteristics of old beech ( <i>Fagus sylvatica</i> L.) trees individually dispersed in spruce monocultures. Journal of Forest Science. 56(9): 406-416.
Dobrowolska 2015	Dobrowolska, D. 2015. Vitality of European beech ( <i>Fagus sylvatica</i> L.) at the limit of its natural range in Poland. Polish Journal of Ecology. 63: 260-272.
Drobyshev et al. 2010	Drobyshev, I., Övergaard, R., Saygin, I., Niklasson, M., Hickler, T., Karlsson, M. and M. T. Sykes. 2010. Masting behaviour and dendrochronology of European beech ( <i>Fagus sylvatica</i> L.) in southern Sweden. Forest Ecology and Management. 259(11): 2160-2171.
Drobyshev et al. 2014	Drobyshev, I., Niklasson, M., Mazerolle, M. J., and Y. Bergeron. 2014. Reconstruction of a 253-year long mast record of European beech reveals its association with large scale temperature variability and no long-term trend in mast frequencies. Agricultural and Forest Meteorology. 192: 9-17.
Agricoltore	La Domenica dell'Agricoltore VII(24), 1932.
EAN	EAN - European Aerobiological Network - Austria
EGI	EGI - Edward Grey Institute of Field Ornithology, University of Oxford
Eichhorn et al. 2008	Eichhorn, J., Dammann, I., Schönfelder, E., Albrecht, M., Beck, W. and U. Paar. 2008. Untersuchungen zur Trockenheitstoleranz der Buche am Beispiel des witterungsextremen Jahres 2003. Beiträge aus der BW-FVA. Band 3.
Emborg 1998	Emborg, J. 1998. Understorey light conditions and regeneration with respect

	to the structural dynamics of a near-natural temperate deciduous forest in Denmark. <i>Forest Ecology and Management</i> . 106(2): 83-95.
Erbrecht 1960	Erbrecht, J. 1960. Buchennaturverjüngung aus dem Mastjahr 1958. <i>Allgemeine Forstzeitung</i> . 15: 289-291.
Eriksson 1984	Eriksson, M. 1984. Winter breeding in three rodent species, the bank vole <i>Clethrionomys glareolus</i> , the yellow-necked mouse <i>Apodemus flavicollis</i> and the wood mouse <i>Apodemus sylvaticus</i> in southern Sweden. <i>Ecography</i> . 7(4): 428-429.
Feichtner 1998	Feichtner, B. 1998. Ursachen der Streckenschwankungen beim Schwarzwild im Saarland. <i>Zeitschrift für Jagdwissenschaft</i> . 44(3): 140-150.
Fietz 2010	Fietz, J. 2010. Reproduktionsdynamik des Siebenschläfers ( <i>Glis glis</i> ) als Zeigerfunktion für den Status von Mischwäldern. Institut für Experimentelle Ökologie. Universität Ulm.
Flade & Schwarz 2004	Flade, M. and J. Schwarz. 2004. Ergebnisse des DDA-Monitoringprogramms, Teil II: Bestandsentwicklung von Waldvögeln in Deutschland. <i>Vogelwelt</i> . 125:177-213.
Forester's record 1935	Forester's Record. 1935. A Forester's Record. <i>Quarterly Journal of Forestry</i> . xxix.2.3:107-204.
Forester's record 1936	Forester's Record. 1936. A Forester's Record. <i>Quarterly Journal of Forestry</i> . xxx.3: 228-234.
Forester's record 1937	Forester's Record. 1937. 'A Forester's Record. <i>Quarterly Journal of Forestry</i> . xxxi.3: 189-195.
Forester's record 1938	Forester's Record. 1938. A Forester's Record. <i>Quarterly Journal of Forestry</i> . xxxii.3: 194-200.
Forester's record	Forester's Record. 1939. A Forester's Record. <i>Quarterly Journal of Forestry</i> .

1939	xxxiii.3: 187-195.
Forester's record 1940	Forester's Record. 1940. A Forester's Record. Quarterly Journal of Forestry. xxxiv.3: 116.
Forst Thuringia	Forst Thuringia unublished data: C. Rösner, Referat Forstsaatgutberatung, forstliches Forschungs- und Kompetenzzentrum Gotha, Referat Monitoring, Klima und Forschung, Gotha (e-mail 2016-05-10)
Gallego Zamorano et al. 2016	Zamorano, J.G., Hokkanen, T., and A. Lehikoinen. 2016. Climate driven synchrony in seed production of masting deciduous and conifer tree species. <i>Journal of Plant Ecology</i> . rtw117.
Gatter 1973	Gatter, W. 1973. Zugplanbeobachtungen an Spechten der Gattung Dendrocopos am Randecker Maar, Schwäbische Alb. <i>Anz. orn. Ges. Bayern</i> . 12: 122-129.
Genet 2010	Genet, H., Bréda, N. and E. Dufrêne. 2010. Age-related variation in carbon allocation at tree and stand scales in beech ( <i>Fagus sylvatica</i> L.) and sessile oak ( <i>Quercus petraea</i> (Matt.) Liebl.) using a chronosequence approach. <i>Tree Physiology</i> . 30(2): 177-192.
Gerasimidis et al. 2006	Gerasimidis, A., Panajiotidis, S., Hicks, S., and N. Athanasiadis. 2006. An eight-year record of pollen deposition in the Pieria mountains (N. Greece) and its significance for interpreting fossil pollen assemblages. <i>Review of Palaeobotany and Palynology</i> 141(3):231-243.
Gloaguen & Touffet 1982	Gloaguen, J. C. and J. Touffet. 1982. Production de litière dans une chênaie- hêtre Atlantique. Relations avec les caractères climatiques. <i>Revue forestière française</i> . 34(2): 108-118.
Granier et al. 2008	Granier, A., Bréda, N., Longdoz, B., Gross, P. and J. Ngao. 2008. Ten years of fluxes and stand growth in a young beech forest at Hesse, North-eastern

	France. Annals of Forest Science. 64(704): 1-13.
Gross 1934	Gross, H. 1934. Die Rotbuche in Ostpreussen. Zeitschrift für Forst-und Jagdwesen. 66/12: 622-651.
Grull 1957	Güll, J. 1957. Waldsamenernteaussicht 1957/58. Holz-Kurier. 12(36): 8-9.
Grulois et al. 2001	Grulois, C., De Meersman, F., Loyen, S., Orfinger, C., Quivy, V., De Vos, B. and B. Van der AA. 2001. Régénération naturelle de la Forêt de Soignes. Rapport Final. Instituut voor Bosbouw en Wildbeheer and Centre de Recherches Agronomiques de Gembloux.
Guggisberg 2002	Guggisberg, M., 2002. Walddynamik der letzten 70 Jahre bei Ponte Tresa/Tessin, Südalpen festgehalten in gewarvten Seesedimenten. MSc Thesis Phil. Nat. Fac. University of Bern, p. 83.
Guitian & da Costa 2005	Gutián, M. A. and F. J. da Costa. 2005. Primeros datos sobre la variabilidad interanual de la producción de semilla de <i>Fagus sylvatica</i> L. en el extremo occidental de la Cornisa Cantábrica. In Proceedings IV Congreso Forestal Nacional. SECF-DGA. 26-27 September Zaragoza, ES.
Guitian et al. 2005	Gutián, M. A., Vilariño, E. C., and F. J. da Costa. 2005. Caracterización biométrica y determinación de la viabilidad de las semillas y crecimiento de las plántulas de <i>Fagus sylvatica</i> L. provenientes de dos hayedos del parque natural de redes (Asturias). In Proceedings IV Congreso Forestal Nacional. SECF-DGA. 26-27 September Zaragoza, ES.
Gunton 1938	Gunton, H. C. 1938. Report on the phenological observations in the British Isles from December, 1936, to November, 1937. Quarterly Journal of the Royal Meteorological Society. 64(274): 135-198.
Gurnell 1993	Gurnell, J. 1993. Tree seed production and food conditions for rodents in an oak wood in southern England. Forestry. 66(3): 291-315.

Hacket-Pain; Personal observation	Hacket-Pain, A.; Personal observation
Hagner 1965	Hagner, S. 1965. Cone crop fluctuations in Scots pine and Norway spruce. Studia Forestalia Suecica 33, ed. Skogshögskolan, Stockholm.
Han et al. 2011	Han, Q., Kabeya, D. and G. Hoch. 2011. Leaf traits, shoot growth and seed production in mature <i>Fagus sylvatica</i> trees after 8 years of CO <sub>2</sub> enrichment. <i>Annals of Botany</i> . 107(8): 1405-1411.
Hansson 1971	Hansson, L. 1971. Small rodent food, feeding and population dynamics. A comparison between granivorous and herbivorous species in Scandinavia. <i>Oikos</i> . 22(2): 183-198.
Haredasht et al. 2013	Haredasht, S. A., Taylor, C. J., Maes, P., Verstraeten, W. W., Clement, J., Barrios, M., Lagrou, K., Van Ranst, M., Coppin, P., Berckmans, D. and J. M. Aerts. 2013. Model-based prediction of Nephropathia epidemica outbreaks based on climatological and vegetation data and bank vole population dynamics. <i>Zoonoses and Public Health</i> . 60(7): 461-477.
Hartig 1889	Hartig, R. 1889. Über den Einfluß der Samenproduktion auf Zuwachsgrösse und Reservestoffvorrathe der Bäume . Allg. Forst Jagdztg. 65, pp. 13–17
Hase 1964	Hase, W. 1964. Die Buchenmast in Schleswig-Holstein und ihre Abhängigkeit von der Witterung. Mitt. Deutsch. Wetterdienst. 31: 31/3-31/45
Hendry 2001	Hendry, S. J., Boswell, R. C. and J. C. Proudfoot. 2001. Forest condition 2000. Forestry Commission Information Note 46. Forestry Commission, Edinburgh.
Herbst et al. 2015	Herbst, M., Mund, M., Tamrakar, R., and A. Knohl. 2015. Differences in

	carbon uptake and water use between a managed and an unmanaged beech forest in central Germany. Forest Ecology and Management 355: 101-108.
Herget; Personal observation	Herget, K.; Personal observation
Hermann 1940	Hermann. 1940. Berichte über die Waldsamenernte. Deutsche Forstzeitung.
Hermansson et al. 2014	Hermansson S., Cato N., and U. Lamberth U. 2014. Skogsstyrelsen. Online: <a href="http://www.skogsstyrelsen.se/Myndigheten/Press-och-information/Pressmeddelanden1/Pressrelease/?releaseId=1643579">http://www.skogsstyrelsen.se/Myndigheten/Press-och-information/Pressmeddelanden1/Pressrelease/?releaseId=1643579</a>
Heroldova et al. 2013	Heroldová, M., Suchomel, J., Purchart, L. and L. Čepelka. 2013. Beech-mast crop evaluation in Kněhyně forest complex (Beskydy Mts. Czech Republic) as a food supply for granivorous rodents. Beskydy. 6(1): 27-32.
Hess 1905	Hess, R. A. 1905. Die Eigenschaften und das forstliche Verhalten der wichtigeren in Deutschland vorkommenden Holzarten: Ein Leitfaden für Studierende, Praktiker und Waldbesitzer. Parey, Berlin.
Heyman et al. 2012	Heyman, P., Thoma, B. R., Marié, J. L., Cochez, C. and S. S. Essbauer. 2012. In search for factors that drive hantavirus epidemics. Frontiers in Physiology. 3(237): 1-23.
Hilton & Packham 2003	Hilton, G. M. and J. R. Packham. 2003. Variation in the masting of common beech ( <i>Fagus sylvatica</i> L.) in northern Europe over two centuries (1800–2001). Forestry. 76(3): 319-328.
Hoch et al. 2013	Hoch, G., Siegwolf, R. T., Keel, S. G., Körner, C. and Q. Han. 2013. Fruit production in three masting tree species does not rely on stored carbon reserves. Oecologia. 171(3): 653-662.
Hoelzl et al. 2015	Hoelzl, F., Bieber, C., Cornils, J. S., Gerritsmann, H., Stalder, G. L., Walzer, C. and T. Ruf. 2015. How to spend the summer? Free-living dormice ( <i>Glis</i>

	glis) can hibernate for 11 months in non-reproductive years. Journal of Comparative Physiology. 185(8): 931-939.
Hofmann et al. 1992	Hofmann, G., Anders, S., Beck, W., Chzron, S. and B. Matthes. 1992. Buchenwälder in der ehemaligen DDR und ihr Vitalitätszustand. NZ NRW-Seminar. 12: 23-34.
Holmsgaard & Olsen 1960	Holmsgaard, E., and H. C. Olsen. 1960. The influence of weather on beech mast. Forstlige Forsøgsvaesen I Danmark. 26: 347-370.
Huss 1964	Huss, J. 1964. Untersuchungen über die natürliche Verjüngung der Buche: die Entwicklung des Aufschlages nach der Mast 1960. Dissertation der Forstwissenschaftlichen Fakultät der Georg-AugustUniversität zu Göttingen in Hann. Münden, 187 S.
Huss et al. 1972	Huss, J., Kratsch, H. D. and E. Röhrig. 1972. Ein Erfahrungsbericht über Maßnahmen zur Förderung der Buchennaturverjüngung, bei der Mast 1970 in acht Forstämtern Südniedersachsens. Forst- und Holzwirt. 27: 365-370.
Hyde 1963	Hyde, H. A. 1963. Pollen-fall as a means of seed prediction in certain trees. Grana. 4(2): 217-230.
Ihrig 1860	Ihrig. 1860. Ueber Wiederkehr der Mastjahre. Natürliche Verjüngung und Eckernertrag in Buchenhochwaldungen. Allgemeine Forst- und Jagdzeitung, 36. Jg., H. September, S. 341-350
Innes 1994	Innes, J. L. 1994. The occurrence of flowering and fruiting on individual trees over 3 years and their effects on subsequent crown condition. Trees. 8(3): 139-150.
Jacamon 1987	Jacamon, M. 1987. Le parc de l'École forestière à Nancy (France). Bulletin de la Société Botanique de France. Lettres Botaniques. 134(1): 29-34.
Jacobsen 2001	Jacobsen, E. M. 2001. Punkttællinger af ynglefugle i eng, by og skov 2000:

	Naturovervågning. Danmarks Miljøundersøgelser, Aarhus Universitet.
Jazewitsch 1953	Jazewitsch von, W. 1953. Jahrringchronologie der Spessart-Buchen. Forstwissenschaftliches Centralblatt. 72(7):234-247.
Jenni 1987	Jenni, L. 1987. Mass concentration of Bramblings <i>Fringilla montifringilla</i> in Europe 1900–1983: Their dependence upon beech mast and the effect of snow cover. <i>Ornis Scandinavica</i> . 18: 84-94.
Jensen 1982	Jensen, T. S. 1982. Seed production and outbreaks of non-cyclic rodent populations in deciduous forests. <i>Oecologia</i> . 54(2): 184-192.
Jezik et al. 2011	Ježík, M., Blaženec, M., Střelcová, K. and L. Ditmarová. 2011. The impact of the 2003–2008 weather variability on intra-annual stem diameter changes of beech trees at a submontane site in central Slovakia. <i>Dendrochronologia</i> . 29(4): 227-235.
Jonard et al. 2008	Jonard, M., André, F., Dambrine, E., Ponette, Q. and E. Ulrich. 2009. Temporal trends in the foliar nutritional status of the French, Walloon and Luxembourg broad-leaved plots of forest monitoring. <i>Annals of Forest Science</i> . 66(4): 1-10.
Jones 1952	Jones, E. W. 1952. Natural regeneration of beech abroad and in England. <i>Quarterly Journal of Forestry</i> . 46: 75-91.
Jutte 2013	Jütte, F. 2013. <i>Forstliches Rechnen: Ein Übungs- und Aufgabensammlungsbuch für angehende Forstleute</i> . Springer-Verlag, Berlin.
Kager & Fietz 2009	Kager, T. and J.Fietz. 2009. Food availability in spring influences reproductive output in the seed-preying edible dormouse ( <i>Glis glis</i> ). <i>Canadian Journal of Zoology</i> . 87(7): 555-565.
Kallander 1993	Källander, H. 1993. Food caching in the European nuthatch <i>Sitta europaea</i> . <i>Ornis Svecica</i> 3(2): 49-58.

Kampfer-Lauenstein & Lederer 2010	Kämpfer-Lauenstein, A. and W. Lederer. 2010. Populationsdynamik des Raufußkauzes <i>Aegolius funereus</i> im Arnsberger Wald. Charadrius. 46(1-2): 69-78.
Kantorowicz	Forest Research Institute (Poland)
Kaplunovskyy 1972	Kaplunovskyy C. 1972. Особенности плодоношения буковых лесов. Лесоведение 1:51-61.
Karlsson 2001	Karlsson, M. 2001. Natural Regeneration of Broadleaved Tree Species in Southern Sweden. P.h.D. Thesis, Swedish University of Agricultural Sciences, Alnarp.
Kasprzyk et al. 2014	Kasprzyk, I., Ortyl, B. and A. Dulska-Jeż. 2014. Relationships among weather parameters, airborne pollen and seed crops of <i>Fagus</i> and <i>Quercus</i> in Poland. Agricultural and Forest Meteorology. 197: 111-122.
Keller 1875	Keller, H. 1875. Waldsamenernte von 1874/75. Allgemeine Forst- und Jagdzeitung. 33.
Kihl et al. 2011	Khil, L., Samwald, O., Tiefenbach, A. u. M. and H. Pacher. 2011. Der Massenschlafplatz von Bergfinken ( <i>Fringilla montifringilla</i> ) in Österreich im Winter 2008/2009. Limicola. 25(2):81-100.
Kishazi 1982	Kisházi, Z. 1982. Makktermés utáni vizsgálatok véghasználati bükkösökben. Erdészeti Lapok. 31 (117.): 393-396.
Kleef & Wijsman 2015	Kleef, H. L. and H.J. Wijsman. 2015. Mast, mice and pine marten ( <i>Martes martes</i> ): the pine marten's reproductive response to wood mouse ( <i>Apodemus sylvaticus</i> ) fluctuations in the Netherlands. Lutra. 58(1): 23-33.
Klhare & Roth 1996	Klhare J, Roth H.U. 1996. Nutritional preferences of Abruzzo brown bears before hibernation. Proceedings of the II Conferenze of the Italian Association of Teriologia, III Italian Simposium on Carnivorous. Perugia,

	October 1996.
Konnert et al. 2014	Konnert, M., Schneck, D., and Zollner, A. (2014). Blühen und Fruktifizieren unserer Waldbäume in den letzten 60 Jahren. LWF Wissen, 74, 37-45.
Konnert; Personal observation	Konnert, M.; Personal observation
Krynytskyy; Personal observation	Krynytskyy, V.; Personal observation
Kulzer et al. 1993	Kulzer, E., von Lindeiner-Wildau, A. and I.-M. Wolters. 1993. Säugetiere im Naturpark Schönbuch. Landesanstalt für Umweltschutz Baden-Württemberg, Karlsruhe.
Latte et al. 2016	Latte, N., Lebourgeois, F. and H. Claessens. 2016. Growth partitioning within beech trees ( <i>Fagus sylvatica</i> L.) varies in response to summer heat waves and related droughts. Trees. 30: 189–201.
Lauprecht 1874	Lauprecht, G . 1874. Buchen- und Eichen-Samenjahre im Vergleich mit der Witterung. Zeitschrift für Forst- und Jagdwesen. 7: 246-266
Lavnyy; Personal observation	Lavnyy, M.; Personal observation
Le Tacon et al. 1976	Le Tacon, F., Oswald, H., Perrin, R., Picard, J. F. and J.P. Vincent. 1976. Les causes de l'échec de la régénération naturelle du hêtre à la suite de la fainée de 1974. Revue forestière française. 28(6): 426-446.
Lebl et al. 2010	Lebl, K., Kürbisch, K., Bieber, C. and T. Ruf. 2010. Energy or information? The role of seed availability for reproductive decisions in edible dormice. Journal of Comparative Physiology B. 180(3): 447-456.
Lebl et al. 2011	Lebl, K., Kürbisch, K., Bieber, C., and T. Ruf. 2010. Energy or information?

	The role of seed availability for reproductive decisions in edible dormice. Journal of Comparative Physiology B. 180(3): 447-456.
Lebret & Forgeard 2001	Lebret, M., Nys, C. and F. Forgeard. 2001. Litter production in an Atlantic beech ( <i>Fagus sylvatica</i> L.) time sequence. Annals of Forest Science. 58(7): 755-768.
Leemans 1991	Leemans, R. 1991. Canopy gaps and establishment patterns of spruce ( <i>Picea abies</i> (L.) Karst.) in two old-growth coniferous forests in central Sweden. Vegetatio. 93(2): 157-165.
LFD	LFD - Landesforstdirektion Südtirol
Liesebach 2012	Liesebach, M. 2012. Der Internationale Herkunftsversuch mit Rot-Buche von 1993/95—Beschreibung der ausgewählten sechs Herkünfte und zwei Versuchsflächen. Landbauforschung. 62(4): 159-168.
Lithner & Johnson 2002	Lithner, S., and K.I. Jonsson. 2002. Abundance of owls and bramblings <i>Fringilla montifringilla</i> in relation to mast seeding in south-eastern Sweden. Ornis Svecica. 12(1-2): 35-45.
Litschauer 2000	Litschauer, R. 2001. Blüh-und Fruktifikationsverhalten der Waldbäume. FBVA—Berichte. 123: 45-66.
Ljungstrom et al. 1990	Ljungström, M., Gyllin, M. and B. Nihlgård. 1990. Effects of liming on soil acidity and beech ( <i>Fagus sylvatica</i> L.) regeneration on acid soils in south Swedish beech forests. Scandinavian Journal of Forest Research. 5(1-4): 243-254.
Louarn & Schmitt 1972	Le Louarn, H. and A. Schmitt. 1972. Relations observées entre la production de Faines et la dynamique de population du Mulot, <i>Apodemus sylvaticus</i> L. en forêt de Fontainebleau. In Annales des Sciences Forestieres. 29(2): 205-214.

Lupke 2007	Lüpke von, B.V. 2008. Einfluss unterschiedlicher Hiebsformen auf die Naturverjüngung eines Traubeneichen-Buchen Mischbestandes. Forsarchiv. 79: 4-14.
MastWeb	Forschungsanstalt für Wald, Schnee und Landschaft WSL <a href="http://www.wsl.ch/mastweb/resultate/index_DE">http://www.wsl.ch/mastweb/resultate/index_DE</a>
Madsen 1995	Madsen, P. 1995. Effects of seedbed type on wintering of beech nuts ( <i>Fagus sylvatica</i> ) and deer impact on sprouting seedlings in natural regeneration. Forest Ecology and Management. 73(1): 37-43.
Majer 1982	Majer, A. 1982. A bükkösök makk termésének időszakossága. Erdészeti Lapok. 31(9): 388-392.
Mancini et al. 2016	Mancini N.M., Mancini G.M., Travaglini D., Nocentini S. and R. Giannini. 2016. Prime osservazioni sulla struttura e la produzione di seme dei boschi cacuminali di faggio nei Monti della Laga (Parco Nazionale del Gran Sasso e dei Monti della Laga). L’Italia Forestale e Montana. 71 (1): 31-47
Margary 1936	Margary, I. D. 1936. Report on the phenological observations in the British Isles from December, 1934, to November, 1935. No. 45. Quarterly Journal of the Royal Meteorological Society. 62(265): 299-358.
Maringer; Personal observation	Maringer, J.; Personal observation
Marteau & Sara 2015	Marteau, M. and M. Sarà. 2015. Habitat preferences of edible dormouse, <i>Glis glis italicus</i> : implications for the management of arboreal mammals in Mediterranean forests. Folia Zoologica 64 (2): 136-150
Martin 2009	Martin, H. 2009. Caractérisation des fructifications des chênaies et hêtraies du réseau Renecofor. Diplôme de l’École Pratique des Hautes Études.

Martinez et al. 2009	Martínez, I. and F. González-Taboada. 2009. Seed dispersal patterns in a temperate forest during a mast event: performance of alternative dispersal kernels. <i>Oecologia</i> . 159(2): 389-400.
Matic et al. 2003	Matic, S., Orsanic, I. and I. Anic. 2003. Establishing forests of Common beech. In: Common beech ( <i>Fagus sylvatica</i> L.) in Croatia. Academy of Forestry Sciences. 326-339.
Matthysen 1989	Matthysen, E. 1989. Nuthatch <i>Sitta europaea</i> demography, beech mast, and territoriality. <i>Ornis Scandinavica</i> . 20(4):278-282.
Maurer & Tabel 2000	Maurer, W. D. and U. Tabel. 2000. Einrichtung und Bewirtschaftung forstlicher Generationsbestände am Beispiel der Buche ( <i>Fagus sylvatica</i> L.) in Rheinland-Pfalz (Deutschland). <i>Forest, Snow &amp; Landscape Research</i> . 75(1/2): 219-231.
Maurer 1964	Maurer, E. 1964. Buchen-und Eichensamenjahre in Unterfranken während der letzten 100 Jahre. <i>Allgemeine Forstzeitschrift</i> . 31: 469-470.
Meiffren 1988	Meiffren, I. 1988. Airborne pollen of Toulouse, southern France: comparison with Bordeaux and Montpellier. <i>Grana</i> . 27(3):183-201.
Melgar et al. 2012	Melgar, M., Trigo, M. M., Recio, M., Docampo, S., García-Sánchez, J. and B. Cabezudo. 2012. Atmospheric pollen dynamics in Münster, north-western Germany: a three-year study (2004–2006). <i>Aerobiologia</i> . 28(4): 423-434.
Mellstrom 1918	Mellström, G. 1918. Skogsträdens frösättning år 1917. <i>Statens Skogsforstoksanstalt</i> . 15 :43-68.
Mencuccini et al. 1995	Mencuccini, M., Piussi, P., and A. Z. Sulli. 1995. Thirty years of seed production in a subalpine Norway spruce forest: patterns of temporal and spatial variation. <i>Forest Ecology and Management</i> . 76(1): 109-125.
Mancuccini &	Mencuccini, M., and P. Piussi. 1995. Production of seed and cones and

Piussi 1995	consequences for radial increment in Norway spruce ( <i>Picea abies</i> (L.) Karst.). Giornale Botanico Italiano. 129(3): 797-812.
MeteoSwiss	Federal Office of Meteorology and Climatology (MeteoSwiss), Switzerland
Mezzavilla 2014	Mezzavilla F. 2014. Il faggio e la fauna. Indagini ecologiche nella riserva naturale biogenetica campo di mezzo - Pian Parrocchia. Foresta del Cansiglio. Ministero delle Politiche Agricole e Forestali.
Michaelis 1911	Michaelis, 1911. Einiges zur Buchenmast 1909. Zeitschrift f. Forst- und Jagdwesen. 43: 267-283.
Milleron et al. 2012	Millerón, M., De Heredia, U. L., Lorenzo, Z., Perea, R., Dounavi, A., Alonso, J., Gil, L. and N. Nanos. 2012. Effect of canopy closure on pollen dispersal in a wind-pollinated species ( <i>Fagus sylvatica</i> L.). Plant Ecology. 213(11): 1715-1728.
Morris & Morris 2010	Morris, P. A. and M.J. Morris. 2010. A 13-year population study of the edible dormouse <i>Glis glis</i> in Britain. Acta Theriologica. 55(3): 279-288.
Moskava & Adamik 2012	Mašková, P. and P. Adamík. 2012. Poznámky o výskytu arboreálních hlodavců (Mammalia: Rodentia) v budkách na Sovinecku, Nízký Jeseník. Zprávy Vlastivědného muzea v Olomouci. 303: 13–21.
Mountford et al. 2006	Mountford, E. P., Savill, P. S. and D.P. Bebb. 2006. Patterns of regeneration and ground vegetation associated with canopy gaps in a managed beechwood in southern England. Forestry. 79(4): 389-408.
Muller 1952	Müller, W.L . 1952. Über die Witterungsabhängigkeit von Samenerträgen bei Buchen und Eichen. Bericht d. dt. Wetterdienstes in der US-Zone 58.
Muller-Haubold et al. 2015	Müller-Haubold, H., Hertel, D. and C. Leuschner. 2015. Climatic drivers of mast fruiting in European Beech and resulting C and N allocation shifts. Ecosystems. 18(6): 1083-1100.

Mund et al. 2010	Mund, M., Kutsch, W. L., Wirth, C., Kahl, T., Knohl, A., Skomarkova, M. V. and E.D. Schulze. 2010. The influence of climate and fructification on the inter-annual variability of stem growth and net primary productivity in an old-growth, mixed beech forest. <i>Tree Physiology.</i> 30: 689-704.
Nather 1962	Nather, H. 1962. Waldsamen-Ernteaussichten für das Jahr 1962. <i>Allgemeine Forstzeitung.</i> 73(60): 17-18.
Nather 1963	Nather, H. 1963. Waldsamen-Ernteaussichten für das Jahr 1963. <i>Holz-Kurier.</i> 18(40): 7.
Nather 1966	Nather, H. 1966. Ernteaussichten für Waldsamen im Jahre 1966. <i>Holz-Kurier.</i> 21 (41): 7-8.
Neckarova 2011	Neckařová, M. 2012. Velkoplošná variabilita plodování buku lesního: semenné roky a prostorová synchronie. <i>Univerzita Palackého v Olomouci.</i>
Newson 1963	Newson, R. 1963. Differences in numbers, reproduction and survival between two neighboring populations of bank voles ( <i>Clethrionomys glareolus</i> ). <i>Ecology.</i> 44(1): 110-120.
Nielsen 1977	Nielsen, B. O. 1977. Beech seeds as an ecosystem component. <i>Oikos.</i> 29(2): 268-274.
Nielsen 2005	Nielsen, J. T. 2005. Yngletidspunktets betydning for produktionen af unger og deres overlevelse hos Spurvehøgen <i>Accipiter nisus</i> i Vendsyssel 1977-97. <i>Dansk Orn. Foren. Tidsskr.</i> 99: 107-114.
Nielsen et al. 2010	Nielsen, A. B., Møller, P. F., Giesecke, T., Stavngaard, B., Fontana, S. L. and R. H. Bradshaw. 2010. The effect of climate conditions on inter-annual flowering variability monitored by pollen traps below the canopy in Draved Forest, Denmark. <i>Vegetation History and Archaeobotany.</i> 19(4): 309-323.
Niklasson 2003	Niklasson, M. 2003. En undersökning av trädåldrar i halländska

	skogsreservat. Information från. Länsstyrelsen halland.
Nilsson & Wastljung 1987	Nilsson, S. G. and U. Wastljung. 1987. Seed predation and cross-pollination in mast-seeding beech ( <i>Fagus sylvatica</i> ) patches. <i>Ecology</i> . 68(2): 260-265.
Nilsson 1985	Nilsson, S. G. 1985. Ecological and evolutionary interactions between reproduction of beech <i>Fagus sylvatica</i> and seed eating animals. <i>Oikos</i> . 44(1): 157-164.
Nopp-Mayr et al. 2012	Nopp-Mayr, U., Kempter, I., Muralt, G. and G. Gratzer. 2012. Seed survival on experimental dishes in a central European old-growth mixed-species forest—effects of predator guilds, tree masting and small mammal population dynamics. <i>Oikos</i> . 121(3):337-346.
Novakova et al. 2011	Nováková, P., Štípek, K., Ježek, M., Červený, J. and V. Ešner. 2011. Effect of diet supply and climatic conditions on population dynamics of the wild boar ( <i>sus scrofa</i> ) in the křivoklát region (Central Bohemia,Czech Republic). <i>Scientia Agriculturae Bohemica</i> . 42(1): 24-30.
Nussbaumer et al. 2014	Nussbaumer, A., Waldner, P. and S. Etzold. 2014. Fruiting occurrence of Beech, Spruce, Pine and Oak trees in Europe. Report I. Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf.
Nussbaumer et al. 2016	Nussbaumer, A., Waldner, P., Etzold, S., Gessler, A., Benham, S., Thomsen, I. M., Jørgensen, B.B., Timmermann, V., Verstraeten, A., Sioen, g., Rautio, P., Ukonmaanaho, L., Skudnik, M., Apuhtin, V., Braun, S. and A. Wauer. 2016. Patterns of mast fruiting of common beech, sessile and common oak, Norway spruce and Scots pine in Central and Northern Europe. <i>Forest Ecology and Management</i> . 363: 237-251.
Nussbaumer presentation	Nussbaumer, A., Etzold, S., Waldner, P. and Dobbertin, M. Carbon allocation to fruits and seeds in European forests as a function of climate,

	atmospheric deposition and nutrient supply. COST Action FP0903: Climate Change and Forest Migration and Adaption in a Polluted Environment. ( <a href="http://baw.ac.at/cms_stamm/430/pdf/ffcc/EP14/07_Nussbaumer.pdf">http://baw.ac.at/cms_stamm/430/pdf/ffcc/EP14/07_Nussbaumer.pdf</a> ) Access 2016-07-01
Oddou-Muratorio 2011	Oddou-Muratorio, S., Bontemps, A., Klein, E. K., Chybicki, I., Vendramin, G. G. and Y. Suyama. 2010. Comparison of direct and indirect genetic methods for estimating seed and pollen dispersal in <i>Fagus sylvatica</i> and <i>Fagus crenata</i> . Forest Ecology and Management. 259(11): 2151-2159.
Olesen & Madsen 2008	Olesen, C. R., and P. Madsen. 2008. The impact of roe deer ( <i>Capreolus capreolus</i> ), seedbed, light and seed fall on natural beech ( <i>Fagus sylvatica</i> ) regeneration. Forest Ecology and Management. 255(12): 3962-3972.
Oosterbaan 1982	Oosterbaan, A. 1982. Voorbereidingen voor het benutten van de beukenmast 1982. Nederlands bosbouw Tijdschrift. 54(9): 284-287.
Ott et al. 2003	Ott, E., Conceprio, F. and A. Pedrini. 2003. Prime valutazioni sull'introduzione della rinnovazione naturale nel bosco ceduo di castagno misto a faggio nella foresta sperimentale e didattica della SPF di Zurigo a Novaggio, Cantone Ticino. Schweizerische Zeitschrift fur Forstwesen. 154(2):51-67.
Overgaard et al. 2007	Overgaard, R., Gemmel, P. and M. Karlsson. 2007. Effects of weather conditions on mast year frequency in beech ( <i>Fagus sylvatica</i> L.) in Sweden. Forestry. 80(5): 555-565.
Overgaard; Personal observation	Overgaard, R.; Personal observation
Paar et al. 2011	Paar, U., Guckland, A., Dammann, I., Albrecht, M. and J. Eichhorn. 2011.

	Häufigkeit und Intensität der Fruktifikation der Buche. AFZ-Der Wald. 6: 26-29.
Pakenham 1996	Pakenham, R. 1996. Natural regeneration of beech in the Chilterns. Quarterly Journal of Forestry. 90(2): 143-149.
Palaghianu; Personal observation	Palaghianu, C.; Personal observation
Parmigiani 2007	Parmigiani, S. 2007. Spatial behaviour of the yellow-necked mouse (Apodemus flavicollis, melchior 1834) at contrasting population density and resource availability. P.h.D Thesis, University of Parma.
Perdeck et al. 2000	Perdeck, A. C., Visser, M. E. and J. H. Van Balen. 2000. Great tit <i>Parus</i> major survival and the beech-crop. <i>Ardea</i> . 88: 99-106.
Perrin 1978	Perrin, R. 1978. Etude de la sporulation de <i>Nectria Ditissima</i> Tul. Agent du chancre du hêtre. <i>Annales des Sciences Forestières</i> . 35(3): 213-228.
Perrins 1965	Perrins, C. M. 1965. Population fluctuations and clutch-size in the Great Tit, <i>Parus major</i> L. <i>The Journal of Animal Ecology</i> . 34(3): 601-647.
Petty et al. 1995	Petty, S. J., Patterson, I. J., Anderson, D. I. K., Little, B., and M. Davison. 1995. Numbers, breeding performance, and diet of the sparrowhawk <i>Accipiter nisus</i> and merlin <i>Falco columbarius</i> in relation to cone crops and seed-eating finches. <i>Forest ecology and management</i> . 79(1): 133-146.
Pidek et al. 2010	Pidek, I. A., Svitavská-Svobodová, H., van der Knaap, W. O., Noryśkiewicz, A. M., Filbrandt-Czaja, A., Noryśkiewicz, B., Filbrandt-Czaja, A., Noryśkiewicz, B., Latałowa, M., Zimny, M., Święta-Musznicka, J., Bozilova, E., Tonkov, S., Filipova-Marinova, M., Poska, A., Giesecke, T. and A. Gikov. 2010. Variation in annual pollen accumulation rates of <i>Fagus</i>

	along a N–S transect in Europe based on pollen traps. <i>Vegetation History and Archaeobotany</i> . 19(4): 259-270.
Pidek et al. 2015	Pidek, I. A., Poska, A. and B. M. Kaszewski. 2015. Taxon-specific pollen deposition dynamics in a temperate forest zone, SE Poland: the impact of physiological rhythmicity and weather controls. <i>Aerobiologia</i> 31(2): 219-238.
Pilastro et al. 2003	Pilastro, A., Tavecchia, G. and G. Marin. 2003. Long living and reproduction skipping in the fat dormouse. <i>Ecology</i> . 84(7): 1784-1792.
Pilegaard et al. 2011	Pilegaard, K., Ibrom, A., Courtney, M. S., Hummelshøj, P. and N. O. Jensen. 2011. Increasing net CO <sub>2</sub> uptake by a Danish beech forest during the period from 1996 to 2009. <i>Agricultural and Forest Meteorology</i> . 151(7): 934-946.
Piovesan & Bernabei 1997	Piovesan, G. and M. Bernabei. 1997. L'influenza delle precipitazioni estive sulla crescita e la riproduzione del faggio ( <i>Fagus sylvatica</i> L.) in una stazione meridionale dell'areale. <i>Italia Forestale e Montana</i> . 6: 444-459.
Piussi; unpublished data	Pietro, P., Istituto di Selvicoltura, Facoltà di Agraria, Università di Firenze; unpublished data
Poska & Pidek 2010	Poska, A. abd I. A. Pidek. 2010. Pollen dispersal and deposition characteristics of <i>Abies alba</i> , <i>Fagus sylvatica</i> and <i>Pinus sylvestris</i> , Roztocze region (SE Poland). <i>Vegetation History and Archaeobotany</i> . 19(2): 91-101.
Potena et al. 2008	Potena G., Sammarone L., Panella M. and M. Romano. 2008. Progetto Life Natura 99/NAT/IT/006244. Conservazione dell'orso bruno nell'Appennino centrale. Azione C1: Linee guida di gestione forestale. Pg. 1-25
Potena et al. 2009	Potena, G., Di Marzio. M., Panella, M., Sammarone, L., Altea, T., Posillico, M., Roman, M., and M. Consalvo. 2009. Il monitoraggio delle produzioni di faggiola: una risorsa trofica critica per l'orso bruno ( <i>Ursus arctos</i> ). VII

	Conference of the Italian Society of Silviculture and Forest Ecology. Isernia, Pesche, 29 September, 2009.
Prach et al. 1996	Prach, K., J. Lepš, and J. Michalek. 1996. Establishment of <i>Picea abies</i> seedlings in a central European mountain grassland: an experimental study. Journal of Vegetation Science. 7: 681-684.
Pucek et al. 1993	Pucek, Z., Jędrzejewski, W., Jędrzejewska, B., and M. Pucek. 1993. Rodent population dynamics in a primeval deciduous forest (Białowieża National Park) in relation to weather, seed crop, and predation. Acta Theriologica. 38(2): 199-232.
Pukkala 2010	Pukkala, T., Hokkanen, T., and T. Nikkanen. 2010. Prediction models for the annual seed crop of Norway spruce and Scots pine in Finland. Silva Fennica. 44(4): 629-642.
Regierungspräsidi um Freiburg	Regierungspräsidium Freiburg
Relazioni annuali della Sezione Forestale	Relazioni annuali della Sezione Forestale del Canton Ticino. Dipartimento del territorio, Repubblica e Cantone del Ticino, Bellinzona.
Renecofor	French National Forest Office, National Network for Long-term FORest ECOSystem Monitoring
Rohle et al. 2009	Röhle, H., Gerold, D. and R. Gemballa. 2010. Beziehungen zwischen Klima und Zuwachs, dargestellt am Beispiel von Fichte, Kiefer und Buche in Sachsen. Allgemeine Forst und Jagdzeitung. 181: 21-35.
Rohrig et al. 1978	Röhrig, E., Bartels, H., Gussone, H. A. and B. Ulrich. 1978. Untersuchungen zur natürlichen Verjüngung der Buche ( <i>Fagus sylvatica</i> ). Forstwissenschaftliches Centralblatt. 97/1: 121-131.

Romania Survey	Romanian National Forest Administration - Romsilva, Suceava County Division
Rossa 1992	Rossa, G. 1992. Geschichtliche Entwicklung der Laubholzbestände. Heimatland Lippe. 85(3): 90.
Ruf et al. 2006	Ruf, T., Fietz, J., Schlund, W. and C. Bieber. 2006. High survival in poor years: Life history tactics adapted to mast seeding in the edible dormouse. Ecology. 87(2): 372-381.
Salmaso et al. 2009	Salmaso, F., Molinari, A., Di Pierro, E., Ghisla, A., Martinoli, A., Preatoni, D., Serino, G., Tosi, G., Bertolino, S. and L. A. Wauters. 2009. Estimating and comparing food availability for tree-seed predators in typical pulsed-resource systems: Alpine conifer forests. Plant Biosystems. 143(2): 258-267.
Scharnweber et al. 2011	Scharnweber, T., Manthey, M., Criegee, C., Bauwe, A., Schröder, C. and M. Wilmking. 2011. Drought matters—Declining precipitation influences growth of <i>Fagus sylvatica</i> L. and <i>Quercus robur</i> L. in North-Eastern Germany. Forest Ecology and Management. 262(6): 947-961.
Schlund et al. 2002	Schlund, W., Scharfe, F. and J. U. Ganzhorn. 2002. Long-term comparison of food availability and reproduction in the edible dormouse ( <i>Glis glis</i> ). Mammalian Biology-Zeitschrift für Säugetierkunde. 67(4): 219-232.
Schmidt 2006	Schmidt, W. 2006. Zeitliche Veränderung der Fruktifikation bei der Rotbuche ( <i>Fagus sylvatica</i> ) in einem Kalkbuchenwald (1981-2004). Allgemeine Forst- u. Jagd-Zeitung. 177: 9-19.
Schneck; unpublished data	Schneck, R.; unpublished data
Schwappach 1895	Schwappach, A. 1895. Die Samenproduktion der wichtigsten Waldholzarten in Preussen. Zeitschrift für Forst- und Jagdwesen. 27: 147-174.

Schwarz 1870	Königlich Preußischer Forstmeiser Schwarz von Erfurt. 1870. Ein Beitrag zur Buchenwirtschaft im höherem Gebirge des Thüringer Waldes. Allgemeine Forst- und Jagdzeitung. 55-95.
Schwarz et al. 2009	Schwarz, A. C., Ranft, U., Piechotowski, I., Childs, J. E. and S. O. Brockman. 2009. Risk factors for human infection with Puumala virus, Southwestern Germany. Emerging Infectious Diseases. 15(7): 1032-1039.
Seeger 1913	Seeger. 1913. Ein Beitrag zur Samenproduktion der Waldbäume im Großherzogtum Baden. Naturwissenschaftliche Zeitschrift für Forst- und Landwirtschaft. 11: 529-554.
Seifert & Muller-Starck 2009	Seifert, T. and G. Müller-Starck. 2009. Impacts of fructification on biomass production and correlated genetic effects in Norway spruce ( <i>Picea abies</i> [L.] Karst.). European Journal of Forest Research. 128(2): 155-169.
Selas et al. 2002	Selås, V., Framstad, E., and T. K. Spidsø. 2002. Effects of seed masting of bilberry, oak and spruce on sympatric populations of bank vole ( <i>Clethrionomys glareolus</i> ) and wood mouse ( <i>Apodemus sylvaticus</i> ) in southern Norway. Journal of Zoology. 258(4): 459-468.
Silva et al. 2012	Silva, D. E., Mazzella, P. R., Legay, M., Corcket, E. and J. L. Dupouey. 2012. Does natural regeneration determine the limit of European beech distribution under climatic stress? Forest Ecology and Management. 266: 263-272.
Simoleit et al. 2016	Simoleit, A., Wachter, R., Gauger, U., Werchan, M., Werchan, B., Zuberbier, T. and K. C. Bergmann. 2016. Pollen season of European beech ( <i>Fagus sylvatica</i> L.) and temperature trends at two German monitoring sites over a more than 30-year period. Aerobiologia (in press). DOI:10.1007/s10453-016-9421-y.

Sioen et al. 2008	Sioen, G., Roskams, P., Verschelde P., Van der Aa, B. and A. Verstraeten. 2008. Monitoring the masting behaviour of beech ( <i>Fagus sylvatica</i> ) in Flanders (Belgium). In Proceedings of the Forest Adaptation Conference, FAO. Session 1. Physiological responses of trees to climate. Poster presentation. Umeå, Sweden from 25-28 August 2008.
Slovenian Forest Service	Slovenian Forest Service (Zavod za Gozdovje Slovenije). 2016. Poročilo zavoda za gozdove slovenije o gozdovih. Za leto 2015. Ljubljana.
Sohn et al. 2013	Sohn, J. A., Gebhardt, T., Ammer, C., Bauhus, J., Häberle, K. H., Matyssek, R., and T. E. Grams. 2013. Mitigation of drought by thinning: short-term and long-term effects on growth and physiological performance of Norway spruce ( <i>Picea abies</i> ). Forest Ecology and Management. 308: 188-197.
Spek et al. 2015	Spek, G.J. and A. Vliet. 2015. Veel eikels aan de bomen, record aantal zwijnen in 2016 verwacht. ( <a href="https://www.naturetoday.com/intl/nl/nature-reports/message/?msg=21765">https://www.naturetoday.com/intl/nl/nature-reports/message/?msg=21765</a> ) Access Date 2016-07-01.
SRS	Survey of the Public Forest Enterprise "Sume Republike Srpske" (Forests of the Republic of Srpska; <a href="http://www.sumers.org/portal/index.php">http://www.sumers.org/portal/index.php</a> )
Staatsklenge Forstamt Nagold	Staatsklenge Forstamt Nagold
Standovar & Kenderes 2003	Standovár, T. and K. Kenderes. 2003. A review on natural stand dynamics in beechwoods of East Central Europe. Applied Ecology and Environmental Research. 1(1): 19-46.
Suchomel 2007	Suchomel, J. 2014. A study of the synusia of small terrestrial mammals (Insectivora, Rodentia) of the Kelečská pahorkatina Upland–Czech Republic. Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis. 55(5): 165-170.

Suchomel et al. 2014	Suchomel, J., Purchart, L., Čepelka, L. and M. Heroldová. 2014. Structure and diversity of small mammal communities of mountain forests in Western Carpathians. European Journal of Forest Research. 133(3): 481-490.
Svardson 1957	Svardson, G. 1957. The "invasion" type of bird migration. British Birds. 50: 314-343.
Szewczyk & Szwalgrzyk 2010	Szewczyk, J. and J. Szwalgrzyk. 2010. Spatial and temporal variability of natural regeneration in a temperate old-growth forest. Annals of Forest Science. 67(2): 202.
Szwagrzyk et al. 2015	Szwagrzyk, J., Gratzer, G., Stepniewska, H., Szewczyk, J. and B. Veselinovic. 2015. High reproductive effort and low recruitment rates of European beech: Is there a limit for the superior competitor?. Polish Journal of Ecology. 63: 198-212.
Tacon & Malphettes 1974	Tacon, F. Le and C. V. Malphettes. 1974. Germination et comportement de semis de hêtre sur six stations de la Forêt domaniale de Villers-Cotterets. Revue Forestière Française. 26: 111-123.
Teissier 1977	Teissier, D. C. 1977. Étude de la variabilité du Hêtre ( <i>Fagus sylvatica</i> L.). Revue Forestière Française. 29(5): 355-362.
Tersago et al. 2009	Tersago, K., Verhagen, R., Servais, A., Heyman, P., Ducoffre, G. and H. Leirs. 2009. Hantavirus disease (nephropathia epidemica) in Belgium: effects of tree seed production and climate. Epidemiology and Infection. 137(02): 250-256.
Thissen 2010	Thissen, J. 2010. Knaagdiermonitoring in Nederland. Zoogdiervereniging Rapport 2010. 36. Zoogdiervereniging, Nijmegen.
Tirén 1935	Tirén, L., 1935. Om granens kottsättning, dess periodicitet och samband med temperatur och nederbörd (On the fruit setting of spruce, its periodicity and

	relation to temperature and precipitation). Meddelanden Från Statens Skogsforsöksanst Häfte. 28(4): 413-521.
Tollefsrud 2015	Tollefsrud, M. M. 2015. Blomstring hos gran gir frø til skogbruket. Norsk institutt for biookonomi. <a href="http://www.skogoglandskap.no/nyheter/2015/blomstring_hos_gran_gir_fro_til_skogbruket">(<a href="http://www.skogoglandskap.no/nyheter/2015/blomstring_hos_gran_gir_fro_til_skogbruket">http://www.skogoglandskap.no/nyheter/2015/blomstring_hos_gran_gir_fro_til_skogbruket</a>)</a> . Date of access 2016-06-27.
Topoliantz & Ponge 2000	Topoliantz, S. and J. F. Ponge. 2000. Influence of site conditions on the survival of <i>Fagus sylvatica</i> seedlings in an old-growth beech forest. Journal of Vegetation Science. 11(3): 369-374.
Tosoni et al. 2014	Tosoni, E., Gentile, D., Altea, T., Latini, R. and P. Ciucci. 2014. Conta cumulativa delle unità familiari di orso bruno marsicano per la stima della produttività della popolazione: estate 2014. UE Project LifeNAT/IT/000160 "ARCTOS", Dept. Biology and Biotechnologies, University of Rome "La Sapienza", Roma, Italy.
Trauboldt; Personal observation	Trauboth, V.; Personal observation
UK survey 1980-2007	Packham, J. R., Thomas, P. A., Lageard, J. G. A., and G. M. Hilton. 2008. The English beech masting survey 1980–2007. Variation in the fruiting of the common beech ( <i>Fagus sylvatica</i> L.) and its effects on woodland ecosystems. Arboricultural Journal 31(3):189-214.
UK survey 2008-2016	UK survey - Lageard, J.G.A and Thomas, P.
Vacchiano; Personal	Vacchiano, G.; Personal observation

observation	
Vacek & Hejcman 2012	Vacek, S. and M. Hejcman. 2012. Natural layering, foliation, fertility and plant species composition of a <i>Fagus sylvatica</i> stand above the alpine timberline in the Giant (Krkonoše) Mts., Czech Republic. European Journal of Forest Research. 131(3): 799-810.
Vacek & Jurasek 1986	Vacek, S. and A. Jurasek. 1986. Fruktifikace bukovych porostů pod vlivem imis. Opera Corcontica. 23: 111-142.
van der Knaap et al. 2010	Van der Knaap, W.O., van Leeuwen, J.F., Svitavská-Svobodová, H., Pidek, I. A., Kvavadze, E., Chichinadze, M., Giesecke, T., Kaszewski, B.M., Oberli, F., Kalnina, L., Pardoe, H. S., Tinner, W. and B. Ammann. S. 2010. Annual pollen traps reveal the complexity of climatic control on pollen productivity in Europe and the Caucasus. Vegetation History and Archaeobotany. 19(4): 285-307.
van der Maaten 2012	Van der Maaten, E. 2012. Climate sensitivity of radial growth in European beech ( <i>Fagus sylvatica</i> L.) at different aspects in southwestern Germany. Trees. 26(3): 777-788.
Vanha-Majamaa et al. 1996	Vanha-Majamaa, I., Tuittila, E. S., Tonteri, T., and R. Suominen. 1996. Seedling establishment after prescribed burning of a clear-cut and a partially cut mesic boreal forest in southern Finland. Silva Fennica. 30(1): 31-45.
Vieghofer 1952	Vieghofer, L. 1952. Die Rotbuchenverjüngung des Wienerwaldes in ihrer Abhängigkeit vom Boden. Bundesforschungszentrum für Wald, Wien.
Viel et al. 2010	Viel, J. F., Lefebvre, A., Marianneau, P., Joly, D., Giraudoux, P., Upegui, E., Tordo, N., Hoen, B. 2011. Environmental risk factors for haemorrhagic fever with renal syndrome in a French new epidemic area. Epidemiology and Infection. 139(6): 867-874.

Wachter 1964	Wachter, H. 1964. Über die Beziehungen zwischen Witterung und Buchenmastjahren. Forstarchiv. 35(4): 69-78.
Watt 1923	Watt, A. S. 1923. On the ecology of British beechwoods with special reference to their regeneration. Journal of Ecology. 11(1): 1-48.
Watt 1925	Watt, A. S. 1925. On the ecology of British beechwoods with special reference to their regeneration: part II, sections II and III. The development and structure of beech communities on the Sussex Downs. Journal of Ecology. 13(1): 27-73.
Wauters et al. 2007	Wauters, L. A., Vermeulen, M., Van Dongen, S., Bertolino, S., Molinari, A., Tosi, G. and E. Matthysen. 2007. Effects of spatio-temporal variation in food supply on red squirrel <i>Sciurus vulgaris</i> body size and body mass and its consequences for some fitness components. Ecography. 30(1): 51-65.
Wesolowski et al. 2015	Wesołowski, T., Rowiński, P., and M. Maziarz. 2015. Interannual variation in tree seed production in a primeval temperate forest: does masting prevail?. European Journal of Forest Research. 134(1): 99-112.
Westergren et al.; unpublished data	Westergren, M., Bozic, G., Brus, R., Grech, Z., Minic, M., and H. Kraigher; unpublished data
Witteczek 2009	Witteczek, K. 2009. Cone and seed entomofauna of Norway spruce <i>Picea abies</i> (L.) Karst. In the selected stands of the Gorce National Park. Part II. Influence of altitude and of variable spruce stands masting on the chosen insect species. Acta Agraria et Silvestria. 47: 73-92.
WSL	Swiss Federal Institute for Forest, Snow and Landscape Research (WSL)
Wu et al. 2013	Wu, J., Larsen, K. S., van der Linden, L., Beier, C., Pilegaard, K. and A. Ibrom. 2013. Synthesis on the carbon budget and cycling in a Danish, temperate deciduous forest. Agricultural and Forest Meteorology. 181: 94-

	107.
Zingg & Brang 2003	Zingg, A. and P. Brang. 2003. Sterben Buchen wegen der Trockenheit?. Wald und Holz. 9(03): 44-46.
Zwander H.; Pollenwarndienst – Kärnten	Zwander H., Klagenfurt; EAN-Datenbank Wien. Bearbeitung: R. Litschauer, Genetik, BFW-Wien
Zwolak et al. 2016	Zwolak, R., Bogdziewicz, M. and L. Rychlik. 2016. Beech masting modifies the response of rodents to forest management. Forest Ecology and Management. 359: 268-276.

429

430 10. Comments

431 For some data, a brief comment provided by the data compiler may be included in the 18<sup>th</sup> column.

432 In the case of continuous data from scientific and the grey literature, we report the table or figure  
433 number (if available) from which we extracted data values.

434

435 CLASS V. SUPPLEMENTARY DESCRIPTORS

436 A. Data acquisition

437

438 Data forms: n/a

439 Location of completed data forms: n/a

440 Data entry/verification procedures: Data were introduced in a spreadsheet from published  
441 references and unpublished series. The main compilers (Ascoli and Maringer) reviewed all  
442 individual series to homogenise criteria and to detect any inconsistencies.

443

444 B. Quality assurance/quality control procedures: see Authentication procedure (Class IV).

445

446 C. Related material: n/a

447

448 D. Computer programs and data processing algorithms:

449 The file can be read using different statistical, database or spreadsheet software. The command line  
450 to read it in R version 3.2.5 (R Development Core Team 2016) reads:

451 `dataFrameName <- read.csv("MASTREE_2016.11.csv")`

452

453 E. Archiving: n/a

454

455 F. Publications using the data set: The full data set has not yet been used in any publication. Several  
456 papers using the database are in preparation by the same authors.

457

458 G. History of data set usage: n/a (the data has not yet been used by any secondary user).

459

## 460 ACKNOWLEDGEMENTS

461 We thank following colleagues and Institutions for providing further data: Rudolf Litschauer, BFW  
462 Wien; Corpo Forestale dello Stato - Ministero delle Politiche Agricole, Alimentari e Forestali,  
463 Ufficio Territoriale per la Biodiversità di Verona, Centro Nazionale Biodiversità Forestale di Peri.  
464 Edward Grey Institute, University of Oxford; Andreas Wurm helped with interpreting the Ihrig  
465 1860 categories; prof. Alessandra Zanzi Sulli, sig. Alberto Pierguidi and Ilario Cavada for the  
466 analysis of seed data from the Paneveggio Forest and dott. Ilario Cavada for monitoring cone  
467 production. We thanks all unknown foresters who collected valuable masting data through time.

468

## 469 LITERATURE CITED

- 470 Ascoli, D., Vacchiano, G., Maringer, J., Bovio, G. and M. Conedera. 2015. The synchronicity of  
471 masting and intermediate severity fire effects favors beech recruitment. Forest Ecology and  
472 Management. 353: 126-135.
- 473 Bogdziewicz, M. and J. Szymkowiak. 2016. Oak acorn crop and Google search volume predict  
474 Lyme disease risk in temperate Europe. Basic and Applied Ecology. 17(4): 300-307.
- 475 Bohn, U., Neuhäusl, R., Gollub, G., Hettwer, C., Neuhäuslová, Z., Schlüter, H., and H. Weber.  
476 2003. Map of the Natural Vegetation of Europe, ed. Landwirtschaftsverlag, Münster.
- 477 Boutin, S., Wauters, L. A., McAdam, A. G., Humphries, M. M., Tosi, G., and A. A. Dhondt. 2006.  
478 Anticipatory reproduction and population growth in seed predators. Science. 314: 1928-1930.
- 479 Crone, E. E., and J. M. Rapp. 2014. Resource depletion, pollen coupling, and the ecology of mast  
480 seeding. Annals of the New York Academy of Sciences. 1322: 21-34.
- 481 Drobyshev, I., Niklasson, M., Mazerolle, M. J., and Y. Bergeron. 2014. Reconstruction of a 253-  
482 year long mast record of European beech reveals its association with large scale temperature  
483 variability and no long-term trend in mast frequencies. Agricultural and Forest Meteorology. 192:  
484 9-17.
- 485 Dengler, A. 1944. Waldbau auf ökologischer Grundlage, ed. Parey, Hamburg.
- 486 Hacket-Pain, A. J., Friend, A. D., Lageard, J. G., and P. A. Thomas. 2015. The influence of masting  
487 phenomenon on growth-climate relationships in trees: explaining the influence of previous  
488 summers' climate on ring width. Tree physiology. 35(3): 319-330.
- 489 Hase, W. 1964. Die Buchenmast in Schleswig-Holstein und ihre Abhängigkeit von der Witterung.  
490 Vol. 31. Mitteilungen des Deutschen Wetterdienst, Offenbach a.M.3
- 491 Hilton, G. M., and J. R. Packham. 2003. Variation in the masting of common beech (*Fagus*  
492 *sylvatica* L.) in northern Europe over two centuries (1800–2001). Forestry. 76(3): 319-328.
- 493 Kasprzyk, I., Ortył, B. and A. Dulska-Jeż. 2014. Relationships among weather parameters, airborne  
494 pollen and seed crops of *Fagus* and *Quercus* in Poland. Agricultural and Forest Meteorology. 197:  
495 111-122.

- 496 Kelly, D. and V. L. Sork. 2002. Mast seeding in perennial plants: why, how, where? Annual  
497 Review of Ecology and Systematics. 33: 427-447.
- 498 Kelly, D., Geldenhuys, A., James, A., Penelope Holland, E., Plank, M. J., Brockie, R. E., Cowan, P.  
499 E., Harper, G. A., Lee, W. G., Maitland, M. J., Mark, A. F., Mills, J. A., Wilson, P. R. and A. E.  
500 Byrom. 2013. Of mast and mean: differential-temperature cue makes mast seeding insensitive to  
501 climate change. Ecology Letters. 16(1): 90-98.
- 502 Koenig, W. D. and J. M. Knops. 1998. Scale of mast-seeding and tree-ring growth. Nature.  
503 396(6708): 225-226.
- 504 Koenig, W. D., and Knops, J. M. 2000. Patterns of annual seed production by northern hemisphere  
505 trees: a global perspective. The American Naturalist 155(1):59-69.
- 506 Koenig, W. and J. Knops. 2005. The Mystery of Masting in Trees Some trees reproduce  
507 synchronously over large areas, with widespread ecological effects, but how and why? American  
508 Scientist. 93(4): 340-347.
- 509 Koenig, W. D., Knops, J. M., Carmen, W. J., and I. S. Pearse. 2015. What drives masting? The  
510 phenological synchrony hypothesis. Ecology. 96(1): 184-192.
- 511 Herrera, C. M., Jordano, P., Guitián, J. and A. Traveset. 1998. Annual variability in seed production  
512 by woody plants and the masting concept: reassessment of principles and relationship to pollination  
513 and seed dispersal. The American Naturalist. 152(4):576-594.
- 514 Jenni, L. 1987. Mass concentration of Bramblings *Fringilla montifringilla* in Europe 1900–1983:  
515 Their dependence upon beech mast and the effect of snow cover. Ornis Scandinavica. 18: 84-94.
- 516 Maurer, E. 1964. Buchen-und Eichensamenjahre in Unterfranken während der letzten 100 Jahre.  
517 Allgemeine Forstzeitschrift. 31: 469-470.
- 518 Mencuccini, M., and P. Piussi. 1995. Production of seed and cones and consequences for radial  
519 increment in Norway spruce (*Picea abies* (L.) Karst.). Giornale Botanico Italiano. 129(3): 797-812.

- 520 Monks, A., Monks, J. M. and A. J. Tanentzap. 2016. Resource limitation underlying multiple  
521 masting models makes mast seeding sensitive to future climate change. *New Phytologist*. 210:419-  
522 430.
- 523 Nather, H. 1962. Waldsamen-Ernteaussichten für das Jahr 1962. *Allgemeine Forstzeitung*. 73: 17-  
524 18.
- 525 Nilsson, S. G. and U. Wastljung. 1987. Seed predation and cross-pollination in mast-seeding beech  
526 (*Fagus sylvatica*) patches. *Ecology*. 68(2): 260-265.
- 527 Norton, D. A. and D. Kelly. 1988. Mast seeding over 33 years by *Dacrydium cupressinum*  
528 Lamb.(rimu)(Podocarpaceae) in New Zealand: the importance of economies of scale. *Functional*  
529 *ecology*. 2: 399-408.
- 530 Pearse, I. S., Koenig, W. D. and J. M. Knops. 2014. Cues versus proximate drivers: testing the  
531 mechanism behind masting behavior. *Oikos*. 123(2): 179-184.
- 532 Pearse, I. S., Koenig, W. D., and D. Kelly. 2016. Mechanisms of mast seeding: resources, weather,  
533 cues, and selection. *New Phytologist* (in press).
- 534 Perrins, C. M. 1965. Population fluctuations and clutch-size in the Great Tit, *Parus major* L. *The*  
535 *Journal of Animal Ecology* 34(3): 601-647.
- 536 Pesendorfer, M. B., Koenig, W. D., Pearse, I. S., Knops, J. M. and K. A. Funk. 2016. Individual  
537 resource-limitation combined with population-wide pollen availability drives masting in the valley  
538 oak (*Quercus lobata*). *Journal of Ecology*. 104: 637-645.
- 539 Pidek, I. A., Svitavska-Svobodova, H., van der Knaap, W. O., Noryskiewicz, A. M., Filbrandt-  
540 Czaja, A., Noryskiewicz, B., Latalowa, M., Zimny, M., Swieta-Musznicka, J., Bozilova, E.,  
541 Tonkov, S., Filipova-Marinova, M., Poska, A., Giesecke, T. and A. Gikov. 2010. Variation in  
542 annual pollen accumulation rates of *Fagus* along a N-S transect in Europe based on pollen traps.  
543 *Vegetation History and Archaeobotany*. 19: 259-270.
- 544 R Development Core Team. 2016. *The R Project for Statistical Computing*. R Foundation for  
545 Statistical Computing, Vienna, Austria. URL: <http://www.R-project.org>.

- 546 Reil, D., Imholt, C., Eccard, J. A. and J. Jacob. 2015. Beech Fructification and Bank Vole
- 547 Population Dynamics-Combined Analyses of Promoters of Human Puumala Virus Infections in
- 548 Germany. *PloS One*. 10(7): e0134124.
- 549 Schauber, E. M., Kelly, D., Turchin, P., Simon, C., Lee, W. G., Allen, R. B., Payton, I. J., Wilson,
- 550 P. R., Cowan, P.E. and R. E. Brockie. 2002. Masting by eighteen New Zealand plant species: the
- 551 role of temperature as a synchronizing cue. *Ecology*. 83(5): 1214-1225.
- 552 Snell, R. S., Huth, A., Nabel, J. E. M. S., Bocedi, G., Travis, J. M. J., Gravel, D., Bugmann, H.,
- 553 Gutiérrez, A. G., Hickler, T., Higgins, S. I., reineking, B., Schwestjanoi, M., Zurbriggen, N. and H.
- 554 Lischke. 2014. Using dynamic vegetation models to simulate plant range shifts. *Ecography*. 37(12):
- 555 1184-1197.