


Please cite the Published Version

Luiselli, L, Dendi, D, Eniang, EA, Fakae, BB, Akani, GC and Fa, JE  (2017) State of knowledge of research in the Guinean forests of West Africa region. *Acta Oecologica*, 94. pp. 3-11. ISSN 1146-609X

DOI: <https://doi.org/10.1016/j.actao.2017.08.006>

Publisher: Elsevier

Version: Accepted Version

Downloaded from: <https://e-space.mmu.ac.uk/619127/>

Usage rights:  [Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0](#)

Additional Information: This is an Author Accepted Manuscript of a paper accepted for publication in *Acta Oecologica*, published by and copyright Elsevier.

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

ACCEPTED *Acta Oecologica*

**Understanding the influence of non-wealth factors in determining
bushmeat consumption: results from four West African countries**

Luca Luiselli^{a,b}, Emmanuel M. Hema^c, Gabriel Hoinssoudé Segniagbeto^d, Valy
Ouattara^e, Edem A. Eniang^f, Massimiliano Di Vittorio^g, Nioking Amadi^b, Gnoumou
Parfait^c, Nic Pacini^{h,i}, Godfrey C. Akani^b, Djidama Sirima^c, Wendengoudi Guenda^c,
Barineme B. Fakae^{a,b}, Daniele Dendi^{a,b}, John E. Fa^j

^a IDECC - Institute for Development, Ecology, Conservation and Cooperation, via G.
Tomasi di Lampedusa 33, I-00144 Rome, Italy.

^b Department of Applied and Environmental Biology, Rivers State University of
Science and Technology, P.M.B. 5080, Port Harcourt, Nigeria.

^c Université Ouaga 1 Professeur Joseph KI-ZERBO/CUPD, Laboratoire de Biologie
et Ecologie Animales, 09 B.P. 848 Ouagadougou 09 - Burkina Faso.

^d Department de Zoologie, Université de Lomé, Lomé, Togo.

^e Groupe des Expert en Gestion des Eléphants et de la Biodiversité de l'Afrique, de
l'Ouest, Ouagadougou, Burkina Faso.

^f Department of Forestry and Wildlife, University of Uyo, Akwa-Ibom State, Nigeria

^g Ecologia Applicata Italia s.r.l., via E. Jenner 50, Rome, Italy.

^h Department of Environmental and Chemical Engineering, University of Calabria,
Arcavacata di Rende (CS), Italy.

- 24 ⁱ Centre for Landscape and Climate Research, University of Leicester, UK.
- 25 ^j Division of Biology and Conservation Ecology, School of Science and the
- 26 Environment, Manchester Metropolitan University, Manchester M1 5GD, UK.

27 **Abstract**

28 The meat of wild animals (bushmeat) is consumed extensively in many tropical regions.
29 Over the past few decades bushmeat consumption has greatly increased, threatening the
30 survival of some hunted species and the supply of animal protein to countless numbers of
31 people. Understanding patterns of bushmeat consumption is thus vital to ensure the
32 sustainable use of this resource. Although the economic drivers of bushmeat consumption
33 have been well studied, non-wealth correlates have been poorly considered. Here, we
34 analyse how variables such as age and gender may influence bushmeat consumption in
35 four West African countries, within the Guinean forests (Togo and Nigeria) and Sahel
36 (Burkina Faso and Niger). We interviewed a total of 2,453 persons (1,253 urban, 1,200 in
37 rural areas) to determine frequency of consumption of bushmeat as well as the main
38 species eaten. We found significant differences in bushmeat consumption between rural
39 and urban areas in all four countries. In particular, the proportion of persons not
40 consuming any bushmeat was highest in urban areas. Gender differences in bushmeat
41 consumption were not generally important but young people consistently avoided eating
42 bushmeat, especially in Togo and Nigeria, and in urban areas. The complicated interplay
43 between tradition and evolution of social systems (especially the trends towards
44 westernization) may explain the different perceptions that people may have towards
45 consuming bushmeat in the four studied countries. In addition, we found considerable
46 variation in types of bushmeat eaten, with antelopes and large rodents eaten by the great
47 majority of interviewees, but bats, monkeys, and snakes being avoided, especially in urban
48 settlements.

49 *Key words:* Age; gender; Togo; Burkina Faso; Nigeria; Niger; wildlife; species eaten;
50 frequency.

51 **1. Introduction**

52 Terrestrial wild vertebrates are central to the nutritional wellbeing of many rural
53 people, particularly those inhabiting the world's tropical regions (Fa et al., 2002; Golden et
54 al., 2011). This reliance on wild meat is as much a consequence of the lack of alternative
55 domestic meat resources (Mainka & Trivedi 2002; Nasi et al. 2008), as much as it is an
56 attribute of centuries-old cultural traditions (Milner-Gulland et al., 2003). However,
57 although wild animals have been hunted for millennia, their consumption has greatly
58 increased over the past few decades (Maxwell et al. 2016). In West and Central Africa,
59 commercial hunting, especially to supply large urban centres, has risen dramatically, largely
60 driven by a human population growth of 2–3% per year (Nasi et al., 2011). Such
61 intensification of demand for bushmeat will have fatal consequences for many species but
62 particularly large-bodied and slow-growing species if extraction exceeds their replacement
63 rate (Wilkie et al., 2001). Indeed, the decline of some species as a consequence of
64 bushmeat extraction has already been documented for tortoises (Luiselli, 2003) and
65 antelopes (Fischer and Linsenmair, 2001; Grande-Vega et al., 2016; Hema et al., 2017). As a
66 consequence, loss of wildlife may threaten the food security of many marginalized forest
67 foragers, and farmer-forager communities that are isolated from markets and depend on
68 bushmeat as their primary protein source (Eves and Ruggiero, 2001).

69 Few studies have centred on understanding why people eat bushmeat. Knowing
70 what motivates people to eat bushmeat can help in developing politically acceptable ways
71 to manage wildlife hunting and trading with the aim of halting unsustainable exploitation.
72 Bushmeat may be eaten because it is cheaper or there are no alternatives available in the
73 market place (Apaza et al., 2002; Wilkie and Godoy, 2001), because consumers prefer the
74 taste of wildlife (Chardonnet et al., 1995; Trefon and de Maret, 1999) or to add variety to

75 the diet and for special social events and occasions (Njiforti, 1996). Despite this variety of
76 possible reasons that may motivate buyers to eat bushmeat, most studies have focused on
77 the socioeconomic background of consumers as the main reason underpinning their choice
78 (e.g. Wilkie and Godoy, 2001; Brashares et al., 2011). Findings show that wealthier
79 households consume more bushmeat in settlements nearer urban areas, but the opposite
80 pattern is typical of more isolated settlements (Brashares et al., 2011). Price and income
81 have significant roles in determining the level of consumption of bushmeat, fish, chicken,
82 and beef (Apaza et al., 2002; Wilkie and Godoy, 2001; Wilkie et al., 2005). Nonetheless, as
83 Brashares et al. (2011) has indicated, household wealth is only weakly linked to wildlife
84 consumption, and thus such a lack of a strong correlation could be explained by the
85 undisclosed importance of other factors. Few investigations have focused on how
86 bushmeat consumption may be affected by non-wealth factors such as age, gender and
87 geographical setting (Hema et al., 2017; Luiselli et al., 2017).

88 Most published studies reporting amounts of bushmeat consumed and preferences
89 have been based on data gathered within households (see Brashares et al. 2011). These
90 have been useful in determining possible socioeconomic characteristics of a community
91 that may be linked to bushmeat consumption patterns, but are generally costly in terms of
92 time and resources. Moreover, because households can include residents of different ages
93 and literacy - attributes generally linked to contrasting lifestyles and points of view
94 including the perception towards bushmeat consumption (Luiselli et al., 2017) - household
95 surveys are less suitable for exploring the influence of non-wealth factors. In this paper,
96 we use face-to-face interviews to examine the possible links between bushmeat
97 consumption frequency and types eaten relative to the age and gender of consumers as
98 well as the influence of the settlement type, ecological and country setting where they live.

We focus on a large sample of inhabitants of four West African countries found in Sahelian and Guinean forests environments.

2. Methods

2.1. Study sites

We interviewed a total of 2,453 individuals (1,253 urban, 1,200 in rural areas) from 27 separate human settlements in Nigeria, Togo, Burkina Faso and Niger (Fig. 1). Study localities in Nigeria and Togo were located within the Guinean Forests of West Africa (GFWA) region; swamp forest and moist rainforest vegetation zones in southern Nigeria (Niger Delta Environmental Survey, 1998; Oates et al., 2004) and in the deciduous moist forest zone of southwestern Togo (Ern, 1979). Sites in Burkina Faso and Niger were found within the Sahel, in Sudanian and Sahel Acacia savannahs (Thiombiano and Kampmann, 2010).

2.2. Interviews

To obtain information on bushmeat use, we conducted face-to-face interviews using a standardized questionnaire. All data were gathered during 2012-2016. We selected interviewees at marketplaces, roadsides, canteens, restaurants, hairdressing salons, food shops, and other gathering places. We stopped the first person encountered after a given time period (in minutes); the time interval randomly generated by a Random Number Generator. Local scientists (VO, NA, GP, DS, WG, EAE and other students) applied all interviews. All interviewees were informed of the aims of the project and their consent was obtained before proceeding. All interviews were conducted in the local language.

We interviewed persons in urban areas (Ouagadougou, Niamey, Lomé, Benin City, Port Harcourt and Calabar, all cities with more than 500,000 residents) as well as in smaller rural settlements (500 to 40,000 inhabitants). We recorded the interviewees' gender (male or female) and age (≤ 25 years, 26-50 years, ≥ 51 years) but not their names (St. John, 2010; Nuno et al., 2014; Luiselli et al., 2017). To avoid non-independence of the data, we never interviewed two persons of the same family or those living in the same house, even if they were not relatives (see also Hema et al., 2017, for similar procedure).

Interviewees were asked the following two main questions: (1) Do you like eating bushmeat? (2) If yes, how often do you eat bushmeat? Interviewees would then be asked if they ate bushmeat frequently (at least once a week), rarely (about once per month or less) or never. Persons who answered that they consumed bushmeat only occasionally were then asked whether they selected the type of animal or whether they would just buy/eat whatever kind of bushmeat was available.

2.3. Statistical analyses

We employed Generalized Linear Models (GLZs) to determine the relationship between bushmeat consumption frequency and site (rural versus urban), gender (male/female) and age classes (three categories) (Hosmer and Lemeshow, 2000). The codes for the variables used in the GLZs are given in Appendix 1. In the model, the response "never eat bushmeat" was the dependent variable (i.e. consumption data were converted into a binary variable, 1 = eat (often or rarely) and 0 = never eat bushmeat) and the identity of the link function and a normal distribution of error were used (McCullagh and Nelder, 1989). Three age categories were used for all analyses: persons aged less than 25, aged less than 50, and aged 51 years or more. In the GLZ models, a stepwise forward regression

procedure was used to test the statistical significance of each variable in turn, and variables were excluded when they did not correlate significantly with the dependent variable (Wald test $P > 0.05$).

To explore deviance and hierarchical partitioning, the selected variables were analyzed in order to determine the comparative influence of each variable (Borcard et al., 1992). The decomposition of the variation into subsets of explanatory variables was carried out by means of a partial regression analysis (Legendre and Legendre, 1998).

Frequency differences between groups of interviewed people were analyzed using the χ^2 test, for comparing both differences among frequently-eating, rarely-eating and non-eating bushmeat respondents, and for determining differences in terms of type of bushmeat eaten. The statistical software PASW 11.0 was used for all analyses, and alpha was set at 5%.

3. Results

3.1. General patterns

A summary of the data gathered for this study is shown in Table 1, the raw dataset is given in Appendix 2. In general terms, bushmeat was consumed more often by rural than urban interviewees in all countries (Fig. 2). An average total of $70.3 \pm 15.7\%$ of rural respondents answered that they ate bushmeat (either eaten rarely or often) in contrast to only $42.8 \pm 19.0\%$ of urban interviewees. In all countries more rural than urban respondents ate bushmeat; 1.59 times more in Niger, 1.26 times more in Nigeria, 0.46 times in Togo and 0.14 times in Burkina Faso.

A general GLZ model using data from all countries pooled and type of bushmeat eaten as the dependent variable showed that, the probability of eating ungulates or birds

was significantly affected by gender or age of the respondents respectively, while the eating of monkeys, bats, carnivores, crocodiles, snakes and turtles was influenced by the age of the respondents and their urban/rural location (Table 2).

Significant differences in responses between interviewees in Guinean forests and the Sahel region appeared. Age classes, followed by urban/rural location, accounted for the strongest pure effect in the Sahelian localities with gender explaining only a very small proportion of the variance (Fig. 3). Within the Guinean forest localities, urban/rural location was the predominant effect, age had a lesser relevance in terms of the explained variance, but gender had almost no effect (Fig. 3).

We found a significant effect of distance (in km) of the interviewee to the nearest urban area; the probability of never-eating bushmeat increased in Sahelian countries, but not in Guinean forest region countries (for Sahel: GLZ estimate = 6.56, standard error = 1.34, Wald = 24.0, $P < 0.0001$; for age classes: estimate = -7.62, standard error = 2.32, Wald = 10.79, $P < 0.001$; for Guinean forests: in all cases $P > 0.165$).

3.2. Country effects

Our GLZ model revealed that effect of country on bushmeat consumed were relatively minor (Table 2). Nonetheless, country had a statistical effect on the consumption of primates, with people from the Guinean Forests countries being more likely to eat monkeys than those in the Sahel (Table 2). Thus, apart from primates, there were no other statistical differences found between Guinean forest and Sahelian country sites in terms of the probability of consuming other types of bushmeat.

Overall, there were no significant differences between countries (in all cases, at least $P > 0.225$ at χ^2 test) in the proportion of respondents who declared that they never

ate bushmeat (Table 3) as well as in those that declared frequently eating bushmeat (Table 4). However, there were clear confounding effects of age, gender and urban/rural location on the pure effect of the country (see below). Overall, patterns for the frequency of 'often-eaten-bushmeat' responses were more consistent among countries than in the 'never-eating-bushmeat' answers (Table 4).

In Togo, there was a significant effect of age in urban and rural areas; the frequency of respondents never-eating bushmeat declined significantly with age in both locations (Table 3). No effect of gender was found, but the differences between rural and urban areas depended on the strength of the frequency decreases of never-eating-bushmeat respondents in these two locations, i.e. rural and urban people in Togo tended to respond similarly. In Nigeria (Table 3), there was no effect of age in urban areas (people do not eat bushmeat in general) but in rural areas (only young people did not eat bushmeat). In addition, there was a significant effect of gender in urban areas, with women avoiding eating bushmeat more than men. The overall differences between rural and urban areas were significant for both gender and age (Tables 3 and 4). In Burkina Faso, there was a significant effect of age in urban areas (more young people did not eat bushmeat) but not for rural areas, where people do generally eat bushmeat independent of their age (Tables 3 and 4). In Niger, there was only a significant effect of age, with more young people responding that they would never eat bushmeat compared to older people, in both urban and rural locations (Table 3).

3.3. *Age effects*

Our GLZ model revealed that the age of the interviewees affected the probability of consuming primates, bats, carnivores, crocodiles, snakes and chelonians, in all cases older people were more likely to consume these animals than younger people (Table 2).

Overall, age had a significantly stronger effect on the likelihood of consuming bushmeat in the Sahelian region compared to the Guinean forest region (Fig. 3). Nonetheless, the tendency was the same in both regions: young people tended to never or very rarely consume bushmeat significantly more than people of >25 years age ($P < 0.001$ at χ^2 test).

3.4. *Gender effects*

Overall, gender effects were negligible in both Sahelian and Guinean forests regions, and contributed little to the hierarchical variance partitioning in the interview dataset (Figure 3). Nonetheless, some effects of gender were detected in the attitude of consuming a few types of bushmeat as well as in a few local contexts. Indeed, although most people ate ungulates and rodents, there were significant effects of gender on the consumption of these animals, with men being more likely to eat them than women (Table 2). In addition, females tended to avoid eating bushmeat more frequently than males in some countries such as Nigeria. However, this was not a pure gender effect, as it was mediated by age and rural/urban condition in a rather complicated way (Tables 3 and 4). Overall, the 'often-eating-bushmeat' response was especially linked to men in either Guinean forests (e.g. Togo) or Sahel (e.g. Burkina Faso) regions.

3.5. *Rural versus urban*

Whether living in rural or urban locations determined the outcome of the interviewees' answers in the Guinean forest region but not in the Sahelian region (Fig. 3). In other words, attitude towards bushmeat of people from Sahelian regions was similar in both rural and urban locations, whereas in the Guinean forest region there were differences between locations. In addition, in terms of frequency of never-eating bushmeat people, statistical differences between rural versus urban conditions were much higher ($P < 0.001$ at χ^2 test) than those occurring between countries (see above). Whether a person lived in an urban or rural location affected the probability of consuming bushmeat much more than their country of residence.

A total of 41.9% of urban and 67.3% of rural respondents stated they consumed bushmeat (Fig. 4); this difference being significant ($\chi^2 = 231.9$, $df = 2$, $P < 0.0001$). According to the different response categories, most interviewees in rural areas mentioned they frequently ate bushmeat ($\chi^2 = 7.3$, $df = 2$, $P < 0.05$), but in urban areas most said they never ate bushmeat ($\chi^2 = 193.4$, $df = 2$, $P < 0.0001$).

Overall, ungulates and rodents were eaten by almost all respondents in either rural or urban areas, but carnivores, monkeys and snakes were eaten rarely (differences significant at $P < 0.00001$ compared to ungulates and rodents, χ^2 test), and mainly in rural areas (Fig. 5). Contingency table analysis showed that there were no significant differences between urban and rural areas in terms of frequency of respondents eating the various bushmeat types ($\chi^2 = 14.48$, $df = 8$, $P = 0.0699$). However, our GLZ model revealed that primates, bats, carnivores, crocodiles, snakes and chelonians were significantly more likely to be eaten by rural than by urban people, with the highest estimates being for monkeys

and bats (Table 2); it was unlikely that people from urban areas, in any of the surveyed countries, ate monkeys and bats.

Differences between urban and rural areas were also strongly mediated by the effects of age and gender (Tables 3 and 4). Overall, there were significant differences in both gender and age between rural and urban areas.

4. Discussion

Previous studies have suggested that bushmeat was universally preferred “due to its superior taste” (King, 1994) and thus African communities therefore preferred and thus primarily ate bushmeat. These statements were not based on empirical evidence until a study reporting on two-choice taste tests showed that consumers in Gabon had only a weak preference for bushmeat and only rural consumers consistently preferred bushmeat over alternatives (Schenck et al., 2006). This result is particularly important given that it manifests that even though basic desires such as hunger and the need for nourishment can influence food choice, availability and cultural norms also affect these. Thus, it is not simply taste that is driving demand for bushmeat, but that price or other culturally mediated factors such as familiarity, tradition, and prestige play a role.

The use of face-to-face interviews for assessing bushmeat consumption has been used in some studies e.g. in a recent one in Liberia by Ordaz-Németh et al. (2017). Like in our study, Ordaz-Németh et al. (2017) did not test for any potential bias in responses by interviewees not responding truthfully. In both studies, hunting is not illegal per se and we did not mention explicitly any particular species in the interview form, but instead used open questions so that the interviewee could respond without any prompting e.g. by naming protected species.

288 Our analyses indicate a very clear and significant difference in bushmeat
 289 consumption among rural and urban peoples in all countries. This effect appeared in 7 out
 290 of 7 models, in all four of the investigated countries. This difference has been
 291 demonstrated in a number of other studies in the African continent (e.g. in the Democratic
 292 Republic of Congo, see Van Vliet et al., 2014) and in Madagascar (Jenkins et al., 2011). This
 293 contrast between rural and urban dwellers is largely explained by the availability of
 294 bushmeat versus alternative protein sources. Rural dwellers are usually restricted in terms
 295 of the availability and accessibility of domestic meats but in a much better position to
 296 option these resources from the wild. By contrast, urban dwellers have greater access to
 297 alternative proteins (Apaza et al., 2002). Nonetheless, cultural complications also explain
 298 the preference of non-bushmeat proteins by urban people (see Luiselli et al., 2017 and
 299 below).

300 Our analyses clearly showed that age was important in most countries; the pure
 301 effect of age was significant in 5 out of 7 models, in Nigeria, Togo and Niger. Younger
 302 interviewees generally ate less bushmeat than older persons. That young people ate less
 303 bushmeat can in part be due to a growing 'westernization' of the lifestyles, especially
 304 among the middle classes. These sector of the community often do not see it as 'socially
 305 acceptable' to consume bushmeat, since this is perceived by them as a sign of 'being very
 306 local' (i.e. not culturally advanced). In contrast the eating of 'fast foods' (hamburgers, pizza,
 307 kebab, etc.) is now the favourite 'social diet' of young people. This pattern is especially
 308 evident in urban Nigeria and Togo (our unpublished observations), where young
 309 interviewees not only declared that they would not eat bushmeat, but even commented
 310 that eating bushmeat was not acceptable because it produces a loss of personal prestige
 311 within their circle of friends. In this regard, it was particularly interesting that, among the

rarely-eating bushmeat urban people, a sample of 7 young (<25 years) persons from Togo and 15 from Nigeria declared that they would never eat bushmeat in public, but that very occasionally they do during private family events, and only when they visit their rural relatives. Thus, among the respondents who declared that they rarely ate bushmeat, many would only consume bushmeat for special occasions. We suggest that in urban areas the lower consumption of bushmeat is not because of lack of access, but that it responds to a more culturally-driven avoidance in response to the changing socio-economic context.

By contrast to the effects of age we observed in our study, the pure effect of gender was only apparent in 1 out of 7 models. In terms of mixed factors, 'Gender X Rural/Urban' were significant in 4/7 models and 'Age X Rural/Urban' were significant in 5/7 models, whereas 'Gender X Age' in 2/7 models. From these results, we conclude that rural/urban and age are much more important than gender in determining the probability for a person to consume bushmeat. The non-effect of gender is probably related to the enhanced equal rights of women and men in West African societies (especially in Nigeria), with young generations being much more equal in terms of gender and lifestyle (see Gender Equality Index database by the African Development Bank, available at www.afdb.org). Thus, since young men and women typically share a similar life-style (especially in urban areas), even their food preferences tend to be very similar.

Since, in all countries, and in urban areas in particular, most of the young respondents stated they never ate bushmeat, this would suggest that bushmeat consumption has been substantially decreasing among the new generations of West Africans, independently on their local culture, religion, ethnicity and level of human development. In Nigeria, where the level of human development (average wealth and scholarization standards) are clearly higher than in the other countries (the country being

the 22th economy of the world; World Bank, 2016), only older people in rural areas (age > 51 years) answered that they consumed bushmeat more regularly (Table 2).

Although our study is the first to cover a broad spectrum of situations, it is important to note that bushmeat trade analysis are much easier to undertake in the Guinean forests region (such as Ivory Coast, Ghana, Nigeria; e.g. see Fa et al., 2002a, 2002b, 2006) than in Sahel. This difference is related to the fact that in the Sahel region there are no open bushmeat markets and people here may be more reluctant to answer interviewers openly because of social norms (Hema et al., 2017, but see Lindsey et al. 2013). This is also possibly linked to the fact that forest can occur close to urban areas in the Guinean Forest region (e.g., Niger Delta forests surrounding Port Harcourt), whereas the same is not true in the Sahel where all the forested or mature savannah sites (from which most of the bushmeat trade does originate) are situated far from larger urban centres (our unpublished observations). Therefore, ‘hub’ markets (Akani et al., 2015) are more likely to be found nearby large cities in the forest zone than in the savannah zone. In conclusion, we argue that the cultural drivers of wildlife use are crucial to take into account when seeking long-term sustainability solutions of wildlife resource extraction (e.g., Luiselli et al., 2017).

Acknowledgements. We are thankful to several organizations, and in particular to the Rivers State University of Science and Technology (Port Harcourt), IDECC – Institute for Development, Ecology, Conservation and Cooperation (Rome), Mohamed Bin Zayed Species Conservation Fund, Conservation International, Turtle Conservation Fund, Andrew Sabin & Family Foundation, T.S.K.J. Nigeria Ltd., IUCN/SSC TFTSG, Aquater s.p.a. and Snamprogetti s.p.a. for having funded our field projects in the studied countries.

References

- Akani, G.C., Amadi, N., Eniang, E.A., Luiselli, L. & Petrozzi, F., 2015. Are mammal communities occurring at a regional scale reliably represented in “hub” bushmeat markets? A case study with Bayelsa State (Niger Delta, Nigeria). *Folia Zoologica*, 64, 79–86.
- Borcard, D., Legendre, P., Drapeau, P., 1992. Partialling out the spatial component of ecological variation. *Ecology*, 73, 1045-1055.
- Brashares, J.S., Golden, C.D., Weinbaum, K.Z., Barrett, C.B., Okello, G.V., 2011. Economic and geographic drivers of wildlife consumption in rural Africa. *Proc. Natl. Acad. Sci. U.S.A.* 108, 13931–13936.
- Ern, H., 1979. Vegetation Togos. Gliederung, Gefährdung, Erhaltung. *Willdenowia* 9, 295-312.
- Fa, J.E., Peres, C., Meeuwig, J., 2002a. Bushmeat exploitation in tropical forests: an intercontinental comparison. *Conserv. Biol.* 16, 232-237.
- Fa, J.E., Seymour, S., Dupain, J., Amin, R., Albrechtsen, L., Macdonald, D., 2006. Getting to grips with the magnitude of exploitation: Bushmeat in the Cross-Sanaga rivers region, Nigeria and Cameroon. *Biol. Conserv.* 129, 497–510.
- Fa, J.E., Juste, J., Burn, R.W., Broad, G., 2002b. Bushmeat consumption and preferences of two ethnic groups in Bioko Island, West Africa. *Human Ecol.* 30, 397–416.
- Fischer, F., Linsenmair, K.E., 2001. Decreases in ungulate population densities. Examples from the Comoé National Park, Ivory Coast. *Biol. Conserv.* 101, 131–135.

- 381 Golden, C.D., Fernald, L.C.H., Brashares, J.S., Rasolofoniaina, B. J. R., Kremen, C., 2011.
 382 Benefits of wildlife consumption to child nutrition in a biodiversity hotspot. *Proc. Natl.*
 383 *Acad. Sci. USA* 108, 19653–19656.
- 384 Grande-Vega, M., Farfan, M.A., Ondo, A., Fa, J.E., 2016. Decline in hunter offtake of blue
 385 duikers in Bioko Island, Equatorial Guinea. *Afr. J. Ecol.* 54, 49–58.
- 386 Hema, E.M., Ouattara, V., Parfait, G., Di Vittorio, M., Sirima, D., Dendi, D., Guenda, W.,
 387 Petrozzi, F., Luiselli, L., 2017. Bushmeat consumption in the West African Sahel of
 388 Burkina Faso based on interview campaigns, and the decline of some consumed
 389 species. *Oryx*, <https://doi.org/10.1017/S0030605316001721>.
- 390 Hosmer, D.W., Lemeshow, S., 2000. *Applied logistic regression analysis*. 2nd ed. John Wiley
 391 and Sons, New York.
- 392 Jenkins, R.K., Keane, A., Rakotoarivelo, A.R., Rakotomboavonjy, V., Randrianandrianina,
 393 F.H., Razafimanahaka, H. et al., 2011. Analysis of patterns of bushmeat consumption
 394 reveals extensive exploitation of protected species in Eastern Madagascar. *PLoS*
 395 *ONE* 6(12), e27570.
- 396 Legendre, P., Legendre, L., 1998. *Numerical ecology*. Elsevier Science. Amsterdam,
 397 Netherlands.
- 398 Lindsey, P., Balme, G., Becker, M., Begg, C., Bento, C., Bocchino, C., Dickman, A., Diggle, R.,
 399 Eves, H., Henschel, P., Lewis, D., Marnewick, K., Mattheus, J., McNutt, J.W.,
 400 McRobb, R., Midlane, N., Milanzi, J., Morley, R., Murphree, M., Nyoni, P., Opyene,
 401 V., Phadima, J., Purchase, N., Rentsch, D., Roche, C., Shaw, J., van der Westhuizen,
 402 H., Van Vliet, N., Zisadza, P., 2012. Illegal hunting and the bush-meat trade in
 403 savanna Africa: drivers, impacts and solutions to address the problem.

- 404 Panthera/Zoological Society of London/Wildlife Conservation Society report, New
405 York.
- 406 Luiselli, L., Petrozzi, F., Akani, G.C., Di Vittorio, M., Amadi, N., Ebere, N., Dendi, D., Amori,
407 G., Eniang, E.A., 2017. Rehashing bushmeat –interview campaigns reveal some
408 controversial issues about the bushmeat trade dynamics in Nigeria. *Rev. Ecol. (Terre*
409 *et Vie)* 72, 3-18.
- 410 Mainka, S. A., Trivedi, M., 2002. Links between biodiversity conservation, livelihoods and
411 food security: the sustainable use of wild species for meat. IUCN, Gland,
412 Switzerland.
- 413 Maxwell, S.L., Fuller, R.A., Brooks T.M., Watson J.E.M., 2016. Biodiversity: The ravages of
414 guns, nets and bulldozers. *Nature* 536:143-145.
- 415 McCullagh, P., Nelder, J.A., 1989. Generalized linear models. Chapman and Hall/CRC,
416 London, United Kingdom.
- 417 Milner-Gulland, E.J., Bennett, E.L., the SCB 2002 Annual Meeting Wild Meat Group, 2003.
418 Wild meat: the bigger picture. *Tr. Ecol. Evol.* 18, 351-357.
- 419 Nasi, R., Brown, D., Wilkie, D., Bennett, E., Tutin, C., van Tol, G., Christophersen, T., 2008.
420 Conservation and use of wildlife-based resources: the bushmeat crisis. Technical
421 Series no. 33, Secretariat of the Convention on Biological Diversity, Montreal, and
422 Center for International Forestry Research (CIFOR), Bogor.
- 423 Nasi, R., Taber, A., Van Vliet, N., 2011. Empty forests, empty stomachs? Bushmeat and
424 livelihoods in the Congo and Amazon Basins. *Internat. For. Rev.* 13, 355-368.

- 425 Niger Delta Environmental Survey (NDES), 1998. Environmental and Socio-Economic
426 characteristics, volume 1. NDES, Port Harcourt, Nigeria.
- 427 Nuno, A., Bunnefeld, N., Naiman, L.C., Milner-Gulland, E.J., 2014. Novel Approach to
428 Assessing the Prevalence and Drivers of Illegal Bushmeat Hunting in the Serengeti.
429 *Conserv. Biol.* 27, 1355–1365.
- 430 Oates, J.F., Bergl, R.A., Linder, J.M., 2004. Africa's Gulf of Guinea forests: biodiversity
431 patterns and conservation priorities. *Adv. Appl. Biodiv. Sci.* 6, 1–91.
- 432 Ordaz-Németh, I., Arandjelovic, M., Boesch, L., Gatiso, T., Grimes, T., Kuehl, H.S., Lormie,
433 M., Stephens, C., Tweh, C., Junker, J., 2017. The socio-economic drivers of bushmeat
434 consumption during the West African Ebola crisis. *PLoS Negl Trop Dis* 11(3):
435 e0005450. <https://doi.org/10.1371/journal.pntd.0005450>
- 436 St John, F.A.V., Gibbons, J.M., Edwards-Jones, G., 2010. Testing novel methods for
437 assessing rule breaking in conservation. *Biol. Conserv.* 143, 1025–1030.
- 438 Thiombiano, A., Kampmann, D., 2010. Atlas de la biodiversité de l'Afrique de l'Ouest. Tome
439 II. Burkina Faso. Ouagadougou and Frankfurt/Main.
- 440 Wilkie, D. S., Godoy, R.A., 2001. Income and price elasticities of bushmeat demand in
441 lowland Amerindian societies. *Conserv. Biol.* 15, 1–9.
- 442 World Bank, 2016. Statistics: countries and economies. World Bank, New York.

443 Table 1. Synopsis of the interview raw data collected during the present surveys in the four studied countries.

	Urban			Total urban	Rural			Total rural
	Often eaten	Rarely eaten	Never eaten		Often eaten	Rarely eaten	Never eaten	
Burkina Faso								
Males (< 25 yr)	0	0	7	7	4	2	2	8
Males (< 50 yr)	7	69	43	119	66	24	7	97
Males (> 51)	7	12	1	20	17	9	3	29
Females (< 25 yr)	2	1	10	13	1	1	21	23
Females (< 50 yr)	6	52	30	88	17	21	33	71
Females (> 51)	1	8	6	15	9	5	3	17
TOTAL SAMPLE	23	142	97	262	114	62	69	245
Niger								
Males (< 25 yr)	2	2	56	60	20	11	45	76
Males (< 50 yr)	4	6	32	42	30	20	39	89
Males (> 51)	5	6	22	33	33	9	37	79
Females (< 25 yr)	1	0	46	47	14	9	44	67
Females (< 50 yr)	4	7	39	50	24	11	30	65
Females (> 51)	7	7	26	40	22	10	31	63
TOTAL SAMPLE	23	28	221	272	143	70	226	439
Togo								
Males (< 25 yr)	11	9	33	53	14	8	21	43
Males (< 50 yr)	12	16	15	43	33	24	2	59
Males (> 51)	14	12	5	31	24	7	1	32
Females (< 25 yr)	0	11	41	52	4	16	26	46
Females (< 50 yr)	7	17	23	47	14	7	4	25
Females (> 51)	16	11	11	38	16	2	2	20
TOTAL SAMPLE	60	76	128	264	105	64	56	225
Nigeria								
Males (< 25 yr)	7	14	56	77	17	31	11	59
Males (< 50 yr)	12	23	44	79	21	23	8	52
Males (> 51)	16	31	39	86	22	41	5	68

Females (< 25 yr)	3	6	62	71	13	43	14	70
Females (< 50 yr)	7	12	46	65	9	11	2	22
Females (> 51)	19	23	35	77	14	5	1	20
TOTAL SAMPLE	64	109	282	455	96	154	41	291
GRAND TOTAL	170	355	728	1253	458	350	392	1200

444 Table 2. Results of the Generalized Linear Model on the probability of eating bushmeat by type of
 445 animals by country, urban/rural locality, age, sex and gender (female/male). Intercepts are included
 446 in all models, and the explained deviance (in %) is also shown. Negative estimates for gender
 447 means a preponderance of male respondents. Positive estimates for age indicate a preponderance
 448 of older age classes respondents.

Variable	Estimate	St. error	Wald	P
Ungulates				
Intercept	211.71	72.41	8.55	0.003
Gender	-1.68	0.82	4.18	0.041
Explained deviance (%)	90.20			
Rodents				
Intercept	295.85	202.43	2.14	0.144
Gender	-5.14	2.29	5.02	0.025
Explained deviance (%)	88.08			
Monkeys				
Intercept	-2079.08	431.87	23.17	0.000001
Country	5.93	2.99	3.92	0.048
Urban/Rural	-31.26	4.89	40.89	0.000001
Age	14.92	2.99	24.85	0.000001
Explained deviance (%)	34.07			
Bats				
Intercept	-1596.00	376.76	17.94	0.000023
Urban/Rural	-29.26	6.03	23.54	0.000001
Age	16.13	3.69	19.08	0.000013
Explained deviance (%)	45.79			
Carnivores				
Intercept	-1408.96	335.10	17.68	0.000026
Urban/Rural	-17.45	5.36	10.58	0.0011
Age	14.22	3.28	18.74	0.000015
Explained deviance (%)	55.90			
Birds				

Intercept	-837.82	493.40	2.88	0.089
Age	7.33	3.42	4.60	0.032
Explained deviance (%)	88.74			

Crocodiles

Intercept	-1439.25	453.05	10.09	0.0015
Urban/Rural	-22.29	5.13	18.89	0.000014
Age	15.28	3.14	23.68	0.000001
Explained deviance (%)	46.41			

Snakes

Intercept	-1330.51	345.56	14.82	0.000118
Urban/Rural	-20.61	5.53	13.87	0.000195
Age	13.41	3.39	15.66	0.000076
Explained deviance (%)	55.05			

Turtles

Intercept	-853.91	296.62	8.29	0.0039
Urban/Rural	-15.25	4.75	10.31	0.0013
Age	9.156	2.9079	9.91504	0.001639
Explained deviance (%)	64.03			

449

450

451 Table 3. Summary of the results of contingency tables on the frequencies of the never-eating bushmeat respondents by country. In this table,
 452 'towards' would indicate the direction of the significant effect. For instance, if in a given area, there was a significantly higher number of 'never-
 453 eating-bushmeat' respondents for young people (< 25 years age), this is highlighted in the table with 'towards young'.

	Differences between gender in rural	Differences between gender in urban	Differences by age in rural	Differences by age in urban	Differences between urban and rural by gender	Differences between urban and rural by age
Togo	P = n.s.	P = n.s.	P < 0.01 (towards young) P < 0.01 (towards old people)	P < 0.01 (towards young)	P = n.s. P < 0.01 (due to men in urban areas)	P = n.s.
Nigeria	P = n.s.	P < 0.05 (towards men)	P = n.s.	P = n.s.	P < 0.0001 (due to opposite signs of differences)	P < 0.05 (due to age in rural areas)
Burkina Faso	P < 0.01 (towards men)	P < 0.05 (towards women)	P < 0.05 (towards young)	P < 0.05 (towards young)		P < 0.05 (due to consistent trends of age: young do not eat bushmeat)
Niger	P = n.s.	P = n.s.	P < 0.05 (towards young)	P < 0.05 (towards young)	P = n.s.	P = n.s.

454


455

456 Table 4. Summary of the results of contingency tables on the frequencies of the often-eating bushmeat respondents by country. In this table,
 457 'towards' would indicate the direction of the significant effect. For instance, if in a given area, there was a significantly higher number of 'never-
 458 eating-bushmeat' respondents for young people (< 25 years age), this is highlighted in the table with 'towards young'.

	Differences between gender in rural	Differences between gender in urban	Differences by age in rural	Differences by age in urban	Differences between urban and rural by gender	Differences between urban and rural by age
Togo	P < 0.05 (towards men)	P < 0.05 (towards men)	P < 0.05 (towards young)	P < 0.01 (towards young)	P = n.s.	P = n.s.
Nigeria	P = n.s.	P = n.s.	P = n.s.	P < 0.01 (towards young)	P = n.s.	P < 0.05 (due to young people responses negative effect)
Burkina Faso	P < 0.001 (towards men)	P < 0.001 (towards men)	P = n.s.	P < 0.0001 (towards young)	P < 0.05 (due to men)	P < 0.05 (due to young people responses negative effect)
Niger	P = n.s.	P = n.s.	P = n.s.	P < 0.05 (towards young)	P = n.s.	P = n.s.

459 Figure 1. Map of West Africa showing the study sites where interviews were carried out

460




461


462

463 Figure 2. Percent interviewees responding whether bushmeat was eaten often, rarely or never in
 464 urban and rural settlements in the four countries studied in West Africa.

465




466 Figure 3. Relative importance of predictors (pure effect), as determined by hierarchical variation
467 partitioning, for the model considering all the interviewees' responses as dependent variable, for
468 the Sahel countries (upper graphic) and for the Guinean forests countries (lower graphic). Spatial =
469 urban/rural.



473 Figure 4. Distribution of the various types of answer by respondents in urban versus rural areas in
474 the four studied countries of West Africa as for whether they would eat bushmeat often, rarely or
475 never. All data from the different countries were pooled for this graphic


476

477



478 Figure 5. Distribution of the various types of answer by respondents in urban versus rural areas in
479 the four studied countries of West Africa as for the type of consumed bushmeat is concerned. All
480 data from the different countries were pooled for this graphic

481



482

483 Appendix 1. Codes for the variables used in the Generalized Linear Models (GLZs).

Country	Class	Locality	Cod Loc	Cod age	Cod sex	Never eat bushmeat
Burkina Faso	Males (< 25 yr)	urban	1	g	1	7
Burkina Faso	Males (< 50 yr)	urban	1	m	1	43
Burkina Faso	Males (> 51)	urban	1	a	1	1
Burkina Faso	Females (< 25 yr)	urban	1	g	0	10
Burkina Faso	Females (< 50 yr)	urban	1	m	0	30
Burkina Faso	Females (> 51)	urban	1	a	0	6
Burkina Faso	Males (< 25 yr)	rural	0	g	1	2
Burkina Faso	Males (< 50 yr)	rural	0	m	1	7
Burkina Faso	Males (> 51)	rural	0	a	1	3
Burkina Faso	Females (< 25 yr)	rural	0	g	0	21
Burkina Faso	Females (< 50 yr)	rural	0	m	0	33
Burkina Faso	Females (> 51)	rural	0	a	0	3
Nigeria	Males (< 25 yr)	urban	1	g	1	77
Nigeria	Males (< 50 yr)	urban	1	m	1	79
Nigeria	Males (> 51)	urban	1	a	1	86
Nigeria	Females (< 25 yr)	urban	1	g	0	71
Nigeria	Females (< 50 yr)	urban	1	m	0	65
Nigeria	Females (> 51)	urban	1	a	0	77
Nigeria	Males (< 25 yr)	rural	0	g	1	59
Nigeria	Males (< 50 yr)	rural	0	m	1	52
Nigeria	Males (> 51)	rural	0	a	1	68
Nigeria	Females (< 25 yr)	rural	0	g	0	70
Nigeria	Females (< 50 yr)	rural	0	m	0	22
Nigeria	Females (> 51)	rural	0	a	0	20
Niger	Males (< 25 yr)	urban	1	g	1	4
Niger	Males (< 50 yr)	urban	1	m	1	4
Niger	Males (> 51)	urban	1	a	1	7
Niger	Females (< 25 yr)	urban	1	g	0	1
Niger	Females (< 50 yr)	urban	1	m	0	9
Niger	Females (> 51)	urban	1	a	0	6
Niger	Males (< 25 yr)	rural	0	g	1	6
Niger	Males (< 50 yr)	rural	0	m	1	5
Niger	Males (> 51)	rural	0	a	1	9
Niger	Females (< 25 yr)	rural	0	g	0	3
Niger	Females (< 50 yr)	rural	0	m	0	7
Niger	Females (> 51)	rural	0	a	0	9
Togo	Males (< 25 yr)	urban	1	g	1	33
Togo	Males (< 50 yr)	urban	1	m	1	15
Togo	Males (> 51)	urban	1	a	1	5
Togo	Females (< 25 yr)	urban	1	g	0	41
Togo	Females (< 50 yr)	urban	1	m	0	23
Togo	Females (> 51)	urban	1	a	0	11
Togo	Males (< 25 yr)	rural	0	g	1	21
Togo	Males (< 50 yr)	rural	0	m	1	2
Togo	Males (> 51)	rural	0	a	1	1
Togo	Females (< 25 yr)	rural	0	g	0	26
Togo	Females (< 50 yr)	rural	0	m	0	4
Togo	Females (> 51)	rural	0	a	0	2

484

485

486 Appendix 2. Summary of the raw data on the types of eaten bushmeat by people in the investigated countries.

		Males (< 25 yr)	Males (< 50 yr)	Males (> 51)	Females (< 25 yr)	Females (< 50 yr)	Females (> 51)	Total interviewees
Nigeria		21	35	47	9	19	42	173
	ungulates	21	34	45	9	19	40	
	rodents	21	31	41	9	19	39	
Urban	monkeys	2	7	11	0	3	6	
	bats	3	7	8	1	3	11	
	carnivores	2	10	15	0	2	5	
	birds	15	26	33	5	15	28	
	crocodiles	8	15	27	1	5	9	
	snakes	1	3	5	0	0	3	
	turtles	16	22	26	6	14	32	
Nigeria		48	44	63	56	20	19	250
	ungulates	48	43	61	56	20	19	
	rodents	48	41	55	56	18	19	
Rural	monkeys	6	23	31	3	3	11	
	bats	7	17	22	7	5	13	
	carnivores	9	14	20	21	6	11	
	birds	39	40	55	45	17	17	
	crocodiles	23	21	38	32	9	17	
	snakes	2	8	19	4	6	9	
	turtles	44	33	51	36	16	19	
Togo		20	28	26	11	23	27	136
	ungulates	19	26	25	11	22	25	
	rodents	13	23	24	11	22	23	
Urban	monkeys	4	6	6	0	1	4	
	bats	4	9	6	0	1	3	
	carnivores	8	6	6	0	2	8	
	birds	15	21	21	10	17	23	
	crocodiles	3	18	8	3	6	11	
	snakes	2	4	4	0	1	6	

Togo	turtles	9	16	16	5	15	24	169
		22	57	31	20	21	18	
	ungulates	22	54	31	20	21	18	
	rodents	22	49	31	20	21	18	
Rural	monkeys	14	25	25	2	11	15	
	bats	13	27	27	0	8	17	
	carnivores	3	11	21	1	6	16	
	birds	16	24	31	17	19	18	
	crocodiles	13	26	28	3	8	14	
	snakes	11	22	20	1	6	11	
	turtles	14	41	27	16	17	16	
Niger		4	10	11	1	11	14	51
	ungulates	4	9	11	1	11	14	
	rodents	4	10	11	1	11	14	
Urban	monkeys	0	1	4	0	4	7	
	bats	0	2	3	0	5	11	
	carnivores	0	1	4	0	8	7	
	birds	4	8	9	0	10	12	
	crocodiles	1	6	6	0	7	8	
	snakes	1	4	6	0	7	5	
	turtles	3	8	9	0	10	9	
Niger		31	50	41	23	35	32	213
	ungulates	30	48	41	22	35	32	
	rodents	29	48	41	23	35	32	
Rural	monkeys	8	21	26	16	20	25	
	bats	12	33	27	14	20	27	
	carnivores	8	19	17	11	16	22	
	birds	21	41	37	18	31	31	
	crocodiles	9	28	33	13	28	28	
	snakes	8	21	27	9	18	25	
	turtles	23	33	38	16	28	30	

