

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

***N'*-(*E*)-2-Chlorobenzylidene]-2-(6-methoxynaphthalen-2-yl)propanohydrazide**Joel T. Mague,^a Shaaban K. Mohamed,^{b,c} Mehmet Akkurt,^d Herman Potgieter^e and Mustafa R. Albayati^{f*}

^aDepartment of Chemistry, Tulane University, New Orleans, LA 70118, USA, ^bChemistry and Environmental Division, Manchester Metropolitan University, Manchester M1 5GD, England, ^cChemistry Department, Faculty of Science, Minia University, 61519 El-Minia, Egypt, ^dDepartment of Physics, Faculty of Sciences, Erciyes University, 38039 Kayseri, Turkey, ^eAnalytical Development Division, Manchester Metropolitan University, Manchester M1 5GD, England, and ^fKirkuk University, College of Science, Department of Chemistry, Kirkuk, Iraq
Correspondence e-mail: shaabankamel@yahoo.com


Received 10 April 2014; accepted 16 April 2014

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.026; wR factor = 0.064; data-to-parameter ratio = 13.7.

In the title compound, $\text{C}_{21}\text{H}_{19}\text{ClN}_2\text{O}_2$, the benzene ring and the naphthalene ring system are oriented at a dihedral angle of $65.24(10)^\circ$. In the crystal, $\text{N}-\text{H}\cdots\text{O}$, $\text{C}-\text{H}\cdots\text{N}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules, forming chains along the b -axis direction. Further $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds link the chains, forming corrugated sheets lying parallel to $(10\bar{1})$.

Related literature

For the use of Naproxen [systematic name: (+)-(*S*)-2-(6-methoxynaphthalen-2-yl)propanoic acid] and hydrazide-hydrazones in the treatment of disease and inflammation, see: Merlet *et al.* (2013); Almasirad *et al.* (2005, 2006). For the harmful side-effects of non-steroidal anti-inflammatory drugs (NSAIDs), see: Uzgören-Baran *et al.* (2012); Tozkoparan *et al.* (2012). For the synthesis of NSAIDs with safer pro-drug profiles and enhanced chromophore efficacy, see: Koopaei *et al.* (2013). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_{21}\text{H}_{19}\text{ClN}_2\text{O}_2$
 $M_r = 366.83$
 Monoclinic, $P2_1$
 $a = 6.5703(2)$ Å
 $b = 8.6166(2)$ Å
 $c = 16.3411(4)$ Å
 $\beta = 98.6850(9)^\circ$
 $V = 914.52(4)$ Å³
 $Z = 2$
 Cu $K\alpha$ radiation
 $\mu = 1.99$ mm⁻¹
 $T = 100$ K
 $0.23 \times 0.09 \times 0.02$ mm

Data collection

Bruker D8 VENTURE PHOTON
 100 CMOS diffractometer
 Absorption correction: multi-scan
 (SADABS; Bruker, 2013)
 $T_{\min} = 0.87$, $T_{\max} = 0.96$
 13492 measured reflections
 3243 independent reflections
 3122 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.030$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.026$
 $wR(F^2) = 0.064$
 $S = 1.05$
 3243 reflections
 237 parameters
 1 restraint
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.20$ e Å⁻³
 $\Delta\rho_{\min} = -0.12$ e Å⁻³
 Absolute structure: Flack
 parameter determined using 1358
 quotients $[(I^+) - (I^-)] / [(I^+) + (I^-)]$
 (Parsons *et al.*, 2013)
 Absolute structure parameter:
 0.056 (5)

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1}\cdots\text{O2}^i$	0.91	1.93	2.829 (2)	169
$\text{C12}-\text{H12}\cdots\text{N2}^i$	1.00	2.48	3.444 (3)	163
$\text{C15}-\text{H15}\cdots\text{O2}^i$	0.95	2.48	3.259 (3)	139
$\text{C19}-\text{H19}\cdots\text{O1}^{ii}$	0.95	2.57	3.486 (3)	163

Symmetry codes: (i) $-x + 1, y - \frac{1}{2}, -z + 1$; (ii) $x + 1, y, z + 1$.

Data collection: APEX2 (Bruker, 2013); cell refinement: SAINT (Bruker, 2013); data reduction: SAINT; program(s) used to solve structure: SHELXT (Sheldrick, 2008); program(s) used to refine structure: SHELXL (Sheldrick, 2008); molecular graphics: DIAMOND (Brandenburg & Putz, 2012); software used to prepare material for publication: SHELXTL (Sheldrick, 2008).

The support of NSF-MRI grant No. 1228232 for the purchase of the diffractometer is gratefully acknowledged.

Supporting information for this paper is available from the IUCr electronic archives (Reference: SJ5398).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Almasirad, A., Hosseini, R., Jalalizadeh, H., Rahimi-Moghaddam, Z., Abaeian, N., Janafrooz, M., Abbaspour, M., Ziaee, V., Dalvandi, A. & Shafiee, A. (2006). *Biol. Pharm. Bull.* **29**, 1180–1185.
- Almasirad, A., Tajik, M., Bakhtiari, D., Shafiee, A., Abdollahi, M., Zamani, M. J. & Esmaily, H. (2005). *J. Pharm. Pharm. Sci.* **8**, 419–425.
- Brandenburg, K. & Putz, H. (2012). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2013). *APEX2, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Koopaei, M. N., Assarzadeh, M. J., Almasirad, A., Ghasemi-Niri, S. F., Amini, M., Kebriaeezadeh, A., Koopaei, N. N., Ghadimi, M. & Tabei, A. (2013). *Iran. J. Pharm. Res.* **12**, 721–727.
- Merlet, N., Busseuil, D., Rhéaume, E. & Tardif, J.-C. (2013). *Anti-Inflamm. Anti-Allergy Agents Med. Chem.* **12**, 24–35.
- Parsons, S., Flack, H. D. & Wagner, T. (2013). *Acta Cryst.* **B69**, 249–259.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Tozkoparan, B., Aytaç, S. P., Gürsoy, S. & Aktay, G. (2012). *Med. Chem. Res.* **21**, 192–201.
- Uzgören-Baran, A., Tel, B. C., Deniz Sarıgöl, D., Öztürk, E. I., Kazkayası, I., Okay, G., Ertand, M. & Tozkoparan, B. (2012). *Eur. J. Med. Chem.* **57**, 398–406.

supplementary materials

Acta Cryst. (2014). E70, o631–o632 [doi:10.1107/S1600536814008629]

***N'*-[(*E*)-2-Chlorobenzylidene]-2-(6-methoxynaphthalen-2-yl)propanohydrazide**

Joel T. Mague, Shaaban K. Mohamed, Mehmet Akkurt, Herman Potgieter and Mustafa R. Albayati

1. Comment

Naproxen ((+)-(S)-2-(6-methoxynaphthalen-2-yl)propanoic acid), among many non-steroidal anti-inflammatory drugs (NSAIDs), is used mainly in the treatment of pain, rheumatoid and inflammatory diseases (Merlet *et al.*, 2013). It was reported that the presence of a carboxylic acid group in the parent drug leads to many undesirable side-effects such as gastrointestinal toxicity and ulceration (Uzgören-Baran *et al.*, 2012; Tozkoparan *et al.*, 2012). Recently, it was found that masking the carboxylic acid residue in the parent drug of NSAIDs led to safer pro-drug profiles and enhanced the chromophore efficacy (Koopaei *et al.*, 2013). On the other hand, hydrazide-hydrazone scaffold compounds have been found to possess significant anti-inflammatory effects (Almasirad *et al.*, 2005; Almasirad *et al.*, 2006). Based on these findings the title compound was designed to be a hydrazone profile incorporating the Naproxen core structure without a carboxylic acid substituent.

The naphthalene ring system of the title compound (I, Fig.-1) is essentially planar [maximum deviation = 0.025 (2) Å for C7 and -0.022 (2) Å for C9]. The dihedral angle between the mean planes of the naphthalene and phenyl groups is 65.24 (10)°. The C1–O1–C2–C11, C6–C7–C12–C13, C7–C12–C14–O2, C7–C12–C14–N1, C12–C14–N1–N2, O2–C14–N1–N2 and C14–N1–N2–C15 torsion angles are -3.6 (3), 28.5 (3), 82.8 (2), -95.9 (2), 177.67 (18), -3.6 (3) and 177.4 (2)°, respectively. In (I), the bond lengths (Allen *et al.*, 1987) and angles are within normal ranges.

In the crystal structure, the N—H···O and C—H···N hydrogen bonds link the molecules, forming chains along the *b*-axis (Table 1 and Fig. 2). However, sensible C—H···O contacts are also present that link molecules into chains along *c* and extend the packing along the *c* axis.

2. Experimental

A mixture of 1 mmol (244 mg) Naproxen acid hydrazide [2-(6-methoxy-2-naphthyl)propanehydrazide] and 1 mmol (141 mg) 2-chlorobenzaldehyde was refluxed in 30 ml ethanol for 5hr in the presence of a few catalytic drops of glacial acetic acid. The mixture was cooled and separated, the solid filtered off, dried under vacuum and recrystallized from ethanol to furnish white crystals in a good quality suitable for X-ray diffraction. Mp 488–451 K.

3. Refinement

H-atoms attached to carbon were placed in calculated positions (C—H = 0.95 - 1.00 Å) while those attached to nitrogen were placed in locations derived from a difference map and their parameters adjusted to give N—H = 0.91 Å. All were included as riding contributions with isotropic displacement parameters 1.2 - 1.5 times those of the attached atoms.




Figure 1

A perspective view of the title molecule with 50% probability ellipsoids.




Figure 2

Crystal packing viewed down the *a* axis showing hydrogen bonds as dotted lines (N—H...O: purple, C—H...O: black, C—H...N: green).

***N'*-[*E*]-2-Chlorobenzylidene]-2-(6-methoxynaphthalen-2-yl)propanohydrazide**

Crystal data

$C_{21}H_{19}ClN_2O_2$

$M_r = 366.83$

Monoclinic, $P2_1$

Hall symbol: P 2yb

$a = 6.5703 (2) \text{ \AA}$

$b = 8.6166 (2) \text{ \AA}$

$c = 16.3411 (4) \text{ \AA}$

$\beta = 98.6850 (9)^\circ$

$V = 914.52 (4) \text{ \AA}^3$

$Z = 2$

$F(000) = 384$

$D_x = 1.332 \text{ Mg m}^{-3}$

Cu $K\alpha$ radiation, $\lambda = 1.54178 \text{ \AA}$
 Cell parameters from 9961 reflections
 $\theta = 2.7\text{--}68.2^\circ$
 $\mu = 1.99 \text{ mm}^{-1}$

$T = 100 \text{ K}$
 Plate, colourless
 $0.23 \times 0.09 \times 0.02 \text{ mm}$

Data collection

Bruker D8 VENTURE PHOTON 100 CMOS
 diffractometer
 Radiation source: INCOATEC $I\mu\text{S}$ micro-focus
 source
 Mirror monochromator
 Detector resolution: $10.4167 \text{ pixels mm}^{-1}$
 ω and ϕ scans
 Absorption correction: multi-scan
 (SADABS; Bruker, 2013)

$T_{\min} = 0.87, T_{\max} = 0.96$
 13492 measured reflections
 3243 independent reflections
 3122 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.030$
 $\theta_{\max} = 68.2^\circ, \theta_{\min} = 2.7^\circ$
 $h = -7 \rightarrow 7$
 $k = -9 \rightarrow 10$
 $l = -19 \rightarrow 19$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.026$
 $wR(F^2) = 0.064$
 $S = 1.05$
 3243 reflections
 237 parameters
 1 restraint
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map

Hydrogen site location: mixed
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0303P)^2 + 0.1372P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.20 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.12 \text{ e \AA}^{-3}$
 Absolute structure: Flack parameter determined
 using 1358 quotients $[(I^+) - (I^-)] / [(I^+) + (I^-)]$
 (Parsons *et al.*, 2013)
 Absolute structure parameter: 0.056 (5)

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement on F^2 for ALL reflections except those flagged by the user for potential systematic errors. Weighted R -factors wR and all goodnesses of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The observed criterion of $F^2 > \sigma(F^2)$ is used only for calculating $-R$ -factor-obs *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	1.26515 (8)	0.49714 (7)	0.64924 (3)	0.0316 (2)
O1	0.4688 (2)	1.0112 (2)	-0.00565 (9)	0.0298 (5)
O2	0.3102 (2)	0.82203 (18)	0.49852 (9)	0.0270 (5)
N1	0.5530 (3)	0.6307 (2)	0.51581 (11)	0.0228 (6)
N2	0.6679 (3)	0.7022 (2)	0.58251 (11)	0.0244 (6)
C1	0.6649 (4)	0.9724 (4)	-0.02662 (17)	0.0422 (9)
C2	0.4163 (3)	0.9427 (3)	0.06423 (14)	0.0246 (7)
C3	0.2255 (3)	0.9925 (3)	0.08485 (12)	0.0251 (6)
C4	0.1563 (3)	0.9358 (3)	0.15342 (14)	0.0255 (7)
C5	0.2726 (3)	0.8254 (3)	0.20542 (13)	0.0217 (6)
C6	0.2070 (3)	0.7663 (3)	0.27802 (14)	0.0227 (7)

C7	0.3221 (3)	0.6596 (2)	0.32769 (14)	0.0220 (6)
C8	0.5082 (3)	0.6063 (3)	0.30327 (14)	0.0245 (7)
C9	0.5762 (3)	0.6609 (3)	0.23431 (14)	0.0246 (7)
C10	0.4627 (3)	0.7741 (2)	0.18331 (14)	0.0227 (6)
C11	0.5318 (3)	0.8356 (3)	0.11192 (14)	0.0235 (6)
C12	0.2624 (3)	0.5983 (3)	0.40812 (13)	0.0231 (7)
C13	0.0309 (3)	0.5967 (3)	0.41077 (14)	0.0272 (7)
C14	0.3743 (3)	0.6955 (2)	0.47855 (14)	0.0226 (7)
C15	0.8290 (3)	0.6268 (3)	0.61428 (14)	0.0236 (7)
C16	0.9609 (3)	0.6871 (3)	0.68768 (14)	0.0243 (7)
C17	0.8874 (4)	0.7958 (3)	0.73949 (14)	0.0294 (8)
C18	1.0113 (4)	0.8504 (3)	0.80960 (15)	0.0353 (8)
C19	1.2109 (4)	0.7965 (3)	0.82972 (16)	0.0387 (9)
C20	1.2888 (4)	0.6892 (3)	0.77994 (16)	0.0330 (8)
C21	1.1634 (4)	0.6351 (3)	0.70945 (15)	0.0276 (7)
H1	0.58750	0.53150	0.50470	0.0270*
H1A	0.77230	0.99770	0.01990	0.0630*
H1B	0.68830	1.03170	-0.07550	0.0630*
H1C	0.66940	0.86110	-0.03860	0.0630*
H3	0.14500	1.06600	0.05070	0.0300*
H4	0.02800	0.97060	0.16660	0.0310*
H6	0.08020	0.80120	0.29270	0.0270*
H8	0.58740	0.53030	0.33610	0.0290*
H9	0.70180	0.62280	0.22000	0.0300*
H11	0.65930	0.80210	0.09730	0.0280*
H12	0.31390	0.48930	0.41570	0.0280*
H13A	0.00370	0.53980	0.45990	0.0410*
H13B	-0.04030	0.54570	0.36090	0.0410*
H13C	-0.01910	0.70350	0.41320	0.0410*
H15	0.86290	0.53170	0.59020	0.0280*
H17	0.75010	0.83280	0.72640	0.0350*
H18	0.95920	0.92500	0.84390	0.0420*
H19	1.29500	0.83360	0.87820	0.0460*
H20	1.42620	0.65270	0.79360	0.0400*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0219 (3)	0.0340 (3)	0.0383 (3)	0.0029 (2)	0.0027 (2)	0.0080 (3)
O1	0.0250 (8)	0.0373 (9)	0.0268 (8)	0.0025 (8)	0.0030 (6)	0.0061 (8)
O2	0.0245 (8)	0.0235 (8)	0.0314 (9)	0.0027 (7)	-0.0005 (6)	0.0002 (7)
N1	0.0203 (10)	0.0203 (9)	0.0262 (10)	0.0012 (7)	-0.0014 (7)	-0.0004 (7)
N2	0.0211 (9)	0.0251 (10)	0.0256 (10)	-0.0019 (7)	-0.0007 (8)	0.0014 (8)
C1	0.0297 (13)	0.0577 (19)	0.0408 (14)	0.0070 (12)	0.0104 (11)	0.0197 (13)
C2	0.0245 (12)	0.0266 (12)	0.0214 (11)	-0.0034 (8)	-0.0003 (9)	-0.0032 (8)
C3	0.0248 (11)	0.0244 (10)	0.0242 (10)	0.0035 (10)	-0.0023 (8)	0.0000 (10)
C4	0.0208 (11)	0.0283 (11)	0.0260 (12)	0.0053 (9)	-0.0012 (9)	-0.0028 (9)
C5	0.0204 (11)	0.0210 (11)	0.0224 (11)	0.0010 (9)	-0.0011 (8)	-0.0044 (9)
C6	0.0168 (11)	0.0247 (11)	0.0258 (12)	0.0018 (8)	0.0009 (9)	-0.0050 (9)
C7	0.0193 (11)	0.0215 (11)	0.0242 (11)	-0.0021 (8)	-0.0003 (9)	-0.0037 (8)

C8	0.0199 (11)	0.0211 (11)	0.0310 (12)	0.0018 (8)	-0.0005 (9)	0.0013 (9)
C9	0.0168 (11)	0.0235 (11)	0.0331 (13)	0.0031 (8)	0.0028 (9)	-0.0031 (9)
C10	0.0198 (11)	0.0213 (11)	0.0258 (11)	-0.0007 (8)	-0.0005 (9)	-0.0065 (8)
C11	0.0195 (11)	0.0240 (11)	0.0267 (11)	0.0008 (9)	0.0023 (9)	-0.0038 (9)
C12	0.0190 (11)	0.0212 (11)	0.0279 (12)	0.0016 (8)	-0.0001 (9)	0.0021 (9)
C13	0.0211 (12)	0.0328 (13)	0.0272 (12)	-0.0011 (9)	0.0016 (10)	0.0029 (10)
C14	0.0194 (11)	0.0225 (11)	0.0261 (12)	0.0002 (8)	0.0039 (9)	0.0026 (9)
C15	0.0215 (11)	0.0234 (11)	0.0258 (12)	-0.0004 (9)	0.0033 (9)	0.0032 (9)
C16	0.0252 (12)	0.0234 (12)	0.0237 (11)	-0.0035 (8)	0.0016 (9)	0.0054 (8)
C17	0.0321 (13)	0.0267 (13)	0.0285 (13)	-0.0022 (9)	0.0019 (10)	0.0045 (9)
C18	0.0487 (16)	0.0289 (14)	0.0279 (13)	-0.0086 (11)	0.0045 (11)	-0.0003 (10)
C19	0.0438 (16)	0.0399 (16)	0.0281 (13)	-0.0181 (12)	-0.0080 (11)	0.0046 (11)
C20	0.0269 (12)	0.0356 (14)	0.0328 (14)	-0.0107 (10)	-0.0071 (10)	0.0107 (10)
C21	0.0250 (12)	0.0283 (13)	0.0285 (12)	-0.0051 (9)	0.0006 (10)	0.0092 (9)

Geometric parameters (Å, °)

C11—C21	1.740 (3)	C16—C17	1.397 (3)
O1—C1	1.422 (3)	C17—C18	1.384 (3)
O1—C2	1.375 (3)	C18—C19	1.383 (4)
O2—C14	1.231 (2)	C19—C20	1.380 (4)
N1—N2	1.374 (3)	C20—C21	1.392 (4)
N1—C14	1.358 (3)	C1—H1A	0.9800
N2—C15	1.283 (3)	C1—H1B	0.9800
N1—H1	0.9100	C1—H1C	0.9800
C2—C11	1.362 (3)	C3—H3	0.9500
C2—C3	1.413 (3)	C4—H4	0.9500
C3—C4	1.362 (3)	C6—H6	0.9500
C4—C5	1.419 (3)	C8—H8	0.9500
C5—C6	1.417 (3)	C9—H9	0.9500
C5—C10	1.422 (3)	C11—H11	0.9500
C6—C7	1.375 (3)	C12—H12	1.0000
C7—C12	1.522 (3)	C13—H13A	0.9800
C7—C8	1.419 (3)	C13—H13B	0.9800
C8—C9	1.358 (3)	C13—H13C	0.9800
C9—C10	1.419 (3)	C15—H15	0.9500
C10—C11	1.417 (3)	C17—H17	0.9500
C12—C13	1.528 (3)	C18—H18	0.9500
C12—C14	1.520 (3)	C19—H19	0.9500
C15—C16	1.465 (3)	C20—H20	0.9500
C16—C21	1.398 (3)		
C1—O1—C2	116.70 (19)	C11—C21—C20	117.8 (2)
N2—N1—C14	120.35 (17)	O1—C1—H1A	109.00
N1—N2—C15	114.50 (18)	O1—C1—H1B	109.00
N2—N1—H1	117.00	O1—C1—H1C	110.00
C14—N1—H1	121.00	H1A—C1—H1B	109.00
O1—C2—C3	114.2 (2)	H1A—C1—H1C	109.00
O1—C2—C11	125.31 (19)	H1B—C1—H1C	110.00
C3—C2—C11	120.5 (2)	C2—C3—H3	120.00

C2—C3—C4	120.4 (2)	C4—C3—H3	120.00
C3—C4—C5	121.10 (19)	C3—C4—H4	119.00
C6—C5—C10	119.4 (2)	C5—C4—H4	119.00
C4—C5—C6	122.54 (19)	C5—C6—H6	119.00
C4—C5—C10	118.05 (19)	C7—C6—H6	119.00
C5—C6—C7	121.66 (19)	C7—C8—H8	119.00
C6—C7—C8	118.1 (2)	C9—C8—H8	119.00
C6—C7—C12	123.46 (18)	C8—C9—H9	119.00
C8—C7—C12	118.43 (18)	C10—C9—H9	119.00
C7—C8—C9	121.8 (2)	C2—C11—H11	120.00
C8—C9—C10	121.1 (2)	C10—C11—H11	120.00
C5—C10—C11	119.72 (19)	C7—C12—H12	108.00
C9—C10—C11	122.36 (19)	C13—C12—H12	108.00
C5—C10—C9	117.92 (19)	C14—C12—H12	108.00
C2—C11—C10	120.28 (19)	C12—C13—H13A	109.00
C7—C12—C14	107.72 (18)	C12—C13—H13B	109.00
C7—C12—C13	114.52 (18)	C12—C13—H13C	110.00
C13—C12—C14	110.68 (18)	H13A—C13—H13B	109.00
N1—C14—C12	113.54 (17)	H13A—C13—H13C	109.00
O2—C14—C12	122.90 (19)	H13B—C13—H13C	110.00
O2—C14—N1	123.55 (19)	N2—C15—H15	120.00
N2—C15—C16	120.1 (2)	C16—C15—H15	120.00
C17—C16—C21	117.6 (2)	C16—C17—H17	119.00
C15—C16—C17	121.2 (2)	C18—C17—H17	119.00
C15—C16—C21	121.2 (2)	C17—C18—H18	120.00
C16—C17—C18	121.1 (2)	C19—C18—H18	120.00
C17—C18—C19	120.1 (2)	C18—C19—H19	120.00
C18—C19—C20	120.5 (2)	C20—C19—H19	120.00
C19—C20—C21	119.1 (2)	C19—C20—H20	120.00
C16—C21—C20	121.7 (2)	C21—C20—H20	120.00
C11—C21—C16	120.48 (19)		
C1—O1—C2—C3	176.4 (2)	C8—C7—C12—C13	-152.7 (2)
C1—O1—C2—C11	-3.6 (3)	C8—C7—C12—C14	83.7 (2)
C14—N1—N2—C15	177.4 (2)	C7—C8—C9—C10	-0.2 (4)
N2—N1—C14—O2	3.6 (3)	C8—C9—C10—C5	-1.7 (3)
N2—N1—C14—C12	-177.67 (18)	C8—C9—C10—C11	178.7 (2)
N1—N2—C15—C16	-177.67 (19)	C5—C10—C11—C2	-0.4 (3)
O1—C2—C3—C4	-178.9 (2)	C9—C10—C11—C2	179.2 (2)
C11—C2—C3—C4	1.0 (4)	C7—C12—C14—O2	82.8 (2)
O1—C2—C11—C10	179.3 (2)	C7—C12—C14—N1	-95.9 (2)
C3—C2—C11—C10	-0.7 (4)	C13—C12—C14—O2	-43.1 (3)
C2—C3—C4—C5	-0.3 (4)	C13—C12—C14—N1	138.2 (2)
C3—C4—C5—C6	178.8 (2)	N2—C15—C16—C17	20.3 (3)
C3—C4—C5—C10	-0.8 (3)	N2—C15—C16—C21	-161.3 (2)
C4—C5—C6—C7	-179.9 (2)	C15—C16—C17—C18	178.8 (2)
C10—C5—C6—C7	-0.4 (3)	C21—C16—C17—C18	0.3 (4)
C4—C5—C10—C9	-178.5 (2)	C15—C16—C21—C11	0.3 (3)
C4—C5—C10—C11	1.1 (3)	C15—C16—C21—C20	-178.7 (2)

C6—C5—C10—C9	2.0 (3)	C17—C16—C21—C11	178.83 (19)
C6—C5—C10—C11	-178.4 (2)	C17—C16—C21—C20	-0.2 (4)
C5—C6—C7—C8	-1.5 (3)	C16—C17—C18—C19	-0.5 (4)
C5—C6—C7—C12	177.2 (2)	C17—C18—C19—C20	0.6 (4)
C6—C7—C8—C9	1.8 (3)	C18—C19—C20—C21	-0.4 (4)
C12—C7—C8—C9	-177.0 (2)	C19—C20—C21—C11	-178.8 (2)
C6—C7—C12—C13	28.5 (3)	C19—C20—C21—C16	0.2 (4)
C6—C7—C12—C14	-95.1 (2)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1 \cdots O2 ⁱ	0.91	1.93	2.829 (2)	169
C12—H12 \cdots N2 ⁱ	1.00	2.48	3.444 (3)	163
C15—H15 \cdots O2 ⁱ	0.95	2.48	3.259 (3)	139
C19—H19 \cdots O1 ⁱⁱ	0.95	2.57	3.486 (3)	163

Symmetry codes: (i) $-x+1, y-1/2, -z+1$; (ii) $x+1, y, z+1$.