

**Please cite the Published Version**

Beeton, ML, Alves, DR, Enright, Mark and Jenkins, ATA (2015) Assessing phage therapy against *Pseudomonas aeruginosa* using a *Galleria mellonella* infection model. *International Journal of Antimicrobial Agents*, 46 (2). pp. 196-200. ISSN 0924-8579

**DOI:** <https://doi.org/10.1016/j.ijantimicag.2015.04.005>

**Publisher:** Elsevier

**Version:** Accepted Version

**Downloaded from:** <https://e-space.mmu.ac.uk/620969/>

**Usage rights:** © In Copyright

**Additional Information:** The Accepted Version is the author's final submitted typescript version that we accepted for publication before it was edited and typeset.

**Enquiries:**

If you have questions about this document, contact [openresearch@mmu.ac.uk](mailto:openresearch@mmu.ac.uk). Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

1 Assessing phage therapy against *Pseudomonas aeruginosa* using a *Galleria mellonella* infection model

2

3 Beeton M. L.<sup>1\*</sup>, Alves D. R.<sup>2\*</sup>, Enright M. C.<sup>2</sup> and Jenkins A. T. A.<sup>2</sup>

4

5 <sup>1</sup>Cardiff Metropolitan University, Western Avenue, Cardiff CF5 2YB, United Kingdom.

6 <sup>2</sup>Department of Chemistry, University of Bath, Claverton Down, Bath BA2 7AY, United Kingdom.

7 \*both authors contributed equally

8

9 Corresponding author. E-mail. mbeeton@cardiffmet.ac.uk

10

11 Keywords. *Pseudomonas aeruginosa*, phage therapy, *Galleria mellonella*, infection model

12

## 13 Synopsis

### 14 Objectives

15 To develop a *Galleria mellonella* infection model to assess the *in vivo* efficacy of phage therapy against  
16 laboratory and clinical strains of *Pseudomonas aeruginosa*.

### 17 Methods

18 *Galleria* were infected with *P. aeruginosa* PA01 and treated with varying multiplicity of infection of  
19 phage either two hours post-infection (treatment) or two hours pre-infection (prevention) via  
20 injection into the haemolymph. To address the kinetics of infection larvae were bled over a period of

21 24 hours for quantification of bacteria and phage. Finally, clinical strains from acute and chronic cystic  
22 fibrosis infections were used in the prevention model to further validate the model.

## 23 **Results**

24 Survival rates at 24 hours when infected with 10 cells/larvae were greater with the prevention model  
25 versus treatment model (47 vs 40% MOI 10, 47 vs 20 % MOI 1 and 33 vs 7% MOI 0.1). This pattern  
26 also held true when infected with 100 cells/larvae (87% vs 20% MOI 10; 53% vs 13 % MOI 1; 67% vs  
27 7% MOI 0.1). By 24 hours post infection phage had kept bacterial numbers in the haemolymph to  
28 1000-fold lower than the non-treated group. Phage increased survival of *Galleria* when infected with  
29 both the acute (0% vs 85%) and chronic (80% vs 100%) clinical *P. aeruginosa* strains.

## 30 **Conclusions**

31 Here we present data for the use of *G. mellonella* as a simple, robust and cost-effective model for the  
32 initial *in vivo* examination *P. aeruginosa* targeted phage therapy which may be applied to other  
33 pathogens with similarly low infective doses.

34

## 35 **Introduction**

36 Multi-drug resistant bacterial pathogens pose an ever increasing threat to human health. This  
37 problem is in part due to a lack of novel antibiotics approved for use over the last few decades resulting  
38 in an urgent need to identify new avenues for treating bacterial infections, especially those caused by  
39 gram-negative pathogens<sup>1</sup> *Pseudomonas aeruginosa* is an opportunistic pathogen that is a leading  
40 cause of infection among burns victim and patients with cystic fibrosis. It is also responsible for a large  
41 number of health-care associated infections. To make matters worse, *P. aeruginosa* is associated with  
42 hypermutability and due to high antibiotic selective pressure has given rise to the emergence of multi-  
43 drug resistant strains in the population and concerns about available effective treatments are  
44 growing.<sup>2, 3</sup> In the UK resistance to two or more antibiotics among *P. aeruginosa* isolated from the

45 lungs of cystic fibrosis (CF) patients has risen to 40%.<sup>4</sup> This is a worrying statistic as colonisation of the  
46 CF lung with *P. aeruginosa* is a predictor of poor prognosis and associated with a 2-3 fold increased  
47 chance of death over an eight year period.<sup>5</sup> For this reason novel anti-infectives are needed.

48

49 Facing such a scenario interest in phage therapy in Western society has experienced resurgence after  
50 research into this area fell out of favour following the discovery of antibiotics. Bacteriophages (or  
51 phages) are viral particles able to infect bacterial cells with high specificity, taking over cellular function  
52 to replicate its genome. Upon maturation is complete the bacterial cell wall is lysed to release viral  
53 progeny.

54 Phage therapy can be broadly subdivided in to four main categories.<sup>6</sup> (1) Conventional phage therapy  
55 principally uses lytic phage to lyse target bacterial species. (2) Modified phage therapy utilizes  
56 genetically altered phage with favourable properties such as non-lytic replication to avoid the  
57 possibility of endotoxin shock, when bacterial cells are lysed. (3) Treatment with enzymes derived  
58 from phage such as administration of endolysins to selectively degrade the bacterial peptidoglycan  
59 cell wall. (4) Finally is the concept of combination therapy with phage and antibiotics where phage  
60 exhibit properties to degrade polysaccharide components of biofilms therefore allowing antibiotics to  
61 penetrate and elicit an action.<sup>7</sup>

62 Although *in vitro* systems allow for a reductionist approach to examining phage interactions with  
63 target bacteria, it does not take into account a more complex *in vivo* system. Mammalian models are  
64 an excellent means of testing phage therapy, but require ethical approval, significant infrastructure  
65 and funds. The *Galleria mellonella* model fills the void between these two systems providing a cheap,  
66 reliable and ethics free system for testing novel antimicrobials.<sup>8</sup> Here we describe the first use of the  
67 *G. mellonella* model to examine the treatment of *P. aeruginosa* infection with phage therapy.

68

## 69 **Materials and Methods**

### 70 **Bacterial strains and preparation of inoculum**

71 Phage therapy was assessed using *P. aeruginosa* PAO1 and two clinical isolates, PA45291 and  
72 BC00907, isolated from cystic fibrosis patients with acute and chronic infections, respectively. Bacteria  
73 were grown to mid-log phase in LB and washed once in PBS. Cells were resuspended in PBS to give a  
74 final cell number of  $1 \times 10^8$  cfu ml<sup>-1</sup> and diluted accordingly in PBS to give the required inoculum size for  
75 each experiment.

76

### 77 **Phage cocktail preparation and titration**

78 Six distinct phages infecting all the *P. aeruginosa* strains mentioned above were combined to establish  
79 a cocktail suspension. Phage solutions were propagated on PAO1. Briefly, 100 µl of phage lysate and  
80 100 µl of host growing culture were mixed and left for 5 min at room temperature. Following  
81 incubation, 3 ml of LB soft-agar containing 0.65% bacteriological agar was added and poured onto  
82 agar plates. After an overnight incubation at 37°C, plates displaying confluent lysis were selected and  
83 3 ml of SM buffer (5 M NaCl, 1M MgSO<sub>4</sub>, 1 M Tris-HCl [pH 7.5], 0.01% w/v gelatine) and 2% (vol/vol)  
84 chloroform were added before incubation at 37°C for 4h. High-titre phage solution was removed from  
85 the plates, centrifuged (8,000 x g, 10 min) to remove cell debris, and then filter sterilized (pore size,  
86 0.22 µm). A PEG purification step was further added to remove any possible bacterial remainings from  
87 the suspensions and the final solutions were stored at 4°C. All the necessary dilutions were performed  
88 in SM buffer. For the titration of the bacteriophage content in the haemolymph, a similar methodology  
89 to the propagation was followed. The several dilutions were mixed with the host bacterial cells and 3  
90 ml of soft agar was added and poured onto agar plates. After an overnight incubation plaques were  
91 counted to determine phage titre.

92

### 93 ***G. mellonella* phage therapy assay**

94 Larvae of *G. mellonella* were obtained from Livefood UK Ltd (Somerset, UK). Larvae were stored at 4  
95 °C and used within 1 week of receipt. A modified methodology developed by Peleg was used to infect  
96 each *Galleria*, but in brief.<sup>8</sup> *Galleria* were surface sterilized with a FASTAID pre-injection swab  
97 containing 70 % ethanol. Using a pair of tweezers each *Galleria* was restrained and with a 26 gauge  
98 Terumo syringe 10 µl of inoculum containing either 100 or 10 cells of *P. aeruginosa* was delivered into  
99 the larval haemolymph behind the last proleg. For the treatment model phage suspension was  
100 delivered behind the last proleg on the opposite side to the infection two hours post-infection and for  
101 the prevention experiment phage suspension was given two hours pre-infection. All experiments used  
102 15 larvae per treatment. A positive control group, where the larvae were infected and treated with  
103 PBS solution, and two negative control groups were also included: one group injected with PBS only,  
104 assessing the impact of any negative effect from the injection process, and one group injected with  
105 phage suspension. Larvae were placed into petri dishes and incubated at 37 °C for 48 hours, being  
106 examined regularly and recorded as dead when they did not move in response to touch.

107

### 108 **Bleeding larvae haemolymph**

109 The prevention model was used to follow the kinetics of bacteria and phage interactions within the  
110 larval haemolymph over time. The phage cocktail, or PBS, was administered two hours prior to  
111 infection with phage initially quantified within the haemolymph at time of infection (time zero).  
112 *Galleria* were infected with 100 cells of *P. aeruginosa* PA01 and at time points of eight and 24 hours  
113 three *Galleria* were sacrificed and bled following incision made with forceps to quantify phage and *P.*  
114 *aeruginosa* in both phage and PBS treated *Galleria*. Titrations of haemolymph were made in SM buffer  
115 for phage counts and for PA01 counts titrations were made in 10 mM of ferrous ammonium sulfate  
116 (FAS) (Sigma Aldrich, UK) for inactivation of extracellular phage prior to viable bacterial counts.

### 117 **Statistical analysis**

118 Kaplan-Meier survival curves were plotted using GraphPad Prism (GraphPad Software, Inc., La Jolla,  
119 CA, USA).

120

## 121 **Results**

### 122 **Treatment of infection**

123 In this study two models of phage and infection interactions were examined. The first was a treatment  
124 whereby *Galleria* were infected with either 10 or 100 cells of *P. aeruginosa* PA01 and left to allow an  
125 infection to establish for two hours. Varying MOIs of phage were then administered and death was  
126 observed over 48 hours. No death was seen in the PBS controls and *Galleria* which were treated with  
127 PBS died quicker when infected with 100 cells versus 10 cells. Administration of phage prolonged the  
128 survival of the *Galleria* in a dose dependent manner, but 0% survival was eventually seen in all groups  
129 by 30 hours [Figure 1a]. At 24 hours there was 100% mortality in the infected and untreated *Galleria*,  
130 but 40% survival for those infected with 10 cells and treated with an MOI of 10 compared with 20%  
131 survival with those infected with 100 cells at the same MOI [Figure 1b].

132

### 133 **Prevention of infection**

134 The second model examined the effect of prevention of infection whereby *Galleria* were given a  
135 prophylactic dose of phage two hours prior to infection with *P. aeruginosa* PA01. Similarly to the  
136 treatment experiment, *Galleria* infected with 100 cells died quicker than those infected with 10 cells  
137 when given PBS two hours before infection. [Figure 2a]. At 24 hours survival ranged from 80% in  
138 *Galleria* given an MOI of 100 to 35% in those given an MOI of 0.1 [Figure 2b]. Compared with 10 cells  
139 there was greater survival at 24 hours for all comparable MOIs when infection was established with  
140 100 cells. Survival ranged from 90% to 60% in *Galleria* given MOIs of 100 and 1, respectively.

141

## 142 Kinetics of *P. aeruginosa* infection and effect of phage treatment

143 To understand the kinetics of a *Pseudomonas* infection within *Galleria*, larvae were infected with 100  
144 cells using the prevention model of infection. Bacteria and phage were quantified at set time points  
145 by bleeding the haemolymph. No endogenous *Pseudomonas* or phage with lytic activity against *P.*  
146 *aeruginosa* PA01 were detected in the uninfected controls. For the *Galleria* which were given *P.*  
147 *aeruginosa* PA01 only the numbers of cells isolated from the haemolymph increased over the duration  
148 of the experiment. By 24 hours all *Galleria* were dead and numbers of *P. aeruginosa* were in the order  
149 of  $10^8$  c.f.u/ml. The second group of *Galleria* were given a prophylactic dose of phage 2 hours prior to  
150 infection and phage and bacteria were then quantified over the course of infection. Numbers of *P.*  
151 *aeruginosa* PA01 were comparable to that of the non-treated *Galleria* after 8 hours infection, but were  
152 three orders of magnitude less cells at 24 hours compared with the non-treated *Galleria*. These  
153 *Galleria* were alive at 24 hours. Numbers of phage increased over the duration of the infection  
154 reaching a peak titre at 24 hours of  $10^8$  p.f.u/ml.

155

## 156 Clinical isolates of *P. aeruginosa*

157 To validate the model of phage therapy with *P. aeruginosa* we sought to test the model with clinical  
158 strains isolated from cystic fibrosis patients experiencing acute and chronic episodes of *P. aeruginosa*  
159 infections. With the PA45291 acute strain all infected *Galleria* were dead by 24 hours whereas there  
160 was 60 % survival at 28 hours in the group which were treated with phage at an MOI 10. When *Galleria*  
161 were infected with the BC09007 chronic strain there was little death at 24 hours (90%) when given  
162 PBS as treatment, but 100 % survival in the phage treated group. By 40 hours all *Galleria* were then  
163 dead.

## 164 Discussion

165 To avoid a scenario whereby society is plunged back into a pre-antibiotic era we need to urgently  
166 identify novel anti-bacterial agents. Phage therapy offers a novel non-antibiotic approach to help in  
167 this battle. The benefits of phage therapy include no cross resistance from pre-existing antibiotic  
168 resistant organisms, high selectivity therefore not wiping out the host microbiota unlike antibiotics as  
169 well as being deemed as safe in trials.<sup>9-11</sup>

170 The *G. mellonella* infection model provides a system that can bridge the gap between *in vitro* studies  
171 and more advanced mammalian studies giving initial proof of principle data. Mammalian models are  
172 crucial for testing the efficacy of phage prior to human trials, but drawbacks include the need for  
173 sufficient infrastructure, substantial costs, as well as the need for ethical approval. *Galleria* larvae  
174 have been used to examine numerous host-pathogen interactions ranging from studies of  
175 pathogenicity to antimicrobial activity with a small number of these examining the potential for phage  
176 therapy.<sup>12-14</sup>

177

178 The strain of *P. aeruginosa* PA01 proved to be highly virulent with only 10 cells per *Galleria* required  
179 to result in death at 24 hours. This is a very low infective dose in this model with organism such as *S.*  
180 *aureus* requiring  $10^5$  - $10^6$  cells/*Galleria* for death, *A. baumannii* requires greater than  $10^4$  and for  
181 *Helicobacter pylori*  $10^6$ - $10^7$  cells are required for establishment of infection.<sup>15-17</sup> This low infectious  
182 dose is of particular interest as it reduces the chances of endotoxin shock due to rapid lysis of high  
183 numbers of Gram negative cells.

184 Two models of therapy were examined. The first was a treatment methodology whereby an acute 2  
185 hour infection was allowed to establish prior to administration of phage. At all MOIs of phage there  
186 was prolonged survival of the *Galleria* regardless whether 10 or 100 bacterial cells were used as the  
187 inoculum. Although there was increased survival compared with the control there was a difference in

188 survival depending on the number of cells in the inoculum. Presumably the 10-fold higher inoculum  
189 of 100 cells vs 10 cells had meant that the infection had become more established within the two hour  
190 time frame therefore reducing the efficacy of the phage to prolong survival.

191 The second model examined the ability to prevent infection using a prophylactic administration of  
192 phage two hours prior to infection. When compared with the treatment model, prophylactic  
193 administration of phage resulted in greater survival after 24 hours at all comparable MOI values.  
194 Presumably this increased efficacy was the result of phage being able to distribute throughout the  
195 haemolymph over the two hour period prior to infection, where as in the treatment model the  
196 bacteria will have had opportunity to establish and begin to express toxins. Interesting was the  
197 observation of greater survival among *Galleria* which received the higher inoculum of 100 cells,  
198 compared with 10 cells. This may have been due to the higher number of bacterial cells resulting in  
199 an increased chance of bacteria and phage interaction resulting in a more rapid amplification of the  
200 phage.

201 In both models phage treated *Galleria* eventually succumbed to the infection resulting in death by 30  
202 hours post infection. For this reason we explored the kinetics of both the *P. aeruginosa* infection as  
203 well as effect phage had on bacterial numbers *in vivo*. The most striking observation was the  
204 comparison between numbers of *P. aeruginosa* in the phage treated and untreated *Galleria*. At 24  
205 hours the phage had kept the number of *P. aeruginosa* to 1000-fold less than the non-treated *Galleria*,  
206 but even in the presence of high titres of phage there had still been active growth, and therefore  
207 infection, from the *P. aeruginosa* over the duration of the experiment. We had previously  
208 hypothesised that the reason for eventual death was the lack of available phage for clearance. From  
209 Figure 3 it is clear that this is not the case due to the high titre of phage within the haemolymph,  
210 although the MOI had shifted from 100 to less than 1 by 24 hours. This hypothesis was also ruled out  
211 by an experiment where *Galleria* were given a second dose of phage four hours after an initial dosing,  
212 but there was no difference when compared with the single dose control (data not shown). One

213 possibility for the continual survival of PA01 in the presence of a high titre of phage was the evolution  
214 of phage resistance within the *Galleria*. This was ruled out after observation of no bacterial growth  
215 when co-cultivating *P. aeruginosa* single colonies, recovered at 24 hours after phage treatment, and  
216 a suspension of phage cocktail (data not shown). The final explanation for the survival would be the  
217 intracellular localisation of *Pseudomonas*. In these experiments we only examined bacterial numbers  
218 within extracted haemolymph. *Pseudomonas aeruginosa* is known to have the ability to invade  
219 epithelial cells which would protect from attack from the phage.<sup>18</sup> This highlights one of the  
220 limitations of phage therapy on pathogens which are able to exist and replicate in an intracellular  
221 environment. Perhaps combination therapy with antibiotics which can enter host cells such as a  
222 fluoroquinolone or tetracycline would have aided in clearance, but this was beyond the scope of this  
223 study. This potential intracellular survival strategy would also explain why the prevention model  
224 showed improved survival compared with the treatment model where the *P. aeruginosa* will have had  
225 time to establish within cells before the *Galleria* received a dose of the phage. Although we  
226 hypothesise the lack of *P. aeruginosa* clearance was due to intracellular localisation, there must have  
227 been a degree of extracellular replication of cells within the haemolymph to allow for the observed  
228 propagation of the phage over time.

229 Finally we looked to demonstrate the effectiveness of the phage model on clinical isolates of *P.*  
230 *aeruginosa*. To do this the prevention model was repeated with clinical isolates from an acute and a  
231 chronic CF infection. Here, the acute isolate resulted in rapid death of the *Galleria* within 24 hours,  
232 with 85 % survival when given phage at an MOI of 10. Interestingly the chronic isolate was less virulent  
233 at 24 hours compared with the acute and PA01 strains, but 100 % mortality was then seen by 40 hours.  
234 In conclusion we present data for the use of the *G. mellonella* as a simple, robust and cost-effective  
235 model for initial examination *P. aeruginosa* targeted phage therapy.

236

237 **Acknowledgements**

238 We thank the Engineering and Physical Sciences Research Council (EPSRC) Healthcare Partnership fund  
239 for their funding and support.

240 The authors declare that they have no conflicts of interest.

241 **Transparency declarations**

242 None to declare

243

244

245 **References**

246

- 247 1. Boucher HW, Talbot GH, Bradley JS et al. Bad bugs, no drugs: no ESKAPE! An update from the  
248 Infectious Diseases Society of America. *Clinical infectious diseases : an official publication of the*  
249 *Infectious Diseases Society of America* 2009; **48**: 1-12.
- 250 2. Oliver A, Mena A. Bacterial hypermutation in cystic fibrosis, not only for antibiotic resistance.  
251 *Clinical microbiology and infection : the official publication of the European Society of Clinical*  
252 *Microbiology and Infectious Diseases* 2010; **16**: 798-808.
- 253 3. Barbier F, Wolff M. [Multi-drug resistant *Pseudomonas aeruginosa*: towards a therapeutic  
254 dead end?]. *Medecine sciences : M/S* 2010; **26**: 960-8.
- 255 4. Pitt TL, Sparrow M, Warner M et al. Survey of resistance of *Pseudomonas aeruginosa* from UK  
256 patients with cystic fibrosis to six commonly prescribed antimicrobial agents. *Thorax* 2003; **58**: 794-6.
- 257 5. Emerson J, Rosenfeld M, McNamara S et al. *Pseudomonas aeruginosa* and other predictors of  
258 mortality and morbidity in young children with cystic fibrosis. *Pediatric pulmonology* 2002; **34**: 91-100.
- 259 6. Viertel TM, Ritter K, Horz HP. Viruses versus bacteria-novel approaches to phage therapy as a  
260 tool against multidrug-resistant pathogens. *The Journal of antimicrobial chemotherapy* 2014; **69**:  
261 2326-36.
- 262 7. Bedi M, Verma V, Chhibber S. Amoxicillin and specific bacteriophage can be used together for  
263 eradication of biofilm of *Klebsiella pneumoniae* B5055. *World J Microbiol Biotechnol* 2009; **25**: 1145-  
264 51.
- 265 8. Peleg AY, Monga D, Pillai S et al. Reduced susceptibility to vancomycin influences  
266 pathogenicity in *Staphylococcus aureus* infection. *The Journal of infectious diseases* 2009; **199**: 532-6.
- 267 9. Paul VD, Sundarrajan S, Rajagopalan SS et al. Lysis-deficient phages as novel therapeutic  
268 agents for controlling bacterial infection. *BMC microbiology* 2011; **11**: 195.
- 269 10. Dethlefsen L, Huse S, Sogin ML et al. The pervasive effects of an antibiotic on the human gut  
270 microbiota, as revealed by deep 16S rRNA sequencing. *PLoS biology* 2008; **6**: e280.

- 271 11. Bruttin A, Brussow H. Human volunteers receiving Escherichia coli phage T4 orally: a safety  
272 test of phage therapy. *Antimicrobial agents and chemotherapy* 2005; **49**: 2874-8.
- 273 12. Kamal F, Dennis JJ. Burkholderia cepacia complex Phage-Antibiotic Synergy (PAS): Antibiotics  
274 stimulate lytic phage activity. *Applied and environmental microbiology* 2014.
- 275 13. Seed KD, Dennis JJ. Experimental bacteriophage therapy increases survival of Galleria  
276 mellonella larvae infected with clinically relevant strains of the Burkholderia cepacia complex.  
277 *Antimicrobial agents and chemotherapy* 2009; **53**: 2205-8.
- 278 14. Abbasifar R, Kropinski AM, Sabour PM et al. Efficiency of bacteriophage therapy against  
279 Cronobacter sakazakii in Galleria mellonella (greater wax moth) larvae. *Archives of virology* 2014; **159**:  
280 2253-61.
- 281 15. Gibreel TM, Upton M. Synthetic epidermicin NI01 can protect Galleria mellonella larvae from  
282 infection with Staphylococcus aureus. *The Journal of antimicrobial chemotherapy* 2013; **68**: 2269-73.
- 283 16. Peleg AY, Jara S, Monga D et al. Galleria mellonella as a model system to study Acinetobacter  
284 baumannii pathogenesis and therapeutics. *Antimicrobial agents and chemotherapy* 2009; **53**: 2605-9.
- 285 17. Giannouli M, Palatucci AT, Rubino V et al. Use of larvae of the wax moth Galleria mellonella as  
286 an in vivo model to study the virulence of Helicobacter pylori. *BMC microbiology* 2014; **14**: 228.
- 287 18. Angus AA, Lee AA, Augustin DK et al. Pseudomonas aeruginosa induces membrane blebs in  
288 epithelial cells, which are utilized as a niche for intracellular replication and motility. *Infection and*  
289 *immunity* 2008; **76**: 1992-2001.
- 290

291 **Figure legends**

292 **Figure 1.** Kaplan-Meier survival curves of *G. mellonella* infected with (A) 100 cells or (B) 10 cells of *P.*  
293 *aeruginosa* PA01 and treated with phage at varying multiplicities of infection two hours post-infection.  
294 C. Percentage of *G. mellonella* survival at 24 hours.

295

296 **Figure 2.** Kaplan-Meier survival curves of *G. mellonella* infected with (A) 100 cells or (B) 10 cells of *P.*  
297 *aeruginosa* PA01 and pre-treated with phage at varying multiplicities of infection two hours pre-  
298 infection. C. Percentage of *G. mellonella* survival at 24 hours.

299

300 **Figure 3.** *In vivo* kinetics of *P. aeruginosa* infection within *G. mellonella* with and without phage  
301 treatment.

302

303 **Figure 4.** Kaplan-Meier survival curves of *G. mellonella* infected with 10 cells of (A) acute *P. aeruginosa*  
304 PA45291 or (B) chronic *P. aeruginosa* BC09007 and pre-treated with phage at an MOI 10 two hours  
305 pre-infection.

306