

Please cite the Published Version

Chandler, B, Lovell, H, Boston, C, Lukas, S, Barr, I, Benediktsson, ÍÖ, Benn, D, Clark, C, Darvill, C, Evans, D, Ewertowski, M, Loibl, D, Margold, M, Otto, J-C, Roberts, D, Stokes, C, Storrar, R and Stroeve, A (2018) Glacial geomorphological mapping: A review of approaches and frameworks for best practice. *Earth-Science Reviews*, 185. pp. 806-846. ISSN 0012-8252

DOI: <https://doi.org/10.1016/j.earscirev.2018.07.015>

Publisher: Elsevier

Version: Accepted Version

Downloaded from: <https://e-space.mmu.ac.uk/621339/>

Usage rights:  [Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0](https://creativecommons.org/licenses/by-nc-nd/4.0/)

Additional Information: This article was originally published in *Earth-Science Reviews*, published by and copyright Elsevier.

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

Glacial geomorphological mapping: a review of approaches and frameworks for best practice

Benjamin M.P. Chandler¹*, Harold Lovell², Clare M. Boston², Sven Lukas³, Iestyn D. Barr⁴,
Ívar Örn Benediktsson⁵, Douglas I. Benn⁶, Chris D. Clark⁷, Christopher M. Darvill⁸,
David J.A. Evans⁹, Marek W. Ewertowski¹⁰, David Loibl¹¹, Martin Margold¹², Jan-Christoph Otto¹³,
David H. Roberts⁹, Chris R. Stokes⁹, Robert D. Storrar¹⁴, Arjen P. Stroeven^{15, 16}

¹ School of Geography, Queen Mary University of London, Mile End Road, London, E1 4NS, UK

² Department of Geography, University of Portsmouth, Portsmouth, UK

³ Department of Geology, Lund University, Lund, Sweden

⁴ School of Science and the Environment, Manchester Metropolitan University, Manchester, UK

⁵ Institute of Earth Sciences, University of Iceland, Reykjavík, Iceland

⁶ Department of Geography and Sustainable Development, University of St Andrews, St Andrews, UK

⁷ Department of Geography, University of Sheffield, Sheffield, UK

⁸ Geography, School of Environment, Education and Development, University of Manchester, Manchester, UK

⁹ Department of Geography, Durham University, Durham, UK

¹⁰ Faculty of Geographical and Geological Sciences, Adam Mickiewicz University, Poznań, Poland

¹¹ Department of Geography, Humboldt University of Berlin, Berlin, Germany

¹² Department of Physical Geography and Geoecology, Charles University, Prague, Czech Republic

¹³ Department of Geography and Geology, University of Salzburg, Salzburg, Austria

¹⁴ Department of the Natural and Built Environment, Sheffield Hallam University, Sheffield, UK

¹⁵ Geomorphology & Glaciology, Department of Physical Geography, Stockholm University, Stockholm, Sweden

¹⁶ Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden

*Corresponding author. Email: b.m.p.chandler@qmul.ac.uk

Abstract

Geomorphological mapping is a well-established method for examining earth surface processes and landscape evolution in a range of environmental contexts. In glacial research, it provides crucial data for a wide range of process-oriented and palaeoglaciological reconstruction studies; in the latter case providing an essential geomorphological framework for establishing glacial chronologies. In recent decades, there have been significant developments in remote sensing and Geographical Information Systems (GIS), with a plethora of high-quality remotely-sensed datasets now (often freely) available. Most recently, the emergence of unmanned aerial vehicle (UAV) technology has allowed sub-decimetrescale aerial images and Digital Elevation Models (DEMs) to be obtained. Traditional field mapping methods still have an important role in

glacial geomorphology, particularly in cirque glacier, valley glacier and icefield/ice-cap outlet settings. Field mapping is also used in ice sheet settings, but often takes the form of necessarily highly-selective ground-truthing of remote mapping. Given the increasing abundance of datasets and methods available for mapping, effective approaches are necessary to enable assimilation of data and ensure robustness. In this contribution, we provide a review and assessment of the various glacial geomorphological methods and datasets currently available, with a focus on their applicability in particular glacial settings. We distinguish two overarching ‘work streams’ that recognise the different approaches typically used in mapping landforms produced by ice masses of different sizes: (i) mapping of ice sheet geomorphological imprints using a combined remote sensing approach, with some field checking (where feasible); and (ii) mapping of alpine and plateau-style ice mass (cirque glacier, valley glacier, icefield and ice-cap) geomorphological imprints using remote sensing and considerable field mapping. Key challenges to accurate and robust geomorphological mapping are highlighted, often necessitating compromises and pragmatic solutions. The importance of combining multiple datasets and/or mapping approaches is emphasised, akin to multi-proxy/-method approaches used in many Earth Science disciplines. Based on our review, we provide idealised frameworks and general recommendations to ensure best practice in future studies and aid in accuracy assessment, comparison and integration of geomorphological data. These will be of particular value where geomorphological data are incorporated in large compilations and subsequently used for palaeoglaciological reconstructions. Finally, we stress that robust interpretations of glacial landforms and landscapes invariably requires additional chronological and/or sedimentological evidence, and that such data should be collected as part of a coupled inductive-deductive approach.

Keywords: glacial geomorphology; geomorphological mapping; GIS; remote sensing; field mapping

1. Introduction

1.1 Background and importance

Mapping the spatial distribution of landforms and features through remote sensing and/or field-based approaches is a well-established method in Earth Sciences to examine earth surface processes and landscape evolution (e.g. Kronberg, 1984; Hubbard and Glasser, 2005; Smith et al., 2011). Moreover, geomorphological mapping is utilised in numerous applied settings, such as natural hazard assessment, environmental planning and civil engineering (e.g. Kienholz, 1977, Finke, 1980; Paron and Claessens, 2011; Marc and Hovius, 2015; Griffiths and Martin, 2017).

Two overarching traditions exist in geomorphological mapping: Firstly, the classical approach involves mapping all geomorphological features in multiple thematic layers (e.g. landforms, breaks of slope, slope angles, drainage), regardless of the range of different processes responsible for forming the landscape. This approach to geomorphological mapping has been particularly widely used in mainland Europe and has resulted in the creation of national legends to record holistic geomorphological data that may be comparable across much larger areas or between studies (Demek, 1972; van Dorsser and Salomé, 1973; Leser and Stäblein, 1975; Klimaszewski, 1990; Schoeneich, 1993; Kneisel et al., 1998; Gustavsson et al., 2006; Rączkowska and Zwoliński, 2015). The second approach involves more detailed, thematic geomorphological mapping commensurate with particular research questions; for example, the map may have an emphasis on mass movements or glacial and periglacial landforms and processes. Such a reductionist approach is helpful in ensuring a map is not ‘cluttered’ with less relevant data that may in turn make a multi-layered map unreadable (e.g. Kuhle, 1990; Robinson et al., 1995; Kraak and Ormeling, 2006). In recent years, the second approach has become much more widespread due to increasing specialisation and thus forms the basis for this review, which focuses on geomorphological mapping in glacial environments.

In glacial research, the production and analysis of geomorphological maps provide a wider context and basis for various process-oriented and palaeoglaciological studies, including:

- (1) analysing glacial sediments and producing process-form models (e.g. Price, 1970; Benn, 1994; Lukas, 2005; Benediktsson et al., 2016);
- (2) quantitatively capturing the pattern and characteristics (‘metrics’) of landforms to understand their formation and evolution (e.g. Spagnolo et al., 2014; Ojala et al., 2015; Ely et al., 2016a);
- (3) devising glacial landsystem models that can be used to elucidate former glaciation styles or inform engineering geology (e.g. Eyles, 1983; Evans et al., 1999; Evans, 2017; Bickerdike et al., 2018);

- (4) reconstructing the extent and dimensions of former or formerly more extensive ice masses (e.g. Dyke and Prest, 1987a; Kleman et al., 1997; Houmark-Nielsen and Kjær, 2003; Benn and Ballantyne, 2005; Glasser et al., 2008; Clark et al., 2012);
- (5) elucidating glacier and ice sheet dynamics, including advance/retreat cycles, flow patterns/velocities and thermal regime (e.g. Kjær et al., 2003; Kleman et al., 2008, 2010; Evans, 2011; Boston, 2012a; Hughes et al., 2014; Darvill et al., 2017);
- (6) identifying sampling locations for targeted numerical dating programmes and ensuring robust chronological frameworks (e.g. Owen et al., 2005; Barrell et al., 2011, 2013; Garcia et al., 2012; Kelley et al., 2014; Stroeven et al., 2014; Blomdin et al., 2016a; Gribenski et al., 2016, 2017);
- (7) calculating palaeoclimatic variables for glaciated regions, namely palaeotemperature and palaeoprecipitation (e.g. Kerschner et al., 2000; Bakke et al., 2005; Stansell et al., 2007; Mills et al., 2012; Boston et al., 2015)
- (8) providing parameters to constrain and test numerical simulations of ice masses (e.g. Kleman et al., 2002; Napieralski et al., 2007a; Golledge et al., 2008; Stokes and Tarasov, 2010; Livingstone et al., 2015; Seguinot et al., 2016; Patton et al., 2017a).

Thus, accurate representation of glacial and associated landforms is crucial to producing geomorphological maps of subsequent value in a wide range of glacial research. This is exemplified in glacial geochronological investigations, where a targeted radiometric dating programme first requires a clear geomorphological (and/or stratigraphic) framework and understanding of the relationships and likely relative ages of different sediment-landform assemblages. In studies that ignore this fundamental principle, it can be challenging to then reconcile any scattered or anomalous numerical ages with a realistic geomorphological interpretation, as the samples have been obtained without a clear genetic understanding of the landforms being sampled (see Boston et al., 2015, for further discussion).

The analysis of geomorphological evidence has been employed in the study of glaciers and ice sheets for over 150 years, with the techniques used in geomorphological mapping undergoing a number of significant developments in that time. The earliest geomorphological investigations involved intensive field surveys (e.g. Close, 1867; Penck and Brückner, 1901/1909; De Geer, 1910; Trotter, 1929; Caldenius, 1932; Raistrick, 1933), before greater efficiency was achieved through the development of aerial photograph interpretation from the late 1950s onwards (e.g. Lueder, 1959; Price, 1963; Welch, 1967; Howarth, 1968; Prest et al., 1968; Sugden, 1970; Sissons, 1977a; Prest, 1983; Kronberg, 1984; Mollard and Janes, 1984). Satellite imagery and digital elevation models (DEMs) have been in widespread usage since their development in the late 20th Century and have, in particular, helped revolutionise our understanding of palaeo-ice sheets (e.g. Barents-Kara Ice Sheet: Winsborrow et al.,

2010; British Ice Sheet: Hughes et al., 2014; Cordilleran Ice Sheet: Kleman et al., 2010; Fennoscandian Ice Sheet: Stroeve et al., 2016; Laurentide Ice Sheet: Margold et al., 2018; Patagonian Ice Sheet: Glasser et al., 2008). In recent times, increasingly higher-resolution DEMs have become available due to the adoption of Light Detection and Ranging (LiDAR) technology (e.g. Salcher et al., 2010; Jónsson et al., 2014; Miller et al., 2014; Dowling et al., 2015; Hardt et al., 2015; Putniņš and Henriksen, 2017) and Unmanned Aerial Vehicles (UAVs) (e.g. Chandler et al., 2016a; Evans et al., 2016a; Ewertowski et al., 2016; Tonkin et al., 2016; Ely et al., 2017). Aside from improvements to remote sensing technologies, the last decade has seen a revolution in data accessibility, with the proliferation of freely available imagery (e.g. Landsat data), freeware mapping platforms (e.g. *Google Earth*) and open-source Geographical Information System (GIS) packages (e.g. *QGIS*). As a result, tools for glacial geomorphological mapping are becoming increasingly accessible, both practically and financially.

Field mapping remains a key component of the geomorphological mapping process, principally in the context of manageable study areas relating to alpine- and plateau-style ice masses, i.e. cirque glaciers, valley glaciers, icefields and ice-caps (e.g. Bendle and Glasser, 2012; Boston, 2012a, b; Jónsson et al., 2014; Pearce et al., 2014; Gribenski et al., 2016; Lardeux et al., 2016; Chandler and Lukas, 2017). This approach is also employed in ice sheet settings, but typically in the form of selective ground checking of mapping from remotely-sensed data or focused mapping of regional sectors (e.g. Stokes et al., 2013; Bendle et al., 2017a; Pearce et al., 2018). Frequently, field mapping is conducted in tandem with sedimentological investigations (see Evans and Benn, 2004, for methods), providing a means of testing preliminary interpretations and identifying problems for specific (and more detailed) studies. This interlinked approach is particularly powerful and enables robust interpretations of genetic processes, glaciation styles and/or glacier dynamics (e.g. Benn and Lukas, 2006; Evans, 2010; Benediktsson et al., 2010, 2016; Gribenski et al., 2016). In this context, it is worth highlighting the frequent use of the term ‘sediment-landform assemblage’ (or ‘landform-sediment assemblage’) as opposed to ‘landform’ in glacial geomorphology, underlining the importance of studying both surface form and internal composition (e.g. Evans, 2003a, 2017; Benn and Evans, 2010; Lukas et al., 2017).

Geomorphological mapping using a combination of field mapping and remotely-sensed data interpretation (hereafter ‘remote mapping’), or a number of remote sensing methods, permits a holistic approach to mapping, wherein the advantages of each method/dataset can be combined to produce an accurate map with robust genetic interpretations (e.g. Boston, 2012a, b; Darvill et al., 2014; Storrar and Livingstone, 2017). As such, approaches are required that allow the accurate transfer and assimilation of data from these various sources, particularly where data are transferred from analogue (e.g. hard-copy aerial photographs) to digital format. Apart from a few recent exceptions for specific locations (e.g. the Scottish Highlands: Boston, 2012a, b; Pearce et al., 2014), there has been limited

explicit discussion of the approaches used to integrate geomorphological data in map production (i.e. the relative contributions of different methods and/or datasets and their associated uncertainties), with many contributions simply stating that the maps were produced through fieldwork and/or remote sensing (e.g. Ballantyne, 1989; Lukas, 2007a; Evans et al., 2009a; McDougall, 2013). Given the diversity of scales, data sources and research questions inherent in glacial geomorphological research, and the increasing abundance of high-quality remotely-sensed datasets, finding the most cost- and time- effective approach is difficult, especially for researchers new to the field.

1.2 Aims and scope

In this contribution, we review the wide range of approaches and datasets available to practitioners and students for geomorphological mapping in glacial environments. The main aims of this review are to (i) synthesise scale-appropriate mapping approaches that are relevant to particular glacial settings, (ii) devise frameworks that will help ensure best practice when mapping, and (iii) encourage clear communication of details on mapping methods used in glacial geomorphological studies. This will ensure transparency and aid data transferability against a background of growing demand to collate geomorphological (and chronological) data in regional compilations (e.g. the BRITICE project: Clark et al., 2004, 2018a; the DATED-1 database: Hughes et al., 2016). A further aim of this contribution is to emphasise the continued and future importance of field mapping in geomorphological research, despite the advent of very high-resolution remotely-sensed datasets in recent years (e.g. UAV-captured imagery).

The following two sections of this review focus on field mapping (Section 2) and remote mapping (Section 3), respectively. We consider these methods in a broadly chronological order to provide historical context and illustrate the evolution of geomorphological mapping in glacial environments. Section 4 discusses the errors associated with each mapping method, an important issue that often receives limited attention within geomorphological studies. Within this discussion, we highlight approaches that can help manage and minimise residual errors. Subsequently, we review the mapping methods used in particular glacial environments (Section 5) and synthesise frameworks to help ensure best practice when mapping (Section 6).

For the purposes of this review, we distinguish two overarching ‘work streams’: (i) mapping of palaeo-ice sheet geomorphological imprints using a combined remote sensing approach, with some field checking (where feasible); and (ii) mapping of alpine- and plateau-style ice mass geomorphological imprints using a combination of remote sensing and considerable field mapping/checking. The second workstream incorporates a spatial continuum of glacier morphologies, namely cirque glaciers, valley glaciers, icefields and ice-caps (cf. Sugden and John, 1976; Benn and

Evans, 2010). The rationale for this subdivision is fourfold: Firstly, the approaches are governed by the size of the (former) glacial systems and thus feasibility of using particular mapping methods in certain settings (cf. Clark, 1997; Storrar et al., 2013). Secondly, there is a greater overlap of spatial and temporal scales (i.e. more detailed records are preserved) in areas glaciated by smaller ice masses that respond more rapidly to climate (cf. Lukas, 2005, 2012; Bradwell et al., 2013; Boston et al., 2015; Chandler et al., 2016b). Thirdly, the different mapping methodologies reflect the difficulties in identifying vertical limits, thickness distribution and surface topography of palaeo-ice sheets (i.e. emphasis often on mapping bed imprints) (cf. Stokes et al., 2015). Finally, the overarching methods employed to map glacial landforms in alpine and plateau settings do not differ fundamentally with ice mass morphology, i.e. most studies in these environments employ a combination of field mapping and remote sensing. In Section 5.3, we also specifically consider geomorphological mapping in modern glacial environments to highlight important issues relating to the temporal resolution of remotely-sensed data and landform preservation potential. We emphasise the importance of utilising multiple datasets and/or mapping approaches in an iterative process in all glacial settings (multiple remotely-sensed datasets in the case of ice sheet-scale geomorphology) to increase accuracy and robustness, akin to multi-proxy methodologies used in many Earth Science disciplines.

2. Field mapping methods

2.1 Background and applicability of field mapping

Traditionally, glacial geomorphological mapping has been undertaken through extensive field surveys, an approach that dates back to the late 19th Century and early 20th Century (e.g. Close, 1867; Goodchild, 1875; Partsch, 1894; Sollas, 1896; Penck and Brückner, 1901/1909; Kendall, 1902; Wright, 1912; Hollingworth, 1931; Caldenius, 1932). Field mapping involves traversing the study area and recording pertinent landforms onto (enlarged) topographic base maps (Figure 1). Typically, field mapping is conducted at cartographic scales of ~1: 10,000 (e.g. Leser and Stäblein, 1975; Rupke and De Jong, 1983; Thorp, 1986; Ballantyne, 1989; Evans, 1990; Benn et al., 1992; Mitchell and Riley, 2006; Rose and Smith, 2008; Boston, 2012a, b) or 1: 25,000 (e.g. Leser, 1983; Ballantyne, 2002, 2007a, b; Benn and Ballantyne, 2005; Lukas and Lukas, 2006). Occasionally, it is conducted at even larger scales, such as 1: 1,000 to 1: 5,000, but this is most appropriate for small areas or project-specific purposes (e.g. Kienholz, 1977; Leser, 1983; Lukas et al., 2005; Coray, 2007; Graf, 2007; Reinardy et al., 2013).

With improvements in technology, the widespread availability of remotely-sensed datasets, and a concomitant ease of access to high-quality printing facilities, alternative approaches to the traditional, *purely* field mapping method have also been employed, including (i) documenting sediment-landform

assemblages during extensive field campaigns both prior to and after commencing remote mapping (e.g. Dyke et al., 1992; Krüger 1994; Lukas and Lukas, 2006; Kjær et al. 2008; Boston, 2012a, b; Jónsson et al., 2014; Schomacker et al. 2014; Everest et al., 2017), (ii) mapping directly onto or annotating print-outs of imagery (e.g. aerial photographs) in the field (e.g. Lovell, 2014), (iii) recording the locations of individual landforms using a (handheld) Global Navigation Satellite System (GNSS) device (e.g. Bradwell et al., 2013; Brynjólfsson et al., 2014; Lovell, 2014; Małecki et al., 2018), or (iv) digitally mapping landforms in the field using a ruggedised tablet PC with built-in GNSS and GIS software (e.g. Finlayson et al., 2011; Pearce et al., 2014). These approaches to field mapping are particularly useful where large-scale topographic maps are unavailable or obsolete.

Detailed field mapping is typically restricted to alpine- and plateau-style ice masses due to logistical and financial constraints (Clark, 1997; Storrar et al., 2013). When conducted at the ice sheet scale, field mapping is (or historically was) undertaken either as part of long-term campaigns by national geological surveys in conjunction with surficial geology mapping programmes (e.g. Barrow et al., 1913; Flint et al., 1959; Krygowski, 1963; Campbell, 1967a, b; Hodgson et al., 1984; Klassen, 1993; Priamonosov et al., 2000; Follestad and Bergstrøm, 2004) or necessarily highly-selective ground-truthing of remote mapping (e.g. Kleman et al., 1997, 2010; Golledge and Stoker, 2006; Stokes et al., 2013; Darvill et al., 2014; Stroeven et al., 2016; Pearce et al., 2018).

2.2 The field mapping process

Field mapping should ideally begin with systematic traverses of the study area – sometimes referred to as a ‘walk-over’ (e.g. Demek, 1972; Otto and Smith, 2013) – to get a sense of the scale of the study area and ensure that subtle features of importance, such as the location and orientation of ice-flow directional indicators (e.g. flutes, striae, roches moutonnées and ice-moulded bedrock), are not missed. In a palaeo-ice sheet context, mapping the location and orientation of striae in the field may be of most interest as these can provide information on multiple (local) ice flow directions, of which not all are recorded in the pattern of elongated bedforms mappable from remotely-sensed data (cf. Kleman, 1990; Smith and Knight, 2011). Similarly, in a contemporary outlet glacier context, flutes are an important indicator of ice flow direction – sometimes of annual ice flow trajectories of glacier margins (cf. Chandler et al., 2016a; Evans et al., 2017) – but due to their subtlety they may only be identifiable in the field (e.g. Jónsson et al. 2014).

Traversing should ideally start from higher ground, where an overview can be gained, and proceed by crossing a valley axis (or a cirque floor, for example) many times to enable the viewing and assessment of landforms from as many perspectives, angles and directions as possible (cf. Demek, 1972). In addition to systematic traverses, landform assemblages in, for example, individual

valleys/basins should ideally be viewed from a high vantage point in low light (e.g. Benn, 1990). Depending on the location and orientation of landforms, it may be beneficial to see the same area either (i) early in the morning or late in the afternoon/evening due to longer shadows, or (ii) both in the morning and afternoon/evening due to the changing position of longer shadows. These procedures ensure that apparent dimensions and orientations, which are influenced by perception under different viewing angles and daylight conditions, can be taken into account in descriptions and interpretations. This approach circumvents potential complications relating to subtle features that may only be visible from one direction or certain angles.

The location of features should be recorded on field maps or imagery (e.g. aerial photograph) extracts with reference to ‘landmarks’ that are clearly identifiable both in the field and on the base maps/imagery, such as distinct changes in contour-line inflection, river bends, confluences, prominent bedrock exposures and large ridges or mounds (Lukas and Lukas, 2006; Boston, 2012a, b). Where geomorphological features are small, background relief is low and/or conspicuous reference points are absent, a network of mapped reference points can be established by either taking a series of cross-bearings on prominent features using a compass (e.g. Benn, 1990) or by verifying locations using a handheld GNSS (e.g. Lukas and Lukas, 2006; Boston, 2012a, b; Brynjólfsson et al., 2014; Jónsson et al. 2014; Lovell, 2014; Pearce et al., 2014; van der Bilt et al., 2016). The latter is useful for recording the location of point-data such as striae, erratic or glacially-transported boulders, and sediment exposures (cf. Lukas and Lukas, 2006; Boston, 2012a, b; Pearce et al., 2014). Additional information between known reference points can then be interpolated and marked on the geomorphological map.

Establishing the size of landforms and features and plotting them on the map as accurately as possible is of crucial importance, and in addition to the inflections of contours (which may mark the location and boundaries of prominent ridges, for example), the mapper may pace out and/or estimate lengths, heights and widths. For larger landforms, or those masked by forest, walking around the perimeter of landforms and establishing a GNSS-marked ‘waypoint-trail’ is a good first approximation.

The strategy outlined above offers a broad perspective on the overall landform pattern and ensures accurate representation of landforms on field maps. To ensure accurate genetic interpretation of individual landforms, and the landscape as a whole, this field mapping strategy should ideally form part of an iterative process of observation and interpretation whilst still in the field (see Section 2.3, below).

2.3 Interpreting glacial landforms

In the preceding section, we focused on the technical aspects of field mapping and the means of recording glacial landforms. However, geomorphological mapping typically forms the foundation of process-oriented and palaeoglaciological reconstruction studies (see Section 1.1) and should, therefore, be embedded within a process of observation and interpretation. Definitive interpretation of glacial landforms, and glacial landscapes as a whole, can rarely be made on the basis of surface morphology alone. Additional strands of field evidence may become highly relevant, if not essential, depending on the objectives of the individual project: sedimentological data are crucial to interpreting processes of landform formation and glacier dynamics (e.g. Lukas, 2005; Benn and Lukas, 2006; Benediktsson et al., 2010, 2016; Chandler et al., 2016a), whilst chronological data are fundamental to robust palaeoglaciological reconstructions and related palaeoclimatic studies (e.g. Finlayson et al., 2011; Gribenski et al., 2016; Hughes et al., 2016; Stroeven et al., 2016; Bendle et al., 2017b; Darvill et al., 2017). Moreover, time and resources are limited and pragmatism necessary. Thus, observations must be targeted efficiently and effectively, in line with the research aims.

Much field-based research adopts an inductive approach, in which observations are collected and used to argue towards a particular conclusion. This is a valid approach at the exploratory stage of research, but deeper understanding of a landscape requires a more iterative process, in which data collection is conducted within a framework of hypothesis generation and testing. For this reason, it is useful to adopt a number of alternative working hypotheses (Chamberlin, 1897) that can be tested and gradually eliminated, following the principle of falsification (Popper, 1972). This process is best conducted in the field while it is possible to make key observations to test an interpretation, especially if the field site is remote and expensive to re-visit.

Following initial data collection, preliminary interpretations can be used to predict the outcome of new observations, which can then be used to test and refine the interpretation. Well-framed hypotheses allow an investigator to anticipate other characteristics of a glacial landscape, then to test those predictions by targeted investigation of key localities (see Benn, 2006). For example, the presence of a certain group of landforms (e.g. moraines trending downslope into a side valley) can be used to formulate hypotheses (e.g. blockage of the side-valley by glacier ice), which in turn can be used to predict the presence of other sediment-landform associations in a particular locality (e.g. lacustrine sediments or shoreline terraces in the side-valley). Further detailed geomorphological mapping (and sedimentological analyses) in that area would then allow testing and falsification of the alternative working hypotheses. Iterations of this process during field mapping enable an increasingly detailed and robust understanding of the glacier system to be constructed. This coupled inductive-deductive approach is much more powerful than a purely inductive process: narratives that ‘explain’ a set of observations can appear very persuasive, even self-evident, but there may be other narratives that are also consistent with the same observations (cf. Popper, 1972).

Process-form models are useful tools in this inductive-deductive approach to landscape interpretation. In particular, *landsystem* or *facies models* make explicit links between landscape components and genetic processes, providing structure and context for data collection and interpretation (e.g. Eyles, 1983; Brodzikowski and van Loon, 1991; Evans, 2003a; Benn and Evans, 2010). At best, process-form models are not rigid templates or preconceived categories into which observations are forced, but a flexible set of possibilities that can guide, shape and enrich investigations (Benn and Lukas, 2006). For example, preliminary remote mapping may reveal features that suggest former glacier lobes may have surged (e.g. Lovell et al., 2012). Systematic study of sediment-landform assemblages, sediment exposures and other evidence, with reference to modern analogues (e.g. Evans and Rea, 2003), allows this idea to be rigorously evaluated in a holistic context (e.g. Darvill et al., 2017). This then opens up new avenues for research in a creative and open-ended process.

This inductive-deductive approach to interpreting glacial landscapes and events should be embedded as part of the geomorphological mapping process (see Section 6). When dealing with palaeo-ice sheets, such field-based investigations may be guided by (existing) remote mapping. In alpine and plateau-style ice mass settings, sedimentological and chronological investigations should ideally form an integral part of field surveys.

3. Remote mapping methods

In the following sections, we review the principal remote mapping approaches employed in glacial geomorphological research, with analogue (or hard-copy) remote mapping (Section 3.1) and digital remote mapping (Section 3.2) considered separately. We give an overview of a number of datasets used for digital remote (i.e. GIS-based) mapping, namely satellite imagery (see Section 3.2.2.1), aerial photographs (see Section 3.2.2.2), digital elevation models (see Section 3.2.2.3), freeware virtual globes (see Section 3.2.2.4) and UAV-captured imagery (see Section 3.2.2.5). Each individual section provides a brief outline of the historical background and development of the methods, and we discuss the individual approaches in a broadly chronological order. Section 3.3 then provides an overview of image processing techniques, highlighting that pragmatic solutions are often required.

We focus principally on remotely-sensed datasets relevant to terrestrial (onshore) glacial settings in the following sections, since submarine (bathymetric) datasets and mapping of submarine glacial landforms have been subject to recent reviews elsewhere (see Dowdeswell et al., 2016; Batchelor et al., 2017). Nevertheless, we do acknowledge that the emergence of geophysical techniques to investigate submarine (offshore) glacial geomorphology is a major development over the last two decades. Similarly, the emergence of geophysical datasets of sub-ice geomorphology in the last

decade or so has been revolutionary, particularly in relation to subglacial bedforms (see Stokes, 2018). Many of the issues we discuss in relation to mapping from DEMs are transferable to those environments.

3.1 Analogue remote mapping

3.1.1 Background and applicability of analogue remote mapping

Geomorphological mapping from analogue (hard-copy) aerial photographs became a mainstream approach in glacial geomorphology in the 1960s and 1970s, with early proponents including, for example, the Geological Survey of Canada (e.g. Craig, 1961, 1964; Prest et al., 1968) and UK-based researchers examining the Quaternary geomorphology of upland Britain (e.g. Price, 1961, 1963; Sissons, 1967, 1977a, b, 1979a, b; Sugden, 1970) and contemporary glacial landsystems (e.g. Petrie and Price, 1966; Price, 1966; Welch, 1966, 1967, 1968; Howarth, 1968; Howarth and Welch, 1969a, b). The latter research on landsystems in Alaska and Iceland was particularly pioneering in that it exploited a combination of aerial photograph interpretation, surveying techniques and early photogrammetry (see Evans, 2009, for further details).

Despite continued development of remote sensing technologies and the availability of digital aerial photographs (see Section 3.2.2.2), analogue stereoscopic aerial photographs are still used for glacial geomorphological mapping (e.g. Hättestrand, 1998; Benn and Ballantyne, 2005; Lukas et al., 2005; Boston, 2012a, b; Evans and Orton, 2015). Additionally, the availability of high-quality photogrammetric scanners means that archival, hard-copy aerial photographs can be scanned at high resolutions, processed using digital photogrammetric methods and subsequently used for on-screen digitisation (Section 3.2; e.g. Bennett et al., 2010; Jónsson et al., 2014). As with field mapping, the interpretation of analogue aerial photographs is primarily used for mapping alpine- and plateau-style ice mass geomorphological imprints. Historically, analogue aerial photograph interpretation was extensively used for mapping palaeo-ice sheet geomorphological imprints, particularly by the Geological Survey of Canada, who combined aerial photograph interpretation with detailed ground checking and helicopter-based surveys (e.g. Craig, 1961, 1964; Hodgson et al., 1984; Aylsworth and Shilts, 1989; Dyke et al., 1992; Klassen, 1993; Dyke and Hooper, 2001). This approach has largely been superseded by satellite imagery and DEM interpretation in palaeo-ice sheet settings (see Section 5.1) but is applied in palaeo-ice sheet contexts for more detailed mapping of selected/complex areas (e.g. Dyke, 1990; Kleman et al., 2010; Stokes et al., 2013; Storrar et al., 2013; Darvill et al., 2014; Evans et al., 2014).

3.1.2 Mapping from analogue datasets

For glacial geomorphological mapping purposes, vertical panchromatic aerial photographs have traditionally been employed, with pairs of photographs (stereopairs) viewed in stereo using a stereoscope (with magnification) (e.g. Melander, 1975; Horsfield, 1983; Krüger, 1994; Kleman et al., 1997; Hättestrand, 1998; Evans and Twigg, 2002; Benn and Ballantyne, 2005; Lukas and Lukas, 2006; Boston, 2012a, b; Chandler and Lukas, 2017). During aerial surveys, longitudinally-overlapping photographs along the flight path (endlap $\geq 60\%$) are captured in a series of laterally-overlapping parallel strips (sidelap $\geq 30\%$), with the two different viewing angles of the same area resulting in the stereoscopic effect (due to the principle of parallax; see Lillesand et al., 2015, for further details). This form of aerial photograph interpretation has been demonstrated to be a particularly valuable tool for determining the exact location, shape and planform of small features in glaciated terrain (e.g. Ballantyne, 1989, 2002, 2007a, b; Bickerton and Matthews, 1992, 1993; Lukas and Lukas, 2006; Boston, 2012a, b), provided the photographs are of appropriate scale, quality and tonal contrast (cf. Benn, 1990; Benn et al., 1992).

Mapping from hard-copy aerial photographs is undertaken by drawing onto acetate sheets (transparency films) whilst viewing the aerial photographs through a stereoscope, with the acetate overlain on one photograph of a stereopair (Figure 2). Ideally, mapping should be conducted using a super-fine pigment liner with a nib size of 0.05 mm to enable small features to be mapped. Even so, it may still be necessary to compromise on the level of detail mapped; for example, meltwater channels between ice-marginal moraines have been left off maps in some studies due to map scale, with the associated text describing chains of moraines interspersed with meltwater channels (e.g. Benn and Ballantyne, 2005; Lukas, 2005).

Examining stereopairs from multiple sorties ('flight missions') in parallel or in combination with digital aerial photographs may be beneficial and help alleviate issues such as localised cloud cover, snow cover, poor tonal contrast, afforestation and anthropogenic developments (e.g. Horsfield, 1983; Bennett, 1991; McDougall, 2001). Additionally, it is advantageous to examine stereopairs multiple times – preferably before and after field mapping – to increase feature identification and improve the accuracy of genetic interpretations (Lukas and Lukas, 2006; Sahlin and Glasser, 2008). When conducting mapping over a large area with multiple stereopairs, examining stereopairs from a sortie 'out of sequence' (i.e. not mapping from consecutive pairs of photographs) may provide a means of internal corroboration and ensure objectivity and robustness (Bennett, 1991).

In order to reduce geometric distortion, which increases towards the edges of aerial photographs due to the central perspective (Lillesand et al., 2015), it is advisable to keep the areas mapped onto the acetate as close as possible to the centre of one aerial photograph of a stereopair (Kronberg, 1984; Lukas, 2002, 2005a; Evans and Orton, 2015). These hand-drawn overlays can subsequently be

scanned at high resolutions and then georeferenced and digitised using GIS and graphics software (see also Section 3.3.1; e.g. Lukas and Lukas, 2006; Boston, 2012a, b).

3.2 Digital remote mapping

3.2.1 Background and applicability of digital remote mapping

The development of GIS software packages (e.g. commercial: *ArcGIS*; open source: *QGIS*) and the proliferation of digital imagery, particularly freely available satellite imagery, have undoubtedly been the most significant developments in glacial geomorphological mapping. GIS packages have provided platforms and tools for visualising, maintaining, manipulating and analysing vast quantities of remotely-sensed and geomorphological data (cf. Gustavsson et al., 2006, 2008; Napieralski et al., 2007b). Their use in combination with digital imagery allows geomorphological features to be mapped directly in GIS software (Figure 3), with individual vector layers created for each geomorphological feature. Moreover, the availability of digital imagery enables the mapper to alter the viewing scale instantaneously and switch between various datasets/types, allowing for a flexible but systematic approach.

Digital mapping (on-screen digitisation) also provides georeferenced geomorphological data, which has two important benefits: Firstly, these data can easily be used to derive landform metrics (e.g. Hättestrand et al., 2004; Clark et al., 2009; Spagnolo et al., 2010, 2014; Storrar et al., 2014; Ojala et al., 2015; Dowling et al., 2016; Ely et al., 2016a, 2017a); and, secondly, these data can be seamlessly incorporated into wider, regional-scale GIS compilations (e.g. Bickerdike et al., 2016; Stroeve et al., 2016; Clark et al., 2018a). Additionally, digital remote mapping allows the user to record attribute data (e.g. data source) tied to individual map (vector) layers, which can be useful for large compilations of previously published mapping (e.g. Bickerdike et al., 2016; Clark et al., 2018a). Such compendia help to circumvent issues relating to the often-fragmented nature of geomorphological evidence (i.e. numerous spatially separate studies) and identify gaps in the mapping record. Once assembled across large areas, they also enable evidence-based reconstructions of entire ice sheets and regional ice sheet sectors (see Clark et al., 2004, 2018a). Indeed, the ongoing open access data revolution in academia and the increasing publication/availability of mapping output (in the form of GIS files; e.g. Finlayson et al., 2011; Darvill et al., 2014; Bickerdike et al., 2016; Bendle et al., 2017a), means that geomorphological mapping can have wider impact beyond individual local to regional studies.

3.2.2 Datasets for digital remote mapping

There is now a plethora of remotely-sensed datasets covering a wide range of horizontal resolutions (10^{-2} to 10^2 m), enabling the application of digital mapping (in some form) to all glacial settings. We

provide an overview of the principal datasets used in digital mapping below, with mapping approaches in specific glacial settings reviewed in Section 5.

3.2.2.1 Satellite imagery. The development of satellite-based remote sensing in the 1970s and subsequent advances in technology have revolutionised understanding of glaciated terrain, particularly with respect to palaeo-ice sheet geomorphology and dynamics (see Section 5.1; Clark, 1997; Stokes, 2002; Stokes et al., 2015). The potential of satellite imagery was first demonstrated by the pioneering work of Sugden (1978), Andrews and Miller (1979) and Punkari (1980), with the availability of large-area view (185 km x 185 km) Landsat Multi-Spectral Scanner (MSS) images affording a new perspective of glaciated regions. These allowed a single analyst to systematically map ice sheet-scale (1:45,000 to 1: 1,000,000) glacial geomorphology (e.g. Boulton and Clark, 1990a, b) in a way that previously would have required the painstaking mosaicking of thousands of aerial photographs (e.g. Prest et al., 1968). Since the 1980s, there has been an explosion in the use of satellite imagery for glacial geomorphological mapping and there is now a profusion of datasets available (Table 1). Importantly, many of these sensors capture multispectral data, which can enhance landform detection through image processing and the use of different band combinations (see Section 3.3.2). The uptake of satellite imagery has coincided with improvements in the availability and spatial and spectral resolution of satellite datasets globally, with Landsat (multispectral: 30 m; panchromatic: 15 m), ASTER (15 m), Sentinel-2 (10 m) and SPOT (up to 1.5 m) images proving the most popular. More recently, satellite sensor advancements have enabled the capture of satellite images with resolutions comparable to aerial photographs (Figure 4; e.g. SPOT6/7, QuickBird, WorldView). Thus, these datasets are also suitable for mapping typically smaller and/or complex glacial landforms produced by cirque glaciers, valley glaciers and icefield/ice-cap outlets (e.g. Chandler et al., 2016a; Evans et al., 2016b; Ewertowski et al., 2016; Gribenski et al., 2016; Małeckki et al., 2018).

In general, as better-resolution imagery has become more widely available at low to no cost, older, coarser-resolution datasets (e.g. Landsat MSS: 60 m) have largely become obsolete. Nevertheless, Landsat data (TM, ETM+ and OLI: 15 to 30 m) are still the standard data source for ice-sheet scale mapping, with the uptake of high-resolution commercial satellite imagery still relatively slow in such studies. This is primarily driven by the cost of purchasing high-resolution commercial datasets, making freely-available imagery such as Landsat a valuable resource. In addition, archival satellite data afford time-series of multi-spectral images that may facilitate assessments of geomorphological changes through time; for example, fluctuations in highly dynamic (surging or rapidly retreating) glacial systems (e.g. Flink et al., 2015; Jamieson et al., 2015). Conversely, for smaller research areas (e.g. for a single valley or foreland), high-resolution satellite imagery is becoming an increasingly viable option, with prices for georeferenced and orthorectified products comparable to those for digital aerial photographs (see Section 3.2.2.2). This also has the benefit of saving time on

photogrammetric processing, with many vendors providing consumers with various processing options. Consequently, on-demand, high-resolution (commercial) satellite imagery will inevitably come into widespread usage, where costs are not prohibitive. Alternatively, freeware virtual globes and web mapping services (e.g. *Bing Maps*, *Google Earth*) offer valuable resources for free visualisation of such high-resolution imagery (see Section 3.2.2.4).

3.2.2.2 Digital aerial photographs. With improvements in technology, high-resolution (ground resolution <0.5 m per pixel) digital copies of aerial photographs have become widely available and used for glacial geomorphological mapping (e.g. Brown et al., 2011a; Bradwell et al., 2013; Brynjólfsson et al., 2014; Jónsson et al., 2014; Pearce et al., 2014; Schomacker et al., 2014; Chandler et al., 2016a; Evans et al., 2016c; Lardeux et al., 2016; Lønne, 2016; Allaart et al., 2018). Indeed, digital aerial photographs, along with scanned copies of archival aerial photographs, are now more widely used than hard-copy stereoscopic aerial photographs, particularly in modern glacial settings. Additionally, the introduction of UAV technology in recent years has allowed sub-decimetre resolution aerial photographs to be captured on demand (see Section 3.2.2.5). Aside from their typically superior resolution/scale compared with hard-copy aerial photographs, a key advantage of aerial photographs in digital format is the ability to produce orthorectified aerial photograph mosaics (or ‘orthophotographs’) and DEMs with low root mean square errors (RMSEs <1 m; see Section 4.4), when combined with ground control points collected using surveying equipment (e.g. Kjær et al. 2008; Bennett et al., 2010; Schomacker et al., 2014; Chandler et al., 2016b; Evans et al., 2017). These photogrammetric products can then be used for on-screen digitisation and generation of georeferenced geomorphological mapping (Figure 5), as outlined above.

Typically, digital aerial photographs are captured by commercial surveying companies (e.g. Loftmyndir ehf, Iceland; Getmapping, UK), meaning that they may be expensive to purchase and costs may be prohibitive for large study areas. This is in contrast to hard-copy (archival) aerial photographs that are often freely available for viewing in national collections. Additionally, digital aerial photographs are not readily viewable in stereo with a standard desktop setup, although on-screen mapping in stereoscopic view is possible on workstations equipped with stereo display and software such as *BAE Systems SOCET SET* (e.g. Kjær et al., 2008; Benediktsson et al., 2009). However, this approach is not applicable to orthophotographs. An alternative approach is to visualise orthophotographs in 3D by draping them over a DEM (see Section 3.2.2.3) in GIS software such as *ESRI ArcScene* or similar (Figure 6; e.g. Benediktsson et al., 2010; Jónsson et al., 2014; Schomacker et al., 2014; van der Bilt et al., 2016). Three-dimensional assessment in *ArcScene*, parallel to mapping in *ArcMap*, may aid in landform detection, delineation and interpretation.

3.2.2.3 *Digital Elevation Models*. Over the last ~15 years there has been increasing use of DEMs in glacial geomorphology, particularly for mapping at the ice-sheet scale (e.g. Glasser and Jansson, 2008; Hughes et al., 2010; Ó Cofaigh et al., 2010; Evans et al., 2014, 2016d; Principato et al., 2016; Ojala, 2016; Stokes et al., 2016a; Mäkinen et al., 2017; Norris et al., 2017). DEMs are raster-based models of topography that record absolute elevation, with each pixel or grid cell in a DEM representing the average height for the area it covers (Clark, 1997; Smith et al., 2006). Terrestrial DEMs can be generated by a variety of means, including from surveyed contour data, directly from stereo imagery (aerial photographs, satellite and UAV-captured imagery), or from air- and space-borne radar and LiDAR systems (Smith and Clark, 2005). An important recent development in this regard has been the Surface Extraction with TIN-based Search-space Minimization (SETSM) algorithm for automated extraction of DEMs from stereo satellite imagery (Noh and Howat, 2015), which has been used to generate the ArcticDEM dataset (<https://www.pgc.umn.edu/data/arcticdem/>). However, SETSM DEMs may contain systematic vertical errors that require correction (e.g. Carrivick et al., 2017; Storrar et al., 2017).

The majority of DEMs with national- to international-scale coverage (Table 2) typically have a coarser spatial resolution than aerial photographs and satellite imagery and represent surface elevations rather than surface reflectance. As a result, it may be difficult to identify glacial landforms produced by relatively small ice masses (cirque glaciers, valley glaciers and icefield outlets), precluding detailed mapping of their planforms (cf. Smith et al., 2006; Hughes et al., 2010; Brown et al., 2011a; Boston, 2012a, b; Pearce et al., 2014). Conversely, these DEMs can be particularly valuable for mapping glacial erosional features (e.g. glacial valleys, meltwater channels), as well as major glacial depositional landforms produced by larger ice masses (e.g. Greenwood and Clark, 2008; Heyman et al., 2008; Livingstone et al., 2008; Hughes et al., 2010; Morén et al., 2011; Barr and Clark, 2012; Stroeven et al., 2013; Turner et al., 2014a; Margold et al., 2015a; Blomdin et al., 2016a, b; Lindholm and Heyman, 2016; Mäkinen et al., 2017; Storrar and Livingstone, 2017). However, the recent development of UAV (see section 3.2.2.5) and LiDAR technologies have allowed the generation of very high resolution DEMs (<0.1 m), enabling the application of DEMs to map small glacial landforms (e.g. Evans et al., 2016a; Ewertowski et al., 2016; Ely et al., 2017). We anticipate national-scale LiDAR DEMs becoming widely-used in the future, with a number of nations recently releasing or currently capturing/processing high horizontal resolution (≤ 2 m) LiDAR data (Table 2; e.g. Dowling et al. 2013; Johnson et al. 2015).

Although the principal focus of this contribution is terrestrial/onshore glacial geomorphological mapping, it is worth highlighting here that the availability of spatially-extensive bathymetric charts, such as the General Bathymetric Chart of the Oceans (GEBCO) and International Bathymetric Chart of the Arctic Ocean (IBCAO: Jakobsson et al., 2012), and high-resolution, regional (often industry-

acquired) bathymetric data has been an important development in submarine/offshore glacial geomorphological mapping. This has enabled the gridding of DEMs to map submarine glacial geomorphological imprints (see Dowdeswell et al., 2016), markedly enhancing understanding of palaeo-ice sheets in marine sectors (e.g. Ottesen et al., 2005, 2008a, 2016; Bradwell et al., 2008; Winsborrow et al., 2010, 2012; Livingstone et al., 2012; Ó Cofaigh et al., 2013; Hodgson et al., 2014; Stokes et al., 2014; Margold et al., 2015a, b; Greenwood et al., 2017) and modern tidewater (often surging) glaciers (e.g. Ottesen and Dowdeswell, 2006; Ottesen et al., 2008b, 2017; Robinson and Dowdeswell, 2011; Dowdeswell and Vazquez, 2013; Flink et al., 2015; Streuff et al., 2015; Allaart et al., 2018). In addition, recent years have seen the production of DEMs of sub-ice topography from geophysical datasets (radar and seismics) at spatial resolutions suitable for identifying and mapping bedforms (see King et al., 2007, 2009, 2016a; Smith et al., 2007; Smith and Murray, 2009). This work has advanced understanding of the evolution of bedforms beneath Antarctic ice streams, providing important genetic links between the formation of landforms beneath modern ice sheets and those left behind by palaeo-ice masses (Stokes, 2018). The interested reader is directed to recent reviews (Livingstone et al., 2012; Ó Cofaigh, 2012; Stokes et al., 2015; Dowdeswell et al., 2016; Batchelor and Dowdeswell, 2017; Stokes, 2018) for further discussion on the importance of geophysical evidence for understanding ice sheet extent and dynamics.

3.2.2.4 Freeware virtual globes. The advent of freeware virtual globes (e.g. *Google Earth*, *NASA Worldwind*) and web mapping services (e.g. *Bing Maps*, *Google Maps*) have provided platforms for free visualisation of imagery from various sources and low-cost mapping resources. A key benefit of virtual globes is the ability to visualise imagery and terrain in 3D and from multiple viewing angles, which may aid landform detection when used in conjunction with other datasets and software (e.g. Heyman et al., 2008; Bendle et al., 2017a). Moreover, a number of virtual globes and web mapping services have the ability to link with other freeware and open-source programmes; for example, free plugins are available to import *Google Earth* and *Bing Maps* imagery into the open-source GIS software package *QGIS*. Thus, a mapper can combine freely available, often high-resolution (e.g. SPOT6/7, QuickBird, WorldView), imagery and the capabilities of GIS technology without the expense associated with commercial imagery and software (see Sections 3.2.2.1 and 3.2.2.2).

The most widely-used virtual globe is *Google Earth*, with a ‘professional’ version (*Google Earth Pro*) freely available since 2015 (see Mather et al., 2015, for a review). An increasing number of glacial geomorphological studies are noting the use of *Google Earth* (but not necessarily the imagery type) as a mapping tool (see Table 1), principally to cross-check mapping conducted from other imagery. However, some studies have also utilised the built-in digitising tools for mapping (e.g. Margold and Jansson, 2011; Margold et al., 2011; Fu et al., 2012). There is a compromise on the functionality of freeware virtual globes and digitisation tools are often not as flexible and/or user-friendly, but these

can be counteracted by the option to import imagery into GIS software. In the case of *Google Earth*, it is also possible to export Keyhole Markup Language (KML) files that can be used for subsequent analyses and map production in GIS software (following file conversion). Open access remotely-sensed datasets are also available through commercial GIS software, with high resolution satellite imagery (e.g. GeoEye-1, SPOT-5, WorldView) available for mapping through the in-built ‘World Imagery’ service in *ESRI ArcGIS* (e.g. Bendle et al., 2017a).

Despite the benefits, some caution is necessary when using freeware virtual globes as there may be substantial errors in georeferencing of imagery, which users cannot account/correct for. Moreover, dating of imagery is not necessarily clear or accurate (Mather et al., 2015; Wyshnytzky, 2017). The latter may not be a concern if mapping in a palaeoglaciological setting, whilst any georeferencing errors may not be as significant if mapping broad patterns at the ice sheet scale. Conversely, errors associated with freeware mapping may be significant when comparing imagery from different times and/or when mapping in highly dynamic, contemporary glacial environments. Aside from these potential issues, limitations are imposed by pre-processing of imagery, with no option to, for example, modify band combinations to enhance landform detection (see Section 3.3.2).

3.2.2.5 UAV-captured imagery. The recent emergence of UAV technology provides an alternative method for the acquisition of very high-resolution (<0.1 m per pixel) geospatial data that circumvents some of the issues associated with more established approaches, particularly in relation to temporal resolution and the high-cost of acquiring commercial remotely-sensed data (see also Smith et al., 2016a). This method provides a rapid, flexible and relatively inexpensive (following the initial acquisition of the UAV and associated software) means of acquiring up-to-date imagery at an unprecedented spatial resolution and is becoming increasingly employed in glacial research (Figure 7; Rippin et al., 2015; Ryan et al., 2015; Chandler et al., 2016b; Ewertowski et al., 2016; Tonkin et al., 2016; Westoby et al., 2016; Ely et al., 2017; Allaart et al., 2018). UAV-captured images are processed using Structure-from-Motion (SfM) photogrammetry techniques, with *Agisoft Photoscan* being the most common software in use at present (e.g. Chandler et al., 2016b; Evans et al., 2016a; Ely et al., 2017; Allaart et al., 2018). This methodology has enabled the production of sub-decimetres resolution orthophotographs and DEMs with centimetre-scale error values (RMSEs <0.1 m; see Section 4.4) for glacial geomorphological mapping (e.g. Evans et al., 2016a; Ely et al., 2017). Although surveying of ground control points is still preferable for processing UAV-captured imagery, a direct georeferencing workflow (see Turner et al., 2014b, for further details) is capable of producing reliable geospatial datasets from imagery captured using consumer-grade UAVs and cameras, without the need for expensive survey equipment (see Carbonneau and Dietrich, 2017).

The use of UAVs will be valuable in future glacial geomorphological research due to their flexibility and low-cost. In particular, UAVs open up the exciting possibility of undertaking repeat surveys at high temporal (sub-annual to annual) resolutions in modern glacial settings (Immerzeel et al., 2014; Chandler et al., 2016b; Ely et al., 2017). Multi-temporal UAV imagery will enable innovative geomorphological studies on issues such as (i) the modification and preservation potential of landforms over short timescales (Ely et al., 2017), (ii) the frequency of ice-marginal landform formation, particularly debates on sub-annual to annual landform formation (Chandler et al., 2016b), and (iii) changes in process-form regimes at contemporary ice-margins (Evans et al., 2016a).

Using UAVs to capture aerial imagery is not without challenges, particularly in relation to the challenge of intersecting suitable weather conditions in modern glacial environments: many UAVs are unable to fly in high windspeeds, whilst rain can infiltrate electrical components and create hazy imagery (Ely et al., 2017). Flight times and areal coverage are also limited by battery life, with some battery packs permitting as little as 10 minutes per flight. There are also legal considerations, with the use of UAVs prohibited in some localities/countries or requiring licenses/permits. Moreover, there may be restrictions on flying heights and UAVs may need to be flown in visual line of sight, further limiting areal coverage. Nevertheless, we envisage UAV technology becoming more widespread and a key tool in high-resolution glacial geomorphological investigations, especially if future technological developments can increase the range of conditions in which UAVs can be flown. In future, it is likely that UAV technology will be primarily used for investigating short-term changes across relatively small areas.

3.3 Image processing for mapping

An important part of geomorphological mapping is processing remotely-sensed datasets in preparation for mapping, but this is often given limited prominence in glacial geomorphological studies. Crucially, processing of remotely-sensed data aids the identification of glacial landforms and ensures accurate transfer of geomorphological data from the imagery. In the sections below, we provide a brief overview of image processing solutions for aerial photographs (Section 3.3.1), satellite imagery (Section 3.3.2) and DEMs (Section 3.3.3). Reference is made to common processing techniques used to remove distortion and displacement evident in aerially-captured imagery (see Campbell and Wynne (2011) and Lillesand et al. (2015) for further details), but these are not discussed in detail for reasons of brevity and clarity. However, a detailed workflow diagram outlining the potential procedures for a range of scenarios (depending on data, resources and time) is available as Supplementary Material. We emphasise that compromises and pragmatic solutions are necessary, particularly in the case of aerial photographs, as the ‘idealised’ scenario is frequently not an option due to data limitations or logistical constraints.

3.3.1 Aerial photograph processing

Aerial photographs contain varying degrees of distortion and displacement owing to their central (or perspective) projection. Geometric distortion is related to radial lens distortion and refraction of light rays in the atmosphere. Additional displacement occurs as a result of the deviation of the camera from a vertical position (caused by roll, pitch and yaw of the aircraft), relief and curvature of the Earth. Non-corrected aerial photographs are therefore characterised by relief displacement and scale variations, which increase towards the edges of the photograph (see Campbell and Wynne (2011) and Lillesand et al. (2015) for further details). Thus, it is necessary to apply geometric corrections to aerial photographs before geomorphological mapping.

Ideally, aerial photographs would be corrected using stereoscopic (or conventional) photogrammetric processing in software packages such as *Imagine Photogrammetry* (formerly *Leica Photogrammetry Suite*, or *LPS*). This approach involves the extraction of quantitative elevation data from stereoscopic (overlapping) imagery to generate DEMs and orthorectified imagery (see also Section 3.2). Internal and external parameters, along with the location of GCPs, are used to establish the relationship between the position of the images and a ground coordinate system (e.g. Kjær et al., 2008; Bennett et al., 2010). However, this approach may be impractical and unsuitable in many glacial settings; for example, it is unrealistic to collect GCPs using (heavy) survey equipment (e.g. RTK-GPS) in former plateau icefield and ice-cap settings due to their location (remote, upland environments) and the size of the study area (and thus quantity of aerial photographs and GCPs required). Moreover, camera calibration data (focal length, fiducial marks, principal point coordinates and lens distortion) are frequently unavailable or incomplete for archive datasets, and the process is also not applicable to acetate overlays. Thus, orthorectification of imagery – correction of geometric distortions in both the horizontal (x and y coordinates) and vertical (z coordinates) dimensions – is typically precluded over larger areas, although it may be possible to employ this approach for individual cirque basins, valleys and glacier forelands (e.g. Wilson, 2005; Bennett et al., 2010; Chandler et al., 2016a). Consequently, pragmatic solutions are required for georectification of imagery, i.e. the process of transforming and projecting imagery to a (local) planar coordinate system. Several approaches have been used to overcome this and we briefly outline these below in relation to analogue aerial photographs (Section 3.3.1.1) and digital aerial photographs (Section 3.3.1.2).

3.3.1.1 Analogue aerial photograph processing. A potential pragmatic solution to correcting analogue (hard-copy) aerial photographs is to georeference scanned copies of acetate overlays or the original aerial photographs to reference points on other forms of (coarser) georeferenced digital imagery (if available; e.g. DEMs, orthorectified radar images, satellite images). The scanned images can then be georectified and resampled using the georeferencing functions within GIS and remote sensing

programmes such as *ArcGIS* or *Erdas Imagine* (cf. Boston, 2012a, for further details). This approach is particularly useful where hard-copy aerial photographs are used in combination with (coarser) digital imagery. Using this procedure, georeferenced acetate overlays of Quaternary features in the Scottish Highlands have been produced with RMSE values ranging between 2.71 m and 7.82 m (Boston, 2012a), comparable to archival aerial photographs that have been processed using conventional photogrammetric techniques (e.g. Bennett et al., 2010).

The above georectification method works best if relatively small areas are mapped on one acetate. This is because radial distortion increases towards the edges of aerial photographs and, as such, presents a significant problem to matching reference points when large areas have been mapped. From our experience, we estimate the maximum effective area that can be corrected without the danger of mismatches at $\sim 6 \text{ km}^2$. However, this figure depends on the terrain conditions and would have to be smaller in high mountain areas where relief distortion is increased due to greater differences between valleys and adjacent peaks (Lillesand et al., 2015). The mapped area could, conversely, be somewhat larger in low-relief terrain as objects are roughly equally far away from the camera lens over larger areas and thus subject to less distortion (Kronberg, 1984; Lillesand et al., 2015). The aforementioned constraints might seem to make georectification from hard-copy aerial photographs a laborious process, but this is counterbalanced by being able to record small landforms in great detail due to the high-resolution 3D visualisation allowed by stereopairs.

3.3.1.2 Digital aerial photograph processing. Digital aerial photographs can be georeferenced within GIS and remote sensing software following a similar process to that outlined in Section 3.3.1.1, i.e. digital aerial photographs can be georeferenced to other forms of (coarser) georeferenced imagery. Alternatively, SfM photogrammetry can be used to produce orthophotographs and DEMs from digital aerial photographs, which partly circumvents issues relating to incomplete or absent camera calibration data (e.g. Chandler et al., 2016a; Evans et al., 2016e, 2017; Tonkin et al., 2016; Midgley and Tonkin, 2017; Mertes et al., 2017). SfM photogrammetry functions under the same basic principles as stereoscopic photogrammetry but there are fundamental differences: the geometry of the ‘scene’, camera positions and orientation are solved automatically in an arbitrary ‘image-space’ coordinate system without the need to specify either the 3D location of the camera or a network of GCPs with known ‘object-space’ coordinates (cf. Westoby et al., 2012; Carrivick et al., 2016; Smith et al., 2016a, for further details). However, positional data (GCPs) are still required to process the digital photographs for geomorphological mapping, i.e. to assign the SfM models to an ‘object-space’ coordinate system. Ideally, this should be conducted through ground control surveys (see above), but a potential pragmatic solution is to utilise coordinate data from freeware virtual globes such as *Bing Maps* (see also Supplementary Material). Position information (‘object-space’ coordinates) is

introduced after model production, with the benefit that errors in GCPs will not propagate in the DEM.

3.3.2 Satellite imagery processing

Satellite imagery products are typically available in georectified form as standard and therefore do not require geometric correction prior to geomorphological mapping. With respect to high-resolution, commercial satellite imagery (e.g. WorldView-4 captured imagery; 0.31 m GSD), these products are often available for purchase as either georeferenced and orthorectified products (with consumers able to define the processing technique used) at comparable prices to commercial aerial photographs, thereby removing the need for photogrammetric processing. Alternatively, it is possible to purchase less expensive ‘ortho-ready’ imagery and perform orthorectification (where DEM or GCP data are available), thus providing greater end-user control on image processing (e.g. Chandler et al., 2016a; Ewertowski et al., 2016).

Although satellite imagery does not typically require geometric correction for mapping, it is important to consider the choice of band combinations when using multispectral satellite imagery (e.g. Landsat, ASTER; Table 1). Since the detection of glacial landforms from optical satellite imagery relies on the interaction of reflected radiation with topography, different combinations of spectral bands can be employed to optimise landform identification (see Jansson and Glasser, 2005). Manipulating the order of bands with different spectral wavelengths allows the generation of various visualisations, or false-colour composites, of the terrain. For example, specific band combinations may be particularly useful for detecting moraine ridges (7, 5, 2 and 5, 4, 2), mega-scale glacial lineations (4, 5, 6) and meltwater channels (4, 3, 2) from Landsat TM and ETM+ imagery (Jansson and Glasser, 2005; Lovell et al., 2011; Morén et al., 2011). This is principally due to the change in surface vegetation characteristics (e.g. type, density and degree of development) between different landforms, between landforms and the surrounding terrain. For example, former meltwater channels typically appear as overly-wide corridors (relative to any modern drainage) of lush green vegetation and stand out clearly as bright red when using a near-infrared false-colour composites (bands 4, 3, 2: Landsat TM and ETM+), since the green colour of surface vegetation is strongly reflected in near-infrared bands (band 4: Landsat TM and ETM+). In addition to the manipulation of band combinations during the mapping process, it can also be beneficial to use satellite image derivatives based on ratios of band combinations, such as vegetation indices (see Walker et al., 1995) and semi-automated image classification techniques (e.g. Smith et al., 2000, 2016b).

Aside from manipulating spectral band combinations, it may also be beneficial to use the higher-resolution panchromatic band as a semi-transparent layer alongside the multispectral bands to aid landform detection (e.g. Morén et al., 2011; Stroeven et al., 2013; Lindholm and Heyman, 2016), or to

merge the pixel resolutions of the panchromatic and multispectral bands through pan-sharpening techniques (e.g. Glasser and Jansson, 2008; Greenwood and Clark, 2008; Storrar et al., 2014; Chandler et al., 2016a; Ewertowski et al., 2016). Pan-sharpening can be particularly valuable when it is desirable to have both multispectral capabilities (e.g. different band combinations to differentiate between features with varying surface characteristics) and higher-spatial resolutions to help determine the extent and morphology of individual landforms.

3.3.3 Digital Elevation Model processing

Various processing techniques are available that can be beneficial when identifying and mapping glacial landforms from DEMs (Bolch and Loibl, 2017). For mapping and analytical purposes, DEM data are typically converted into ‘hillshaded relief models’ (Figure 8), whereby different solar illumination angles and azimuths are simulated within GIS software to produce the shaded DEMs. This rendition provides a visually realistic representation of the land surface, with shadows improving detection of surface features. Consequently, hillshaded relief models should be generated using a variety of illumination azimuths (direction of light source) and angles (elevation of light source) to alleviate the issue of ‘azimuth bias’, the notion that some linear landforms are less visible when shaded from certain azimuths (see Lidmar-Bergström et al., 1991; Smith and Clark, 2005). An illumination angle of 30° and azimuths set at orthogonal positions of 45° and 315° have been suggested as optimal settings for visualisation (Smith and Clark, 2005; Hughes et al., 2010). Vertical exaggeration of these products (e.g. three to four times) can also aid landform identification (e.g. Hughes et al., 2010). Semi-transparent DEMs can be draped over shaded-relief images to accentuate topographic contrasts (Figure 9), or a semi-transparent satellite image can be draped over a DEM to achieve both a multispectral and topographic assessment of a landscape (e.g. Jansson and Glasser, 2005). First- and second-order DEM-derivatives, including surface gradient (slope) and curvature, have also been found to be useful for mapping (e.g. Smith and Clark, 2005; Evans, 2012; Storrar and Livingstone, 2017).

4. Assessment of mapping errors and uncertainties

In this section, we provide an overview of the main errors and uncertainties associated with the various geomorphological mapping methods introduced in the preceding sections. Consideration and management of mapping errors should be an important part of glacial geomorphological mapping studies, since any errors/uncertainties incorporated in the geomorphological map will propagate into subsequent palaeoglaciological and palaeoclimatic reconstructions. This is of most relevance to small ice masses (cirque glaciers, valley glaciers, outlet glaciers), e.g. metre-scale geolocation errors would have significant implications for studies aiming to establish ice-margin retreat rates at the order of tens of metres (e.g. Krüger, 1995; Lukas and Benn, 2006; Lukas, 2012; Bradwell et al., 2013;

Chandler et al., 2016b). Conversely, mapping errors might be negligible in the context of continental-scale ice sheet reconstructions (e.g. Hughes et al., 2014; Stroeve et al., 2016; Margold et al., 2018).

The overall ‘quality’ of any geomorphological map is a function of three interlinked factors: mapping resolution, accuracy and precision. It is important to highlight that, irrespective of the mapping method employed (field or remote-based), the accuracy and precision of the mapping reflects two related factors: (i) the skill, philosophy and experience of the mapper; and (ii) the detectability of the landforms (Smith and Wise, 2007; Otto and Smith, 2013; Hillier et al., 2015). Mapper philosophy concerns issues such as how landforms are mapped (e.g. generalised mapping vs. mapping the intricate details of individual landforms) and interpreted (e.g. differences in terminology and landform classification), which will partly vary with background and training. The significance of the skill, philosophy and experience in mapping is exemplified by the stark differences across boundaries of British Geological Survey (BGS) map sheets that have been mapped by different surveyors, for example (cf. Clark et al., 2004).

A key determinant of landform detectability is resolution, generally defined as the finest element that can be distinguished during survey/observation (Lam and Quattrochi, 1992). In geomorphological mapping it may be, for example, the smallest distinguishable landform that is visible from remotely-sensed data or that can be drawn on a field map. The accuracy of geomorphological mapping relates to positional accuracy (i.e. difference between ‘true’ and mapped location of the landform), geometric accuracy (i.e. difference between ‘true’ and mapped shape of the landform), and attribute accuracy (i.e. deviation between ‘true’ and mapped landform types) (Smith et al., 2006). For spatial data, it is usually not possible to obtain absolute ‘true’ data, due to limitations such as the ‘resolution’ of remotely-sensed data and the accuracy of instruments/surveying equipment. Precision is often used to express the reproducibility of surveys, which is controlled by random errors. These are errors that are innate in the survey/observation process which cannot be removed (Butler et al., 1998). We now outline the specific uncertainties associated with field mapping (Section 4.1), analogue remote mapping (Section 4.2) and digital remote mapping (Section 4.3).

4.1 Field mapping errors and uncertainty

The correct positioning, orientation and scale of individual geomorphological features on field maps is dependent on the skill of the mapper and the ability to correctly interpret and record landforms. If a handheld GNSS device is used to locate landforms in the field, the positional accuracy is usually restricted to several metres and related to three factors: (i) the quality of the device (e.g. antenna, number of channels, ability to use more than one GNSS); (ii) the position of satellites; and (iii) the characteristics of the surrounding landscapes and space weather (solar activity can affect signal

quality). Higher accuracy (cm- or even mm-scale) can only be achieved when supplemented by measurements using additional surveying (e.g. differential Global Positioning Systems (dGPS), real time kinematic (RTK-) GPS or total station). Alongside positioning errors, the horizontal resolution (and, consequently, accuracy) of the field map is related to line thickness on the field map (Knight et al., 2011; Boston, 2012a, b; Otto and Smith, 2013). A pencil line has a thickness of between 0.20 and 0.50 mm on a field map; therefore, individual lines represent a thickness of between 2 and 5 m on 1: 10,000 scale maps, rendering the maps accurate to this level at best (Raisz, 1962; Robinson et al., 1995; Boston, 2012a). This necessitates some element of selection during field mapping of relatively small landforms formed by alpine- and plateau-style ice masses, as not all the information that can be seen in the field can be mapped, even at a large scale such as 1: 10,000. In terms of the vertical accuracy of field maps, it should be recognised that the mapping is only as accurate as the resolution of the source elevation data: if the topographic base map has contours at 10 m intervals, the mapping has a vertical resolution, and thus accuracy, of 10 m at best, irrespective of the (perceived) skill of the cartographer. As with positional accuracy, higher vertical accuracy necessitates the use of geodetic-grade surveying equipment.

4.2 Analogue remote mapping errors and uncertainty

Accurate detection and mapping of individual landforms from analogue (hard-copy) aerial photographs is influenced by factors such as the scale or resolution of the photographs, shadow length (shadows may obscure the ‘true’ planform or landforms altogether), the presence/absence of vegetation, cloud cover and tonal contrast (photographs may appear ‘flat’, thus limiting landform detection). The resolution of analogue remotely-sensed datasets is associated with scale, which results from the altitude of the plane and camera lens focal length, and the optical resolution of the lens and sensor (Wolf et al., 2013). It should also be recognised that the accuracy of the (non-rectified) mapping, as with field mapping, is limited by the thickness of the pen used for drawing on the acetate sheets. Super-fine pens typically have a nib size of 0.05–0.20 mm; thus, lines on an acetate overlay typically represent thicknesses between ~1.25 m and 5.00 m at a common aerial photograph average scale of 1: 25,000. Despite being particularly useful for detailed mapping of small features and complex landform patterns, the level of accuracy achievable using this methodology is therefore ~1.25 m at best. However, further errors will be introduced to the geomorphological mapping once the raw, non-rectified acetates are georectified (see Section 3.3.1).

4.3 Digital remote mapping errors and uncertainty

A key influence on landform detectability from digital remotely-sensed data is the scale of the feature relative to the resolution of the digital dataset, with a particular challenge being the mapping of

features with a scale close to or smaller than the resolution of the imagery. Conversely, mapping exceptionally large ('mega-scale') glacial landforms can be challenging, depending on the remotely-sensed dataset employed (e.g. Greenwood and Kleman, 2010). Unlike analogue mapping (both in the field and remotely), the thickness of digital lines is not typically a problem for digital mapping, so landform detection and recording are fundamentally linked to spatial resolution. Spatial resolution of digital remotely-sensed data refers to the capability to distinguish between two objects, typically expressed as either (i) pixel size or grid cell size or (ii) ground sampled distance. Pixel/grid size refers to the projected ground dimension of the smallest element of the digital image (Figure 10), whilst ground sampled distance (GSD) refers to the ground distance between two measurements made by the detector (the value of measurement is subsequently assigned to a pixel) (Figure 10; Duveiller and Defourny, 2010). In practice, the spatial resolution of digital imagery is lower than the pixel size (Figure 10).

Landform detectability from raster images (i.e. remotely-sensed data) can be considered with reference to the Nyquist-Shannon sampling theorem, since they comprise discrete sampled values. According to this theorem, the *intrinsic resolution* is twice the sampling distance of the measured values, whereas the *nominal resolution* is twice the pixel/grid size (cf. Pipaud et al., 2015, and references therein). The *effective resolution* and, consequently, the minimum landform footprint/planform that can be unambiguously sampled are defined by the smaller of these two values (cf. Pike, 1988). Where the Nyquist-Shannon criterion is not satisfied for either the intrinsic or nominal resolution, landforms with footprints below the critical value may be visible but are rendered ambiguously in digital imagery, i.e. their boundaries are not clearly definable and mappable (cf. Cumming and Wong, 2005). Further factors that influence landform identification from digital imagery include the strength of the landform signal relative to background terrain, and the azimuth bias introduced by differences in the orientation of linear features and the illumination angle of the sun (Smith and Wise, 2007), along with localised issues such as cloud cover, snow cover and vegetation. The timing of data collection is also a key factor, particularly in the case of modern glacial environments (see Section 5.3).

Aside from the factors outlined above, (raw) remotely-sensed data will contain distortion and/or geometric artefacts of varying degrees. Distortions inherent in raw aerial photographs can be partially or almost fully removed during georeferencing of acetate sheets or photogrammetric processing of aerial photographs (see Section 3.3.1). Raw satellite imagery will contain biases related to attitude, ephemeris and drift errors as well as displacements related to the relief, which, similarly to aerial photographs, is more visible in mountainous areas than in lowland settings (Grodecki and Dial, 2003; Shean et al., 2016). With respect to DEMs, some datasets captured using air- and space-borne radar approaches may contain a number of artefacts (Clark, 1997; Figure 11), with geometric artefacts

particularly significant in upland settings. Geometric artefacts, such as foreshortening and layover, are corrected during image processing by stretching high terrain into the correct position, which can result in a smoothed region on steep slopes (Figure 12). In other parts of upland terrain, information will be lost on the leeward side of slopes, away from the sensor, where high ground prevents the radar beam from reaching the lower ground beneath it (Figure 11). Such issues can be alleviated, at least partly, by examining multiple complementary remotely-sensed datasets and mapping at a variety of scales.

4.4. Assessment and management of uncertainties

Due to the subjective nature of geomorphological mapping, assessing mapping precision is not an easy task. One possible approach is to compare results of mapping using different datasets/methods with a perceived more ‘truthful’ dataset (i.e. field-based survey) (Smith et al., 2006). The number, size and shape of mapped landforms in comparison with a ‘true’ dataset can be used as an approximation of mapping reliability. Precision and accuracy of the produced geomorphological map can also be estimated based on the quality of the source data. Most of the datasets are delivered with at least some assessment of uncertainties, often expressed as accuracy; for example, the SRTM DEM has a horizontal accuracy of ± 20 m and a vertical accuracy of ± 16 m (Rabus et al., 2003). Alternatively, some remotely-sensed datasets have an associated *total root mean square error* (RMSE), which indicates displacement between ‘true’ control points and corresponding points on the remotely-sensed data (Wolf et al., 2013). However, both are measures of the overall (‘global’) quality of the dataset. Thus, these errors may be deceptive because such ‘global’ measures ignore spatial patterns of errors and local terrain characteristics (cf. Lane et al., 2005; James et al., 2017). For example, DEM errors will typically be more pronounced on steep slopes, where even a small horizontal shift will incur large differences in elevation.

Ideally, remotely-sensed datasets should be evaluated independently by the mapper to establish their geolocation accuracy (accuracy of x , y and z coordinates). If feasible, surveys of ground control points (GCPs) should be conducted using geodetic-grade surveying equipment (e.g. RTK-GPS, total station). A sub-sample of this GCP dataset can be used for photogrammetric processing, and RMSE values calculated. Subsequently, the remaining GCPs (i.e. those not used for photogrammetric processing) can be used to perform a further quality check, by quantifying deviations from the coordinates of the GCPs and the corresponding points on the generated DEM (e.g. Carrivick et al., 2017). An additional approach, in geomorphologically stable areas, is to compare the location of individual data points from the DEM (or raw point cloud) being used for mapping with those on a reference DEM (or raw point cloud) (e.g. King et al., 2016b; Carrivick et al., 2017; James et al., 2017; Midgley and Tonkin, 2017; Mertes et al., 2017). Parameters such as the mean deviation, standard deviation and relative standard deviation between the two datasets can then be calculated to perform a quantitative

assessment of quality and accuracy of the DEM (e.g. King et al., 2016b; Mertes et al., 2017). Performing these assessments may then facilitate correction of the processed datasets (e.g. Nuth and Kääb, 2011; Carrivick et al., 2017; King et al., 2017).

To some extent, residual uncertainties relating to the skill, philosophy and experience of the mapper may be reduced by developing a set of clear criteria for identifying and mapping particular landforms (e.g. Barrell et al., 2011; Darvill et al., 2014; Bendle et al., 2017a; Lovell and Boston, 2017). That said, there are currently no ‘agreed’ genetic classification schemes for interpreting glacial sediment-landform assemblages, despite the development of facies and landsystem models for particular glacial environments (e.g. Eyles, 1983; Brodzikowski and van Loon, 1991; Evans, 2003a; Benn and Evans, 2010). Indeed, terminologies are inconsistently used in glacial geomorphological research, as different ‘schools’ or traditions still exist. Thus, it is probably most appropriate to select a scheme that has been in frequent use in a given area (to enable ready comparison) or to develop one suited for a particular area or problem. Notwithstanding potential discrepancies relating to genetic classification or terminology, this will at least ensure transparency in future use and analysis of the geomorphological mapping.

Given the influence of the individual mapper on accuracy and precision, it may be beneficial and desirable for multiple mappers to complete (initially) independent field surveys and examination of remotely-sensed datasets to enhance reliability and reproducibility (cf. Hillier et al., 2015; Ewertowski et al., 2017). However, this approach would only be applicable in collaborative efforts and may be impractical due to various factors (e.g. study area size, data access restrictions). The level of detection of individual landforms might be improved by employing multiple methods to enhance landform detectability, whilst the genetic interpretation of landforms (landform classification) can be tested by detailed sedimentological investigations (see Section 2.3). Some uncertainties associated with the quality of data source (e.g. shadows, artefacts) can be alleviated, at least partly, by examining multiple complementary remotely-sensed datasets and mapping at a variety of scales.

5. Scale-appropriate mapping approaches

The following sections place the main geomorphological mapping methods (see Sections 2 and 3) in the spatial and temporal context of the glacial settings in which they are commonly used, demonstrating that particular methods are employed depending on factors such as the size of the study area, former glacial system and landforms (Table 3). We focus on three broad glacial settings for the purposes of this discussion: palaeo-ice sheets (Section 5.1), alpine- and plateau-style ice masses (Section 5.2) and modern glacier forelands (Section 5.3). Although geomorphological mapping in modern glacial settings follows the same general procedures as in former alpine and plateau-style ice

mass settings (see Section 6.2), specific consideration of contemporary glacier forelands is warranted due to important issues relating to the temporal resolution of remotely-sensed data and landform preservation potential, which are not as significant in palaeoglaciological settings.

5.1 Palaeo-ice sheet settings

The continental-scale of palaeo-ice sheets typically necessitates a mapping approach that enables systematic mapping of a large area in a time and cost-effective manner while still allowing accurate identification of landform assemblages at a variety of scales. However, the nature of the approach will differ depending on the aim of the investigation, as this fundamentally determines *what* needs to be mapped and *how* it should be mapped. Palaeo-ice sheet reconstructions have been produced at a range of scales, from entire ice sheets (e.g. Dyke and Prest, 1987a, b, c; Kleman et al., 1997, 2010; Boulton et al., 2001; Glasser et al., 2008; Clark et al., 2012; Livingstone et al., 2015) to regional/local sectors (e.g. Hättestrand, 1998; Jansson et al., 2003; Stokes and Clark, 2003; Ó Cofaigh et al., 2010; Astakhov et al., 2016; Darvill et al., 2017). Depending on the aim of the study, some investigations may focus specifically on mapping particular landforms. For example, studies of ice-sheet flow patterns frequently focus on mapping subglacial bedforms, such as drumlins (e.g. Boulton and Clark, 1990a, b; Kleman et al., 1997, 2010; Stokes and Clark, 2003; Hughes et al., 2010). Nonetheless, cartographic reduction is often still required to manage the volume of information, resulting in the grouping of similarly-orientated bedforms into flow-sets (occasionally termed fans or swarms) (e.g. Jansson et al., 2002, 2003; De Angelis and Kleman, 2007; Greenwood and Clark, 2009a, b; Stokes et al., 2009; Hughes et al., 2014; Atkinson et al., 2016).

In many cases, studies incorporate all or most of the landforms across ice sheet scales to derive palaeoglaciological reconstructions (e.g. Kleman et al., 1997, 2010; Stroeven et al., 2016). The rationale for this is that glaciation styles and processes (e.g. ice-marginal, subglacial) can be inferred from particular combinations of landforms in landform assemblages (e.g. Clayton et al., 1985; Stokes and Clark, 1999; Evans, 2003b; Kleman et al., 2006; Evans et al., 2008, 2014; Darvill et al., 2017; Norris et al., 2018). Establishing relationships between landforms is therefore valuable, not only in understanding glaciation styles, but also in helping decipher the relative sequence of formation (e.g. Clark, 1993; Kleman and Borgström, 1996) that may lay the foundations for absolute dating. Typically, ice sheet investigations are focused on the spatial and temporal evolution of these various aspects, requiring the robust integration of geomorphological mapping with absolute dating techniques (see Stokes et al., 2015). For example, following pioneering palaeoglaciological studies of the Fennoscandian ice sheet (e.g. Kleman, 1990, 1992; Kleman and Stroeven, 1997; Kleman et al., 1997), cosmogenic nuclide exposure dating offered a means to quantify dates and rates (e.g. Fabel et al., 2002, 2006; Stroeven et al., 2002a, b, 2006; Harbor et al., 2006). Such data are crucial to tune and

validate numerical models used to reconstruct evolving ice sheet limits, flow configurations and subglacial processes (e.g. Boulton and Clark, 1990a, b; Näslund et al., 2003; Evans et al., 2009b; Hubbard et al., 2009; Stokes and Tarasov, 2010; Kirchner et al., 2011; Livingstone et al., 2015; Stokes et al., 2016b; Patton et al., 2017a, b).

5.1.1 Manual mapping of palaeo-ice sheet geomorphological imprints

Satellite imagery and DEMs are the prevailing remotely-sensed datasets used for mapping ice sheet-scale landforms, and these datasets have been at the forefront of key developments in the understanding of palaeo-ice sheets (cf. Stokes, 2002; Stokes et al., 2015). Notably, the use of satellite imagery resulted in the identification of hitherto-unrecognised mega-scale glacial lineations (MSGs; Boulton and Clark, 1990a, b; Clark, 1993), which are now recognised as diagnostic geomorphological evidence of ice streams within palaeo-ice sheets (see Stokes and Clark, 1999, 2001, and references therein). This has allowed tangible links to be made between the behaviours of former Quaternary ice sheets and present-day ice sheets (e.g. King et al., 2009; Stokes and Tarasov, 2010; Stokes et al., 2016b). Aerial photograph interpretation and field mapping are also used in some studies (e.g. Hättestrand and Clark, 2006; Kleman et al., 2010; Darvill et al., 2014), but satellite imagery and DEMs are in wider usage for practical reasons (see also Section 3.2). In recent years, the development of LiDAR datasets has led to their increasing application for high resolution mapping of landforms formed by palaeo-ice sheets, particularly in Scandinavia (e.g. Dowling et al., 2015; Greenwood et al., 2015; Ojala et al., 2015; Ojala, 2016; Mäkinen et al., 2017; Peterson et al., 2017). We expect this to be a major area of growth in future mapping studies of former ice sheets.

Mapping glacial landforms from remotely-sensed data typically involves manual on-screen digitisation using one of two main approaches: (i) creating polylines along the crestline or thalweg of landforms or (ii) digitising polygons that delineate the breaks of slope around landform margins (i.e. digitising the planform). The approach employed will depend on the requirements of the study; for example, flow-parallel bedforms (e.g. drumlins and MSGs) have variously been mapped as polylines (e.g. Kleman et al., 1997, 2010; Stokes and Clark, 2003; De Angelis and Kleman, 2007; Storrar and Stokes, 2007; Livingstone et al., 2008; Brown et al., 2011b) and polygons (e.g. Hättestrand and Stroeven, 2002; Hättestrand et al., 2004; Hughes et al., 2010; Spagnolo et al., 2010, 2014; Stokes et al., 2013; Ely et al., 2016a; Bendle et al., 2017a) (Figure 13). The rationale behind mapping flow-parallel bedforms as linear features is that dominant orientations of a population provide sufficient information when investigating ice sheet-scale flow patterns and organisation, although image resolution may also be a determining factor. Mapping polygons allows the extraction of individual landform metrics (e.g. elongation ratios) that can provide insights into subglacial processes (e.g. Ely et al., 2016a) and regional variations in ice sheet flow dynamics (e.g. Stokes and Clark, 2002, 2003; Hättestrand et al., 2004; Spagnolo et al., 2014), but it is far more time-consuming than digitising

linear features. Increasingly, it is being recognised that the population metrics and spectral characteristics of the subglacial bedform ‘field’ as a whole are most important for quantifying bedforms and deciphering subglacial processes and conditions (see Hillier et al., 2013, 2016; Spagnolo et al., 2017; Clark et al., 2018b; Ely et al., 2018; Stokes, 2018).

5.1.2 Automated mapping of palaeo-ice sheet geomorphological imprints

Comprehensive mapping of palaeo-ice sheet geomorphological imprints, and particularly of bedforms, typically entails the identification and mapping of large numbers (in some cases >10,000) of the same, or very, similar types of features (e.g. Clark et al., 2009; Kleman et al., 2010; Storrar et al., 2013). The manual digitisation of such large numbers of landforms is a time-consuming process. Consequently, semi-automated and automated mapping techniques are increasingly being applied to glacial geomorphology (e.g. Napieralski et al., 2007b; Saha et al., 2011; Maclachlan and Eyles, 2013; Eisank et al., 2014; Robb et al., 2015; Yu et al., 2015; Jorge and Brennand, 2017a, b), particularly given that features of a single landform type (e.g. drumlins or MSGs) will have fairly uniform characteristics (orientation, dimensions and morphology). Automated and semi-automated mapping techniques typically use either a pixel- or object-based approach (see Robb et al., 2015, and references therein). Thus far, automated and semi-automated approaches have primarily focused on mapping drumlins or MSGs from medium- to high-resolution DEMs. Several methods have been used, including multi-resolution segmentation (MRS) algorithms (Eisank et al., 2014), a Curvature Based Relief Separation (CBRS) technique (Yu et al., 2015), Object Based Image Analysis (OBIA) (Saha et al., 2011; Robb et al., 2015), and clustering algorithms (Smith et al., 2016b).

Most recently, 2D discrete Fourier transformations have been applied to automatically quantify MSGs (see Spagnolo et al., 2017). In contrast to traditional mapping approaches, this new method analyses the whole topography simultaneously to identify the wavelength and amplitude of periodic features (i.e. waves or ripples across the topography) without the need to manually digitise them. This automated approach is in its infancy but is likely to provide quantitative data that are useful for (i) testing and parameterising models of subglacial processes and landforms (e.g. Barchyn et al., 2016; Stokes, 2018) and (ii) facilitating comparison between subglacial bedforms and other bedforms (e.g. Fourrière et al., 2010; Kocurek et al., 2010; Murray et al., 2014).

5.2 Alpine and plateau glacial settings

Mapping the geomorphological imprints of former alpine- and plateau-style ice masses (cirque glaciers, valley glaciers, icefields and ice-caps) is particularly significant, since the geomorphological imprints of such discrete ice masses facilitate reconstructions of their three-dimensional form (extent,

morphology and thickness). By contrast, establishing the vertical limits, thickness distribution and surface topography of palaeo-ice sheets is challenging (cf. Stokes et al., 2015). Importantly, three-dimensional glacier reconstructions permit the calculation of palaeoclimatic boundary conditions for glaciated regions (e.g. Kerschner et al., 2000; Bakke et al., 2005; Stansell et al., 2007; Mills et al., 2012; Boston et al., 2015), data that cannot be obtained from point-source palaeoenvironmental records in distal settings (e.g. lacustrine archives). Empirical palaeoclimatic data derived from glacier reconstructions are important for three reasons. Firstly, these data facilitate analyses of wind patterns across loci of former glaciers and, in a wider context, regional precipitation gradients and atmospheric circulation patterns (e.g. Ballantyne, 2007a, b). Secondly, the data allow glaciodynamic conditions reconstructed from sediment-landform assemblages (e.g. moraines) to be directly linked to climatic regimes, thereby providing insights into glacier-climate interactions at long-term timescales (e.g. Benn and Lukas, 2006; Lukas, 2007a). Finally, independent, empirical information on climatic boundary conditions is fundamental to parameterising and testing numerical models used to simulate past glacier-climate interactions (e.g. Golledge et al., 2008). Thus, the geomorphological records of alpine and plateau-style ice masses are powerful proxies for understanding the interactions of such ice masses with climate.

Alpine- and plateau-style ice masses encompass a broad spatial spectrum of glacier morphologies (cf. Sugden and John, 1976; Benn and Evans, 2010), but geomorphological mapping of glacial landforms in alpine and plateau settings generally follows a similar approach that combines remote sensing and considerable field mapping/checking (Figure 14; e.g. Federici et al., 2003, 2017; Bakke et al., 2005; Lukas and Lukas, 2006; Reuther et al., 2007; Hyatt, 2010; Bendle and Glasser, 2012; Pearce et al., 2014; Borsellino et al., 2017). Hence, alpine- and plateau-style ice masses are considered collectively here. The similarities in mapping approaches across a wider range of spatial scales partly reflect the fact that, in both alpine and plateau settings, the majority of (preserved) glacial landforms are confined to spatially- and/or topographically-restricted areas (e.g. glaciated valleys), i.e. glacial landforms relating to plateau-style ice masses (i.e. plateau icefields, ice-caps) are dominantly formed by outlet glaciers. Conversely, an important component of mapping in upland environments is often assessing any glacial geomorphological evidence for connections between supposed valley glaciers and plateau surfaces/rounded summits, i.e. alpine vs. plateau styles of glaciation (e.g. McDougall, 2001; Boston et al., 2015). The recognition of any plateau-based ice has significant implications for studies aiming to assess glacier dynamics and regional palaeoclimate (see Rea et al., 1999; Boston, 2012a, and references therein). Consequently, it is important to utilise a versatile geomorphological mapping approach in alpine and plateau settings that allows mapping of glacial landforms at a wide range of spatial scales and potentially across very large areas ($>500 \text{ km}^2$), whilst also providing sufficiently high resolution imagery to map planforms of individual, small landforms (e.g. moraines).

5.2.1 Remote mapping of alpine and plateau settings

Glacial geomorphological mapping from remotely-sensed data in alpine and plateau ice mass settings typically involves interpretation of either analogue or digital aerial photographs (see Sections 3.1 and 3.2.2.2; e.g. Bickerton and Matthews, 1993; Boston, 2012a; Finlayson et al., 2011; Lukas, 2012; Izagirre et al., 2018). This reflects the superior resolution required to map in detail the frequently smaller glacial landforms produced by alpine and plateau-style ice masses, by contrast to the coarser resolution satellite imagery and DEMs predominantly used in ice sheet settings (see Section 5.1). The use of analogue (hard-copy) and digital aerial photographs varies in alpine and plateau settings, depending on data availability and the preference of individual mappers. For example, hard-copy, panchromatic aerial photographs have been widely used in conjunction with stereoscopes (see Section 3.1) for mapping Younger Dryas glacial landforms in Scotland, owing to their excellent tonal contrast (e.g. Benn and Ballantyne, 2005; Lukas and Lukas, 2006; Boston, 2012a, b). Indeed, depending on the environment and quality/resolution of available remotely-sensed imagery, panchromatic, stereoscopic aerial photographs can provide the most accurate approach (in terms of landform identification), with photographs of this format having superior tonal contrast than their digital (colour) counterparts: digital colour aerial photographs may appear ‘flat’ (i.e. shadows are absent or less pronounced) making it more difficult to pick out subtle features, particularly in the absence of *SOCET SET* stereo display software and equipment (see Section 3.2.2.2). Nevertheless, mapping from digital aerial photographs has the advantage of providing georeferenced data and avoiding the duplication of effort, with hand-drawing on acetate overlays necessitating subsequent digitisation (see Sections 3.1 and 3.2). Although panchromatic aerial photographs are invariably older, temporality usually presents no issue in palaeoglaciological (non-glacierised) settings, with the critical factor being image quality.

Irrespective of the type of aerial photographs used for geomorphological mapping, georectification is required to ensure accurate depiction of glacial landforms on the final maps (Section 3.3). This is important for minimising potential geospatial errors that will propagate into any subsequent glacier reconstructions and analyses of glacier-climate interactions. Ideally, georectification would involve stereoscopic photogrammetry, as discussed in Section 3.3, but this approach is impractical for larger ice masses (i.e. plateau icefields and plateau ice-caps). Thus, it is necessary to apply the pragmatic solutions described in Section 3.3.1.1, namely georectifying the aerial photographs or acetate overlays to other (coarser) georeferenced digital imagery or topographic data. Conversely, geomorphological studies at the scale of individual cirque basins, valley glaciers or glacier forelands would be appropriate for topographic surveys and hence stereoscopic photogrammetry, provided (i) the accessibility of the study area permits the use of surveying equipment and (ii) camera calibration data are available (see Section 3.3).

In some locations, coarse to medium resolution satellite imagery may be sufficient to map the geomorphological imprint of former or formerly more extensive icefields and ice-caps (Figure 15; e.g. Glasser et al., 2005; Heyman et al., 2008; Barr and Clark, 2009, 2012; Morén et al., 2011; Hochreuther et al., 2015; Loibl et al., 2015; Gribenski et al., 2016). However, these coarse remotely-sensed datasets may only allow for mapping of broad landform arrangements and patterns, rather than the intricate details of individual landforms, and preclude mapping of small features (cf. Barr and Clark, 2012; Fu et al., 2012; Stroeve et al., 2013; Blomdin et al., 2016b). The emergence of high-resolution (commercial) satellite imagery may result in more widespread use of satellite imagery for mapping in alpine and plateau settings, although the benefits of increased resolution may be counteracted by prohibitive costs for large study areas (see Section 3.2.2.1).

5.2.2 Field mapping in alpine and plateau settings

Detailed field mapping, following the procedures outlined in Section 2.2, has been widely applied as part of geomorphological studies focused on alpine- and plateau-style ice masses (e.g. Benn, 1992; Federici et al., 2003, 2017; Lukas, 2007a; Reuther et al., 2007; Boston, 2012a; Małeckı et al., 2018). Although field mapping is widely used in such settings, many studies do not explicitly report whether this entails field mapping *sensu stricto* (i.e. the procedure outlined in Section 2.2), or verification of landforms mapped from remotely-sensed data by direct ground observations ('ground truthing'). We reaffirm the points raised in Sections 2.2 and 2.3 that, whenever possible, field mapping should be combined with remote mapping in cirque glacier, valley glacier, icefield and ice-cap settings in order to identify subtle glacial landforms and test interpretations of ambiguous features. While we advocate the application of detailed field mapping, we recognise that logistical and/or financial issues may preclude this and that ground truthing (of selected areas) may only be possible. Nevertheless, some form of field survey is important in alpine and plateau settings to (i) circumvent potential issues with the quality/resolution of remotely-sensed data (e.g. poor tonal contrast) and (ii) arrive at definitive of glacial landforms and landscapes (see also Section 2.3)

5.3 Modern glacial settings

Many contemporary glacier forelands are rapidly evolving and new landscapes are emerging. This is largely due to changes resulting from the current retreat of ice masses and exposure of previously-glaciated terrain, leading to destabilisation of some landforms (e.g. Krüger and Kjær, 2000; Kjær and Krüger, 2001; Lukas et al., 2005; Lukas, 2011), erosion by changing meltwater routes and remoulding or complete obliteration of extant landforms in areas following a glacier re-advance or surge (e.g. Evans et al., 1999; Evans and Twigg, 2002; Evans, 2003b; Evans and Rea, 2003; Benediktsson et al., 2008). Glaciofluvial processes on active temperate glacier forelands (e.g. Iceland) often make these

environments unfavourable for preservation of (small) landforms (e.g. Evans and Twigg, 2002; Evans, 2003b, Kirkbride and Winkler, 2012; Evans and Orton, 2015; Evans et al., 2016a). In addition, de-icing and sediment re-working processes prevalent in many modern glacial environments (e.g. Iceland, Svalbard) typically result in substantial ice-marginal landscape modification and topographic inversion (e.g. Etzelmüller et al., 1996; Krüger and Kjær, 2000; Kjær and Krüger, 2001; Lukas et al., 2005; Schomacker, 2008; Bennett and Evans, 2012; Ewertowski and Tomczyk, 2015). Anthropogenic activity can also have considerable implications for glacial systems (Jamieson et al., 2015; Evans et al., 2016b). The rapidity, ubiquity and efficacy of these censoring processes (cf. Kirkbride and Winkler, 2012, for further details) in contemporary glacial environments should be key considerations in geomorphological mapping studies; in particular, the recognition that ice-cored features mapped at a given interval in time are not the ‘final’ geomorphological products (cf. Krüger and Kjær, 2000; Kjær and Krüger, 2001; Everest and Bradwell, 2003; Lukas et al., 2005, 2007; Lukas, 2007b).

In addition to landform preservation potential, spatial and temporal scales will be key determinants in the approaches used in mapping of ice-marginal landscapes, with studies in such settings often focused on the formation of small features (<3 m in height) on recent, short-term timescales (0–30 years) (e.g. Beedle et al., 2009; Lukas, 2012; Bradwell et al., 2013; Reinardy et al., 2013; Chandler et al., 2016b) and/or evolution of the glacier foreland over a given time period (e.g. Bennett et al., 2010; Bennett and Evans, 2012; Ewertowski, 2014; Jamieson et al., 2015; Chandler et al., 2016a, b; Evans et al., 2016a). Thus, the approach to geomorphological mapping discussed in Section 5.2 require some modification, as discussed below. It is also worth noting that geomorphological mapping usually forms part of process-oriented studies in modern glacial settings (Figure 16), often with the intention of providing modern analogues for palaeo-ice masses and their geomorphological imprints (e.g. Evans et al., 1999; Evans, 2011; Schomacker et al., 2014; Benediktsson et al., 2016).

Geophysical surveying methods can also strengthen links between modern and ancient landform records through surveying of the internal architecture of landforms that can be directly linked to depositional processes, as well as glaciological and climatic conditions (e.g. Bennett et al., 2004; Benediktsson et al., 2009, 2010; Lukas and Sass, 2011; Midgley et al., 2013, 2018). Recent advances in geophysical imaging of sub-ice geomorphology has allowed links to be made between modern and palaeo-ice sheets (see Section 3.2.2.3), and we expect this to a growth area going forward (see also Stokes, 2018). More broadly, geophysical methods can also be used to identify the extent of buried ice, allowing an assessment of the geomorphological stability of contemporary glacier forelands (e.g. Everest and Bradwell, 2003).

5.3.1 Remote mapping of modern glacial settings

The spatial resolution of remotely-sensed data is of critical importance in modern glacial settings: spatial resolutions commensurate with the size of the landforms being mapped and the scope of the research are required. Typically, aerial photographs or satellite imagery with GSDs of <0.5 m are used in modern glacial settings to enable mapping of small features (e.g. Benediktsson et al., 2010; Lukas, 2012; Bradwell et al., 2013; Brynjólfsson et al., 2014; Lovell, 2014; Schomacker et al., 2014; Chandler et al., 2016a; Ewertowski et al., 2016; Lovell et al., 2018). LiDAR or UAV-derived DEMs are also becoming increasingly used for mapping in modern glacial environments (e.g. Brynjólfsson et al., 2014, 2016; Jónsson et al. 2014, 2016; Benediktsson et al., 2016; Chandler et al., 2016a; Ewertowski et al., 2016; Everest et al., 2017; Lovell et al., 2018). Despite the high-resolution of the imagery, some compromise on the level of detail may be necessary, such as deciding on a maximum mapping scale (e.g. 1:500–1:1000, Schomacker et al., 2014) to prevent too detailed mapping or by simplifying the mapping of certain features. In studies of low-amplitude (annual) moraines, the crestlines rather than the planforms are typically mapped, reflecting a combination of image resolution and data requirements: annual moraine sequences are often used to calculate ice-margin retreat rates and mapping of crestlines is sufficient detail for this purpose (Figure 17; Krüger, 1995; Beedle et al., 2009; Lukas, 2012; Bradwell et al., 2013; Chandler et al., 2016a, b). Moreover, this approach can actually ‘normalise’ the data for subsequent analyses, removing the variability of, for example, moraine-base widths that result from gravitational processes during or after moraine formation.

The temporality (both month and year) of imagery takes on greater significance in modern glacial environments. Depending on the purpose of the research, either the most recent high resolution remotely-sensed dataset available or a series of images from a number of intervals during a given time period are commonly required (e.g. Benediktsson et al., 2010; Bennett et al., 2010; Bradwell et al., 2013; Reinardy et al., 2013; Chandler et al., 2016a; Evans et al., 2016b; Ewertowski et al., 2016). In exceptional circumstances, the research may require an annual temporal resolution; for example, aerial photographs are commonly captured annually at the beginning and end of the ablation season in many forelands of the European Alps (cf. Lukas, 2012; Zemp et al., 2015). The burgeoning use of UAVs provides very high-resolution imagery (<0.1 m GSD) of contemporary glacier forelands and the option to capture up-to-date imagery during every visit to the site, circumventing issues relating to temporal resolution. This approach is likely to come into greater usage for studies examining short-term ice-marginal landscape evolution and preservation potential.

Photogrammetric image processing (see Section 3.3) is arguably of most importance in contemporary glacial environments, particularly where the purpose of the mapping is to investigate small variations of the order of metres to tens of metres at short-term (0–30 years) timescales (cf. Evans, 2009). However, such constraints are not necessarily applicable where broader landsystem mapping is

conducted (e.g. Evans, 2009; Evans and Orton, 2015; Evans et al., 2016a). Ideally, digital aerial photographs should be processed using stereoscopic photogrammetry techniques using GCPs collected during topographic surveys to enable the production of DEMs and orthorectified imagery with low error values (RMSEs <2 m; see Section 3.3). It is preferable to survey GCPs and capture imagery contemporaneously, with surveyed GCPs appearing in the captured aerial imagery (e.g. Evans and Twigg, 2002; Evans et al., 2006, 2012; Schomacker et al., 2014), but imagery often pre-dates the geomorphological investigations and topographic surveys (e.g. Bennett et al., 2010; Bradwell et al., 2013; Chandler et al., 2016b). Alternatively, the digital aerial photographs could be processed using SfM photogrammetry methods (see Section 3.3.1.2).

5.3.2 Field mapping in modern glacial settings

The rapidly-changing nature of modern glacier forelands presents a number of challenges when using topographic base maps (see Section 2). Firstly, in relation to spatial limitations, topographic maps available in many settings (typically at scales of 1: 25,000 or 1: 50,000) may offer insufficient spatial resolution for mapping: (i) the relief of the small geomorphological features ubiquitous in contemporary glacial environments is often less than the contour intervals depicted on the maps; and (ii) many forelands, such as those of southeast Iceland, have limited elevation changes across the foreland (cf. Evans and Twigg, 2002; Evans et al., 2016a).

Publicly-available topographic maps are rarely updated frequently enough to be useful for mapping the often rapid (annual to decadal-scale) changes exhibited by modern glacier margins and proglacial landscapes. Instead, it is desirable to undertake geodetic-grade surveying (i.e. using an RTK-GPS) of landforms and measurement of high-resolution topographic profiles, where conditions allow a safe approach towards the glacier margin (e.g. Benediktsson et al., 2008; Bradwell et al., 2013). Indeed, conducting detailed surveying with geodetic-grade equipment is essential for quantifying small changes in ice-marginal/proglacial landscapes (e.g. Schomacker and Kjær, 2008; Ewertowski and Tomczyk, 2015; Korsgaard et al., 2015) and obtaining metre-scale ice-margin retreat rates from the geomorphological record (e.g. Bradwell et al., 2013; Chandler et al., 2016a). This level of detail and accuracy may be unnecessary for some glacial geomorphological studies (e.g. those focused on the overall glacial landsystem), and annotation of aerial photograph extracts may be sufficient. There are still potential temporal limitations with these approaches, namely (a) limitations imposed by the date/year of image capture when mapping on print-outs and (b) difficulties with correlating survey data with imagery, depending on the time difference and rapidity of landscape changes. In localities where (parts of) the ice-marginal/proglacial landscape cannot be satisfactorily or safely traversed, imagery and elevation control from remotely-sensed sources will be necessary (e.g. Evans et al., 2016e).

6. Frameworks for best practice

Based on our review of the various mapping approaches, we here synthesise *idealised* frameworks for mapping palaeo-ice sheet geomorphological imprints (Section 6.1) and alpine and plateau-style ice mass (cirque glaciers, valley glaciers, ice-fields and ice-caps) geomorphological imprints (Section 6.2). The aim is to provide frameworks for best practice in glacial geomorphological mapping, ensuring robust and systematic geomorphological mapping programmes. The templates outlined can be modified as necessary, depending on the study area size and project scope, along with the datasets, software and time available.

Before outlining the idealised frameworks, we offer four general recommendations for undertaking and reporting glacial geomorphological mapping that are applicable at all scales of investigation:

- (1) The methods, datasets and equipment employed in mapping should be clearly stated, including the resolution and format of remotely-sensed data.
- (2) Any processing methods and imagery rectification errors (RMSEs) should be reported, as well as mapping uncertainties (both in terms of the location of the landforms and their identification/classification). Where remotely-sensed datasets are obtained as pre-processed, georeferenced products, this should also be stated.
- (3) Establishing and reporting criteria for identifying and mapping different landforms is desirable (e.g. Barrell et al., 2011; Darvill et al., 2014; Bendle et al., 2017a; Lovell and Boston, 2017). As a minimum, this could take the form of a brief definition of the mapped landform (e.g. Storrar and Livingstone, 2017).
- (4) GIS software (e.g. *ArcGIS*, *QGIS*) should be used for geomorphological mapping and digitisation to provide georeferenced geomorphological data.

Following the above general recommendations will provide transparency about how the mapping was compiled and what considerations were made during the process, aiding accuracy assessment, comparison and integration of geomorphological data. This is particularly valuable for the incorporation of the geomorphological mapping in large compilations (Bickerdike et al., 2016; Stroeve et al., 2016; Clark et al., 2017a) and subsequently using the data for palaeoglaciological reconstruction and/or testing numerical ice sheet models (Stokes et al., 2015; Margold et al., 2018).

In relation to software (recommendation 4), some practitioners may prefer to use graphics software packages (e.g. *Adobe Illustrator*, *Canvas X*, *CorelDRAW*) for producing final glacial geomorphological maps (e.g. Brynþólfsson et al., 2014; Darvill et al., 2014; Chandler et al., 2016a; Bendle et al., 2017a; Norris et al., 2017). Such graphics software can provide greater functionality

than current GIS packages for fine adjustments of the final cartographic design. However, any modification in graphics software should be kept to a minimum in order to avoid compromising the transferability of the data for other users (e.g. as shapefiles), with the focus instead on adjustments to the map symbology and ensuring optimal map presentation.

6.1 Palaeo-ice sheet geomorphological imprints

For mapping of palaeo-ice sheet geomorphological imprints we recommend the use of multiple remotely-sensed datasets in a synergistic and systematic process, subject to data availability and coverage (Figure 18). As a minimum, remote-sensing investigations should involve reconnaissance-level mapping using multiple remotely-sensed datasets to establish the most suitable dataset (e.g. Stokes et al., 2016a). However, mapping often benefits from utilising a range of imagery types and resolutions, enabling the advantages of each respective method/dataset to be integrated to produce an accurate geomorphological map (see below). At the outset of the mapping, a decision should be made on the level of mapping detail required for particular landforms (i.e. polyline or polygon mapping), in line with the aims and requirements of the study (see Section 5.1.1).

Initially, mapping should involve an assessment of the study area using remotely-sensed data in conjunction with existing maps and literature to identify gaps in the mapping record and localities for focused mapping. Following this reconnaissance stage, the mapper may proceed with mapping from both DEMs and satellite imagery, adding increasing levels of detail with increasingly higher resolution datasets. Recommended techniques for processing the satellite images and DEMs are outlined in Sections 3.3.2 and 3.3.3, including the generation of false-colour composites with different spectral band combinations to aid landform identification (e.g. Jansson and Glasser, 2005; Lovell et al., 2011; Storrar and Livingstone, 2017).

DEMs may provide a superior source of imagery as they directly record the shape of landforms, rather than the interaction of reflected radiation and topography, and therefore allow for more accurate and intuitive mapping. For example, DEMs are often particularly useful for identifying and mapping meltwater channels (e.g. Greenwood et al., 2007; Storrar and Livingstone, 2017). Specific features may also only be identifiable on satellite imagery, such as low-relief corridors of glaciofluvial deposits, due to their distinctive spectral signatures (e.g. Storrar and Livingstone, 2017). Moreover, the typically superior resolution of satellite imagery may enhance landform detectability and allow for more detailed mapping. Many glacial landforms are also clearly distinguishable in one or more sets of remotely-sensed data (or through using a combination of datasets).

To ensure that all landforms are mapped from remotely-sensed data, the datasets should be viewed at a variety of scales and mapping conducted through multiple passes of the area, enabling the addition of increasing levels of detail to and/or refinement of initial mapping with each pass (Norris et al., 2017). It may be advantageous to perform a final check at a small cartographic scale (e.g. 1:500,000) to ensure there are no errors in the mapping, such as duplication of landforms at image overlaps (e.g. De Angelis, 2007). The mapping should be iterative, with repeated consultations of various remotely-sensed datasets throughout the process recommended.

In this contribution, we have focused on the use of satellite imagery and DEMs for mapping palaeo-ice sheet geomorphological imprints, since these are the most widely used for practical reasons. However, aerial photograph interpretation and fieldwork should not be abandoned altogether in palaeo-ice sheet settings. Aerial photographs, where available, can be used to add further detail and refine the mapping, whilst fieldwork enables ground-truthing of remote mapping (e.g. Hättestrand and Clark, 2006; Kleman et al., 2010; Darvill et al., 2014; Evans et al., 2014). Furthermore, mapping from satellite imagery and DEMs can direct fieldwork, highlighting areas for sedimentological and stratigraphic investigations. Such studies can provide invaluable data on landform genesis, subglacial processes and ice dynamics (e.g. Livingstone et al., 2010; Evans et al., 2015; Spagnolo et al., 2016; Phillips et al., 2017; Norris et al., 2018). Remote mapping of palaeo-ice sheet geomorphology also guides targeted dating for chronological investigations and should be an essential first phase in such studies (e.g. Darvill et al., 2014, 2015).

6.2 Alpine and plateau-style ice mass geomorphological imprints

Our idealised framework for mapping alpine and plateau-style ice mass geomorphological imprints is an iterative process involving several consultations of remotely-sensed data and field mapping (Figures 19 and 20). This methodology provides a robust approach to mapping that has been broadly used in previous studies (e.g. Benn and Ballantyne, 2005; Lukas and Lukas, 2006; Kjær et al., 2008; Boston, 2012a, b; Brynjólfsson et al., 2014; Jónsson et al., 2014; Pearce et al., 2014; Schomacker et al., 2014; Chandler et al., 2016a; Chandler and Lukas, 2017). This framework is also applicable to modern glacial settings as the overarching methods do not differ fundamentally, but practitioners should be aware of issues relating to the temporal resolution of remotely-sensed data (see Section 5.3).

In the initial preparatory stage, the mapper should consult topographic, geological and extant geomorphological maps (where available), and ideally undertake mapping of the study area using remotely-sensed data, at least at a reconnaissance level. This essential phase familiarises the mapper with the study area prior to fieldwork and enables the identification of significant areas for targeted,

detailed field mapping (or ground verification) and sedimentological investigations of specific landforms. Conversely, the reconnaissance investigations may also clarify which areas are less important for a field visit and aid route planning. Importantly, this enables a systematic approach to mapping, and is particularly important in previously-unmapped areas (e.g. Boston, 2012a, b). During the initial stage, it may also be desirable to establish a legend/mapping system in readiness for subsequent field mapping (Otto and Smith, 2013).

Following the preparatory/reconnaissance stage, detailed field mapping, or at a minimum some ground verification, should ideally be conducted to avoid overlooking (subtle) landforms and misinterpreting others. Depending on the nature of the project and accessibility limitations, ground verification may be done during a single (and relatively short) field visit (e.g. Lukas, 2012; Chandler et al., 2016a), whilst detailed field mapping would usually require longer field visits or even repeated, long-term field campaigns (e.g. Kjær et al. 2008; Boston, 2012a, b; Schomacker et al., 2014; Evans et al., 2016a). During field surveys, consultation of initial remote mapping helps to ensure accurate representation of landforms on field maps and allows verification of all features identified remotely (e.g. Boston, 2012a, b; Pearce et al., 2014).

Following field mapping, which may be an intermittent and ongoing process in the case of large study areas and long-term research projects, it is ideal to finalise the geomorphological mapping using high-resolution imagery (i.e. aerial photographs, satellite imagery, LiDAR DEMs, UAV-derived imagery). This allows complex patterns of landforms, such as Scottish ‘hummocky moraine’ (e.g. Lukas and Lukas, 2006; Boston, 2012b), crevasse-squeeze ridges (e.g. Kjær et al., 2008), drumlin fields (e.g. Benediktsson et al., 2016), and sawtooth ‘annual’ moraines (e.g. Chandler et al., 2016a; Evans et al., 2016a), to be mapped with high spatial accuracy, following landform identification and interpretation in the field. Again, during this stage, previous mapping from DEMs and field maps should be consulted. As highlighted in the scale-appropriate examples, the procurement of remotely-sensed data with appropriate spatial and temporal resolution is important (see Sections 5.2 and 5.3).

Depending on the type of imagery used (hard-copy or digital), the rectification of imagery/overlays may precede or follow aerial photograph mapping: where digital format aerial photographs are used, rectification will be undertaken before mapping (Figure 19), whilst acetate overlays will be corrected *after* mapping from hard-copy aerial photographs (Figure 20) (see also Supplementary Material). Subsequently, acetate overlays can be checked against digital imagery (if available) before being digitised, either in a GIS software package (e.g. *ArcMap*, *QGIS*) or in a graphics software package.

In our view, geomorphological mapping in cirque glacier, valley glacier and icefield/ice-cap outlet settings should not be reliant solely on the morphological characteristics of features and should ideally

be combined with detailed sedimentological investigations of available exposures as part of an inductive-deductive process, using standard procedures (cf. Evans and Benn, 2004; Lukas et al., 2013, and references therein). This reflects the fact that these glacier systems occupy more manageable study areas and therefore sedimentological analyses can be more readily applied. By combining geomorphological mapping and sedimentology, issues relating to equifinality (Chorley, 1962; Möller and Dowling, 2018) will be avoided, which is important when attempting to establish the wider palaeoglaciological and palaeoclimatic significance of the geomorphological evidence (cf. Benn and Lukas, 2006). This multi-proxy, process-form approach ensures accurate genetic interpretations on geomorphological maps.

7. Conclusions

Geomorphological mapping forms the basis of a wide range of process-oriented, glacial chronological and palaeoglaciological studies. As such, it is imperative that effective approaches are used to ensure robust assimilation of data and that errors and uncertainties are explicitly reported. This is particularly the case where field mapping and analogue data are transferred to digital format and combined with digital remotely-sensed data.

In general, specific methods and datasets are often applied to particular glacial settings: (i) a mixture of satellite imagery (e.g. Landsat) and DEMs (e.g. ASTER GDEM, SRTM) are typically used for mapping in palaeo-ice sheet settings; and (ii) a combination of aerial photographs and field mapping are widely employed for mapping alpine and plateau-style ice mass geomorphological imprints. Increasingly, UAV-captured aerial imagery and high resolution DEMs (derived from UAV-captured imagery and LiDAR) are being utilised for mapping of modern glacial environments and are likely to be a growth area in future geomorphological mapping studies, enabling high resolution, multi-temporal remotely-sensed datasets to be obtained at relatively low cost. The usage of particular methods reflects the spatial and temporal resolution of remotely-sensed datasets, along with the practicality of their application (both in terms of time and finance).

In this contribution, we have highlighted that compromises and pragmatic solutions are often necessary in glacial geomorphological mapping, particularly with respect to processing techniques and the level of mapping detail. For example, detailed GNSS surveys using geodetic-grade equipment are desirable for photogrammetric processing of aerial photographs, but this is impractical for the large areas covered by icefields, ice-caps and ice sheets. Thus, pragmatic approaches may be used, such as georeferencing analogue-derived mapping to existing (coarser) georeferenced datasets (e.g. satellite imagery, DEMs or orthophotographs). In relation to the level of mapping detail, it is often

necessary to map particular landforms as linear features (e.g. subglacial bedforms, moraines) or define a maximum scale during mapping, due to image resolution and/or study requirements.

Based on our review, we have outlined idealised frameworks and general recommendations to ensure best practice in future studies. In particular, we emphasise the importance of utilising multiple datasets or mapping approaches in synergy, akin to multi-proxy/-method approaches used in many Earth Science disciplines; multiple remotely-sensed datasets in the case of ice sheet-scale geomorphology and a combination of remote sensing and field mapping for cirque glaciers to ice-caps. Further key recommendations are the clear reporting of (i) the methods, datasets and equipment employed in mapping, (ii) any processing methods employed and imagery rectification errors (RMSEs) associated with imagery, along with mapping uncertainties, (iii) the criteria for identifying and mapping different landforms. We also recommend that mapping is conducted in GIS software to provide georeferenced geomorphological data. Finally, we advocate sedimentological investigations of available exposures as part of an inductive-deductive process during fieldwork to ensure accurate genetic interpretations of the geomorphological record as part of a holistic approach. Following these recommendations will aid in comparison, integration and accuracy assessment of geomorphological data, particularly where geomorphological data are incorporated in large compilations and subsequently used for palaeoglaciological reconstruction.

Acknowledgements

We are grateful to numerous colleagues for informal discussions that have directly or indirectly helped shape this paper. Alex Clayton is thanked for kindly supplying the UAV imagery and DEM for the Skálafellsjökull foreland, whilst Jon Merritt is thanked for providing CMB and SL with access to aerial photographs at the British Geological Survey in Edinburgh. We are also grateful to Jacob Bendle, Natacha Gribenski and Sophie Norris for kindly providing figures for inclusion in this contribution. The NEXTMap Great Britain™ data for Ben More Coigach was licensed to BMPC by the NERC Earth Observation Data Centre under a Demonstration Use License Agreement. CMB and HL obtained access to aerial photographs and NEXTMap Great Britain™ data through NERC Earth Observation Data Centre whilst in receipt of NERC Algorithm studentships NE/G52368X/1 (CMB) and NE/I528050/1 (HL). This contribution was written whilst BMPC was in receipt of a Queen Mary Natural and Environmental Science Studentship, which is gratefully acknowledged. We thank Richard Waller and an anonymous reviewer for constructive comments that helped improve the clarity of this contribution, along with Ian Candy for editorial handling.

References

- Alexanderson, H., Adrielsson, L., Hjort, C., Möller, P., Antonov, O., Eriksson, S., Pavlov, M., 2002. Depositional history of the North Taymyr ice- marginal zone, Siberia—a landsystem approach. *J. Quat. Sci.* 17(4), 361–382.
- Allaart, L., Friis, N., Ingólfsson, Ó., Håkansson, L., Noormets, R., Farnsworth, W.R., Mertes, J., Schomacker, A., 2018. Drumlins in the Nordenskiöldbreen forefield, Svalbard. *Gff*, in press. doi: 10.1080/11035897.2018.1466832
- Andrews, J.T., Miller, G.H., 1979. Glacial erosion and ice sheet divides, northeastern Laurentide Ice Sheet, on the basis of the distribution of limestone erratics. *Geology* 7(12), 592–596.
- Astakhov, V., Shkatova, V., Zastrozhnov, A., Chuyko, M., 2016. Glaciomorphological Map of the Russian Federation. *Quat. Int.* 420, 4–14.
- Atkinson, N., Pawley, S., Utting, D.J., 2016. Flow- pattern evolution of the Laurentide and Cordilleran ice sheets across west- central Alberta, Canada: implications for ice sheet growth, retreat and dynamics during the last glacial cycle. *J. Quat. Sci.* 31(7), 753–768.
- Aylsworth, J.M., Shilts, W.W., 1989. Glacial features around the Keewatin Ice Divide: Districts of Mackenzie and Keewatin. Geological Survey of Canada paper 88-24.
- Bakke, J., Dahl, S.O., Paasche, Ø., Løvlie, R., Nesje, A., 2005. Glacier fluctuations, equilibrium-line altitudes and palaeoclimate in Lyngen, northern Norway, during the Lateglacial and Holocene. *The Holocene* 15, 518–540.
- Ballantyne, C.K., 1989. The Loch Lomond Readvance on the Isle of Skye, Scotland: glacier reconstruction and palaeoclimatic implications. *J. Quat. Sci.* 4, 95–108.
- Ballantyne, C.K., 2002. The Loch Lomond Readvance on the Isle of Mull, Scotland: glacier reconstruction and palaeoclimatic implications. *J. Quat. Sci.* 17, 759–771.
- Ballantyne, C.K., 2007a. The Loch Lomond Readvance on north Arran, Scotland: glacier reconstruction and palaeoclimatic implications. *J. Quat. Sci.* 22, 343–359.
- Ballantyne, C.K., 2007b. Loch Lomond Stadial glacier on North Harris, Outer Hebrides, North-West Scotland: glacier reconstruction and palaeoclimatic implications. *Quat. Sci. Rev.* 26, 3134–3149.
- Barchyn, T.E., Dowling, T.P.F., Stokes, C.R., Hugenholtz, C.H., 2016. Subglacial bed form morphology controlled by ice speed and sediment thickness. *Geophys. Res. Lett.* 43, 7572–7580.
- Barr, I.D., Clark, C.D., 2009. Distribution and pattern of moraines in Far NE Russia reveal former glacial extent. *J. Maps* 5, 186–193.
- Barr, I.D., Clark, C.D., 2012. An updated moraine map of Far NE Russia. *J. Maps* 8, 431–436.
- Barrell, D.J.A., Andersen, B.G., Denton, G.H. and Lyttle, B.S., 2011. Glacial geomorphology of the central South Island, New Zealand. GNS Science Monograph 27. GNS Science, Lower Hutt, 81 pp + map (5 sheets).
- Barrell, D.J.A., Andersen, B.G., Denton, G.H. and Lyttle, B.S., 2013. Glacial geomorphology of the central South Island, New Zealand – digital data. GNS Science Monograph 27a. GNS Science, 17 pp (GIS digital data files + explanatory notes).
- Barrow, G., Hinxman, L.W., Cunningham Craig, E.H., 1913. The Geology of Upper Strathspey, Gaick, and the Forest of Atholl (Explanation of Sheet 64). Memoirs of the Geological Survey, HMSO, Edinburgh, Scotland.
- Batchelor, C., Dowdeswell, J.A., Ottesen, D., 2017. Submarine Glacial Landforms. In: Micallef, A., Krastel, S., Savini, A. (Eds.), *Submarine Geomorphology*. Springer, Cham, Switzerland: pp. 207–234.
- Beedle, M.J., Menounos, B., Luckman, B.H., Wheate, R., 2009. Annual push moraines as climate proxy. *Geophys. Res. Lett.* 36, L20501.
- Bendle, J.M., Glasser, N.F., 2012. Palaeoclimatic reconstruction from Lateglacial (Younger Dryas Chronozone) cirque glaciers in Snowdonia, North Wales. *Proceedings of the Geologists' Association* 123, 130–145.
- Bendle, J.M., Thorndycraft, V.R., Palmer, A.P., 2017a. The glacial geomorphology of the Lago Buenos Aires and Lago Pueyrredón ice lobes of central Patagonia. *J. Maps* 13, 654–673.
- Bendle, J.M., Palmer, A.P., Thorndycraft, V.R., Matthews, I.P., 2017b. High-resolution chronology for deglaciation of the Patagonian Ice Sheet at Lago Buenos Aires (46.5° S) revealed through varve chronology and Bayesian age modelling. *Quat. Sci. Rev.* 177, 314–339.

- Benediktsson, Í.Ö., Möller, P., Ingólfsson, Ó., van der Meer, J.J.M., Kjær, K.H., Krüger, J., 2008. Instantaneous end moraine and sediment wedge formation during the 1890 glacier surge of Brúarjökull, Iceland. *Quat. Sci. Rev.* 27, 209–234.
- Benediktsson, Í.Ö., Ingólfsson, Ó., Schomacker, A., Kjaer, K.H., 2009. Formation of submarginal and proglacial end moraines: implications of ice-flow mechanism during the 1963–64 surge of Brúarjökull, Iceland. *Boreas* 38, 440–457.
- Benediktsson, Í.Ö., Schomacker, A., Lokrantz, H., Ingólfsson, Ó., 2010. The 1890 surge end moraine at Eyjabakkajökull, Iceland: a re-assessment of a classic glaciotectonic locality. *Quat. Sci. Rev.* 29, 484–506.
- Benediktsson, Í.Ö., Schomacker, A., Johnson, M.D., Geiger, A.J., Ingólfsson, Ó., Guðmundsdóttir, E.R., 2015. Architecture and structural evolution of an early Little Ice Age terminal moraine at the surge-type glacier Múlajökull, Iceland. *J. Geophys. Res.: Earth Surf.* 120, 1895–1910.
- Benediktsson, Í.Ö., Jónsson, S.A., Schomacker, A., Johnson, M.D., Ingólfsson, Ó., Zoet, L., Iverson, N.R., Stötter, J., 2016. Progressive formation of modern drumlins at Múlajökull, Iceland: stratigraphical and morphological evidence. *Boreas*, 45, 567–583.
- Benn, D.I., 1990. Scottish Lateglacial moraines: debris supply, genesis and significance. Unpublished PhD thesis, University of St Andrews.
- Benn, D.I., 1994. Fluted moraine formation and till genesis below a temperate valley glacier: Slettmarkbreen, Jotunheimen, southern Norway. *Sedimentology* 41, 279–292.
- Benn, D.I., 2006. Interpreting glacial sediments. In: Knight, P. (Ed.) *Glacier Science and Environmental Change*. Oxford, Blackwell, pp. 434–439.
- Benn, D.I., Ballantyne, C.K., 2005. Palaeoclimatic reconstruction from Loch Lomond Readvance glaciers in the West Drumochter Hills, Scotland. *J. Quat. Sci.* 20, 577–592.
- Benn, D.I., Evans, D.J.A., 2010. *Glaciers and Glaciation* (2nd Edition). Hodder Education, London, 802 pp.
- Benn, D.I., Lukas, S., 2006. Younger Dryas glacial landsystems in North West Scotland: an assessment of modern analogues and palaeoclimatic implications. *Quat. Sci. Rev.* 25, 2390–2408.
- Benn, D.I., Lowe, J.J., Walker, M.J.C., 1992. Glacier response to climatic change during the Loch Lomond Stadial and early Flandrian: geomorphological and palynological evidence from the Isle of Skye, Scotland. *J. Quat. Sci.* 7, 125–144.
- Bennett, G.L., Evans, D.J.A., 2012. Glacier retreat and landform production on an overdeepened glacier foreland: the debris-charged glacial landsystem at Kvíárjökull, Iceland. *Earth Surf. Process. Land.* 37, 1584–1602.
- Bennett, G.L., Evans, D.J.A., Carbonneau, P., Twigg, D.R., 2010. Evolution of a debris-charged glacier landsystem, Kvíárjökull, Iceland. *J. Maps* 6, 40–67.
- Bennett, M.R., 1991. Scottish "hummocky moraine": its implications for the deglaciation of the North West Highlands during the Younger Dryas or Loch Lomond Stadial. University of Edinburgh, Unpublished PhD thesis, 362 pp.
- Bennett, M.R., Huddart, D., Waller, R.I., Cassidy, N., Tomio, A., Zukowskyj, P., Midgley, N.G., Cook, S.J., Gonzalez, S., Glasser, N.F., 2004. Sedimentary and tectonic architecture of a large push moraine: a case study from Hagafellsjökull-Eystri, Iceland. *Sedi. Geol.* 172, 269–292.
- Bickerdike, H.L., Evans, D.J.A., Ó Cofaigh, C., Stokes, C.R., 2016. The glacial geomorphology of the Loch Lomond Stadial in Britain: a map and geographic information system resource of published evidence. *J. Maps* 12(5), 1178–1186.
- Bickerdike, H.L., Ó Cofaigh, C., Evans, D.J.A., Stokes, C.R., 2018. Glacial landsystems, retreat dynamics and controls on Loch Lomond Stadial (Younger Dryas) glaciation in Britain. *Boreas* 47, 202–224.
- Bickerton, R.W., Matthews, J.A., 1992. On the accuracy of lichenometric dates: an assessment based on the 'Little Ice Age' moraine sequence of Nigardsbreen, southern Norway. *The Holocene* 2, 227–237.
- Bickerton, R.W., Matthews, J.A., 1993. 'Little ice age' variations of outlet glaciers from the jostedalbreen ice-cap, Southern Norway: A regional lichenometric-dating study of ice-marginal moraine sequences and their climatic significance. *J. Quat. Sci.* 8, 45–66.

1731 Blomdin, R., Stroeve, A.P., Harbor, J.M., Lifton, N.A., Heyman, J., Gribenski, N., Petrakov, D.A.,
1732 Caffee, M.W., Ivanov, M.N., Hättestrand, C., Rogozhina, I., Usubaliev, R., 2016a. Evaluating
1733 the timing of former glacier expansions in the Tian Shan: A key step towards robust spatial
1734 correlations. *Quat. Sci. Rev.* 153, 78–96.

1735 Blomdin, R., Heyman, J., Stroeve, A.P., Hättestrand, C., Harbor, J.M., Gribenski, N., Jansson, K.N.,
1736 Petrakov, D.A., Ivanov, M.N., Alexander, O., Rudoy, A.N., Walther, M., 2016b. Glacial
1737 geomorphology of the Altai and Western Sayan Mountains, Central Asia. *J. Maps* 12(1), 123–
1738 136.

1739 Bolch, T., Loibl, D., 2017. GIS for Glaciers and Glacial Landforms. Reference Module in Earth
1740 Systems and Environmental Sciences.

1741 Borsellino, R., Shulmeister, J., Winkler, S., 2017. Glacial geomorphology of the Barbizon & Butler
1742 Downs, Rangitata Valley, South Island, New Zealand. *J. Maps* 13, 502–510.

1743 Boston, C.M., 2012a. A Lateglacial plateau icefield in the Monadhliath Mountains, Scotland:
1744 reconstruction, dynamics and palaeoclimatic implications. Unpublished PhD thesis, Queen
1745 Mary University of London, 295 pp.

1746 Boston, C.M., 2012b. A glacial geomorphological map of the Monadhliath Mountains, Central
1747 Scottish Highlands. *J. Maps* 8(4), 437–444.

1748 Boston, C.M., Lukas, S., Carr, S.J. 2015. A Younger Dryas plateau icefield in the Monadhliath,
1749 Scotland, and implications for regional palaeoclimate. *Quat. Sci. Rev.* 108, 139–162.

1750 Boulton, G.S., Clark, C.D., 1990a. A highly mobile Laurentide ice sheet revealed by satellite images
1751 of glacial lineations. *Nature* 346(6287), 813–817.

1752 Boulton, G.S., Clark, C.D., 1990b. The Laurentide ice sheet through the last glacial cycle: the
1753 topology of drift lineations as a key to the dynamic behaviour of former ice sheets. *Earth*
1754 *Environ. Sci. Trans. R. Soc. Edinb.* 81(4), 327–347.

1755 Boulton, G.S., Dongelmans, P., Punkari, M., Broadgate, M., 2001. Palaeoglaciology of an ice sheet
1756 through a glacial cycle: the European ice sheet through the Weichselian. *Quat. Sci. Rev.*
1757 20(4), 591–625.

1758 Bradwell, T., Stoker, M.S., Golledge, N.R., Wilson, C.K., Merritt, J.W., Long, D., Everest, J.D.,
1759 Hestvik, O.B., Stevenson, A.G., Hubbard, A.L., Finlayson, A.G., Mathers, H.E., 2008. The
1760 northern sector of the last British Ice Sheet: maximum extent and demise. *Earth-Sci. Rev.* 88,
1761 207–226.

1762 Bradwell, T., Sigurðsson, O., Everest, J., 2013. Recent, very rapid retreat of a temperate glacier in SE
1763 Iceland. *Boreas* 42, 959–973.

1764 Brodzikowski, K., van Loon, A.J., 1991. *Glacigenic Sediments*. Elsevier, Amsterdam, 674 pp.

1765 Brown, V.H., Evans, D.J.A., Evans, I.S., 2011a. The Glacial Geomorphology and Surficial Geology
1766 of the South-West English Lake District. *J. Maps* 7(1), 221–243.

1767 Brown, V.H., Stokes, C.R., Ó Cofaigh, C., 2011b. The Glacial Geomorphology of the North-West
1768 sector of the Laurentide Ice Sheet. *J. Maps* 7, 409–428.

1769 Brynjólfsson, S., Schomacker, A., Ingólfsson, Ó., 2014. Geomorphology and the Little Ice Age extent
1770 of the Drangajökull ice cap, NW Iceland, with focus on its three surge-type outlets.
1771 *Geomorphology* 213, 292–304.

1772 Brynjólfsson, S., Schomacker, A., Korsgaard, N.J., Ingólfsson, Ó. 2016. Surges of outlet glacier from
1773 the Drangajökull ice cap, northwest Iceland. *Earth Planet. Sci. Lett.* 450, 140–151.

1774 Butler, J., Lane, S., Chandler, J., 1998. Assessment of DEM quality for characterizing surface
1775 roughness using close range digital photogrammetry. *The Photogramm. Rec.* 16(92), 271–
1776 291.

1777 Caldenius, C.C., 1932. Las glaciaciones cuaternarias en la Patagonia y Tierra del Fuego. *Geogr. Ann.*
1778 14, 1–164.

1779 Campbell, J.B., Wynne, R.H., 2011. *Introduction to remote sensing* (5th Edition). Taylor and Francis,
1780 London, 667 pp.

1781 Campbell, R.B., 1967a. Geology of Glenlyon map-area, Yukon Territory. Geological Survey of
1782 Canada, Memoir 352, 92 pp.

1783 Campbell, R.B., 1967b. Surficial Geology, Glenlyon, Yukon Territory. Geological Survey of Canada,
1784 "A" Series Map 1222A, 1: 253,440.

- Carbonneau, P.E., Dietrich, J.T., 2017. Cost-effective non-metric photogrammetry from consumer-grade sUAS: implications for direct georeferencing of structure from motion photogrammetry. *Earth Surf. Proc. Land.* 42, 473–486.
- Carrivick, J.L., Smith, M.W., Quincey, D.J., 2016. *Structure from Motion in the Geosciences*. New Analytical Methods in Earth and Environmental Science Series. John Wiley & Sons, Chichester, UK, 208 pp.
- Carrivick, J.L., Yde, J., Russell, A.J., Quincey, D.J., Ingeman-Nielsen, T., Mallalieu, J., 2017. Ice-margin and meltwater dynamics during the mid-Holocene in the Kangerlussuaq area of west Greenland. *Boreas* 46, 369–387.
- Chamberlin, T.C., 1897/1965. The method of multiple working hypotheses. *Science* 148, 745–759.
- Chandler, B.M.P., Lukas, S., 2017. Reconstruction of Loch Lomond Stadial (Younger Dryas) glaciers on Ben More Coigach, NW Scotland, and implications for reconstructing palaeoclimate using small ice masses. *J. Quat. Sci.* 32(4), 475–492.
- Chandler, B.M.P., Evans, D.J.A., Roberts, D.H., Ewertowski, M.W., Clayton, A.I., 2016a. Glacial geomorphology of the Skálafellsjökull foreland, Iceland: A case study of ‘annual’ moraines. *J. Maps* 12(5), 904–916.
- Chandler, B.M.P., Evans, D.J.A., Roberts, D.H., 2016b. Characteristics of recessional moraines at a temperate glacier in SE Iceland: Insights into patterns, rates and drivers of glacier retreat. *Quat. Sci. Rev.* 135, 171–205.
- Chorley, R.J., 1962. *Geomorphology and General Systems Theory*. United States Geological Survey, Professional Paper 500B.
- Clark, C.D., 1993. Mega- scale glacial lineations and cross- cutting ice- flow landforms. *Earth Surf. Proc. Land.* 18(1), 1–29.
- Clark, C.D. 1997. Reconstructing the evolutionary dynamics of former ice sheets using multi-temporal evidence, remote sensing and GIS. *Quat. Sci. Rev.* 16, 1067–1092.
- Clark, C.D., Stokes, C.R., 2001. Extent and basal characteristics of the M’Clintock Channel Ice Stream. *Quat. Int.* 86, 81–101.
- Clark, C.D., Knight, J.K., Gray, J.T., 2000. Geomorphological reconstruction of the Labrador sector of the Laurentide Ice Sheet. *Quat. Sci. Rev.* 19(13), 1343–1366.
- Clark, C.D., Evans, D.J.A., Khatwa, A., Bradwell, T., Jordan, C.J., Marsh, S.H., Mitchell, W.A., Bateman, M.D., 2004. Map and GIS database of glacial landforms and features related to the last British Ice Sheet. *Boreas* 33, 359–375.
- Clark, C.D., Hughes, A.L.C., Greenwood, S.L., Spagnolo, M., Ng, F.S.L., 2009. Size and shape characteristics of drumlins, derived from a large sample, and associated scaling laws. *Quat. Sci. Rev.* 28, 677–692.
- Clark, C.D., Hughes, A.L.C., Greenwood, S.L., Jordan, C., Sejrup, H.P., 2012. Pattern and timing of retreat of the last British-Irish Ice Sheet. *Quat. Sci. Rev.* 44, 112–146.
- Clark, C.D., Ely, J.C., Greenwood, S.L., Hughes, A.L.C., Meehan, R., Barr, I.D., Bateman, M.D., Bradwell, T., Doole, J., Evans, D.J.A., Jordan, C.J., Monteys, X., Pellicer, X.M., Sheehy, M., 2018a. BRITICE Glacial Map, version 2: a map and GIS database of glacial landforms of the last British–Irish Ice Sheet. *Boreas*, 47, 11–27.
- Clark, C.D., Ely, J.C., Spagnolo, M., Hahn, U., Hughes, A.L.C., Stokes, C.R., 2018b. Spatial organization of drumlins. *Earth Surf. Proc. Land.*, 43, 499–513.
- Clayton, L., Teller, J.T., Attig, J.W., 1985. Surging of the southwestern part of the Laurentide Ice Sheet. *Boreas* 14(3), 235–241.
- Close, M.H. 1867. Notes on the general glaciation of Ireland. *J. R. Geogr. Soc. Lond. Dublin* 1, 207–242.
- Coray, S., 2007. Glazialsedimentologische Untersuchungen an Flutes im Vorfeld des Findelengletschers. Unpublished MSc Thesis, Universität Bern, 116 pp.
- Coronato, A., Seppälä, M., Ponce, J.F., Rabassa, J., 2009. Glacial geomorphology of the Pleistocene lake Fagnano ice lobe, Tierra del Fuego, southern South America. *Geomorphology* 112(1), 67–81.
- Craig, B.G., 1961. *Surficial geology of northern District of Keewatin, Northwest Territories*. Geological Survey of Canada, Map 7-1961, Scale 1: 1,013,760.

- Craig, B.G., 1964. Surficial geology of east-central District of Mackenzie. Geological Survey of Canada, Bulletin, 99.
- Cumming, I.G., Wong, F.H., 2005. Digital Processing of Synthetic Aperture Radar Data: Algorithms and Implementation. Artech House, Boston.
- da Rosa, K.K., Vieira, R., Fernandez, G.B., Simoes, F.L., Simoes, J.C., 2011. Formas glaciales y procesos glaciológicos del glaciar templado Wanda, Shetland del Sur. *Investig. Geogr.* 43, 3–16.
- da Rosa, K.K., 2013a. The landforms and pattern of deglaciation of the Dragon glacier, King George Island, South Shetlands, Antarctica. *Rev. Geogr. (Recife)* 30(2), 1–16.
- da Rosa, K.K., Vieira, R., Júnior, C.W.M., de Souza Júnior, E., Simões, J.C., 2013b. Compilation of geomorphological map for reconstructing the deglaciation of ice-free areas in the Martel Inlet, King George Island, Antarctica. *Rev. Bras. Geomorfol.* 14(2), 181–187.
- Darvill, C.M., Stokes, C.R., Bentley, M.R., Lovell, H., 2014. A glacial geomorphological map of the southernmost ice lobes of Patagonia: the Bahía Inútil – San Sebastián, Magellan, Otway, Skyring and Río Gallegos lobes. *J. Maps* 10(3), 500–520.
- Darvill, C.M., Bentley, M.J., Stokes, C.R., Hein, A.S., Rodés, Á., 2015. Extensive MIS 3 glaciation in southernmost Patagonia revealed by cosmogenic nuclide dating of outwash sediments. *Earth Planet. Sci. Lett.* 429, 157–169.
- Darvill, C.M., Stokes, C.R., Bentley, M.J., Evans, D.J.A., Lovell, H., 2017. Dynamics of former ice lobes of the southernmost Patagonian Ice Sheet based on a glacial landsystems approach. *J. Quat. Sci.* 32, 857–876.
- De Angelis, H., 2007. Glacial geomorphology of the east-central Canadian Arctic. *J. Maps* 3, 323–341.
- De Angelis, H., Kleman, J., 2007. Palaeo-ice streams in the Foxe/Baffin sector of the Laurentide Ice Sheet. *Quat. Sci. Rev.* 26(9), 1313–1331.
- De Geer, G. 1910. Den svenska Setsbergsexkursionen 1910 för deltagare i den 11:te internationella geologkonferensen i Stockholm. *Ymer* 30, 305-310.
- Demek, J., 1972. Manual of detailed geomorphological mapping. IUG, Prague, 344 pp.
- Dowdeswell, J.A., Vásquez, M., 2013. Submarine landforms in the fjords of southern Chile: implications for glacial processes and sedimentation in a mild glacier-influenced environment. *Quat. Sci. Rev.* 64, 1–19.
- Dowdeswell, J.A., Canals, M., Jakobsson, M., Todd, B. J., Dowdeswell, E.K., Hogan, K.A., 2016. Atlas of Submarine Glacial landforms: Modern, Quaternary and Ancient. Geological Society, London, Memoirs, No. 46.
- Dowling, T.P.F., Alexanderson, H., Möller, P., 2013. The new high-resolution LiDAR digital height model ('Ny Nationell Höjdmodell') and its application to Swedish Quaternary geomorphology. *GFF* 135(2), 145–151.
- Dowling, T.P.F., Spagnolo, M., Möller, P., 2015. Morphometry and core type of streamlined bedforms in southern Sweden from high resolution LiDAR. *Geomorphology* 236, 54–63.
- Dowling, T.P.F., Möller, P., Spagnolo, M., 2016. Rapid subglacial streamlined bedform formation at a calving bay margin. *J. Quat. Sci.* 31, 879–892.
- Dunstone, R.B., 2014. Testing the groove-ploughing theory for mega-scale glacial lineation (MSGL) formation, using a large dataset of their morphology. Unpublished MSc thesis, University of Durham, 127 pp. [Available at Durham e-theses: <http://etheses.dur.ac.uk/9457/>]
- Duveiller, G., Defourny, P., 2010. A conceptual framework to define the spatial resolution requirements for agricultural monitoring using remote sensing. *Remote Sens. Environ.* 114(11), 2637-2650.
- Dyke, A.S. 1990. Quaternary geology of the Frances Lake map area, Yukon and Northwest Territories. Geological Survey of Canada, Memoir 426.
- Dyke, A.S., Hooper, J.M.G., 2001. Deglaciation of northwest Baffin Island, Nunavut. Geological Survey of Canada Map 1999A, Scale 1: 500,000.
- Dyke, A.S., Prest, V.K., 1987a. Late Wisconsinan and Holocene History of the Laurentide Ice Sheet. *Géogr. Phys. Quat.* 41, 237–263.
- Dyke, A.S., Prest, V.K., 1987b. Late Wisconsinan and Holocene retreat of the Laurentide Ice Sheet, Map 1702A. Geological Survey of Canada, Ottawa.

- 1894 Dyke, A.S., Prest, V.K., 1987c. Paleogeography of northern North America, 18 000 - 5 000 years ago,
1895 Map 1703A. Geological Survey of Canada, Ottawa.
- 1896 Dyke, A.S., Morris, T.F., Green, D.E.C., England, J., 1992. Quaternary geology of Prince of Wales
1897 Island, arctic Canada. Geological Survey of Canada Memoir 433.
- 1898 Eisank, C., Smith, M., Hillier, J., 2014. Assessment of multiresolution segmentation for delimiting
1899 drumlins in digital elevation models. *Geomorphology* 214, 452–464.
- 1900 Ely, J.C., Clark, C.D., Spagnolo, M., Stokes, C.R., Greenwood, S.L., Hughes, A.L.C., Dunlop, P.,
1901 Hess, D., 2016a. Do subglacial bedforms comprise a size and shape continuum?
1902 *Geomorphology* 257, 108–119.
- 1903 Ely, J.C., Gribble, E.A., Clark, C.D., 2016b. The glacial geomorphology of the western cordilleran ice
1904 sheet and Ahklun ice cap, Southern Alaska. *J. Maps* 12(Sup. 1), 415–424.
- 1905 Ely, J.C., Graham, C., Barr, I.D., Rea, B.R., Spagnolo, M., Evans, J., 2017. Using UAV acquired
1906 photography and structure from motion techniques for studying glacier landforms: application
1907 to the glacial flutes at Isfallsglaciären. *Earth Surf. Proc. Land.* 42(6), 877–888.
- 1908 Ely, J.C., Clark, C.D., Spagnolo, M., Hughes, A.L., Stokes, C.R., 2018. Using the size and position of
1909 drumlins to understand how they grow, interact and evolve. *Earth Surf. Proc. Land.* 43(5),
1910 1073–1087.
- 1911 Ercolano, B., Coronato, A., Tiberi, P., Corbella, H., Marderwald, G., 2016. Glacial geomorphology of
1912 the tableland east of the Andes between the Coyle and Gallegos river valleys, Patagonia,
1913 Argentina. *J. Maps* 12(1), 304–313.
- 1914 Espinoza, J.M., Glacial geomorphology and paleoglacial behaviour estimation in the Sierra Baguales
1915 (50°S): Paleoclimatic factors that controlled glacier variations within the Pleistocene –
1916 Holocene regional context. Unpublished PhD thesis, Universidad de Chile.
- 1917 Etzelmüller, B., Hagen, J.O., Vatne, G., Ødegård, R.S., Sollid, J.L., 1996. Glacier debris accumulation
1918 and sediment deformation influenced by permafrost: examples from Svalbard. *Ann. Glaciol.*
1919 22, 53–62.
- 1920 Evans, D.J.A. (Ed.), 2003a. *Glacial Landsystems*. Arnold, London, 532 pp.
- 1921 Evans, D.J.A., 2003b. Ice-Marginal Terrestrial Landsystems: Active Temperate Glacier Margins. In:
1922 Evans, D.J.A. (Ed.), *Glacial Landsystems*. Arnold, London, pp. 12–43.
- 1923 Evans, D.J.A., 2009. Glacial Geomorphology at Glasgow. *Scottish Geographical Journal* 125, 285–
1924 320.
- 1925 Evans, D.J.A., 2010. Controlled moraine development and debris transport pathways in polythermal
1926 plateau icefields: examples from Tunghnafellsjökull, Iceland. *Earth Surf. Proc. Land.* 35,
1927 1430–1444.
- 1928 Evans, D.J.A., 2011. Glacial landsystems of Satujökull, Iceland: A modern analogue for glacial
1929 landsystem overprinting by mountain icecaps. *Geomorphology* 129, 225–237.
- 1930 Evans, D.J.A., 2017. Chapter 4 – Conceptual glacial ground models: British and Irish case studies. In:
1931 Griffiths, J.S., Martin, C.J. (Eds.), *Engineering Geology and Geomorphology of Glaciated
1932 and Periglaciated Terrains: Engineering Group Working Party Report*. Engineering Geology
1933 Special Publication 28. The Geological Society, London, pp. 369–500.
- 1934 Evans, D.J.A., Benn, D.I. (Eds.), 2004. *A Practical Guide to the Study of Glacial Sediments*. Arnold,
1935 London, 266 pp.
- 1936 Evans, D.J.A., Orton, C., 2015. Heinabergsjökull and Skalafellsjökull, Iceland: Active Temperate
1937 Piedmont Lobe and Outwash Head Glacial Landsystem. *J. Maps* 11(3), 415–431.
- 1938 Evans, D.J.A., Rea, B.R., 2003. Surging Glacier Landsystem. In: Evans, D.J.A. (Ed.), *Glacial
1939 Landsystems*. Arnold, London, pp. 259–288.
- 1940 Evans, D.J.A., Twigg, D.R., 2002. The active temperate glacial landsystem: a model based on
1941 Breiðamerkurjökull and Fjallsjökull, Iceland. *Quat. Sci. Rev.* 21, 2143–2177.
- 1942 Evans, D.J.A., Lemmen, D.S., Rea, B.R., 1999. Glacial landsystems of the southwest Laurentide Ice
1943 Sheet: modern Icelandic analogues. *J. Quat. Sci.* 14, 673–691.
- 1944 Evans, D.J.A., Twigg, D.R., Shand, M., 2006. Surficial geology and geomorphology of the þórisjökull
1945 plateau icefield, west-central Iceland. *J. Maps* 2, 17–29.
- 1946 Evans, D.J.A., Clark, C.D., Rea B.R., 2008. Landform and sediment imprints of fast glacier flow in
1947 the southwest Laurentide Ice Sheet. *J. Quat. Sci.* 23(3), 249–272.

- 1948 Evans, D.J.A., Twigg, D.R., Rea, B.R., Orton, C., 2009a. Surging glacier landsystem of
1949 Tungnaárjökull, Iceland. *J. Maps* 5, 134–151.
- 1950 Evans, D.J.A., Livingstone, S.J., Vieli, A., Ó Cofaigh, C., 2009b. The palaeoglaciology of the central
1951 sector of the British and Irish Ice Sheet: reconciling glacial geomorphology and preliminary
1952 ice sheet modelling. *Quat. Sci. Rev.* 28, 739–757.
- 1953 Evans, D.J.A., Strzelecki, M., Milledge, D.G., Orton, C., 2012. Hørbyebreen polythermal glacial
1954 landsystem, Svalbard. *J. Maps* 8, 146–156.
- 1955 Evans, D.J.A., Young, N.J.P., Ó Cofaigh, C., 2014. Glacial geomorphology of terrestrial-terminating
1956 fast flow lobes/ice stream margins in the southwest Laurentide Ice Sheet. *Geomorphology*
1957 204, 86–113.
- 1958 Evans, D.J.A., Roberts, D.H., Cofaigh, C.Ó., 2015. Drumlin sedimentology in a hard-bed, lowland
1959 setting, Connemara, western Ireland: implications for subglacial bedform generation in areas
1960 of sparse till cover. *J. Quat. Sci.* 30, 537–557.
- 1961 Evans, D.J.A., Ewertowski, M., Orton, C., 2016a. Fláajökull (north lobe), Iceland: active temperate
1962 piedmont lobe glacial landsystem. *J. Maps* 12(5), 777–789.
- 1963 Evans, D.J.A., Ewertowski, M., Jamieson, S.S.R., Orton, C., 2016b. Surficial geology and
1964 geomorphology of the Kumtor Gold Mine, Kyrgyzstan: human impacts on mountain glacier
1965 landsystems. *J. Maps* 12(5), 757–769.
- 1966 Evans, D.J.A., Ewertowski, M., Orton, C., 2016c. Eiríksjökull plateau icefield landsystem, Iceland. *J.*
1967 *Maps* 12(5), 747–756.
- 1968 Evans, D.J.A., Storrar, R.D., Rea, B.R., 2016d. Crevasse-squeeze ridge corridors: Diagnostic features
1969 of late-stage palaeo-ice stream activity. *Geomorphology* 258, 40–50.
- 1970 Evans, D.J.A., Ewertowski, M., Orton, C., Harris, C., Guðmundsson, S., 2016e. Snæfellsjökull
1971 volcano-centred ice cap landsystem, West Iceland. *J. Maps* 12(5) 1128–1137.
- 1972 Evans, D.J.A., Ewertowski, M., Orton, C., 2017. Skaftafellsjökull, Iceland: glacial geomorphology
1973 recording glacier recession since the Little Ice Age. *J. Maps* 13, 358–368.
- 1974 Evans, I.S., 1990. Cartographic techniques in geomorphology. In: Goudie, A.S. (Ed.),
1975 *Geomorphological techniques* (2nd Edition). Routledge, London, pp. 97–108.
- 1976 Evans I.S., 2012, Geomorphometry and landform mapping: What is a landform? *Geomorphology* 137,
1977 94–106.
- 1978 Everest, J.D., Bradwell, T., 2003. Buried glacier ice in southern Iceland and its wider significance.
1979 *Geomorphology* 52, 347–358.
- 1980 Everest, J., Bradwell, T., Jones, L., Hughes, L., 2017. The geomorphology of Svínafellsjökull and
1981 Virkisjökull-Falljökull glacier forelands, southeast Iceland. *J. Maps* 13, 936–945.
- 1982 Ewertowski, M.W., 2014. Recent transformations in the high- Arctic glacier landsystem,
1983 Ragnarbreen, Svalbard. *Geogr. Ann.* 96A(3), 265–285.
- 1984 Ewertowski, M.W., Tomczyk, A.M., 2015. Quantification of the ice-cored moraines' short-term
1985 dynamics in the high-Arctic glaciers Ebbabreen and Ragnarbreen, Petuniabukta, Svalbard.
1986 *Geomorphology* 234, 211–227.
- 1987 Ewertowski, M.W., Evans, D.J.A., Roberts, D.H., Tomczyk, A.M., 2016. Glacial geomorphology of
1988 the terrestrial margins of the tidewater glacier, Nordenskiöldbreen, Svalbard. *J. Maps* 12(Sup.
1989 1), 476–487.
- 1990 Ewertowski M.W., Kijowski A., Szuman I., Tomczyk A.M., Kasprzak L., 2017. Low-altitude remote
1991 sensing and GIS-based analysis of cropmarks: classification of past thermal-contraction-crack
1992 polygons in central western Poland. *Geomorphology* 293B, 418–432.
- 1993 Eyles, N. (Ed.), 1983. *Glacial Geology: An Introduction for Engineers and Earth Scientists*.
1994 Pergamon, Oxford, 409 pp.
- 1995 Fabel, D., Stroeve, A.P., Harbor, J., Kleman, J., Elmore, D., Fink, D., 2002. Landscape preservation
1996 under Fennoscandian ice sheets determined from in situ produced ¹⁰Be and ²⁶Al. *Earth*
1997 *Planet. Sci. Lett.* 201, 397–406.
- 1998 Fabel, D., Fink, D., Fredin, O., Harbor, J., Land, M., Stroeve, A.P., 2006. Exposure ages from relict
1999 lateral moraines overridden by the Fennoscandian ice sheet. *Quat. Res.* 65, 136–146.
- 2000 Federici, P.R., Pappalardo, M., Ribolini, A., 2003. *Geomorphological Map of the Maritime Alps*
2001 *Natural Park and surroundings (Argentera Massif, Italy)*, 1: 25000 scale. Selca, Florence.

- 2002 Federici, P.R., Ribolini, A., Spagnolo, M., 2017. Glacial history of the Maritime Alps from the Last
2003 Glacial Maximum to the Little Ice Age. *Geological Society, London, Special Publications*
2004 433(1), 137–159.
- 2005 Finke, L., 1980. Anforderungen aus der Planungspraxis an ein geomorphologisches Kartenwerk. In:
2006 Barsch, D., Liedtke, H. (Eds.), *Methoden und Anwendbarkeit geomorphologischer*
2007 *Detailkarten*. Freie Universität Berlin, Berlin; pp. 75–81.
- 2008 Finlayson, A., Merritt, J., Browne, M., Merritt, J., McMillan, A., Whitbread, K., 2010. Ice sheet
2009 advance, dynamics, and decay configurations: evidence from west central Scotland. *Quat. Sci.*
2010 *Rev.* 29, 969–988.
- 2011 Finlayson, A.G., Gollledge, N., Bradwell, T., Fabel, D., 2011. Evolution of a Lateglacial mountain
2012 icecap in northern Scotland. *Boreas* 40, 536–554.
- 2013 Flink, A.E., Noormets, R., Kirchner, N., Benn, D.I., Luckman, A., Lovell, H., 2015. The evolution of
2014 a submarine landform record following recent and multiple surges of Tunabreen glacier,
2015 Svalbard. *Quat. Sci. Rev.* 108, 37–50.
- 2016 Flint, R.F., Colton, R.B., Goldthwait, R.P., Willman, H.B., 1959. Glacial map of the United States
2017 east of the Rocky Mountains. Geological Society of America.
- 2018 Follestad, B., Bergström, B., 2004. Otta 1718-IV. Quaternary geology map with description.
2019 Norwegian Geological Survey. 1:50,000.
- 2020 Fourrière, A., Claudin, P., Andreotti, B., 2010. Bedforms in a turbulent stream: Formation of ripples
2021 by primary linear instability and of dunes by nonlinear pattern coarsening, *J. Fluid Mech.* 649,
2022 287–328.
- 2023 Fu, P., Heyman, J., Hättestrand, C., Stroeve, A.P., Harbor, J.M., 2012. Glacial geomorphology of the
2024 Shaluli Shan area, southeastern Tibetan Plateau. *J. Maps* 8, 48–55.
- 2025 Garcia, J.L., Kaplan, M.R., Hall, B.L., Schaefer, J.M., Vega, R.M., Schwartz, R., Finkel, R., 2012
2026 Glacier expansion in southern Patagonia throughout the Antarctic cold reversal. *Geology* 40,
2027 859–862
- 2028 Glasser, N.F., Jansson, K.N., 2005. Fast-flowing outlet glaciers of the last glacial maximum
2029 Patagonian Icefield. *Quat. Res.* 63(2), 206–211.
- 2030 Glasser, N., Jansson, K., 2008. The Glacial Map of southern South America. *J. Maps* 4, 175–196.
- 2031 Glasser, N.F., Jansson, K.N., Harrison, S., Rivera, A., 2005. Geomorphological evidence for
2032 variations of the North Patagonian Icefield during the Holocene. *Geomorphology* 71(3-4),
2033 263–277.
- 2034 Glasser, N.F., Jansson, K.N., Harrison, S., Kleman, J., 2008. The glacial geomorphology and
2035 Pleistocene history of South America between 38 S and 56 S. *Quat. Sci. Rev.* 27(3), 365–390.
- 2036 Gollledge, N.R., Stoker, M.S., 2006. A palaeo-ice-stream of the British Ice Sheet in eastern Scotland.
2037 *Boreas* 35, 231–243.
- 2038 Gollledge, N.R., Hubbard, A., Sugden, D.E., 2008. High-resolution numerical simulation of Younger
2039 Dryas glaciation in Scotland. *Quat. Sci. Rev.* 27, 888–904.
- 2040 Goodchild, J.G., 1875. The glacial phenomena of the Eden Valley and the western part of the
2041 Yorkshire-dale District. *Quart. J. Geol. Soc. Lond.* 31, 55–99.
- 2042 Graf, A., 2007. Genese alpiner Seitenmoränen am Beispiel des Findelengletschers bei Zermatt (VS).
2043 Unpublished MSc Thesis, Universität Bern, 126 pp.
- 2044 Greenwood, S.L., Clark, C.D., 2008. Subglacial bedforms of the Irish Ice Sheet. *J. Maps* 4, 332–357.
- 2045 Greenwood, S.L., Clark, C.D., 2009a. Reconstructing the last Irish Ice Sheet 1: changing flow
2046 geometries and ice flow dynamics deciphered from the glacial landform record. *Quat. Sci.*
2047 *Rev.* 28, 3085–3100.
- 2048 Greenwood, S.L., Clark, C.D., 2009b. Reconstructing the last Irish Ice Sheet 2: a geomorphologically-
2049 driven model of ice sheet growth, retreat and dynamics. *Quat. Sci. Rev.* 28(27), 3101–3123.
- 2050 Greenwood, S.L., Kleman, J., 2010. Glacial landforms of extreme size in the Keewatin sector of the
2051 Laurentide Ice Sheet. *Quat. Sci. Rev.* 29(15), 1894–1910.
- 2052 Greenwood, S.L., Clark, C.D., Hughes, A.L.C., 2007. Formalising an inversion methodology for
2053 reconstructing ice-sheet retreat patterns from meltwater channels: application to the British
2054 Ice Sheet. *J. Quat. Sci.* 22, 637–645.

2055 Greenwood, S.L., Clason, C.C., Mikko, H., Nyberg, J., Peterson, G., Smith, C.A., 2015. Integrated
2056 use of lidar and multibeam bathymetry reveals onset of ice streaming in the northern Bothnian
2057 Sea. *GFF* 137, 284–292.

2058 Greenwood, S.L., Clason, C.C., Nyberg, J., Jakobsson, M., Holmlund, P., 2017. The Bothnian Sea ice
2059 stream: early Holocene retreat dynamics of the south- central Fennoscandian Ice Sheet.
2060 *Boreas* 46, 346–362.

2061 Gribenski, N., Jansson, K.N., Lukas, S., Stroeven, A.P., Harbor, J.M., Blomdin, R., Ivanov, M.N.,
2062 Heyman, J., Petrakov, D.A., Rudoy, A., Clifton, T., Lifton, N.A., Caffee, M.W., 2016.
2063 Complex patterns of glacier advances during the late glacial in the Chagan Uzun Valley,
2064 Russian Altai. *Quat. Sci. Rev.* 149, 288–305.

2065 Gribenski, N., Lukas, S., Stroeven, A.P., Jansson, K.N., Harbor, J.M., Blomdin, R., Ivanov, M.N.,
2066 Heyman, J., Petrakov, D.A., Rudoy, A., Clifton, T., Lifton, N.A., Caffee, M.W., 2017. Reply
2067 to comment received from J. Herget et al. regarding “Complex patterns of glacier advances
2068 during the late glacial in the Chagan Uzun Valley, Russian Altai” by Gribenski et al. (2016),
2069 *Quaternary Science Reviews* 149, 288–305. *Quat. Sci. Rev.* 168, 219–221.

2070 Griffiths, J.S., Martin, C.J. (Eds.), 2017. *Engineering Geology and Geomorphology of Glaciated and*
2071 *Periglaciated Terrains: Engineering Group Working Party Report. Engineering Geology*
2072 *Special Publication 28. The Geological Society, London, 953 pp.*

2073 Grodecki, J., Dial, G., 2003. Block Adjustment of High-Resolution Satellite Images Described by
2074 Rational Polynomials. *Photogramm. Eng. Remote Sens.* 69(1), 59–68.

2075 Gustavsson, M., Kolstrup, E., Seijmonsbergen, A.C., 2006. A new symbol-and-GIS based detailed
2076 geomorphological mapping system: Renewal of a scientific discipline for understanding
2077 landscape development. *Geomorphology* 77(1–2), 90–111.

2078 Gustavsson, M., Seijmonsbergen, A.C., Kolstrup, E., 2008. Structure and contents of a new
2079 geomorphological GIS database linked to a geomorphological map – With an example from
2080 Liden, central Sweden. *Geomorphology* 95, 335–349.

2081 Harbor, J., Stroeven, A.P., Fabel, D., Clarhäll, A., Kleman, J., Li, Y.K., Elmore, D., Fink, D., 2006.
2082 Cosmogenic nuclide evidence for minimal erosion across two subglacial sliding boundaries of
2083 the late glacial Fennoscandian ice sheet. *Geomorphology* 75, 90–99.

2084 Hardt, J., Hebenstreit, R., Lüthgens, C., Böse, M., 2015. High-resolution mapping of ice-marginal
2085 landforms in the Barnim region, northeast Germany. *Geomorphology* 250, 41–52.

2086 Hättestrand, C., 1998. The glacial geomorphology of central and northern Sweden. *Sver. Geol.*
2087 *Unders. Ca* 85, 1–47.

2088 Hättestrand, C., Clark, C.D., 2006. The glacial geomorphology of Kola Peninsula and adjacent areas
2089 in the Murmansk Region, Russia. *J. Maps* 2, 30–42.

2090 Hättestrand, C., Stroeven, A.P., 2002. A relict landscape in the centre of Fennoscandian glaciation:
2091 Geomorphological evidence of minimal Quaternary glacial erosion. *Geomorphology* 44(1),
2092 127–143.

2093 Hättestrand, C., Kosche, S., Näslund, J.-O., Fabel, D., Stroeven, A.P., 2004. Drumlin formation time -
2094 evidence from northern and central Sweden. *Geogr. Ann.* 86A, 155–167.

2095 Heiser, P.A., Roush, J.J., 2001. Pleistocene glaciations in Chukotka, Russia: moraine mapping using
2096 satellite synthetic aperture radar (SAR) imagery. *Quat. Sci. Rev.* 20(1), 393–404.

2097 Hess, D.P., Briner, J.P., 2009. Geospatial analysis of controls on subglacial bedform morphometry in
2098 the New York Drumlin Field – implications for Laurentide Ice Sheet dynamics. *Earth Surf.*
2099 *Proc. Land.* 34, 1126–1135.

2100 Heyman, J., Hättestrand, C., Stroeven, A.P., 2008. Glacial geomorphology of the Bayan Har sector of
2101 the NE Tibetan Plateau. *J. Maps* 4(1), 42–62.

2102 Hillier, J.K., Smith, M.J., Clark, C.D., Stokes, C.R., Spagnolo, M., 2013. Subglacial bedforms reveal
2103 an exponential size–frequency distribution. *Geomorphology* 190, 82–91.

2104 Hillier, J.K., Smith, M.J., Armugam, R., Barr, I., Boston, C.M., Clark, C.D., Ely, J., Frankl, A.,
2105 Greenwood, S.L., Gosselin, L., Hättestrand, C., Hogan, K., Hughes, A.L.C., Livingstone, S.J.,
2106 Lovell, H., McHenry, M., Munoz, Y., Pellicer, X.M., Pellitero, R., Robb, C., Roberson, S.,
2107 Ruther, D., Spagnolo, M., Standell, M., Stokes, C.R., Storrar, R., Tate, N.J., Wooldridge, K.,
2108 2015. Manual mapping of drumlins in synthetic landscapes to assess operator effectiveness. *J.*
2109 *Maps* 11, 719–729.

2110 Hillier, J.K., Kougioumtzoglou, I.A., Stokes, C.R., Smith, M.J., Clark, C.D., Spagnolo, M.S., 2016.
2111 Exploring Explanations of Subglacial Bedform Sizes Using Statistical Models. *PLoS One* 11,
2112 e0159489.

2113 Hochreuther, P., Loibl, D., Wernicke, J., Zhu, H., Grieflinger, J., Bräuning, A., 2015. Ages of major
2114 Little Ice Age glacier fluctuations on the southeast Tibetan Plateau derived from tree-ring-
2115 based moraine dating. *Palaeogeogr., Palaeoclimatol., Palaeoecol.* 422, 1–10.

2116 Hodgson, D.A., Vincent, J.S., Fyles, J.G., 1984. Quaternary geology of central Melville Island,
2117 Northwest Territories. Geological Survey of Canada paper 83-16.

2118 Hodgson, D.A., Graham, A.G.C., Griffiths, H.J., Roberts, S.J., Ó Cofaigh, C., Bentley, M.J., Evans,
2119 D.J.A., 2014. Glacial history of sub-Antarctic South Georgia based on the submarine
2120 geomorphology of its fjords. *Quat. Sci. Rev.* 89, 129–147.

2121 Hollingworth, S.E., 1931. The glaciation of western Edenside and adjoining areas and the drumlins of
2122 Edenside and the Solway Basin. *Quart. J. Geol. Soc. Lond.* 87, 281–359.

2123 Horsfield, B.R., 1983. The deglaciation pattern of the western Grampians of Scotland. Unpublished
2124 PhD thesis, University of East Anglia.

2125 Houmark-Nielsen, M., Kjær, K.H. 2003. Southwest Scandinavia, 40-15 kyr BP: Palaeogeography and
2126 environmental change. *J. Quat. Sci.* 18, 769–786.

2127 Howarth, P.J., 1968. Geomorphological and Glaciological Studies, Eastern Breiðamerkurjökull,
2128 Iceland, Unpublished PhD thesis, University of Glasgow.

2129 Howarth, P.J., Welch, R., 1969a. Breiðamerkurjökull, South-east Iceland, August 1945, 1:30,000
2130 scale map. University of Glasgow.

2131 Howarth, P.J., Welch, R., 1969b. Breiðamerkurjökull, South-east Iceland, August 1965, 1:30,000
2132 scale map. University of Glasgow.

2133 Hubbard, A., Bradwell, T., Golledge, N., Hall, A., Patton, H., Sugden, D., Cooper, R., Stoker, M.,
2134 2009. Dynamic cycles, ice streams and their impact on the extent, chronology and
2135 deglaciation of the British–Irish ice sheet. *Quat. Sci. Rev.* 28, 758–776.

2136 Hubbard, B., Glasser, N.F., 2005. Field techniques in glaciology and glacial geomorphology. John
2137 Wiley and Sons, Chichester, 400 pp.

2138 Hughes, A.L.C., Clark, C.D., Jordan, C.J., 2010. Subglacial bedforms of the last British Ice Sheet. *J.*
2139 *Maps* 6, 543–563.

2140 Hughes, A.L.C., Clark, C.D., Jordan, C.J., 2014. Flow-pattern evolution of the last British Ice Sheet.
2141 *Quat. Sci. Rev.* 89, 148–168.

2142 Hughes, A.L.C., Gyllencreutz, R., Lohne, Ø.S., Mangerud, J., Svendsen, J.I., 2016. The last Eurasian
2143 ice sheets - a chronological database and time-slice reconstruction, DATED-1. *Boreas* 45, 1-
2144 45.

2145 Hyatt, O.M., 2010. Insights into New Zealand Glacial Processes from studies of glacial
2146 geomorphology and sedimentology in Rakaia and other South Island Valleys. Unpublished
2147 PhD thesis, University of Canterbury, 251 pp.

2148 Immerzeel, W.W., Kraaijenbrink, P.D.A., Shea, J.M., Shrestha, A.B., Pellicciotti, F., Bierkens,
2149 M.F.P., de Jong, S.M., 2014. High-resolution monitoring of Himalayan glacier dynamics
2150 using unmanned aerial vehicles. *Remote Sens. Environ.* 150, 93–103.

2151 Izagirre, E., Darvill, C.M., Rada, C., Aravena, J.C., 2018. Glacial geomorphology of the Marinelli and
2152 Pigafetta glaciers, Cordillera Darwin Icefield, southernmost Chile. *J. Maps* 14, 269–281.

2153 Jakobsson, M., Mayer, L., Coakley, B., Dowdeswell, J.A., Forbes, S., Fridman, B., Hodnesdal, H.,
2154 Noormets, R., Pedersen, R., Rebesco, M., Schenke, H.W., Zarayskaya, Y., Accettella, D.,
2155 Armstrong, A., Anderson, R.M., Bienhoff, P., Camerlenghi, A., Church, I., Edwards, M.,
2156 Gardner, J.V., Hall, J.K., Hell, B., Hestvik, O., Kristoffersen, Y., Marcussen, C., Mohammad,
2157 R., Mosher, D., Nghiem, S.V., Pedrosa, M.T., Travaglini, P.G., Weatherall, P., 2012. The
2158 International Bathymetric Chart of the Arctic Ocean (IBCAO) Version 3.0. *Geophys. Res.*
2159 *Lett.* 39, L12609.

2160 James, M.R., Robson, S., Smith, M.W., 2017. 3-D uncertainty-based topographic change detection
2161 with structure-from-motion photogrammetry: precision maps for ground control and directly
2162 georeferenced surveys. *Earth Surf. Process. Land.* 42, 1769–1788.

2163 Jamieson, S.S.R., Ewertowski, M.W., Evans, D.J.A., 2015. Rapid advance of two mountain glaciers
2164 in response to mine-related debris loading, *J. Geophys. Res.: Earth Surf.* 120, 1418–1435.

- 2165 Jansson, K.N., Glasser, N.F., 2005. Using Landsat 7 ETM+ imagery and Digital Terrain Models for
2166 mapping glacial lineaments on former ice sheet beds. *Int. J. Remote Sens.* 26(18), 3931–3941.
- 2167 Jansson, K.N., Kleman, J., Marchant, D.R., 2002. The succession of ice-flow patterns in north-central
2168 Québec-Labrador, Canada. *Quat. Sci. Rev.* 21(4), 503–523.
- 2169 Jansson, K.N., Stroeve, A.P., Kleman, J., 2003. Configuration and timing of Ungava Bay ice
2170 streams, Labrador–Ungava, Canada. *Boreas* 32(1), 256–262.
- 2171 Johnson, M.D., Fredin, O., Ojala, A.E.K., Peterson, G., 2015. Unravelling Scandinavian
2172 geomorphology: the LiDAR revolution. *GFF* 137, 245–251.
- 2173 Jónsson, S.A., Schomacker, A., Benediktsson, Í.Ö., Ingólfsson, Ó., Johnson, M.D., 2014. The drumlin
2174 field and the geomorphology of the Múlajökull surge-type glacier, central Iceland.
2175 *Geomorphology* 207, 213–220.
- 2176 Jónsson, S.A., Benediktsson, Í.Ö., Ingólfsson, Ó., Schomacker, A., Bergsdóttir, H.L., Jacobsson,
2177 W.R., Linderson, H., 2016. Submarginal drumlin formation and late Holocene history of
2178 Fláajökull, southeast Iceland. *Ann. Glaciol.* 57, 128–141.
- 2179 Jorge, M.G., Brennand, T.A., 2017a. Measuring (subglacial) bedform orientation, length, and
2180 longitudinal asymmetry—Method assessment. *PloS one*, 12(3), e0174312.
- 2181 Jorge, M.G., Brennand, T.A., 2017b. Semi-automated extraction of longitudinal subglacial bedforms
2182 from digital terrain models—Two new methods. *Geomorphology*, 288, 148–163.
- 2183 Juyal, N., Thakkar, P.S., Sundriyal, Y.P., 2011. Geomorphic evidence of glaciations around Mount
2184 Kailash (Inner Kora): implication to past climate. *Current Science* 100(4), 535–541.
- 2185 Kassab, C., Wang, J., Harbor, J., 2013. Glacial geomorphology of the Dalijia Shan region,
2186 northeastern Tibetan Plateau. *J. Maps* 9(1), 98–105.
- 2187 Kelley, S.E., Kaplan, M.R., Schaefer, J.M., Andersen, B.G., Barrell, D.J.A., Putnam, A.E., Denton,
2188 G.H., Schwartz, R., Finkel, R.C., Doughty, A.M., 2014. High-precision ^{10}Be chronology of
2189 moraines in the Southern Alps indicates synchronous cooling in Antarctica and New Zealand
2190 42,000 years ago. *Earth Planet. Sci. Lett.* 405, 194–206.
- 2191 Kendall, P.F. 1902. A system of glacier lakes in the Cleveland Hills. *Quart. J. Geol. Soc. Lond.* 58,
2192 471–571.
- 2193 Kerschner, H., Kaser, G., Sailer, R., 2000. Alpine Younger Dryas glaciers as palaeo-precipitation
2194 gauges. *Ann. Glaciol.* 31, 80–84.
- 2195 Kienholz, H., 1977. Kombinierte geomorphologische Gefahrenkarte 1:10,000 von Grindelwald.
2196 *Geographica Bernensia*, G4: 1–204.
- 2197 King, E.C., Woodward, J., Smith, A. M., 2007. Seismic and radar observations of subglacial bed
2198 forms beneath the onset zone of Rutford Ice Stream, Antarctica. *J. Glaciol.* 53(183), 665–672.
- 2199 King, E.C., Hindmarsh, R.C., Stokes, C.R., 2009. Formation of mega-scale glacial lineations observed
2200 beneath a West Antarctic ice stream. *Nat. Geosci.* 2(8), 585–588.
- 2201 King, E.C., Pritchard, H.D., Smith, A.M., 2016a. Subglacial landforms beneath Rutford Ice Stream,
2202 Antarctica: detailed bed topography from ice-penetrating radar. *Earth Syst. Sci. Data* 8(1),
2203 151–158.
- 2204 King, O., Hambrey, M.J., Irvine-Fynn, T.D.L., Holt, T.O., 2016b. The structural, geometric and
2205 volumetric changes of a polythermal Arctic glacier during a surge cycle: Comfortlessbreen,
2206 Svalbard. *Earth Surf. Process. Land.* 41, 162–177.
- 2207 King, O., Quincey, D.J., Carrivick, J.L., Rowan, A.V., 2017. Spatial variability in mass loss of
2208 glaciers in the Everest region, central Himalayas, between 2000 and 2015. *The Cryosphere*
2209 11, 407–426.
- 2210 Kirchner, N., Greve, R., Stroeve, A.P., Heyman, J., 2011. Paleoglaciological reconstructions for the
2211 Tibetan Plateau during the last glacial cycle: evaluating numerical ice sheet simulations
2212 driven by GCM-ensembles. *Quat. Sci. Rev.* 30, 248–267.
- 2213 Kirkbride, M.P., Winkler, S., 2012. Correlation of Late Quaternary moraines: impact of climate
2214 variability, glacier response, and chronological resolution. *Quat. Sci. Rev.* 46, 1–29.
- 2215 Kjær, K.H., Krüger, J., 2001. The final phase of dead-ice moraine development: processes and
2216 sediment architecture, Kotlujökull, Iceland. *Sedimentology* 48, 935–952.
- 2217 Kjær, K.H., Houmark-Nielsen, M., Richardt, N., 2003. Ice-flow patterns and dispersal of erratics at the
2218 southwestern margin of the last Scandinavian Ice Sheet: signature of paleo-ice streams.
2219 *Boreas* 32, 130–148.

- 2220 Kjær, K.H., Korsgaard, N.J., Schomacker, A., 2008. Impact of multiple glacier surges – a
2221 geomorphological map from Brúarjökull, East Iceland. *J. Maps* 4, 5–20.
- 2222 Kłapyta, P., 2013. Application of Schmidt hammer relative age dating to Late Pleistocene moraines
2223 and rock glaciers in the Western Tatra Mountains, Slovakia. *Catena* 111, 104–121.
- 2224 Klassen, R.A., 1993. Quaternary geology and glacial history of Bylot Island, Northwest Territories.
2225 Geological Survey of Canada Memoir 429.
- 2226 Kleman, J., 1990. On the use of glacial striae for reconstruction of palaeo-ice sheet flow patterns.
2227 *Geogr. Ann.* 72A, 217–236.
- 2228 Kleman, J., 1992. The Palimpsest Glacial Landscape in Northwestern Sweden. Late Weichselian
2229 Deglaciation Landforms and Traces of Older West-Centered Ice Sheets. *Geogr. Ann.* 74A(4),
2230 305–325.
- 2231 Kleman, J., Borgström, I., 1996. Reconstruction of Palaeo-Ice Sheets: The Use of Geomorphological
2232 Data. *Earth Surf. Process. Land.* 21, 893–909.
- 2233 Kleman, J., Stroeve, A.P., 1997. Preglacial surface remnants and Quaternary glacial regimes in
2234 northwestern Sweden. *Geomorphology* 19(1–2), 35–54.
- 2235 Kleman, J., Hättestrand, C., Borgström, I., Stroeve, A., 1997. Fennoscandian palaeoglaciology
2236 reconstructed using a glacial geological inversion model. *J. Glaciol.* 43(144), 283–299.
- 2237 Kleman, J., Fastook, J., Stroeve, A.P., 2002. Geologically and geomorphologically constrained
2238 numerical model of Laurentide Ice Sheet inception and build-up. *Quat. Int.* 95–96, 87–98.
- 2239 Kleman, J., Hättestrand, C., Stroeve, A.P., Jansson, K.N., De Angelis, H., Borgström, I., 2006.
2240 Reconstruction of palaeo-ice sheets; inversion of their glacial geomorphological record. In:
2241 Knight, P.G. (Ed.), *Glacier Science and Environmental Change*. Blackwell, Oxford, pp. 192–
2242 199.
- 2243 Kleman, J., Stroeve, A.P., Lundqvist, J., 2008. Patterns of Quaternary ice sheet erosion and
2244 deposition in Fennoscandia. *Geomorphology* 97, 73–90.
- 2245 Kleman, J., Jansson, K., De Angelis, H., Stroeve, A.P., Hättestrand, C., Alm, G., Glasser, N., 2010.
2246 North American Ice Sheet build-up during the last glacial cycle, 115–21 kyr. *Quat. Sci. Rev.*
2247 29, 2036–2051.
- 2248 Klimaszewski, M., 1990. Thirty years of detailed geomorphological mapping. *Geogr. Pol.* 58, 11–18.
- 2249 Kneisel, C., Lehmkuhl, F., Winkler, S., Tresselt, E., Schröder, H., 1998. *Legende für*
2250 *geomorphologische Kartierungen in Hochgebirgen (GMK Hochgebirge)*. Trier. *Geogr. Stud.*
2251 18, 1–24.
- 2252 Knight, J., Mitchell, W., Rose, J., 2011. Geomorphological Field Mapping. In: Smith, M.J., Paron, P.,
2253 Griffiths, J. (Eds.), *Geomorphological Mapping: Methods and Applications*. Developments in
2254 Earth Surface Processes 15. Elsevier, London, pp. 151–188.
- 2255 Kocurek, G., Ewing, R.C., Mohrig, D., 2010. How do bedform patterns arise? New views on the role
2256 of bedform interactions within a set of boundary conditions. *Earth Surf. Process. Land.* 35,
2257 51–63.
- 2258 Korsgaard, N.J., Schomacker, A., Benediktsson, Í.Ö., Larsen, N.K., Ingólfsson, Ó., Kjær, K.H., 2015.
2259 Spatial distribution of erosion and deposition during a glacier surge: Brúarjökull, Iceland.
2260 *Geomorphology* 250, 258–270.
- 2261 Kraak, M.-J., Oreming, F.J., 2010. *Cartography: visualization of spatial data* (3rd Edition). Routledge,
2262 London.
- 2263 Kronberg, P., 1984. *Photogeologie. Eine Einführung in die Grundlagen und Methoden der*
2264 *geologischen Auswertung von Luftbildern*. Enke, Stuttgart.
- 2265 Krüger, J., 1994. Glacial processes, sediments, landforms and stratigraphy in the terminus region of
2266 Mýrdalsjökull, Iceland. *Folia Geographica Danica* 21, 1–233.
- 2267 Krüger, J., 1995. Origin, chronology and climatological significance of annual moraine ridges at
2268 Mýrdalsjökull, Iceland. *The Holocene* 5, 420–427.
- 2269 Krüger, J., Kjær, K.H., 2000. De-icing progression of ice-cored moraines in a humid, subpolar
2270 climate, Kötlujökull, Iceland. *The Holocene* 10, 737–747.
- 2271 Krygowski B. (Ed.), 1963. *Mapa Geomorfologiczna Niziny Wielkopolsko-Kujawskiej*
2272 *[Geomorphological Map of Wielkopolska-Kujawy lowland]*. 1:300 000 Map. Adam
2273 Mickiewicz University.

- 2274 Kuhle, M., 1990. Quantificational reductionism as a risk in geography instanced by the 1:25,000
2275 Geomorphological Map of the Federal Republic of Germany. *Geogr. Pol.* 58, 41–54.
- 2276 Lam, N.S.N., Quattrocchi, D.A., 1992. On the issues of scale, resolution, and fractal analysis in the
2277 mapping sciences. *The Prof. Geogr.* 44(1), 88–98.
- 2278 Lane, S.N., Reid, S.C., Westaway, R.M., Hicks, D.M., 2005. Remotely Sensed Topographic Data for
2279 River Channel Research: The Identification, Explanation and Management of Error. In: R.E.J.
2280 Kelly, Drake, N.A., Barr, S.L. (Eds.), *Spatial Modelling of the Terrestrial Environment*. John
2281 Wiley & Sons, Ltd, Chichester, UK, pp. 113–136.
- 2282 Lardeux, P., Glasser, N., Holt, T., Hubbard, B., 2015. Glaciological and geomorphological map of
2283 Glacier Noir and Glacier Blanc, French Alps. *J. Maps* 12, 582–596.
- 2284 Leser, H., 1983. Anwendung und Auswertung geomorphologischer Kartierungen und Karten. *Mater.*
2285 *Physiogeogr.* 5, 5–13.
- 2286 Leser, H., Stäblein, G., 1975. Geomorphologische Kartierung – Richtlinien zur Herstellung
2287 geomorphologischer Karten 1: 25,000. *Berliner Geographische Abhandlungen Sonderheft*, 1–
2288 33.
- 2289 Levy, L.B., Larsen, N.K., Davidson, T.A., Strunk, A., Olsen, J., Jeppesen, E., 2017. Contrasting
2290 evidence of Holocene ice margin retreat, south-western Greenland. *J. Quat. Sci.* 32, 604–616.
- 2291 Li, Y., Li, Y., Chen, Y., Lu, X., 2016. Presumed Little Ice Age glacial extent in the eastern Tian Shan,
2292 China. *J. Maps* 12(1), 71–78.
- 2293 Lidmar-Bergström, K., Elvhage, C., Ringberg, B., 1991. Landforms in Skane, south Sweden.
2294 *Geografiska Annaler* 73A, 61–91.
- 2295 Lifton, N., Beel, C., Hättestrand, C., Kassab, C., Rogozhina, I., Heermance, R., Oskin, M., Burbank,
2296 D., Blomdin, R., Gribenski, N., Caffee, M., 2014. Constraints on the late Quaternary glacial
2297 history of the Inylchek and Sary-Dzaz valleys from in situ cosmogenic ^{10}Be and ^{26}Al ,
2298 eastern Kyrgyz Tian Shan. *Quat. Sci. Rev.* 101, 77–90.
- 2299 Lillesand, T.M., Kiefer, R.W., Chipman, J.W., 2015. Remote Sensing and Image Interpretation (7th
2300 Edition). John Wiley & Sons, Hoboken, USA, 768 pp.
- 2301 Lindholm, M.S., Heyman, J., 2016. Glacial geomorphology of the Maidika region, Tibetan Plateau. *J.*
2302 *Maps* 12(5), 797–803.
- 2303 Livingstone, S.J., Ó Cofaigh, C., Evans, D.J.A., 2008. The glacial geomorphology of the central
2304 sector of the British-Irish Ice Sheet. *J. Maps* 4, 358–377.
- 2305 Livingstone, S.J., Evans, D.J.A., Ó Cofaigh, C., 2010. Re-advance of Scottish ice into the Solway
2306 Lowlands (Cumbria, UK) during the Main Late Devensian deglaciation. *Quat. Sci. Rev.* 29,
2307 2544–2570.
- 2308 Livingstone, S.J., Ó Cofaigh, C., Stokes, C.R., Hillenbrand, C.-D., Vieli, A., Jamieson, S.S.R., 2012.
2309 Antarctic palaeo-ice streams. *Earth-Sci. Rev.* 111, 90–128.
- 2310 Livingstone, S.J., Storrar, R.D., Hillier, J.K., Stokes, C.R., Clark, C.D., Tarasov, L., 2015. An ice-
2311 sheet scale comparison of eskers with modelled subglacial drainage routes. *Geomorphology*
2312 246, 104–112.
- 2313 Loibl, D., Hochreuther, P., Schulte, P., Hülle, D., Zhu H, Bräuning, A., Lehmkuhl, F., 2015. Toward a
2314 late Holocene glacial chronology for the eastern Nyainqêntanglha Range, southeastern Tibet.
2315 *Quat. Sci. Rev.* 107, 243–259.
- 2316 Lønne, I., 2016. A new concept for glacial geological investigations of surges, based on High-Arctic
2317 examples (Svalbard). *Quat. Sci. Rev.*, 132, 74–100.
- 2318 Lovell, H., 2014. On the ice-sediment-landform associations of surging glaciers on Svalbard.
2319 Unpublished PhD thesis, Queen Mary University of London, 312 pp.
- 2320 Lovell, H., Boston, C.M., 2017. Glacitectonic composite ridge systems and surge-type glaciers: an
2321 updated correlation based on Svalbard, Norway. *arktos* 3, 2.
- 2322 Lovell, H., Stokes, C.R., Bentley, M.J., 2011. A glacial geomorphological map of the Seno Skyring-
2323 Seno Otway-Strait of Magellan region, southernmost Patagonia. *J. Maps* 7, 318–339.
- 2324 Lovell, H., Stokes, C.R., Bentley, M.J., Benn, D.I., 2012. Evidence for rapid ice flow and proglacial
2325 lake evolution around the central Strait of Magellan region, southernmost Patagonia. *J. Quat.*
2326 *Sci.* 27, 625–638.

- 2327 Lovell, H., Benn, D.I., Lukas, S., Spagnolo, M., Cook, S.J., Swift, D.A., Clark, C.D., Yde, J.C.,
2328 Watts, T.P., 2018. Geomorphological investigation of multiphase glacial tectonic composite
2329 ridge systems in Svalbard. *Geomorphology* 300, 176–188.
- 2330 Lueder, D.R., 1959. *Aerial Photographic Interpretation – Principles and Applications*. McGraw-Hill,
2331 New York.
- 2332 Lukas, S., 2002. Geomorphological evidence for the pattern of deglaciation around the Drumochter
2333 Pass, Central Grampian Highlands, Scotland. Unpublished MSc thesis, Ruhr-University of
2334 Bochum, Germany, 115 pp.
- 2335 Lukas, S., 2005. A test of the englacial thrusting hypothesis of 'hummocky' moraine formation – case
2336 studies from the north-west Highlands, Scotland. *Boreas* 34, 287–307.
- 2337 Lukas, S., 2006. Morphostratigraphic principles in glacier reconstruction -a perspective from the
2338 British Younger Dryas. *Prog. Phys. Geogr.* 30, 719–736.
- 2339 Lukas, S., 2007a. Early-Holocene glacier fluctuations in Krundalen, south central Norway: palaeo-
2340 glacier dynamics and palaeoclimate. *The Holocene* 17, 585–598.
- 2341 Lukas, S., 2007b. 'A test of the englacial thrusting hypothesis of "hummocky" moraine formation:
2342 case studies from the northwest Highlands, Scotland': Reply to comments. *Boreas* 36, 108–
2343 113.
- 2344 Lukas, S., 2011. Ice-cored moraines. In: Singh, V., Singh, P. Haritashya, U.K. (Eds.), *Encyclopedia of*
2345 *Snow, Ice and Glaciers*. Springer, Heidelberg, pp. 616–619.
- 2346 Lukas, S., 2012. Processes of annual moraine formation at a temperate alpine valley glacier: insights
2347 into glacier dynamics and climatic controls. *Boreas* 41(3), 463–480.
- 2348 Lukas, S., Benn, D.I., 2006. Retreat dynamics of Younger Dryas glacier in the far NW Scottish
2349 Highlands reconstructed from moraine sequences. *Scott. Geogr. J.* 122, 308–325.
- 2350 Lukas, S., Lukas, T., 2006. A glacial geological and geomorphological map of the far NW Highlands,
2351 Scotland. Parts 1 and 2. *J. Maps* 2, 43–56, 56–58.
- 2352 Lukas, S., Sass, O., 2011. The formation of Alpine lateral moraines inferred from sedimentology and
2353 radar reflection patterns: a case study from Gornergletscher, Switzerland. *Geological Society,*
2354 *London, Special Publications* 354, 77–92.
- 2355 Lukas, S., Nicholson, L.I., Ross, F.H., Humlum, O., 2005. Formation, meltout processes and
2356 landscape alteration of High-Arctic ice-cored moraines—examples from Nordenskiöld Land,
2357 Central Spitsbergen. *Polar Geogr.* 29, 157–187.
- 2358 Lukas, S., Nicholson, L.I., Humlum, O., 2007. Comment on Lønne and Lyså (2005): "Deglaciation
2359 dynamics following the Little Ice Age on Svalbard: Implications for shaping of landscapes at
2360 high latitudes", *Geomorphology* 72, 300–319. *Geomorphology* 84, 145–149.
- 2361 Lukas, S., Benn, D.I., Boston, C.M., Brook, M.S., Coray, S., Evans, D.J.A., Graf, A., Kellerer-
2362 Pirklbauer-Eulenstein, A., Kirkbride, M.P., Krabbendam, M., Lovell, H., Machiedo, M.,
2363 Mills, S.C., Nye, K., Reinardy, B.T.I., Ross, F.H., Signer, M., 2013. Clast shape analysis and
2364 clast transport paths in glacial environments: A critical review of methods and the role of
2365 lithology. *Earth-Sci. Rev.* 121, 96–116.
- 2366 Lukas, S., Preusser, F., Evans, D.J.A., Boston, C.M., Lovell, H., 2017. Chapter 2 – The Quaternary.
2367 In: Griffiths, J.S., Martin, C.J. (Eds.), *Engineering Geology and Geomorphology of Glaciated*
2368 *and Periglaciated Terrains: Engineering Group Working Party Report*. Engineering Geology
2369 Special Publication 28. The Geological Society, London, pp. 31–57.
- 2370 MacLachlan, J.C., Eyles, C.H., 2013. Quantitative geomorphological analysis of drumlins in the
2371 Peterborough drumlin field, Ontario, Canada. *Geogr. Ann.* 95A, 125–144.
- 2372 Mäkinen, J., Kajuutti, K., Palmu, J., Ojala, A., Ahokangas, E., 2017. Triangular-shaped landforms
2373 reveal subglacial drainage routes in SW Finland. *Quat. Sci. Rev.* 164, 37–53.
- 2374 Małeck, J., Lovell, H., Ewertowski, W., Górski, Ł., Kurczba, T., Latos, B., Miara, M., Piniarska, D.,
2375 Płocieniczak, J., Sowada, T., Spiralski, M., Warczowska, A., Rabatel, A., 2018. The
2376 glacial landsystem of a tropical glacier: Charquini Sur, Bolivian Andes. *Earth Surf. Proc.*
2377 *Land.*, in press. doi: 10.1002/esp.4417
- 2378 Marc, O., Hovius, N., 2015. Amalgamation in landslide maps: effects and automatic detection. *Nat.*
2379 *Hazards Earth Syst. Sci.* 15, 723–733.
- 2380 Margold, M., Jansson, K.N., 2011. Glacial geomorphology and glacial lakes of central Transbaikalia,
2381 Siberia, Russia. *J. Maps* 7, 18–30.

2382 Margold, M., Jansson, K.N., 2012. Evaluation of data sources for mapping glacial meltwater features.
2383 Int. J. Remote Sens. 33, 2355–2377.

2384 Margold, M., Jansson, K.N., Kleman, J., Stroeven, A.P., 2011. Glacial meltwater landforms of central
2385 British Columbia. J. Maps 7, 486–506.

2386 Margold, M., Stokes, C.R., Clark, C.D., Kleman, J., 2015a. Ice streams in the Laurentide Ice Sheet: a
2387 new mapping inventory. J. Maps 11, 380–395.

2388 Margold, M., Stokes, C.R., Clark, C.D., 2015b. Ice streams in the Laurentide Ice Sheet: Identification,
2389 characteristics and comparison to modern ice sheets. Earth-Sci. Rev. 143, 117–146.

2390 Margold, M., Stokes, C.R., Clark, C.D., 2018. Reconciling records of ice streaming and ice margin
2391 retreat to produce a palaeogeographic reconstruction of the deglaciation of the Laurentide Ice
2392 Sheet. Quat. Sci. Rev. 189, 1–30.

2393 Mather, A.E., Mills, S., Stokes, M., Fyfe, R., 2015. Ten years on: what can Google Earth offer the
2394 Geoscience community? Geology Today 31(6), 216–221.

2395 May, J.H., Zech, J., Zech, R., Preusser, F., Argollo, J., Kubik, P.W., Veit, H., 2011. Reconstruction of
2396 a complex late Quaternary glacial landscape in the Cordillera de Cochabamba (Bolivia) based
2397 on a morphostratigraphic and multiple dating approach. Quat. Res. 76(1), 106–118.

2398 McDougall, D.A., 2001. The geomorphological impact of Loch Lomond (Younger Dryas) Stadial
2399 plateau icefields in the central Lake District, northwest England. J. Quat. Sci. 16, 531–543.

2400 McDougall, D.A., 2013. Glaciation style and the geomorphological record: evidence for Younger
2401 Dryas glaciers in the eastern Lake District, northwest England. Quat. Sci. Rev. 73, 48–58.

2402 McHenry, M., Dunlop, P., 2016. The subglacial imprint of the last Newfoundland Ice Sheet, Canada.
2403 J. Maps 12(3), 462–483.

2404 Melander, O., 1975. Geomorfologiska kartbladet 29 I Kebnekaise. Statens naturvårdsverk. 78 pp.,
2405 map scale 1:250,000.

2406 Mertes, J.R., Gulley, J.D., Benn, D.I., Thompson, S.S., Nicholson, L.I., 2017. Using structure-from-
2407 motion to create glacier DEMs and orthoimagery from historical terrestrial and oblique aerial
2408 imagery. Earth Surf. Process. Land. 42, 2350–2364.

2409 Midgley, N.G., Tonkin, T.N., 2017. Reconstruction of former glacier surface topography from archive
2410 oblique aerial images. Geomorphology 282, 18–26.

2411 Midgley, N.G., Cook, S.J., Graham, D.J., Tonkin, T.N., 2013. Origin, evolution and dynamic context
2412 of a Neoglacial lateral–frontal moraine at Austre Lovénbreen, Svalbard. Geomorphology 198,
2413 96–106.

2414 Midgley, N.G., Tonkin, T.N., Graham, D.J., Cook, S.J., 2018. Evolution of high-Arctic glacial
2415 landforms during deglaciation. Geomorphology 311, 63–75.

2416 Miller, H., Cotterill, C.J., Bradwell, T., 2014. Glacial and paraglacial history of the Troutbeck Valley,
2417 Cumbria, UK: integrating airborne LiDAR, multibeam bathymetry, and geological field
2418 mapping. Proc. Geol. Assoc. 125, 31–40.

2419 Mills, S.C., Grab, S.W., Rea, B.R., Carr, S.J., Farrow, A., 2012. Shifting westerlies and precipitation
2420 patterns during the Late Pleistocene in southern Africa determined using glacier
2421 reconstruction and mass balance modelling. Quaternary Science Reviews 55, 145–159.

2422 Mitchell, W.A., Riley, J.M., 2006. Drumlin map of the Western Pennines and southern Vale of Eden,
2423 Northern England, UK. J. Maps 2, 10–16.

2424 Mollard, J.D., Janes, J.R., 1984. Airphoto Interpretation and the Canadian Landscape. Energy, Mines
2425 and Resources Canada, Ottawa.

2426 Möller, P., Dowling, T.P.F., 2016. Streamlined subglacial bedforms on the Närke plain, south-central
2427 Sweden – Areal distribution, morphometrics, internal architecture and formation. Quat. Sci.
2428 Rev. 146, 182–215.

2429 Möller, P., Dowling, T.P.F., 2018. Equifinality in glacial geomorphology: instability theory examined
2430 via ribbed moraine and drumlins in Sweden. GFF, in press.

2431 Morén, B., Heyman, J., Stroeven, A.P., 2011. Glacial geomorphology of the central Tibetan Plateau.
2432 J. Maps 7, 115–125.

2433 Murray, A.B., Goldstein, E.B., Coco, G., 2014. The shape of patterns to come: from initial formation
2434 to long-term evolution. Earth Surf. Process. Land. 39, 62–70.

- Napieralski, J., Hubbard, A., Li, Y.K., Harbor, J., Stroeven, A.P., Kleman, J., Alm, G., Jansson, K.N., 2007a. Towards a GIS assessment of numerical ice sheet model performance using geomorphological data. *J. Glaciol.* 53 (180), 71–83.
- Napieralski, J., Harbor, J., Li, Y., 2007b. Glacial geomorphology and geographic information systems. *Earth-Sci. Rev.* 85, 1–22.
- Näslund, J.O., Rodhe, L., Fastook, J.L., Holmlund, P., 2003. New ways of studying ice sheet flow directions and glacial erosion by computer modelling — examples from Fennoscandia. *Quat. Sci. Rev.* 22, 245–258.
- Noh, M.J., Howat, I.M., 2015. Automated stereo-photogrammetric DEM generation at high latitudes: Surface Extraction from TIN-Based Search Minimization (SETSM) validation and demonstration over glaciated regions. *GIScience Remote Sens.* 52(2), 198–217.
- Norris, S.L., Margold, M., Froese, D.G., 2017. Glacial landforms of northwest Saskatchewan. *J. Maps* 13, 600–607.
- Norris, S.L., Evans, D.J.A., Cofaigh, C.Ó., 2018. Geomorphology and till architecture of terrestrial palaeo-ice streams of the southwest Laurentide Ice Sheet: A borehole stratigraphic approach. *Quat. Sci. Rev.* 186, 186–214.
- Nuth, C., Kääb, A., 2011. Co-registration and bias corrections of satellite elevation data sets for quantifying glacier thickness change. *The Cryosphere* 5, 271–290.
- Ó Cofaigh, C., 2012. Ice sheets viewed from the ocean: the contribution of marine science to understanding modern and past ice sheets. *Phil. Trans. R. Soc. A* 370(1800), 5512–5539.
- Ó Cofaigh, C., Evans, D.J.A., Smith, I.R., 2010. Large-scale reorganization and sedimentation of terrestrial ice streams during late Wisconsinan Laurentide Ice Sheet deglaciation. *Geol. Soc. Am. Bull.* 122, 743–756.
- Ó Cofaigh, C., Dowdeswell, J.A., Jennings, A.E., Hogan, K.A., Kilfeather, A., Hiemstra, J.F., Noormets, R.M., Evans, J., McCarthy, D.J., Andrews, J.T., Lloyd, J.M., Moros, M., 2013. An extensive and dynamic ice sheet on the West Greenland shelf during the last glacial cycle. *Geology* 41 (2), 219–222.
- Ojala, A.E.K., 2016. Appearance of De Geer moraines in southern and western Finland — Implications for reconstructing glacier retreat dynamics. *Geomorphology* 255, 16–25.
- Ojala, A.E.K., Putkinen, N., Palmu, J.P., Nenonen, K., 2015. Characterization of De Geer moraines in Finland based on LiDAR DEM mapping. *GFF* 137(4), 304–318.
- Orkhonselenge, A., 2016. Glacial Geomorphology of Mt. Munkh Saridag in the Khuvsgul Mountain Range, Northern Mongolia. *Géomorphologie: relief, processus, environnement* 22(4), 389–398.
- Ottesen, D., Dowdeswell, J.A., 2006. Assemblages of submarine landforms produced by tidewater glaciers in Svalbard. *J. Geophys. Res.* 111, F01016.
- Ottesen, D., Dowdeswell, J.A., Rise, L., 2005. Submarine landforms and the reconstruction of fast-flowing ice streams within a large Quaternary ice sheet: the 2500 km long Norwegian-Svalbard margin (57 degrees – 80 degrees N). *Geol. Soc. Am. Bull.* 117 (7–8), 1033–1050.
- Ottesen, D., Dowdeswell, J.A., 2006. Assemblages of submarine landforms produced by tidewater glaciers in Svalbard. *J. Geophys. Res.* 111, F01016.
- Ottesen, D., Stokes, C.R., Rise, L., Olsen, L., 2008a. Ice-sheet dynamics and ice streaming along the coastal parts of northern Norway. *Quat. Sci. Rev.* 27, 922–940.
- Ottesen, D., Dowdeswell, J.A., Benn, D.I., Kristensen, L., Christiansen, H.H., Christensen, O., Hansen, L., Lebesbye, E., Forwick, M., Vorren, T.O., 2008b. Submarine landforms characteristic of glacier surges in two Spitsbergen fjords. *Quat. Sci. Rev.* 27, 1583–1599.
- Ottesen, D., Stokes, C.R., Bøe, R., Rise, L., Longva, O., Thorsnes, T., Olesen, O., Bugge, T., Lepland, A., Hestvik, O.B., 2016. Landform assemblages and sedimentary processes along the Norwegian Channel Ice Stream. *Sediment. Geol.* 338, 115–137.
- Ottesen, D., Dowdeswell, J.A., Bellec, V.K., Bjarnadóttir, L.R., 2017. The geomorphic imprint of glacier surges into open-marine waters: Examples from eastern Svalbard. *Mar. Geol.* 392, 1–29.

- 2488 Otto, J.-C., Smith, M.J., 2013. Section 2.6: Geomorphological mapping. In: Clarke, L. (Ed.),
2489 Geomorphological Techniques (Online Edition). British Society for Geomorphology, London.
2490 ISSN: 2047-0371.
- 2491 Owen, L.A., Finkel, R.C., Barnard, P.L., Haizhou, M., Asahi, K., Caffee, M.W., Derbyshire, E., 2005.
2492 Climatic and topographic controls on the style and timing of Late Quaternary glaciation
2493 throughout Tibet and the Himalaya defined by ¹⁰Be cosmogenic radionuclide surface
2494 exposure dating. *Quat. Sci. Rev.* 24(12), 1391–1411.
- 2495 Paron, P., Claessens, L., 2011. Makers and Users of Geomorphological Maps. In: Smith, M.J., Paron,
2496 P., Griffiths, J.S. (Eds.), *Geomorphological Mapping: Methods and applications.*
2497 *Developments in Earth Surface Processes*, Volume 15. Elsevier, Oxford; pp. 75–106.
- 2498 Partsch, J., 1894. Die Vergletscherung des Riesengebirges zur Eiszeit. *Forschungen zur Deutschen*
2499 *Landes- und Volkskunde* VIII/2, 103–194.
- 2500 Patton, H., Hubbard, A., Andreassen, K., Auriac, A., Whitehouse, P.L., Stroeve, A.P., Shackleton,
2501 C., Winsborrow, M., Heyman, J., Hall, A.M., 2017a. Deglaciation of the Eurasian ice sheet
2502 complex. *Quat. Sci. Rev.* 169, 148–172.
- 2503 Patton, H., Hubbard, A., Bradwell, T., Schomacker, A., 2017b. The configuration, sensitivity and
2504 rapid retreat of the Late Weichselian Icelandic ice sheet. *Earth-Sci. Rev.* 166, 223–245.
- 2505 Pearce, D., Rea, B.R., Bradwell, T., McDougall, D., 2014. Glacial geomorphology of the Tweedsmuir
2506 Hills, Central Southern Uplands, Scotland. *J. Maps* 10(3), 457–465.
- 2507 Pearce, D.M., Mair, D.W.F., Rea, B.R., Lea, J.M., Schofield, J.E., Kamenos, N., Schoenrock, K.,
2508 2018. The glacial geomorphology of upper Godthåbsfjord (Nuup Kangerlua) in southwest
2509 Greenland. *J. Maps* 14, 45–55.
- 2510 Penck, A., Brückner, E., 1901/1909. *Die Alpen im Eiszeitalter*. Tauchnitz, Leipzig.
- 2511 Peterson, G., Johnson, M.D., Smith, C.A., 2017. Glacial geomorphology of the south Swedish uplands
2512 – focus on the spatial distribution of hummock tracts. *J. Maps* 13, 534–544.
- 2513 Petrie, G., Price, R.J., 1966. Photogrammetric measurements of the ice wastage and morphological
2514 changes near the Casement Glacier, Alaska. *Can. J. Earth Sci.* 3, 827–840.
- 2515 Phillips, E., Evans, D.J.A., Atkinson, N., Kendall, A., 2017. Structural architecture and glaciectonic
2516 evolution of the Mud Buttes cupola hill complex, southern Alberta, Canada. *Quat. Sci. Rev.*
2517 164, 110–139.
- 2518 Pike, R.J., 1988. The geometric signature: quantifying landslide-terrain types from digital elevation
2519 models. *Math. Geol.* 20, 491–511.
- 2520 Pipaud, I., Loibl, D., Lehmkuhl, F., 2015. Evaluation of TanDEM-X elevation data for
2521 geomorphological mapping and interpretation in high mountain environments — A case study
2522 from SE Tibet, China. *Geomorphology* 246, 232–254.
- 2523 Popper, K.R., 1972. *Objective Knowledge*. Oxford University Press, Oxford.
- 2524 Prest, V.K., 1983. Canada's Heritage of Glacial Features. GSC Miscellaneous Report 28.
- 2525 Prest, V.K., Grant, D.R., Rampton, V.N. 1968. Glacial map of Canada. Geological Survey of Canada,
2526 Map 1253A.
- 2527 Priamonoosov, A.P., Kuznetsova, E.J., Abaturova, I.V., 2000. National geological map of the Russian
2528 Federation, Map of Quaternary Formations, Polar-Ural series, map sheet Q-41-XII Kharp.
2529 Ministry of Natural Resources of the Russian Federation. 1:200,000.
- 2530 Price, R.J., 1961. The Deglaciation of the Upper Tweed Basin. Unpublished PhD thesis, University of
2531 Edinburgh.
- 2532 Price, R.J., 1963. A glacial meltwater drainage system in Peebleshire, Scotland. *Scott. Geogr. Mag.*
2533 79, 133–141.
- 2534 Price, R.J., 1966. Eskers near the Casement Glacier, Alaska. *Geogr. Ann.* 48, 111–125.
- 2535 Price, R.J., 1970. Moraines at Fjallsjökull, Iceland. *Arctic Alp. Res.* 2, 27–42.
- 2536 Principato, S.M., Moyer, A.N., Hampsch, A.G., Ipsen, H.A., 2016. Using GIS and streamlined
2537 landforms to interpret palaeo-ice flow in northern Iceland. *Boreas* 45, 470–482.
- 2538 Punkari, M., 1980. The ice lobes of the Scandinavian ice sheet during the deglaciation in Finland.
2539 *Boreas* 9(4), 307–310.
- 2540 Punkari, M., 1995. Glacial flow systems in the zone of confluence between the Scandinavian and
2541 Novaja Zemlya Ice Sheets. *Quat. Sci. Rev.* 14(6), 589–603.

- 2542 Putniņš, A., Henriksen, M., 2017. Reconstructing the flow pattern evolution in inner region of the
2543 Fennoscandian Ice Sheet by glacial landforms from Gausdal Vestfjell area, south-central
2544 Norway. *Quat. Sci. Rev.* 163, 56–71.
- 2545 Rabus, B., Eineder, M., Roth, A., Bamler, R., 2003. The shuttle radar topography mission—a new
2546 class of digital elevation models acquired by spaceborne radar. *ISPRS J. Photogramm.*
2547 *Remote Sens.* 57(4), 241–262.
- 2548 Rączkowska Z., Zwoliński Z., 2015. Digital geomorphological map of Poland. *Geogr. Pol.* 88(2),
2549 205–210.
- 2550 Raistrick, A., 1933. The glacial and post-glacial periods in West Yorkshire. *Proc. Geol. Assoc.* 44,
2551 263–269.
- 2552 Raisz, E.J., 1962. *Principles of cartography*. McGraw-Hill, New York, 315 pp.
- 2553 Rea, B.R., Whalley, W.B., Dixon, T.S., Gordon, J.E., 1999. Plateau icefields as contributing areas to
2554 valley glaciers and the potential impact on reconstructed ELAs: a case study from the Lyngen
2555 Alps, North Norway. *Ann. Glaciol.* 28, 97–102.
- 2556 Reinardy, B.T.I., Leighton, I., Marx, P.J., 2013. Glacier thermal regime linked to processes of annual
2557 moraine formation at Midtdalsbreen, southern Norway. *Boreas* 42(4), 896–911.
- 2558 Reuther, A.U., Urdea, P., Geiger, C., Ivy-Ochs, S., Niller, H.P., Kubik, P.W., Heine, K., 2007. Late
2559 Pleistocene glacial chronology of the Pietrele Valley, Retezat Mountains, Southern
2560 Carpathians constrained by ¹⁰Be exposure ages and pedological investigations. *Quat. Int.*
2561 164, 151–169.
- 2562 Rippin, D.M., Pomfret, A., King, N., 2015. High resolution mapping of supra-glacial drainage
2563 pathways reveals link between microchannel drainage density, surface roughness and surface
2564 reflectance. *Earth Surf. Proc. Land.* 40(10), 1279–1290.
- 2565 Robb, C., Willis, I., Arnold, N., Guðmundsson, S., 2015. A semi-automated method for mapping
2566 glacial geomorphology tested at Breiðamerkurjökull, Iceland. *Remote Sens. Environ.* 163,
2567 80–90.
- 2568 Robinson, A.H., Morrison, J.L., Muehrcke, P.C., Kimerling, A.J., Guptill, S.C., 1995. *Elements of*
2569 *cartography* (6th Edition). Wiley & Sons, Chichester, 674 pp.
- 2570 Robinson, P., Dowdeswell, J.A., 2011. Submarine landforms and the behavior of a surging ice cap
2571 since the last glacial maximum: The open-marine setting of eastern Austfonna, Svalbard. *Mar.*
2572 *Geol.* 286(1), 82–94.
- 2573 Rose, J., Smith, M.J., 2008. Glacial geomorphological maps of the Glasgow region, western central
2574 Scotland. *J. Maps* 4, 399–416.
- 2575 Rossi, C., Rodriguez Gonzalez, F., Fritz, T., Yague-Martinez, N., Eineder, M., 2012. TanDEM-X
2576 calibrated raw DEM generation. *ISPRS J. Photogramm. Remote Sens.* 73, 12–20.
- 2577 Rupke, J., De Jong, M.G.G., 1983. Slope collapse destroying ice-marginal topography in the Walgau
2578 (Vorarlberg, Austria)- an example of the application of a 1:10000 geomorphological mapping
2579 system. *Mater. Physiogeogr.* 5, 33–41.
- 2580 Ryan, J.C., Hubbard, A.L., Todd, J., Carr, J.R., Box, J.E., Christoffersen, P., Holt, T.O., Snooke, N.,
2581 2015. Repeat UAV photogrammetry to assess calving front dynamics at a large outlet glacier
2582 draining the Greenland Ice Sheet. *The Cryosphere* 9, 1–11.
- 2583 Sagredo, E.A., Moreno, P.I., Villa-Martínez, R., Kaplan, M.R., Kubik, P.W., Stern, C.R., 2011.
2584 Fluctuations of the Última Esperanza ice lobe (52 S), Chilean Patagonia, during the last
2585 glacial maximum and termination 1. *Geomorphology* 125(1), 92–108.
- 2586 Saha, K., Wells, N.A., Munro-Stasiuk, M., 2011. An object-oriented approach to automated landform
2587 mapping: A case study of drumlins. *Computers & Geosciences* 37(9), 1324–1336.
- 2588 Sahlin, E.A.U., Glasser, N.F., 2008. A geomorphological map of Cadair Idris, Wales. *J. Maps* 4, 299–
2589 314.
- 2590 Salcher, B.C., Hinsch, R., Wagreich, M., 2010. High-resolution mapping of glacial landforms in the
2591 North Alpine Foreland, Austria. *Geomorphology* 122, 283–293.
- 2592 Schoeneich, P., 1993. *Cartographie géomorphologique en Suisse*, Institut de Géographie Lausanne,
2593 Travaux et recherches, Lausanne, pp. 1–13.
- 2594 Schomacker, A., 2008. What controls dead-ice melting under different climate conditions? A
2595 discussion. *Earth-Sci. Rev.* 90, 103–113.

- Schomacker, A., Kjaer, K.H., 2008. Quantification of dead-ice melting in ice-cored moraines at the high-Arctic glacier Holmströmbreen, Svalbard. *Boreas* 37, 211–225.
- Schomacker, A., Benediktsson, Í.Ö., Ingólfsson, Ó., 2014. The Eyjabakkajökull glacial landsystem, Iceland: Geomorphic impact of multiple surges. *Geomorphology* 218, 98–107.
- Seguinot, J., Rogozhina, I., Stroeve, A.P., Margold, M., Kleman, J., 2016. Numerical simulations of the Cordilleran ice sheet through the last glacial cycle. *The Cryosphere* 10, 639–664.
- Shean, D.E., Alexandrov, O., Moratto, Z.M., Smith, B.E., Joughin, I.R., Porter, C., Morin, P., 2016. An automated, open-source pipeline for mass production of digital elevation models (DEMs) from very-high-resolution commercial stereo satellite imagery. *ISPRS J. Photogramm. Remote Sens.* 116, 101–117.
- Sissons, J.B., 1967. *The Evolution of Scotland's Scenery*. Oliver & Boyd, Edinburgh, 259 pp.
- Sissons, J.B., 1977a. The Loch Lomond Readvance in the northern mainland of Scotland. In: Gray, J.M., Lowe, J.J. (Eds.), *Studies in the Scottish Lateglacial Environment*. Pergamon Press, Oxford, pp. 45–60.
- Sissons, J.B., 1977b. The Loch Lomond Readvance in southern Skye and some palaeoclimatic implications. *Scott. J. Geol.* 13, 23–36.
- Sissons, J.B., 1979a. The limit of the Loch Lomond Advance in Glen Roy and vicinity. *Scott. J. Geol.* 15, 31–42.
- Sissons, J.B., 1979b. The Loch Lomond advance in the Cairngorm Mountains. *Scott. Geogr. Mag.* 95, 66–82.
- Smith, A.M., Murray, T., 2009. Bedform topography and basal conditions beneath a fast-flowing West Antarctic ice stream. *Quat. Sci. Rev.* 28(7), 584–596.
- Smith, A.M., Murray, T., Nicholls, K.W., Makinson, K., Adalgeirsdóttir, G., Behar, A.E., Vaughan, D. G., 2007. Rapid erosion, drumlin formation, and changing hydrology beneath an Antarctic ice stream. *Geology* 35(2), 127–130.
- Smith, G.R., Woodward, J.C., Heywood, D.I., Gibbard, P.L., 2000. Interpreting Pleistocene glacial features from SPOT HRV data using fuzzy techniques. *Computers & Geosciences* 26(4), 479–490.
- Smith, M.J., Clark, C.D., 2005. Methods and visualisation of digital elevation models for landform mapping. *Earth Surf. Process. Land.* 30, 885–900.
- Smith, M.J., Knight, J., 2011. Palaeoglaciology of the last Irish ice sheet reconstructed from striae evidence. *Quat. Sci. Rev.* 30(1–2), 147–160.
- Smith, M.J., Wise, S.M., 2007. Problems of bias in mapping linear landforms from satellite imagery. *Int. J. Appl. Earth Obs. Geoinform.* 9(1), 65–78.
- Smith, M.J., Rose, J., Booth, S., 2006. Geomorphological mapping of glacial landforms from remotely sensed data: An evaluation of the principal data sources and an assessment of their quality. *Geomorphology* 76, 148–165.
- Smith, M.J., Griffiths, J., Paron, P. (Eds.), 2011. *Geomorphological Mapping: Methods and Applications*. Developments in Earth Surface Processes, Volume 15. Elsevier, Oxford, 610 pp.
- Smith, M.J., Anders, N.S., Keesstra, S.D., 2016b. CLustre: semi- automated lineament clustering for palaeo- glacial reconstruction. *Earth Surf. Process. Land.* 41(3), 364–377.
- Smith, M.W., Carrivick, J.L., Quincey, D.J., 2016a. Structure from motion photogrammetry in physical geography. *Prog. Phys. Geogr.* 40(2), 247–275.
- Sollas, W.J., 1896. A map to show the distribution of eskers in Ireland. *The Sci. Trans. R. Dublin Soc.* 5, 785–822.
- Spagnolo, M., Clark, C.D., Hughes, A.L.C., Dunlop, P., Stokes, C.R., 2010. The planar shape of drumlins. *Sediment. Geol.* 232, 119–129.
- Spagnolo, M., Clark, C.D., Ely, J.C., Stokes, C.R., Anderson, J.B., Andreassen, K., Graham, A.G.C., King, E.C., 2014. Size, shape and spatial arrangement of megascale glacial lineations from a large and diverse dataset. *Earth Surf. Process. Land.* 39, 1432–1448.
- Spagnolo, M., Phillips, E., Piotrowski, J.A., Rea, B.R., Clark, C.D., Stokes, C.R., Carr, S.J., Ely, J.C., Ribolini, A., Wysota, W., Szuman, I., 2016. Ice stream motion facilitated by a shallow-deforming and accreting bed. *Nat. Commun.* 7, 10723.

- Spagnolo, M., Bartholomaeus, T.C., Clark, C.D., Stokes, C.R., Atkinson, N., Dowdeswell, J.A., Ely, J.C., Graham, A.G.C., Hogan, K.A., King, E.C., Larter, R.D., Livingstone, S.J., Pritchard, H.D., 2017. The periodic topography of ice stream beds: Insights from the Fourier spectra of mega-scale glacial lineations. *J. Geophys. Res.: Earth Surf.* 122, 1355–1373.
- Stansell, N.D., Polissar, P.J., Abbott, M.B., 2007. Last glacial maximum equilibrium-line altitude and paleo-temperature reconstructions for the Cordillera de Mérida, Venezuelan Andes. *Quat. Res.* 67, 115–127.
- Stokes, C.R., 2002. Identification and mapping of palaeo-ice stream geomorphology from satellite imagery: implications for ice stream functioning and ice sheet dynamics. *Int. J. Remote Sens.* 23(8), 1557–1563.
- Stokes, C.R., 2018. Geomorphology under ice streams: moving from form to process. *Earth Surf. Process. Land.*, 43, 85–123.
- Stokes, C.R., Clark, C.D., 1999. Geomorphological criteria for identifying Pleistocene ice streams. *Ann. Glaciol.* 28(1), 67–74.
- Stokes, C.R., Clark, C.D., 2001. Palaeo-ice streams. *Quat. Sci. Rev.* 20(13), 1437–1457.
- Stokes, C.R., Clark, C.D., 2002. Are long subglacial bedforms indicative of fast ice flow? *Boreas*, 31(3), 239–249.
- Stokes, C.R., Clark, C.D. 2003. The Dubawnt Lake palaeo-ice stream: evidence for dynamic ice sheet behaviour on the Canadian Shield and insights regarding the controls on ice-stream location and vigour. *Boreas* 32, 263–279.
- Stokes, C.R., Tarasov, L., 2010. Ice streaming in the Laurentide Ice Sheet: A first comparison between data-calibrated numerical model output and geological evidence. *Geophys. Res. Lett.* 37, L01501.
- Stokes, C.R., Clark, C.D., Storrar, R., 2009. Major changes in ice stream dynamics during deglaciation of the north-western margin of the Laurentide Ice Sheet. *Quat. Sci. Rev.* 28, 721–738.
- Stokes, C.R., Spagnolo, M., Clark, C.D., Ó Cofaigh, C., Lian, O.B., Dunstone, R.B., 2013. Formation of mega-scale glacial lineations on the Dubawnt Lake Ice Stream bed: 1. size, shape and spacing from a large remote sensing dataset. *Quat. Sci. Rev.* 77, 190–209.
- Stokes, C.R., Corner, G.D., Winsborrow, M.C.M., Husum, K., Andreassen, K., 2014. Asynchronous response of marine-terminating outlet glaciers during deglaciation of the Fennoscandian Ice Sheet. *Geology* 42, 455–458.
- Stokes, C.R., Tarasov, L., Blomdin, R., Cronin, T.M., Fisher, T.G., Gyllencreutz, R., Hättestrand, C., Heyman, J., Hindmarsh, R.C.A., Hughes, A.L.C., Jakobsson, M., Kirchner, N., Livingstone, S.J., Margold, M., Murton, J.B., Noormets, R., Peltier, W.R., Peteet, D.M., Piper, D.J.W., Preusser, F., Renssen, H., Roberts, D.H., Roche, D.M., Saint-Ange, F., Stroeven, A.P., Teller, J.T., 2015. On the reconstruction of palaeo-ice sheets: Recent advances and future challenges. *Quat. Sci. Rev.* 125, 15–49.
- Stokes, C.R., Margold, M., Creyts, T.T., 2016a. Ribbed bedforms on palaeo-ice stream beds resemble regular patterns of basal shear stress (‘traction ribs’) inferred from modern ice streams. *J. Glaciol.* 62, 696–713.
- Stokes, C.R., Margold, M., Clark, C.D., Tarasov, L., 2016b. Ice stream activity scaled to ice sheet volume during Laurentide Ice Sheet deglaciation. *Nature* 530(7590), 322–326.
- Storrar, R.D., Livingstone, S.J., 2017. Glacial geomorphology of the northern Kivalliq region, Nunavut, Canada, with an emphasis on meltwater drainage systems. *J. Maps* 13, 153–164.
- Storrar, R., Stokes, C.R., 2007. A Glacial geomorphological map of Victoria Island, Canadian Arctic. *J. Maps* 3, 191–210.
- Storrar, R.D., Stokes, C.R., Evans, D.J.A., 2013. A map of large Canadian eskers from Landsat satellite imagery. *J. Maps* 9, 456–473.
- Storrar, R.D., Stokes, C.R., Evans, D.J.A., 2014. Morphometry and pattern of a large sample (>20,000) of Canadian eskers and implications for subglacial drainage beneath ice sheets. *Quat. Sci. Rev.* 105, 1–25.
- Storrar, R.D., Jones, A., Evans, D.J.A., 2017. Small-scale topographically-controlled glacier flow switching in an expanding proglacial lake at Breiðamerkurjökull, SE Iceland. *J. Glaciol.* 63(240), 745–750.

2705 Streuff, K., Forwick, M., Szczuciński, W., Andreassen, K., Ó Cofaigh, C., 2015. Submarine landform
2706 assemblages and sedimentary processes related to glacier surging in Kongsfjorden, Svalbard.
2707 *arktos* 1(1), 14.

2708 Stroeven, A.P., Fabel, D., Harbor, J., Hättestrand, C., Kleman, J., 2002a. Quantifying the erosional
2709 impact of the Fennoscandian ice sheet in the Torneträsk-Narvik corridor, northern Sweden,
2710 based on cosmogenic radionuclide data. *Geogr. Ann.* 84A, 275–287.

2711 Stroeven, A.P., Fabel, D., Hättestrand, C., Harbor, J., 2002b. A relict landscape in the centre of
2712 Fennoscandian glaciation: Cosmogenic radionuclide evidence of tors preserved through
2713 multiple glacial cycles. *Geomorphology* 44, 145–154.

2714 Stroeven, A.P., Harbor, J., Fabel, D., Kleman, J., Hättestrand, C., Elmore, D., Fink, D., Fredin, O.,
2715 2006. Slow, patchy landscape evolution in northern Sweden despite repeated ice-sheet
2716 glaciation. *GSA Special Paper* 398, 387–396.

2717 Stroeven, A.P., Hättestrand, C., Heyman, J., Kleman, J., Morén, B.M., 2013. Glacial geomorphology
2718 of the Tian Shan. *J. Maps* 9(4), 505–512.

2719 Stroeven, A.P., Fabel, D., Margold, M., Clague, J.J., Xu, S., 2014. Investigating absolute chronologies
2720 of glacial advances in the NW sector of the Cordilleran Ice Sheet with terrestrial in situ
2721 cosmogenic nuclides. *Quat. Sci. Rev.* 92, 429–443.

2722 Stroeven, A.P., Hättestrand, C., Kleman, J., Heyman, J., Fabel, D., Fredin, O., Goodfellow, B.W.,
2723 Harbor, J.M., Jansen, J.D., Olsen, L., Caffee, M.W., Fink, D., Lundqvist, J., Rosqvist, G.C.,
2724 Strömberg, B., Jansson, K.N., 2016. Deglaciation of Fennoscandia. *Quat. Sci. Rev.* 147, 91–
2725 121.

2726 Sugden, D.E., 1970. Landforms of Deglaciation in the Cairngorm Mountains, Scotland. *Trans. Inst.*
2727 *Br. Geogr.* 51, 201–219.

2728 Sugden, D.E., 1978. Glacial erosion by the Laurentide Ice Sheet. *J. Glaciol.* 20, 367–391.

2729 Sugden, D.E., John, B.S., 1976. *Glaciers and Landscape*. Arnold, London, 376 pp.

2730 Thorp, P.W., 1986. A mountain icefield of Loch Lomond Stadial age, western Grampians, Scotland.
2731 *Boreas* 15, 83–97.

2732 Tonkin, T.N., Midgley, N.G., Cook, S.J., Graham, D.J., 2016. Ice-cored moraine degradation mapped
2733 and quantified using an unmanned aerial vehicle: A case study from a polythermal glacier in
2734 Svalbard. *Geomorphology* 258, 1–10.

2735 Trommelen, M.S., Ross, M., 2010. Subglacial landforms in northern Manitoba, Canada, based on
2736 remote sensing data. *J. Maps* 2010, 618–638.

2737 Trommelen, M.S., Ross, M., 2014. Distribution and type of sticky spots at the centre of a deglacial
2738 streamlined lobe in northeastern Manitoba, Canada. *Boreas* 43, 557–576.

2739 Trotter, F.M., 1929. The glaciation of the eastern Edenside, the Alston Block and the Carlisle Plain.
2740 *Quart. J. Geol. Soc. Lond.* 88, 549–607.

2741 Turner, A.J., Woodward, J., Stokes, C.R., Ó Cofaigh, C., Dunning, S., 2014a. Glacial geomorphology
2742 of the Great Glen Region of Scotland. *J. Maps* 10, 159–178.

2743 Turner, D., Lucieer, A., Wallace, L., 2014b. Direct georeferencing of ultrahigh-resolution UAV
2744 imagery. *IEEE Trans. Geosci. Remote Sens.* 52(5), 2738–2745.

2745 van der Bilt, W.G.M., Bakke, J., Balascio, N.L., 2016. Mapping sediment–landform assemblages to
2746 constrain lacustrine sedimentation in a glacier-fed lake catchment in northwest Spitsbergen. *J.*
2747 *Maps* 12(5), 985–993.

2748 Van Dorsser, J.J., Salomé, A.I., 1973. Different methods of detailed geomorphological mapping.
2749 *K.N.A.G. Geografisch Tijdschrift* VII: 71–74.

2750 Walker, D.A., Auerbach, N.A., Shippert, M.M., 1995. NDVI, biomass, and landscape evolution of
2751 glaciated terrain in northern Alaska. *Polar Record*, 31(177), 169–178.

2752 Welch, R., 1966. A comparison of aerial films in the study of the Breiðamerkur glacier area, Iceland,
2753 *The Photogramm. Rec.* 5, 289–306.

2754 Welch, R., 1967. *The Application of Aerial Photography to the Study of a Glacial Area.*
2755 Breiðamerkur, Iceland, Unpublished PhD thesis, University of Glasgow.

2756 Welch, R., 1968. Color aerial photography applied to the study of a glacial area. In: Smith, J.T. (Ed.),
2757 *Manual of Color Aerial Photography*. American Society of Photogrammetry and Remote
2758 Sensing, Falls Church, VA, pp. 400–401.

- Westoby, M.J., Brasington, J., Glasser, N.F., Hambrey, M.J., Reynolds, J.M., 2012. 'Structure-from-Motion' photogrammetry: A low-cost, effective tool for geoscience applications. *Geomorphology* 179, 300–314.
- Westoby, M.J., Glasser, N.F., Hambrey, M.J., Brasington, J., Reynolds, J.M., Hassan, M.A., 2014. Reconstructing historic Glacial Lake Outburst Floods through numerical modelling and geomorphological assessment: Extreme events in the Himalaya. *Earth Surf. Process. Land.* 39(12), 1675–1692.
- Westoby, M.J., Dunning, S.A., Woodward, J., Hein, A.S., Marrero, S.M., Winter, K., Sugden, D.E., 2016. Interannual surface evolution of an Antarctic blue-ice moraine using multi-temporal DEMs. *Earth Surf. Dyn.* 4, 515–529.
- Wilson, S.B., 2005. Morphological analysis and mapping of Loch Lomond Stadial moraines using digital photogrammetry and geographical information systems. Unpublished PhD thesis, University of Glasgow, 360 pp.
- Winsborrow, M.C.M., Andreassen, K., Corner, G.D., Laberg, J.S., 2010. Deglaciation of a marine-based ice sheet: Late Weichselian palaeo-ice dynamics and retreat in the southern Barents Sea reconstructed from onshore and offshore glacial geomorphology. *Quat. Sci. Rev.* 29(3–4), 424–442.
- Winsborrow, M.C.M., Stokes, C.R., Andreassen, K., 2012. Ice stream flow switching during deglaciation of the southwestern Barents Sea. *Geol. Soc. Am. Bull.* 124(3–4), 275–290.
- Wolf, P.R., DeWitt, B.A., Wilkinson, B.E., 2013. *Elements of Photogrammetry with Application in GIS*, Fourth Edition. McGraw-Hill Education.
- Wright, W.B., 1912. The drumlin topography of south Donegal. *Geol. Mag.* 9, 153–159.
- Wyshnytzky, C.E., 2017. On the mechanisms of minor moraine formation in high-mountain environments of the European Alps. Unpublished PhD thesis, Queen Mary University of London, 329 pp.
- Yu, P., Eyles, N., Sookhan, S., 2015. Automated drumlin shape and volume estimation using high resolution LiDAR imagery (Curvature Based Relief Separation): A test from the Wadena Drumlin Field, Minnesota. *Geomorphology* 246, 589–601.
- Zasadni, J., Kłapyta, P., 2016. From valley to marginal glaciation in alpine-type relief: Lateglacial glacier advances in the Pięć Stawów Polskich/Roztoka Valley, High Tatra Mountains, Poland. *Geomorphology* 253, 406–424.
- Zech, R., Abramowski, U., Glaser, B., Sosin, P., Kubik, P.W. and Zech, W., 2005. Late Quaternary glacial and climate history of the Pamir Mountains derived from cosmogenic ¹⁰Be exposure ages. *Quat. Res.* 64(2), 212–220.
- Zemp, M., Frey, H., Gärtner-Roer, I., Nussbaumer, S.U., Hoelzle, M., Paul, F., Haeberli, W., Denzinger, F., Ahlstrøm, A.P., Anderson, B., Bajracharya, S., Baroni, C., Braun, L.N., Cáceres, B.E., Casassa, G., Cobos, G., Dávila, L.R., Delgado Granados, H., Demuth, M.N., Espizua, L., Fischer, A., Fujita, K., Gadek, B., Ghazanfar, A., Hagen, J.O., Holmlund, P., Karimi, N., Li, Z., Pelto, M., Pitte, P., Popovnin, V.V., Portocarrero, C.A., Prinz, R., Sangewar, C.V., Severskiy, I., Sigurðsson, O., Soruco, A., Usubaliev, R., Vincent, C., 2015. Historically unprecedented global glacier decline in the early 21st century. *J. Glaciology* 61, 745–762.

Figure captions

Figure 1. Digitised versions of two geomorphological maps drawn in the field for (A) Coire Easgainn and (B) Glen Odhar in the Monadhliath, Central Scottish Highlands. These field maps were used in the production of a 1:57,500 geomorphological map for the entire region (Boston, 2012a, b).

Figure 2. The aerial photograph overlay-mapping process using an example from the mountain Arkle, NW Scotland. (A) aerial photograph at an average scale of ~1:25,000 (extract from photo 38 88 087; ©RCAHMS 1988); (B) scan of original overlay mapped through a stereoscope from (A) (see Section 2.2.2 for method description), focusing on moraines, fluted moraines and the approximate upper limit of scree slopes as seen from the aerial photograph; (C) compiled, rectified geomorphological map, incorporating moraines and fluted moraines from (B) and additional data from field mapping, such as the exact upper limits of scree slopes, orientation of striae, solifluction lobes and mountaintop detritus, for example. For description and interpretation of the geomorphology, see Lukas (2006).

Figure 3. Satellite image (A) and geomorphological mapping (B) showing suites of moraines formed by the Lago General Carrera–Buenos Aires ice lobe of the former Patagonian Ice Sheet, located to the east of the present-day Northern Patagonian Icefield. A combination of remotely-sensed datasets and field mapping were used to circumvent issues of localised cloud cover, as visible in A. Where areas were obscured, SPOT-5 and DigitalGlobe images available in *Google Earth* were used. Note the extensive outwash development between moraine suites. The geomorphological map extract is taken from Bendle et al. (2017a).

Figure 4. Comparison of WorldView-2 satellite imagery (June 2012, European Space Imaging) with digital colour aerial photographs (2006, Loftmyndir ehf) for the Skálafellsjökull foreland, SE Iceland. A. Panchromatic satellite image (0.5 m ground sampled distance, GSD). B. Multispectral satellite image (2.0 m GSD). C. Pansharpened three-band natural colour satellite image (0.5 m GSD). D. Digital colour aerial photographs (0.41 m GSD). The satellite imagery is of sufficient resolution to allow mapping of small-scale (<2 m in height) annual moraines (see Chandler et al., 2016a, b).

Figure 5. Geomorphology of the Finsterwalderbreen foreland, Svalbard, mapped in the field and from a digital aerial photograph captured in 2004. Photograph provided by the NERC Earth Observation Data Centre. Modified from Lovell et al. (2018).

Figure 6. Views at various points along the length of the 1890 surge end moraine at Eyjabakkajökull, Iceland, visualised in *ESRI ArcScene* (Benediktsson et al., 2010). Aerial orthophotographs from 2008 are draped over a 3 m grid DEM with 1.5x vertical exaggeration.

Figure 7. High-resolution geomorphological mapping of part of the Fláajökull foreland, Iceland, based on UAV-derived imagery (Evans et al., 2016a). A 1:350 scale version of this map is freely available for download from *Journal of Maps*: <http://dx.doi.org/10.1080/17445647.2015.1073185>.

Figure 8. Geomorphological mapping of lineations in northwest Saskatchewan, Canada, from SRTM imagery (modified from Norris et al., 2017). (A) and (B) Densely spaced drumlins displaying low length to width ratios. (C) and (D) Highly elongated fluting orientated NE–SW east of the Grizzly Bear Hills. Geomorphological map extracts in (C) and (D) show lineations (black lines), eskers (red lines) and meltwater channels (dashed blue lines).

Figure 9. Examples of landforms in relief-shaded DEMs. Red indicates higher elevations and blue lower elevations. (A) Lineations in N Canada shown in 16 m resolution CDED data. (B) De Geer moraines in SW Finland shown in 2 m resolution LiDAR data. (C) Lineations of the Dubawnt Lake Ice Stream shown in 5 m resolution ArcticDEM mosaic data. (D) Esker-fed ice-contact outwash fan in SW Finland shown in 2 m resolution LiDAR data. See Table 2 for DEM data sources.

Figure 10. Conceptual diagrams illustrating the distinction between ground sampled distance (B and E) and pixel size (C and F). The ground distances between two measurements by the detector (i.e. the ground sampled distances) are 30 m and 50 m in (B) and (E), respectively. These ground sample distances are then assigned to pixels in the resulting 30 x 30 m (C) and 50 x 50 m (F) digital images. Note, resultant images may fail to represent accurately the shape of the objects (upper row) or even may fail to reproduce them (lower row), even where the size of the object is the same or larger than the sampling distance.

Figure 11. Geometric artefacts that may be present in space- and air-borne radar captured imagery, resulting from the effects of relief. (A) **Foreshortening**, occurring where the slope of the local terrain is less than the incidence angle (γ). The facing slope, $a - b$, becomes compressed to $a_I - b_I$ in the resulting image. (B) **Layover**, occurring in steep terrain when the slope angle is greater than the

incidence angle. As a mountain-top, b , is closer to the sensor than the base, a , this causes layover in the imagery (an incorrect positioning of b_l relative to a_l). (C) **Radar shadow** in areas of rugged terrain as the illumination is from an oblique source. No data is recorded for the region $b_l - d_l$. (D) In regions of varying topography, a **combination of artefacts** may be present: points b and c will be impacted by layover and will be positioned incorrectly relative to a ; no data will be recorded for the region between c and d due to radar shadow; foreshortening occurs at slope facet $d - e$; further radar shadow occurs at $e - f$; and foreshortening at f and g . After Clark (1997).

Figure 12. Extracts from hillshaded relief models of Ben More Coigach, NW Scottish Highlands, showing the effect of geometric artefacts on the models. The hillshades were generated with azimuths of 45° (A) and 315° (B). Stretching of upland terrain during processing of the DEM data results in blurred regions on the hillshaded relief models. NEXTMap DSM from Intermap Technologies Inc. provided by NERC via the NERC Earth Observation Data Centre.

Figure 13. Example mapping of subglacial bedforms from the Strait of Magellan, Patagonia (A–C), and the Dubawnt Lake Ice Stream (D–F). The bedforms are mapped as polylines along landform crests in B and E, and mapped as polygons delineating lower-break-of-slope in C and F. The Dubawnt Lake Ice Stream polylines (Stokes and Clark, 2003) and polygons (Dunstone, 2014) were mapped by different mappers at different times, which may account for small inconsistencies. For further details on the bedform examples from the Strait of Magellan, see Lovell et al. (2011) and Darvill et al. (2014).

Figure 14. Geomorphological mapping of Coire Easgairn, Monadhliath, Scotland, using a combination of NEXTMap DSMs, analogue aerial photographs and field mapping. Modified from Boston (2012a, b).

Figure 15. Examples of landforms in icefield and valley glacier settings mapped on medium to coarse resolution imagery. Landforms observed in the Chagan Uzun Valley, Russian Altai, displayed on (A) SPOT image and (B) Landsat 7 ETM+ image. (C) Associated geomorphological map extract from Gribenski et al. (2016). Moraines in the Anadyr Lowlands, Far NE Russia, displayed on (D) semi-transparent shaded ViewFinder Panorama (VFP) DEM data (NE solar azimuth) draped over the raw VFP DEM. (E) Associated mapping of moraines (black polygons) from Barr and Clark (2012).

Figure 16. Geomorphological mapping (A) from the Múlajökull foreland, Iceland, completed as part of a process-oriented study examining the internal architecture and structural evolution of a Little Ice Age terminal moraine at this surge-type glacier (Benediktsson et al., 2015). The mapping was

combined with sedimentological investigations (B) to produce a process-form model of moraine formation and evolution (C).

Figure 17. Geomorphological mapping of the foreland of Skálafellsjökull, an active temperate outlet of Vatnajökull, SE Iceland. (A) Digital aerial photographs (2006; 0.41 m GSD; *Loftmyndir ehf*), pan-sharpened WorldView-2 multi-spectral satellite imagery (2012; 0.5 m GSD; *European Space Imaging*), a UAV-derived DEM (2013; 0.09 m GSD) and field mapping were employed to produce the mapping extract (B). A compromise on the level of detail was made, with annual moraines mapped along crestlines due to image resolution and map readability. This mapping detail was sufficient for calculating crest-to-crest moraine spacing (ice-margin retreat rates) shown in (C), which was the principal purpose of the study. Modified from Chandler et al. (2016a, b).

Figure 18. Idealised workflow for mapping palaeo-ice sheet geomorphology. Some pathways in the workflow are optional (grey dashed lines) depending on data availability and the feasibility and applicability of particular methods. Note, where analogue (hard-copy) aerial photographs are used for mapping, processing of acetate overlays would be undertaken *after* mapping from the aerial photographs. Further details on image processing are shown on the processing workflow available as supplementary material.

Figure 19. Idealised workflow for mapping alpine- and plateau-style ice mass geomorphology. In this scenario, digital remotely-sensed datasets are used and this necessitates image processing *before* mapping is undertaken. Ideally, GNSS surveys would be conducted in order to process digital aerial photographs, as depicted in the workflow. Some pathways are optional (grey dashed lines) depending on data availability and the feasibility and applicability of particular methods. Although sedimentology is shown as ‘optional’, it is highly desirable to undertake sedimentological investigations, wherever possible. Alternative image processing solutions are available and readers should consult with the detailed processing workflow which is available as supplementary material.

Figure 20. Idealised workflow for mapping alpine- and plateau-style ice mass geomorphology. In this scenario, analogue (hard-copy) aerial photographs are used and this necessitates image processing *after* mapping is undertaken. Some pathways are optional (grey dashed lines) depending on data availability and the feasibility and applicability of particular methods. Although sedimentology is shown as ‘optional’, it is highly desirable to undertake sedimentological investigations, wherever possible. Alternative image processing solutions are available and readers should consult with the detailed processing workflow which is available as supplementary material.

2956 **Table 1.** Primary satellite imagery types used in glacial geomorphological mapping and example applications. Broadly ordered in terms of spatial resolution.

Satellite	Sensor	Temporal coverage	Spectral bands	Spatial resolution (m)	Source	Example studies
Landsat 1–5	MSS	1972–2013	4	80	USGS Earth Explorer (earthexplorer.usgs.gov)	Clark and Stokes (2001); Stokes and Clark (2002, 2003); Jansson et al. (2003); see also Clark (1997, Table 1)
Landsat 4–5	TM	1982–2013	1 6	120 30	Global Land Cover Facility (landcover.org)	Punkari (1995); Alexanderson et al. (2002); De Angelis (2007); Storrar et al. (2013); Orkhonselenge (2016)
Landsat 7	ETM+	1999–	1 6 1	60 30 15		Kassab et al (2013); Stroeve et al. (2013); Darvill et al. (2014); Blomdin et al. (2016b); Ely et al. (2016b); Ercolano et al. (2016); Lindholm and Heyman (2016); Storrar and Livingstone (2017); see also Clark (1997, Table 1)
Landsat 8	OLI/TIRS	2013–	2 8 1	100 30 15		Espinoza (2016); Carrivick et al. (2017); Storrar and Livingstone (2017)
Terra	ASTER	2000–	5 6 5	90 20 15	LP DAAC (LPDAAC.usgs.gov)	Glasser and Jansson (2005, 2008); Glasser et al. (2005); Lovell et al. (2011); Sagredo et al. (2011); Darvill et al (2014); Ercolano et al. (2016)
ERS 1	SAR	1991–2000	1	30	European Space Agency (earth.esa.int)	Clark et al. (2000); Clark and Stokes (2001); Heiser and Roush (2001); see also Clark (1997, Table 1)
SPOT 1–3	HRV	1986–2009	3 1 1	20 20 10	Airbus Defence and Space (intelligence-airbusds.com)	Smith et al. (2000); Coronato et al. (2009)
SPOT 4	HRVIR	1998–2013	1 3 1	10 20 20		Trommelen and Ross (2010, 2014); Ercolano et al. (2016) [viewed in Google Earth™]; McHenry and Dunlop (2016); Principato et al. (2016)
SPOT 5	HRG/HRS	2002–2015	1 3 1	2.5, 5 10 20		Trommelen and Ross (2010, 2014); Ercolano et al. (2016) [viewed in Google Earth™]; McHenry and Dunlop (2016); Principato et al. (2016); Bendle et al. (2017a)
SPOT 6–7	NAOMI	2012–	1 4	1.5 6		Gribenski et al. (2016)
CORONA/ARGON/LANYARD	KH1–KH6	1959–1972	1	1.8–140	USGS Earth Explorer (earthexplorer.usgs.gov)	Alexanderson et al (2002); Zech et al. (2005); Lifton et al (2014)
IKONOS	HRG	1999–2015	1 4	1 4	DigitalGlobe (digitalglobe.com)	Juyal et al. (2011); Kłapyta (2013); Zasadni and Kłapyta (2016)

COSMO-Skymed	SAR	2008–	1/3/15/16/20	1	e-GEOS (e-geos.it)	da Rosa et al. (2013a)
Quickbird	HRG	2001–	1	0.61	DigitalGlobe	da Rosa et al. (2011, 2013b); May et al (2011);
		2014	4	2.44	(digitalglobe.com)	Lovell et al. (2011)
GeoEye-1		2008–	1	0.46		Westoby et al. (2014)
			4	1.84		
WorldView-2		2009–	1	0.46	European Space Imaging (euspaceimaging.com)	Jamieson et al. (2015); Chandler et al. (2016a);
			8	1.84		Evans et al. (2016e); Ewertowski et al (2016)
Google Earth™ (specific image details not given)	n/a	n/a	n/a	n/a	Google Earth	Margold and Jansson (2011); Margold et al. (2011); Kassab et al (2013); Stroeve et al. (2013); Darvill et al (2014); Blomdin et al. (2016b); Evans et al. (2016d); Li et al (2016); Lindholm and Heyman (2016); Orkhonselenge (2016)

Table 2. Examples of DEM datasets with national- to international-coverage that have been employed in glacial geomorphological map production.

Dataset	Coverage	Spatial resolution (m)	RMSE or CE90 (m)		Data source(s)	Example studies
			Vertical	Horizontal		
SRTM ¹	Global	~90 (3 arc-second) ~30 (1 arc-second)	~5–13	-	Global Land Cover Facility (landcover.org) USGS Earth Resources and Science Center (eros.usgs.gov)	Glasser and Jansson (2008); Barr and Clark (2009); Ó Cofaigh et al. (2010); Morén et al. (2011); Stroeve et al. (2013); Darvill et al. (2014); Evans et al. (2014, 2016d); Trommelen and Ross (2014); Stokes et al. (2016a); Ely et al. (2016b); Lindholm and Heyman (2016)
ASTER GDEM (V2)	Global	~30 (1 arc-second)	~8.7	-	LP DAAC Global Data Explorer (gdex.cr.usgs.gov/gdex) NASA Reverb (reverb.echo.nasa.gov/reverb)	Barr and Clark (2012); Blomdin et al. (2016a, b); Lindholm and Heyman (2016)
Canadian Digital Elevation Dataset (CDED)	Canada	~20 (0.75 arc-second)	-	-	Natural Resources Canada (geogratis.gc.ca)	Margold et al. (2011, 2015a); Evans et al. (2016c); Storrar and Livingstone (2017)
USGS National Elevation Dataset (NED) ²	US	~30 (1 arc-second) ~10 (1/3 arc-second)	~2.4	-	US Geological Survey (ned.usgs.gov)	Hess and Briner (2009); Margold et al. (2015a); Ely et al. (2016a)
TanDEM-X	Global	~12 (0.4 arc-second)	<10	<10	German Aerospace Center (DLR) (tandemx-science.dlr.de)	Pipaud et al. (2015)
NEXMap Britain TM	UK	5	~1	2.5	NERC Earth Observation Data Centre ³ (ceda.ac.uk)	Livingstone et al. (2008); Finlayson et al. (2010, 2011); Hughes et al. (2010); Brown et al. (2011a); Boston (2012a, b); Pearce et al. (2014); Turner et al. (2014a)
ArcticDEM	Arctic	2	2.0	3.8	Polar Geospatial Center (pgc.umn.edu/data/arcticdem)	Levy et al. (2017)
Maanmittauslaitos LiDAR DEM	Finland	2	~0.3	-	National Land Survey of Finland (maanmittauslaitos.fi)	Ojala et al. (2015); Ojala (2016); Mäkinen et al. (2017)
Ny Nationell Höjdmodell	Sweden	2	~0.1	-	Lantmäteriet (lantmateriet.se)	Dowling et al. (2015, 2016); Greenwood et al. (2015); Möller and Dowling (2016); Peterson et al. (2017)
Environment Agency LiDAR DEM	UK (partial)	2, 1, 0.5 and 0.25	0.05 – 0.15	0.4	DEFRA Environment Data (environment.data.gov.uk)	Miller et al. (2014)
Iceland Met Office and Institute of Earth Sciences, University of Iceland, LiDAR DEM ⁴	Iceland (partial)	<5	<0.5	-	Iceland Meteorological Office (en.vedur.is)	Brynjólfsson et al. (2014, 2016); Benediktsson et al. (2016); Jónsson et al. (2016)

¹ SRTM data was only freely available with a spatial resolution of ~90 m (3 arc-seconds) outside of the United States until late 2015 when the highest resolution data were thereafter made available globally (see <http://www2.jpl.nasa.gov/srtm/>)

² The USGS NED dataset has been superseded by the 3D Elevation Program (3DEP), with this data available as seamless 1/3 arc-second, 1 arc-second and 2 arc-second DEMs (see https://nationalmap.gov/3DEP/3dep_prodserv.html)

³ NEXTMap BritainTM data is freely available to NERC staff and NERC-funded researchers, though subsets can be applied for by non-NERC-funded researchers under a Demonstrator User License Agreement (DULA)

⁴ The Icelandic LiDAR DEM data are available at 5 m resolution, but it is possible to derive higher-resolution DEMs (e.g. 2 m) from the point clouds using denser interpolation.

Table 3. Summary of the glacial settings where the main geomorphological mapping methods and remotely-sensed data types are *most* appropriate. ✓ = the method/dataset is appropriate and should be used (where the dataset is available). ● = the method is applicable in certain cases, depending on factors such as the resolution of the *specific* dataset, the size of the study area and landforms, and the accessibility of the study area.

Glacial setting	DEMs	Coarse satellite imagery	LiDAR DEMs	High-resolution satellite imagery	Aerial photographs	UAV imagery	Field mapping
Ice sheets	✓	✓	✓				
Ice sheet sectors/lobes	✓	✓	✓	●	●		●
Ice-caps	●	●	●	✓	✓		✓
Icefields			●	✓	✓		✓
Valley (outlet) glaciers			●	✓	✓	●	✓
Cirque glaciers			●	✓	✓	●	✓
Modern glacier forelands			●	✓	✓	✓	✓