

**Please cite the Published Version**

Collins, NJ, Barton, CJ, Van Middelkoop, M, Callaghan, MJ , Rathleff, MS, Vicenzino, BT, Davis, IS, Powers, CM, Macri, EM, Hart, HF, De Oliveira Silva, D and Crossley, KM (2018) 2018 Consensus statement on exercise therapy and physical interventions (orthoses, taping and manual therapy) to treat patellofemoral pain: Recommendations from the 5th International Patellofemoral Pain Research Retreat, Gold Coast, Australia, 2017. British Journal of Sports Medicine, 52 (18). pp. 1170-1178. ISSN 0306-3674

**DOI:** <https://doi.org/10.1136/bjsports-2018-099397>

**Publisher:** BMJ

**Version:** Accepted Version

**Downloaded from:** <https://e-space.mmu.ac.uk/621353/>

**Additional Information:** This is an Author Accepted Manuscript of a paper accepted for publication in British Journal of Sports Medicine, published by and copyright BMJ.

**Enquiries:**

If you have questions about this document, contact [openresearch@mmu.ac.uk](mailto:openresearch@mmu.ac.uk). Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

# 2018 consensus statement on exercise therapy and physical interventions (orthoses, taping, manual therapy) to treat patellofemoral pain. Recommendations from the 5<sup>th</sup> International Patellofemoral Pain Research Retreat, Gold Coast, Australia, 2017.

Natalie J. Collins<sup>1,2</sup>, Christian J. Barton<sup>2,3</sup>, Marienke van Middelkoop<sup>4</sup>, Michael J. Callaghan<sup>5</sup>, Michael S. Rathleff<sup>6</sup>, Bill Vicenzino<sup>1</sup>, Irene S. Davis<sup>7</sup>, Christopher M. Powers<sup>8</sup>, Erin M. Macri<sup>9,10</sup>, Harvi F. Hart<sup>2,11</sup>, Danilo de Oliveira Silva<sup>2,12</sup>, Kay M. Crossley<sup>2</sup>

<sup>1</sup> School of Health and Rehabilitation Sciences, The University of Queensland, Brisbane, Queensland, Australia

<sup>2</sup> La Trobe Sport and Exercise Medicine Research Centre, College of Science, Health and Engineering, La Trobe University, Melbourne, Victoria, Australia

<sup>3</sup> Department of Surgery, University of Melbourne, St. Vincent's Hospital, Melbourne, Victoria, Australia

<sup>4</sup> Department of General Practice, Erasmus MC University Medical Center, Rotterdam, The Netherlands

<sup>5</sup> Department of Health Professions, Manchester Metropolitan University, Manchester, UK

<sup>6</sup> Research Unit for General Practice in Aalborg, Department of Clinical Medicine, Aalborg University, Aalborg, Denmark

<sup>7</sup> Department of Physical Medicine and Rehabilitation, Spaulding National Running Centre, Harvard Medical School, Cambridge, Massachusetts, USA

<sup>8</sup> Division of Biokinesiology & Physical Therapy, University of Southern California, Los Angeles, California, USA

<sup>9</sup> Department of Physical Therapy, University of Delaware, Newark, Delaware, USA

<sup>10</sup> Centre for Hip Health and Mobility, University of British Columbia, Vancouver, British Columbia, Canada

<sup>11</sup> School of Physical Therapy and Bone and Joint Institute, The University of Western Ontario, London, Ontario, Canada

<sup>12</sup> Department of Physiotherapy, Faculty of Science and Technology, Sao Paulo State University (UNESP), Presidente Prudente, Sao Paulo, Brazil

Corresponding author:

Dr Natalie Collins

School of Health and Rehabilitation Sciences, The University of Queensland, Brisbane, Queensland, 4072 Australia

Email: [n.collins1@uq.edu.au](mailto:n.collins1@uq.edu.au)

Phone: +61 7 3365 2124

## ABSTRACT

Patellofemoral pain affects a large proportion of the population, from adolescents to older adults, and carries a substantial personal and societal burden. An international group of scientists and clinicians meets biennially at the International Patellofemoral Research Retreat to share research findings related to patellofemoral pain conditions, and develop consensus statements using best practice methods. This consensus statement, from the 5<sup>th</sup> International Patellofemoral Research Retreat held in Australia in July 2017, focuses on exercise therapy and physical interventions (e.g. orthoses, taping, manual therapy) for patellofemoral pain. Literature searches were conducted to identify new systematic reviews and randomised clinical trials (RCTs) published since the 2016 Consensus Statement. The methodological quality of included systematic reviews and RCTs was graded using AMSTAR and PEDro, respectively. Evidence-based statements were developed from included papers, and presented to a panel of 41 patellofemoral pain experts for consensus discussion and voting. Recommendations from the expert panel support the use of exercise therapy (especially the combination of hip- and knee-focused exercises), combined interventions, and foot orthoses to improve pain and/or function in people with patellofemoral pain. The use of patellofemoral, knee or lumbar mobilisations in isolation, or electrophysical agents, is not recommended. There is uncertainty regarding the use of patellar taping/bracing, acupuncture/dry needling, manual soft tissue techniques, blood flow restriction training, and gait retraining in patients with patellofemoral pain. In 2017, we launched the International Patellofemoral Research Network ([www.ipfrn.org](http://www.ipfrn.org)), to consolidate and grow our patellofemoral research community, facilitate collaboration, and disseminate patellofemoral pain knowledge to clinicians and the general public. The 6<sup>th</sup> International Patellofemoral Research Retreat will be held in Milwaukee, Wisconsin, United States, in October 2019.

## INTRODUCTION

Patellofemoral pain is a common, chronic musculoskeletal condition, presenting as pain around or behind the patella during patellofemoral joint loading activities (e.g. squatting, stair ambulation, running).<sup>1</sup> Patellofemoral pain has an annual prevalence of approximately 23% of adults and 29% of adolescents in the general population, and affects almost 36% of professional cyclists.<sup>2</sup> Patellofemoral pain tends to persist in about 50% of people<sup>3-5</sup>, in some cases for up to 20 years.<sup>6</sup> Pain and symptoms associated with patellofemoral pain limit participation in daily and occupational tasks, and reduce levels of physical activity.<sup>5 7</sup> Importantly, patellofemoral pain may precede the onset of patellofemoral osteoarthritis,<sup>8 9</sup> which has no cure.

### International Patellofemoral Research Retreat

Every two years, scientists and clinicians who are active researchers in the field of patellofemoral pain meet at the International Patellofemoral Research Retreat. The aims of the Retreat are to: (i) share the latest patellofemoral pain research developments; (ii) discuss the literature to formulate consensus statements to disseminate knowledge; and (iii) develop a future research agenda for patellofemoral pain. The 6<sup>th</sup> International Patellofemoral Research Retreat will be held in Milwaukee, Wisconsin, United States, in October 2019.

In 2017, the International Patellofemoral Research Network (iPFRN) was launched, bringing together past and present Retreat participants into a cohesive, collaborative network of patellofemoral pain researchers from around the world. We also launched the iPFRN website ([www.ipfrn.org](http://www.ipfrn.org)) -- a platform to share news on upcoming Retreats, recruit new members to the research community, facilitate collaboration, and disseminate patellofemoral pain knowledge to clinicians and the general public.

The 5<sup>th</sup> International Patellofemoral Research Retreat (iPFRR17) was held on the Gold Coast, Australia, from July 18-20, 2017. The 51 attendees from 10 countries represented the disciplines of physiotherapy, athletic training, podiatry, biomechanics, imaging, biomedical engineering, and human movement science.

There were two keynote speakers:

- Professor Richard Souza (University of California, San Francisco) shared his expertise on advanced imaging methods for patellofemoral pain and patellofemoral osteoarthritis.
- Associate Professor Thor Besier (The University of Auckland) spoke about how musculoskeletal modelling can be used to improve assessment and treatment of people with patellofemoral pain.

55 peer-reviewed abstracts were accepted for podium presentation. These were grouped into seven categories across nine sessions: (i) imaging; (ii) epidemiology and pain; (iii) muscle and strength; (iv) biomechanics and joint health; (v) biomechanics and patellofemoral joint stress; (vi) treatment; and (vii) psychology and pain.

## Developing this consensus statement

On the final day of the Retreat, we held four structured consensus discussions on key topics. We synthesized current knowledge on patellofemoral pain from published literature, new studies presented at iPFRR17, and expert opinion to develop consensus statements on each topic. Agreement was reached by discussion at the meeting at the Gold Coast, and by email discussion of the draft statement.

Two topics that were discussed in detail were:

1. An evidence-based framework for a pathomechanical model of patellofemoral pain, which was developed after discussions at the 2015 meeting. This paper has been published in the *British Journal of Sports Medicine*.<sup>10</sup>
2. Exercise therapy and physical interventions for patellofemoral pain. This discussion updated and extended the 2016 Patellofemoral Pain Consensus Statement from the 4th International Patellofemoral Pain Research Retreat, Manchester (Part 2).<sup>11</sup>

## METHODS

### Literature review

We repeated the search strategy we had used for the 2016 Consensus Statement<sup>11</sup> to search MEDLINE, EMBASE, CINAHL and Current Contents for systematic reviews and randomised controlled trials (RCTs) (CJB). Searches included literature from June 2015 to July 2017, covering the period since the 2015 meeting. Titles and abstracts were searched using the following patellofemoral pain terms: (patella OR patellofemoral) AND (pain OR syndrome OR dysfunction).<sup>12</sup> To identify systematic reviews, this search was narrowed by combining with the keyword 'review'. To identify RCTs, patellofemoral pain terms were combined with (controlled trial OR clinical trial). Additional searches were performed in the Physiotherapy Evidence Database (PEDro). That list was provided to all authors of this consensus statement, and they were asked to suggest papers they considered were missing. This did not lead to any additional papers being included.

Papers were eligible for inclusion if they: (i) were the most current systematic reviews (including meta-analyses) or RCTs that had been published since the most recent systematic review; (ii) evaluated one or more physical interventions (non-surgical, non-pharmacological), with either no treatment or another physical intervention as the comparator; (iii) evaluated outcome using a measure of pain (e.g. pain visual analogue scale [VAS]), patient-reported improvement (e.g. global rating of change) or function; (iv) randomly allocated participants to intervention groups; and (v) were full-text papers published in English. Two reviewers (CJB, DOS) determined the eligibility status for each paper identified by the search strategy.

### Quality of evidence

The methodological quality of each included paper was rated by three independent reviewers (NJC, CJB, MVM). Systematic reviews were graded using the Assessment of Multiple Systematic

Reviews (AMSTAR) tool<sup>13</sup> (CJB, MVM) (total score /11). PEDro was used to grade the methodological quality of RCTs (NJC, MVM) (total score /10). Discrepancies between raters were discussed at a consensus meeting, with an additional reviewer (KMC) resolving any disagreement.

To maintain consistency with the 2016 Consensus Statement,<sup>11</sup> systematic reviews and RCTs were allocated the following level of evidence, based on AMSTAR and PEDro scores, respectively: high quality  $\geq 7$ ; moderate quality 4–6; and low quality  $\leq 3$ .

### **Data extraction and analysis**

We extracted data from each included paper. Data from low quality systematic reviews was not considered in this evidence update. If no systematic reviews were identified for an intervention, we included all published RCTs related to that intervention. Extracted data were used by members of the consensus group (NJC, MVM, MJC, MSR, KMC) to generate summary tables for the consensus meeting, including study methodology, outcomes and effect sizes. Summary tables also included information on methodological quality. Unless defined otherwise by systematic reviews, we classified duration of treatment effect as follows: (i) short term: <6 months; (ii) medium term: 6-12 months; and (iii) long term: >12 months.<sup>11</sup>

Consistent with the 2016 Consensus Statement,<sup>11</sup> we classified included studies under the following headings: (i) exercise therapy (general; hip-focused; knee-focused); (ii) combined interventions (two or more of the following: exercise therapy, foot orthoses, patellar taping, manual therapy); (iii) foot orthoses; (iv) patellar taping and bracing; and (v) other adjunctive interventions (e.g. manual therapy, soft tissue therapy, acupuncture, gait retraining, electrophysical agents). If a study investigated the effect of adding an adjunct modality to a combined interventions program, compared to combined interventions alone, this was classified as other adjunctive interventions. For an intervention to be considered as a new category, a systematic review on the specific intervention must have been identified by the search strategy.

### **Expert panel voting**

Of the 51 iPFRR17 attendees, 41 comprised the patellofemoral pain expert panel. All panel members were active researchers in the field of patellofemoral pain, and covered disciplines of physiotherapy, athletic training, podiatry, biomechanics, biomedical engineering, and human movement science. A proportion of panel members were also practitioners with experience managing patients with patellofemoral pain. Summary tables developed by the consensus group were provided to the expert panel during the consensus meeting. New evidence was presented alongside each statement from the 2015 Retreat. New statements were formulated where new interventions were identified, based on the evidence, for the outcomes of interest (pain, function, overall improvement).

During the consensus meeting, the consensus group led discussions for each intervention regarding details of included papers, as well as clarification and finalisation of each statement. Following the discussion, the expert panel was instructed to integrate the evidence for each intervention (presented in the summary tables), with their expert opinion and knowledge, to

vote on the appropriateness of new and revised statements. A 10-point scale was used to rate each statement, where 0 was 'not appropriate' and 9 was 'appropriate'.<sup>11 14</sup> Pooled scores were calculated to represent the median appropriateness score for each intervention statement, with interquartile ranges. We considered median scores of 0-3 to represent 'inappropriate', 4-6 as 'uncertain' and 7-9 as 'appropriate'.<sup>11</sup> The consensus group made final recommendations based on statements that had consistency between voting and associated evidence.

## RESULTS

The search strategy yielded 655 papers (311 systematic reviews; 344 RCTs). After title, abstract and full-text review, six systematic reviews and an additional 12 RCTs that were not in the systematic reviews met the inclusion criteria. No additional papers were identified by the consensus panel. There were no new systematic reviews on specific interventions outside of the categories from the 2016 Consensus Statement;<sup>11</sup> thus, no new intervention categories were added.

Quality ratings for included studies are presented in Table 1 (systematic reviews) and Table 2 (RCTs). Figure 1 summarises the updated consensus process. We identified two low quality systematic reviews<sup>15 16</sup> and three moderate quality RCTs<sup>17-19</sup> on exercise therapy; one low quality systematic review<sup>20</sup> and one moderate quality RCT<sup>21</sup> on combined interventions; one high quality<sup>22</sup> and two low quality<sup>23 24</sup> systematic reviews and two RCTs (one high quality<sup>25</sup>, one low quality<sup>26</sup>) on patellar taping and bracing; and two high quality<sup>27 28</sup>, three moderate quality<sup>18 29 30</sup>, and two low quality<sup>31 32</sup> RCTs on other adjunctive interventions. No new systematic reviews or RCTs were identified that related to foot orthoses.

### Evidence update

Ten of the 24 statements voted on by the expert panel from the 2015 Retreat had additional evidence incorporated (Table 3). For eight of these statements, the recommendation did not change despite integration of new evidence. These statements remained unchanged, and therefore were not voted on by the expert panel at the 2017 Retreat.

All voting that took place was for statements regarding other adjunctive interventions. Outcomes are presented in Figure 2. We identified new evidence for two statements that were presented at the 2015 Retreat, relating to acupuncture/dry needling and manual soft tissue techniques. These were voted on again in 2017, based on the new evidence (Table 3).

We also identified new RCT evidence for interventions that were not discussed at the 2015 Retreat. Three new statements relating to other adjunctive interventions were presented to the expert panel. Voting was conducted for recommendations on blood flow restriction training and gait retraining (Table 3). The third statement referred to platelet rich plasma; the expert panel felt that these injections should be excluded from this Consensus Statement as it is more aligned with medical/pharmacological interventions than physical interventions.

### Recommendations from the expert panel

Although the expert panel voted on four statements related to other adjunctive interventions (including two new statements), there was uncertainty associated with the level of evidence for these interventions (acupuncture/dry needling, manual soft tissue techniques, blood flow restriction training, gait retraining). Thus, the integration of new evidence published since the 2015 Retreat did not alter the six key recommendations presented in the 2016 Consensus Statement.<sup>11</sup>

Recommendations from the expert panel on exercise therapy and physical interventions for patellofemoral pain are:

1. Exercise therapy is recommended to reduce pain in the short, medium and long term, and improve function in the medium and long term.
2. Combining hip and knee exercises is recommended to reduce pain and improve function in the short, medium and long term, and this combination should be used in preference to knee exercises alone.
3. Combined interventions are recommended to reduce pain in adults with patellofemoral pain in the short and medium term. Combined interventions as a management program incorporates exercise therapy as well as one of the following: foot orthoses, patellar taping, or manual therapy.
4. Foot orthoses are recommended to reduce pain in the short term.
5. Patellofemoral, knee and lumbar mobilisations are not recommended in isolation.
6. Electrophysical agents are not recommended.

## DISCUSSION

At the 5<sup>th</sup> International Patellofemoral Research Retreat, held in July 2017, we reviewed the consensus-based recommendations for exercise therapy and physical interventions for patellofemoral pain. Despite the integration of new evidence from systematic reviews and RCTs, and voting on two new and two revised statements, the six key recommendations from the expert panel are unchanged from the 4<sup>th</sup> International Patellofemoral Research Retreat Consensus Statement.<sup>11</sup>

These recommendations will provide medical and health practitioners with evidence-based guidelines for managing patients with patellofemoral pain. Considering the widely-held view that patellofemoral pain is a multifactorial condition, as highlighted in our previous consensus statements,<sup>33-35</sup> these recommendations are not intended to be prescriptive for all patients with patellofemoral pain. Rather, in line with evidence-based practice and shared decision-making models, they should be integrated with assessment findings, patient preferences and goals, and clinical expertise.

### Evidence update since 2016 Consensus Statement

In the two years since the 2015 Retreat, we identified six systematic reviews and 13 RCTs that had been published, highlighting the continued output of level 1 and 2 evidence for exercise therapy and physical interventions for patellofemoral pain. Findings of these papers were integrated into existing consensus statements. This new evidence was sufficient to update recommendations for acupuncture and dry needling, and manual soft tissue techniques, which



were subsequently voted on by the expert panel. We also formulated new consensus statements on blood flow restriction training and gait retraining, on which the expert panel also voted. All statements that were voted on were in the category of 'other adjunctive interventions'. The expert panel considered that the recommendation for each of these interventions was uncertain, based on the best available evidence and their clinical expertise. We recommend that practitioners use high-value interventions in the first instance, based on the panel's six key recommendations, and use these adjunctive interventions where indicated for individual patients until high quality evidence supports or refutes their use.

Acupuncture was identified as an area of uncertainty in the 2016 Consensus Statement,<sup>11</sup> based on findings of one moderate quality systematic review.<sup>36</sup> A new high quality RCT evaluating dry needling to trigger points<sup>27</sup> was identified in the 2017 update. While the expert panel acknowledged that acupuncture and trigger point dry needling have key differences in their clinical indication and application, it was decided to group them in a single recommendation on the basis that evidence for both was limited and uncertain. If additional RCTs are published in the future, it is anticipated that acupuncture and dry needling will be considered separately.

There is also uncertainty around the use of manual soft tissue techniques for patellofemoral pain. New studies revealed no significant effect on pain when myofascial techniques were added to hip-focused exercises (vs. exercises alone),<sup>32</sup> as well as small clinically non-meaningful effects on pain and function with ischaemic compression compared to lumbopelvic manipulation.<sup>29</sup> This was considered alongside evidence from the 2015 Retreat, which identified no improvements in pain following ischaemic release to peri- and retro-patellar regions, compared to ischaemic release at the hip.<sup>37</sup> The uncertainty surrounding manual soft tissue techniques reflects expert opinion reported in the mixed methods paper by Barton et al,<sup>12</sup> where massage was considered to add limited value by one individual, while another stated it played a "big role" in patellofemoral pain management.

Blood flow restriction training was added to the consensus discussion after identification of one high quality RCT.<sup>28</sup> The patient performs lower load strengthening exercise with a tourniquet placed around the proximal thigh (e.g. pneumatic cuff, elastic wrap).<sup>28</sup> In musculoskeletal pain conditions such as anterior cruciate ligament reconstruction and knee osteoarthritis, this can be more tolerable than heavy load training, and result in greater strength gains when compared to low load training used in isolation.<sup>38</sup> The RCT evaluated the effects of blood flow restriction in conjunction with low load knee focused exercise therapy, compared to standard high load quadriceps strengthening program.<sup>28</sup> Although there was a significant reduction in pain during activities of daily living after eight weeks, between-group differences were less than the minimal clinically important difference for pain using a visual analogue scale (20mm).<sup>39</sup> There were no effects on worst knee pain or function at eight weeks, and no effects on any outcome measure at six months. On this basis, the expert panel considered there to be uncertainty regarding blood flow restriction training for patellofemoral pain. Considering that this is a new approach to exercise therapy in patellofemoral pain, further high-quality RCTs are needed before evidence-based recommendations for its use or not in patellofemoral pain can be made.

Gait retraining was also incorporated into consensus discussions at the 2017 Retreat, on the basis of two new moderate quality RCTs that met the eligibility criteria.<sup>18 30</sup> Given the current popularity of gait retraining for patellofemoral pain, this generated a large amount of discussion, including whether it should be considered a separate intervention category. Because there was no systematic review evaluating RCT evidence for gait retraining in patellofemoral pain, it was considered to be an adjunctive intervention. One systematic review was excluded from consideration on the basis that it did not identify any published RCTs at the time of their search (April 2015).<sup>40</sup> The expert panel voted that there was uncertainty regarding the short-term effects of gait retraining on pain and function, based on evidence available at the time of the consensus meeting. Esculier et al<sup>18</sup> reported no effects on pain and function over five months when gait retraining (personalised program focused on increasing step rate, landing softer, and adopting a non-rearfoot strike pattern) combined with load management education, was compared to exercise and load management education, or load management education alone, in runners with patellofemoral pain (n=69). In comparison, a small RCT of 16 runners who were self-reported rearfoot strikers found that two weeks of gait retraining (focused on transitioning from rearfoot to forefoot strike pattern) significantly reduced patellofemoral pain immediately post-retraining and at one month follow-up, compared to no intervention.<sup>30</sup> Differences between the two studies extend beyond the selection of runners (e.g. rearfoot strikers<sup>30</sup> vs. any footstrike pattern<sup>18</sup>), choice of running retraining techniques, choice of comparator, and sample size, to approaches used to attain gait retraining effects (e.g. motor retraining principles).<sup>41 42</sup> It is recommended that robust RCTs of gait retraining of runners (and walkers) with patellofemoral pain be conducted. These should focus on the specific gait variables that are associated with PFP,<sup>40 43</sup> include larger samples, and evaluate participants over a longer follow-up period in order to establish the role of gait retraining in patellofemoral pain.

### **Exercise therapy remains the intervention of choice for patellofemoral pain**

Exercise therapy is the intervention of choice for patellofemoral pain, with the largest body of evidence supporting its use to improve pain and function in the short-, medium- and long-term. Our literature search identified two new systematic reviews<sup>15 16</sup> and three new RCTs<sup>17-19</sup> on exercise therapy that were integrated into recommendations from the 2016 Consensus Statement.<sup>11</sup> These new findings strengthened the evidence that underpins our previous recommendations on exercise therapy for patellofemoral pain. Thus, it is recommended that exercise targeting the hip and knee is a key component of management for all patients with patellofemoral pain, particularly when used in combination. Our findings suggest that further studies evaluating the effectiveness of similar exercise therapy programs may not be warranted. Research efforts may be better directed towards determining the optimal exercise dose for patellofemoral pain, especially in light of poor reporting of exercise prescription parameters.<sup>44</sup> This will further facilitate the implementation of RCT findings into clinical practice, and optimise the potential benefits of exercise therapy for patients with patellofemoral pain.

### **Combined interventions and foot orthoses are recommended to reduce pain in the short- to medium-term**

Based on Level 1 evidence and expert opinion, the expert panel recommends the use of combined interventions to reduce pain in the short- and medium-term, and prefabricated foot orthoses to reduce pain in the short-term. We defined combined interventions as a management program incorporating exercise in conjunction with at least one of the following: foot orthoses, patellar taping, or manual therapy. The majority of combined intervention programs studied in RCTs involve exercise as one of the therapeutic components. A combined intervention approach best represents typical clinical management of patellofemoral pain, allowing practitioners to select treatment components that address specific impairments identified in individual patients. It should be noted that there is no evidence supporting combined interventions beyond 12 months for adults with patellofemoral pain. The expert panel voted 'uncertain' regarding the long-term effects of combined interventions on pain in adolescents with patellofemoral pain. Taken together, these conclusions highlight the need to evaluate the effects of combined interventions in both adults and adolescents beyond 12 months, especially considering that more than half of those with patellofemoral pain will continue to experience symptoms after 2-8 years.<sup>4 5</sup>

Prefabricated foot orthoses remain a recommendation for short-term relief of patellofemoral pain. There is no evidence supporting the use of custom fabricated foot orthoses for patellofemoral pain, which are made from a three-dimensional representation of the patient's foot.<sup>45</sup> A consideration when using foot orthoses in the management of patellofemoral pain is the variability in treatment response, demonstrated by wide confidence intervals.<sup>46 47</sup> This may be optimized through selecting patients based on specific characteristics, such as foot posture and mobility characteristics or immediate improvements in functional performance with foot orthoses.<sup>48-50</sup> Further insight into the predictive ability of foot measures on foot orthoses outcomes will be gained from findings of an upcoming RCT.<sup>51</sup>

### **Joint mobilisation and electrophysical agents are not recommended for patellofemoral pain**

The literature search did not identify any new evidence to change recommendations for interventions that are not effective for patellofemoral pain. The expert panel recommends against the use of patellofemoral, knee and lumbar mobilisation, as well as electrophysical agents, as primary interventions when managing patients with patellofemoral pain. It should be noted that patellofemoral mobilisation can be used as a component of a combined intervention approach where appropriate, with evidence and expert opinion in support.<sup>12</sup> Considering the shift towards active, movement-based physical interventions for patellofemoral pain and musculoskeletal pain more broadly, it is advised that these passive interventions should not be the focus of future RCTs.

### **Key areas of uncertainty**

There remain key areas of uncertainty regarding exercise therapy and physical interventions for patellofemoral pain. We found no new evidence to address uncertainty surrounding selection of hip-focused over knee-focused exercise therapy, combined interventions in adolescents with patellofemoral pain, patellar taping and bracing, and the use of other adjunctive interventions such as acupuncture and manual soft tissue therapy.<sup>11</sup> The 2017 consensus meeting also highlighted additional areas of uncertainty, being the use of blood flow

restriction training and gait retraining. It is important to re-iterate that these interventions may still have a place in managing patellofemoral pain, but that further research is required.

## Limitations

Despite only using level 1 and 2 evidence in the consensus process, there were some limitations that should be considered when interpreting recommendations. Firstly, the expert panel consisted of attendees at the 2017 International Patellofemoral Research Retreat, held in Australia. Retreat attendees were required to be active researchers in patellofemoral pain, ensuring that the expert panel were familiar with published literature in the field, and were broadly representative of all current researchers. It is likely that not all active researchers were in attendance, which should be taken in consideration when reading the recommendations. Secondly, the consensus panel was limited to using published literature when updating and composing statements, to ensure adequate availability of study protocols and outcomes and maintain the rigour of the consensus process. Considering that the purpose of the Retreat is to share new knowledge regarding patellofemoral pain and its management, it is plausible that findings of studies presented at the Retreat may have influenced or biased statements presented to the expert panel. We look forward to integrating these findings into future updates on exercise therapy and physical interventions for patellofemoral pain. Finally, it should be highlighted that studies involving medical and surgical interventions were not included in this consensus statement. While it is the opinion of the consensus group that exercise therapy and physical interventions should always be first-line management for patellofemoral pain, there is a need to make evidence-based recommendations for the use of interventions such as pharmacological agents and injectables, to educate practitioners and the public on their use in patellofemoral pain. Ongoing systematic reviews in the field will help to facilitate this.<sup>52 53</sup>

## Future directions

Based on outcomes of the consensus meeting, our consensus group highlights key areas of focus for future studies on exercise therapy and physical interventions for patellofemoral pain. While we are encouraged by the continued output of level 1 and 2 evidence for patellofemoral pain, studies addressing the condition across various stages of the lifespan are required (e.g. adolescents, adults over 40 years of age). Further high-quality RCTs are required for a number of interventions. These should focus primarily on active interventions that have the capacity to induce long-term improvements in pain and function.

Although exercise therapy is the intervention of choice for patellofemoral pain, further work is needed to enhance adherence to exercise programs and understand key components of effective exercise prescription (e.g. load, contraction type, time under tension, rest). This is especially important given findings of a dose-response relationship between exercise frequency and recovery in adolescents with patellofemoral pain.<sup>54</sup> Studies that explore barriers and facilitators to exercise in patients with patellofemoral pain, as well as exercise programs developed in consultation with patients, are encouraged. We also encourage researchers to investigate the role of education in managing patellofemoral pain, as well as interventions aimed at addressing psychosocial impairments that have been identified in people with patellofemoral pain.<sup>55 56</sup>

There is a clear need for minimum reporting standards in patellofemoral pain publications, to allow replication of interventions in clinical practice.<sup>44</sup> These are currently being developed by members of the iPFNRN, and will feature in a future publication. We are also developing consensus around factors that influence treatment response. The next update to the consensus statement on *exercise therapy and physical interventions* will be in four years, at the 7<sup>th</sup> International Patellofemoral Research Retreat in 2021.

## CONCLUSION

Recommendations from the consensus meeting on exercise therapy and physical interventions held at the 5<sup>th</sup> International Patellofemoral Research Retreat (Gold Coast, Australia, 2017), support the use of exercise therapy (especially the combination of hip- and knee-focused exercises), combined interventions, and foot orthoses to improve pain and/or function in people with patellofemoral pain. The use of patellofemoral, knee or lumbar mobilisations in isolation, or electrophysical agents, is not recommended. There is uncertainty regarding the use of patellar taping/bracing, acupuncture/dry needling, manual soft tissue techniques, blood flow restriction training, and gait retraining in patients with patellofemoral pain. The 6<sup>th</sup> International Patellofemoral Research Retreat will be held in Milwaukee, Wisconsin, United States, in October 2019.

## ACKNOWLEDGEMENTS

The authors wish to thank all members of the expert panel who participated in the consensus meeting at the 2017 International Patellofemoral Research Retreat.

## FUNDING

The *British Journal of Sports Medicine* provided an unrestricted grant to support the 5<sup>th</sup> International Patellofemoral Research Retreat.

## REFERENCES

1. Crossley KM, Stefanik JJ, Selfe J, et al. 2016 Patellofemoral pain consensus statement from the 4th International Patellofemoral Pain Research Retreat, Manchester. Part 1: Terminology, definitions, clinical examination, natural history, patellofemoral osteoarthritis and patient-reported outcome measures. *Br J Sports Med* 2016;50(14):839-43. doi: 10.1136/bjsports-2016-096384
2. Smith BE, Selfe J, Thacker D, et al. Incidence and prevalence of patellofemoral pain: A systematic review and meta-analysis. *PLoS One* 2018;13(1):e0190892. doi: 10.1371/journal.pone.0190892 [published Online First: 2018/01/13]
3. Collins NJ, Bierma-Zeinstra SM, Crossley KM, et al. Prognostic factors for patellofemoral pain: a multicentre observational analysis. *Br J Sports Med* 2013;47(4):227-33. doi: 10.1136/bjsports-2012-091696 [published Online First: 2012/12/18]
4. Lankhorst NE, van Middelkoop M, Crossley KM, et al. Factors that predict a poor outcome 5-8 years after the diagnosis of patellofemoral pain: a multicentre observational analysis. *Br J Sports Med* 2016;50(14):881-6. doi: 10.1136/bjsports-2015-094664

5. Rathleff MS, Rathleff CR, Olesen JL, et al. Is Knee Pain During Adolescence a Self-limiting Condition? Prognosis of Patellofemoral Pain and Other Types of Knee Pain. *Am J Sports Med* 2016;44(5):1165-71. doi: 10.1177/0363546515622456
6. Nimon G, Murray D, Sandow M, et al. Natural history of anterior knee pain: a 14- to 20-year follow-up of nonoperative management. *J Pediatr Orthop* 1998;18(1):118-22.
7. Hall R, Barber Foss K, Hewett TE, et al. Sport specialization's association with an increased risk of developing anterior knee pain in adolescent female athletes. *J Sport Rehabil* 2015;24(1):31-5. doi: 10.1123/jsr.2013-0101 [published Online First: 2014/03/14]
8. Thomas MJ, Wood L, Selfe J, et al. Anterior knee pain in younger adults as a precursor to subsequent patellofemoral osteoarthritis: a systematic review. *BMC Musculoskelet Disord* 2010;11:201. doi: 1471-2474-11-201 [pii]10.1186/1471-2474-11-201 [published Online First: 2010/09/11]
9. Crossley KM. Is patellofemoral osteoarthritis a common sequela of patellofemoral pain? *Br J Sports Med* 2014;48(6):409-10. doi: 10.1136/bjsports-2014-093445
10. Powers CM, Witvrouw E, Davis IS, et al. Evidence-based framework for a pathomechanical model of patellofemoral pain: 2017 patellofemoral pain consensus statement from the 4th International Patellofemoral Pain Research Retreat, Manchester, UK: part 3. *Br J Sports Med* 2017;51(24):1713-23. doi: 10.1136/bjsports-2017-098717 [published Online First: 2017/11/08]
11. Crossley KM, van Middelkoop M, Callaghan MJ, et al. 2016 Patellofemoral pain consensus statement from the 4th International Patellofemoral Pain Research Retreat, Manchester. Part 2: recommended physical interventions (exercise, taping, bracing, foot orthoses and combined interventions). *Br J Sports Med* 2016;50(14):844-52. doi: 10.1136/bjsports-2016-096268
12. Barton CJ, Lack S, Hemmings S, et al. The 'Best Practice Guide to Conservative Management of Patellofemoral Pain': incorporating level 1 evidence with expert clinical reasoning. *Br J Sports Med* 2015;49(14):923-34. doi: 10.1136/bjsports-2014-093637
13. Shea BJ, Grimshaw JM, Wells GA, et al. Development of AMSTAR: a measurement tool to assess the methodological quality of systematic reviews. *BMC Med Res Methodol* 2007;7:10. doi: 10.1186/1471-2288-7-10 [published Online First: 2007/02/17]
14. McAlindon TE, Bannuru RR, Sullivan MC, et al. OARSI guidelines for the non-surgical management of knee osteoarthritis. *Osteoarthritis Cartilage* 2014;22(3):363-88. doi: 10.1016/j.joca.2014.01.003 [published Online First: 2014/01/28]
15. Alba-Martin P, Gallego-Izquierdo T, Plaza-Manzano G, et al. Effectiveness of therapeutic physical exercise in the treatment of patellofemoral pain syndrome: a systematic review. *J Phys Ther Sci* 2015;27(7):2387-90. doi: 10.1589/jpts.27.2387 [published Online First: 2015/08/28]
16. Santos TR, Oliveira BA, Ocarino JM, et al. Effectiveness of hip muscle strengthening in patellofemoral pain syndrome patients: a systematic review. *Braz J Phys Ther* 2015;19(3):167-76. doi: 10.1590/bjpt-rbf.2014.0089 [published Online First: 2015/06/04]
17. Bolgia LA, Earl-Boehm J, Emery C, et al. Pain, Function, and Strength Outcomes for Males and Females with Patellofemoral Pain Who Participate in Either a Hip/Core- or Knee-Based Rehabilitation Program. *Int J Sports Phys Ther* 2016;11(6):926-35. [published Online First: 2016/12/03]

18. Esculier JF, Bouyer LJ, Dubois B, et al. Is combining gait retraining or an exercise programme with education better than education alone in treating runners with patellofemoral pain? A randomised clinical trial. *Br J Sports Med* 2017 doi: 10.1136/bjsports-2016-096988
19. Sahin M, Ayhan FF, Borman P, et al. The effect of hip and knee exercises on pain, function, and strength in patients with patellofemoral pain syndrome: a randomized controlled trial. *Turk J Med Sci* 2016;46(2):265-77. doi: 10.3906/sag-1409-66 [published Online First: 2016/08/12]
20. Espi-Lopez GV, Arnal-Gomez A, Balasch-Bernat M, et al. Effectiveness of Manual Therapy Combined With Physical Therapy in Treatment of Patellofemoral Pain Syndrome: Systematic Review. *J Chiropr Med* 2017;16(2):139-46. doi: 10.1016/j.jcm.2016.10.003 [published Online First: 2017/06/01]
21. Banan Khojaste SM, Oliyaie G, Mir MM, et al. The effectiveness of exercise therapy based on Sahrman approach in patients with patella-femoral pain syndrome. *J Babol Univ Med Sci* 2016;18(8):7-13.
22. Smith TO, Drew BT, Meek TH, et al. Knee orthoses for treating patellofemoral pain syndrome. *Cochrane Database Syst Rev* 2015(12):CD010513. doi: 10.1002/14651858.CD010513.pub2 [published Online First: 2015/12/10]
23. Chang WD, Chen FC, Lee CL, et al. Effects of Kinesio Taping versus McConnell Taping for Patellofemoral Pain Syndrome: A Systematic Review and Meta-Analysis. *Evid Based Complement Alternat Med* 2015;2015:471208. doi: 10.1155/2015/471208 [published Online First: 2015/07/18]
24. Logan CA, Bhashyam AR, Tisosky AJ, et al. Systematic Review of the Effect of Taping Techniques on Patellofemoral Pain Syndrome. *Sports Health* 2017;9(5):456-61. doi: 10.1177/1941738117710938 [published Online First: 2017/06/16]
25. Kurt EE, Buyukturan O, Erdem HR, et al. Short-term effects of kinesio tape on joint position sense, isokinetic measurements, and clinical parameters in patellofemoral pain syndrome. *J Phys Ther Sci* 2016;28(7):2034-40. doi: 10.1589/jpts.28.2034 [published Online First: 2016/08/12]
26. Petersen W, Ellermann A, Rembitzki IV, et al. Evaluating the potential synergistic benefit of a realignment brace on patients receiving exercise therapy for patellofemoral pain syndrome: a randomized clinical trial. *Arch Orthop Trauma Surg* 2016;136(7):975-82. doi: 10.1007/s00402-016-2464-2 [published Online First: 2016/05/06]
27. Espi-Lopez GV, Serra-Ano P, Vicent-Ferrando J, et al. Effectiveness of Inclusion of Dry Needling in a Multimodal Therapy Program for Patellofemoral Pain: A Randomized Parallel-Group Trial. *J Orthop Sports Phys Ther* 2017;47(6):392-401. doi: 10.2519/jospt.2017.7389 [published Online First: 2017/05/16]
28. Giles L, Webster KE, McClelland J, et al. Quadriceps strengthening with and without blood flow restriction in the treatment of patellofemoral pain: a double-blind randomised trial. *Br J Sports Med* 2017;51(23):1688-94. doi: 10.1136/bjsports-2016-096329 [published Online First: 2017/05/14]
29. Behrangrad S, Kamali F. Comparison of ischemic compression and lumbopelvic manipulation as trigger point therapy for patellofemoral pain syndrome in young adults: A double-blind randomized clinical trial. *J Bodyw Mov Ther* 2017;21(3):554-64. doi: 10.1016/j.jbmt.2016.08.007 [published Online First: 2017/07/29]

30. Roper JL, Harding EM, Doerfler D, et al. The effects of gait retraining in runners with patellofemoral pain: A randomized trial. *Clin Biomech (Bristol, Avon)* 2016;35:14-22. doi: 10.1016/j.clinbiomech.2016.03.010 [published Online First: 2016/04/26]
31. Orscelik A, Yildiz Y. Comparison of single and triple platelet rich plasma injections in the treatment of patellofemoral pain syndrome. *Turkiye Klinikleri J Med Sci* 2015;35(2):78-87.
32. Telles G, Cristovao DR, Belache FA, et al. The effect of adding myofascial techniques to an exercise programme for patients with anterior knee pain. *J Bodyw Mov Ther* 2016;20(4):844-50. doi: 10.1016/j.jbmt.2016.02.007 [published Online First: 2016/11/07]
33. Davis IS, Powers CM. Patellofemoral pain syndrome: proximal, distal, and local factors, an international retreat, April 30-May 2, 2009, Fells Point, Baltimore, MD. *J Orthop Sports Phys Ther* 2010;40(3):A1-16. doi: 2413 [pii]10.2519/jospt.2010.0302 [published Online First: 2010/03/03]
34. Powers CM, Bolgia LA, Callaghan MJ, et al. Patellofemoral pain: proximal, distal, and local factors, 2nd International Research Retreat. *J Orthop Sports Phys Ther* 2012;42(6):A1-54. doi: 10.2519/jospt.2012.0301
35. Witvrouw E, Callaghan MJ, Stefanik JJ, et al. Patellofemoral pain: consensus statement from the 3rd International Patellofemoral Pain Research Retreat held in Vancouver, September 2013. *Br J Sports Med* 2014;48(6):411-4. doi: 10.1136/bjsports-2014-093450
36. Collins NJ, Bisset LM, Crossley KM, et al. Efficacy of nonsurgical interventions for anterior knee pain: systematic review and meta-analysis of randomized trials. *Sports Med* 2012;42(1):31-49. doi: 10.2165/11594460-000000000-00000 [published Online First: 2011/12/14]
37. Hains G, Hains F. Patellofemoral pain syndrome managed by ischemic compression to the trigger points located in the peri-patellar and retro-patellar areas: a randomized clinical trial. *Clin Chiropr* 2010;13:201-9.
38. Hughes L, Paton B, Rosenblatt B, et al. Blood flow restriction training in clinical musculoskeletal rehabilitation: a systematic review and meta-analysis. *Br J Sports Med* 2017;51(13):1003-11. doi: 10.1136/bjsports-2016-097071 [published Online First: 2017/03/06]
39. Crossley KM, Bennell KL, Cowan SM, et al. Analysis of outcome measures for persons with patellofemoral pain: which are reliable and valid? *Arch Phys Med Rehabil* 2004;85(5):815-22.
40. Neal BS, Barton CJ, Gallie R, et al. Runners with patellofemoral pain have altered biomechanics which targeted interventions can modify: A systematic review and meta-analysis. *Gait Posture* 2016;45:69-82. doi: 10.1016/j.gaitpost.2015.11.018
41. Davis I. Optimising the efficacy of gait retraining. *Br J Sports Med* 2017 doi: 10.1136/bjsports-2017-098297 [published Online First: 2017/09/21]
42. Esculier JF, Dubois B, Roy JS. Response to: 'Optimising the efficacy of gait retraining'. *Br J Sports Med* 2018 doi: 10.1136/bjsports-2017-098609 [published Online First: 2018/03/09]
43. Barton CJ, Bonanno DR, Carr J, et al. Running retraining to treat lower limb injuries: a mixed-methods study of current evidence synthesised with expert opinion. *Br J Sports Med* 2016;50(9):513-26. doi: 10.1136/bjsports-2015-095278 [published Online First: 2016/02/18]



44. Holden S, Rathleff MS, Jensen MB, et al. How can we implement exercise therapy for patellofemoral pain if we don't know what was prescribed? A systematic review. *Br J Sports Med* 2018;52(6):385. doi: 10.1136/bjsports-2017-097547 [published Online First: 2017/11/01]
45. Collins N, Bisset L, McPoil T, et al. Foot orthoses in lower limb overuse conditions: a systematic review and meta-analysis. *Foot Ankle Int* 2007;28(3):396-412.
46. Collins N, Crossley K, Beller E, et al. Foot orthoses and physiotherapy in the treatment of patellofemoral pain syndrome: randomised clinical trial. *BMJ* 2008;337:a1735. doi: 10.1136/bmj.a1735 [published Online First: 2008/10/28]
47. Vicenzino B, Collins N, Cleland J, et al. A clinical prediction rule for identifying patients with patellofemoral pain who are likely to benefit from foot orthoses: a preliminary determination. *Br J Sports Med* 2010;44(12):862-6. doi: bjsm.2008.052613 [pii]10.1136/bjsm.2008.052613 [published Online First: 2008/09/30]
48. Lack S, Barton C, Vicenzino B, et al. Outcome predictors for conservative patellofemoral pain management: a systematic review and meta-analysis. *Sports Med* 2014;44(12):1703-16. doi: 10.1007/s40279-014-0231-5
49. Selfe J, Janssen J, Callaghan M, et al. Are there three main subgroups within the patellofemoral pain population? A detailed characterisation study of 127 patients to help develop targeted intervention (TIPPs). *Br J Sports Med* 2016;50(14):873-80. doi: 10.1136/bjsports-2015-094792 [published Online First: 2016/02/03]
50. Matthews M, Rathleff MS, Claus A, et al. Can we predict the outcome for people with patellofemoral pain? A systematic review on prognostic factors and treatment effect modifiers. *Br J Sports Med* 2017;51(23):1650-60. doi: 10.1136/bjsports-2016-096545 [published Online First: 2016/12/15]
51. Matthews M, Rathleff MS, Claus A, et al. The Foot Orthoses versus Hip eXercises (FOHX) trial for patellofemoral pain: a protocol for a randomized clinical trial to determine if foot mobility is associated with better outcomes from foot orthoses. *J Foot Ankle Res* 2017;10:5. doi: 10.1186/s13047-017-0186-5 [published Online First: 2017/02/01]
52. Macri E, Hart H, Crossley K, et al. PROSPERO International Prospective Register of Systematic Reviews, [Available from: [https://www.crd.york.ac.uk/prospero/display\\_record.php?RecordID=82527](https://www.crd.york.ac.uk/prospero/display_record.php?RecordID=82527) accessed 31 March 2018].
53. Thwaites D, Hart H, Devitt B, et al. PROSPERO International Prospective Register of Systematic Reviews, [Available from: [https://www.crd.york.ac.uk/prospero/display\\_record.php?RecordID=78575](https://www.crd.york.ac.uk/prospero/display_record.php?RecordID=78575) accessed 31 March 2018].
54. Rathleff MS, Roos EM, Olesen JL, et al. Exercise during school hours when added to patient education improves outcome for 2 years in adolescent patellofemoral pain: a cluster randomised trial. *Br J Sports Med* 2015;49(6):406-12. doi: 10.1136/bjsports-2014-093929
55. Maclachlan LR, Collins NJ, Matthews MLG, et al. The psychological features of patellofemoral pain: a systematic review. *Br J Sports Med* 2017;51(9):732-42. doi: 10.1136/bjsports-2016-096705
56. Maclachlan LR, Matthews M, Collins NJ, et al. The psychological features of patellofemoral pain: a cross-sectional study. *Scand J Pain* 2018;accepted manuscript



## FIGURE CAPTIONS

**Figure 1.** Summary of 2017 update of the 2016 Consensus Statement.

**Figure 2.** Consensus voting for new evidence for new and updated recommendations for other adjunctive interventions. Left whisker, quartile 1; orange box, quartile 2; black line, median; blue box, quartile 3.

**Table 1.** Quality ratings of included systematic reviews, evaluated using AMSTAR.

	Score per AMSTAR item:											Total score	Quality	Search date
Systematic review:	i	ii	iii	iv	v	vi	vii	viii	ix	x	xi			
<b>Exercise:</b>														
Alba-Martin <i>et al</i> <sup>15</sup>	0	0	0	0	0	0	1	0	0	0	0	1	low	not reported
Santos <i>et al</i> <sup>16</sup>	0	CA	0	0	0	0	1	0	0	0	0	1	low	September 2014
<b>Combined interventions:</b>														
Espi-Lopez <i>et al</i> <sup>20</sup>	0	CA	0	0	0	0	1	0	0	0	0	1	low	December 2014
<b>Taping and bracing:</b>														
Logan <i>et al</i> <sup>24</sup>	0	0	0	0	0	0	0	0	0	0	0	0	low	April 2015
Smith <i>et al</i> <sup>22</sup>	1	1	1	1	1	1	1	1	1	0	0	9	high	June 2015
Chang <i>et al</i> <sup>23</sup>	0	0	0	0	0	0	1	0	1	1	0	3	low	January 2014

AMSTAR, Assessment of Multiple Systematic Reviews. i, *a priori* design; ii, duplicate study selection and data extraction; iii comprehensive literature search; iv, search for grey literature; v, list of studies included and excluded provided; vi, characteristics of included studies provided; vii, scientific quality assessed; viii, scientific quality used to formulate conclusions; ix, methods to combine study findings appropriate; x, publication bias assessed; xi, conflict of interest. Scoring: 0, no; 1, yes; CA, cannot assess. High quality, total score  $\geq 7$ ; low quality, total score  $\leq 3$ .

**Table 2.** PEDro quality ratings for included RCTs.

	Score per PEDro item											Total score	Quality
Study	i*	ii	iii	iv	v	vi	vii	viii	ix	x	xi		
<b>Exercise:</b>													
Bolglá <i>et al</i> <sup>17</sup>	1	1	0	1	0	0	0	0	1	0	1	4	moderate
Sahin <i>et al</i> <sup>19</sup>	0	1	0	1	0	0	0	1	0	1	1	5	moderate
<b>Combined interventions:</b>													
Banan Khojaste <i>et al</i> <sup>21</sup>	0	1	0	1	0	0	0	0	0	1	1	4	moderate
<b>Taping and bracing:</b>													
Kurt <i>et al</i> <sup>25</sup>	0	1	0	1	1	0	1	1	0	1	1	7	high
Petersen <i>et al</i> <sup>26</sup>	1	1	0	0	0	0	0	1	0	0	1	3	low
<b>Adjunctive interventions:</b>													
Behrangrad <i>et al</i> <sup>29</sup>	1	1	1	0	0	0	0	1	1	1	1	6	moderate
Esculier <i>et al</i> <sup>18</sup>	1	1	1	0	0	0	0	1	1	1	1	6	moderate
Giles <i>et al</i> <sup>28</sup>	1	1	1	1	1	0	1	1	1	1	1	9	high
Espi-Lopez <i>et al</i> <sup>27</sup>	1	1	1	1	0	0	0	1	1	1	1	7	high
Orselik <i>et al</i> <sup>31</sup>	0	0	0	0	0	0	0	0	0	1	1	2	low
Roper <i>et al</i> <sup>30</sup>	0	1	0	0	0	0	0	1	1	1	1	5	moderate
Telles <i>et al</i> <sup>32</sup>	1	1	0	0	0	0	0	0	0	1	1	3	low

PEDro, Physiotherapist Evidence Database. RCTs, randomised controlled trials.

i, eligibility criteria; ii, random allocation; iii, concealed allocation; iv, baseline comparability; v, blind subjects; vi, blind therapists; vii, blind assessors; viii, adequate follow-up; ix, intention-to-treat analysis; x, between-group comparisons; xi, point estimates and variability. Scoring: 0, no or unable to determine; 1, yes; \*item does not contribute to total score.

High quality, total score  $\geq 7$ ; moderate quality, total score 4-6; low quality, total score  $\leq 3$ .


**Table 3.** Summary of consensus-based recommendations from the 2015 and 2017 International Patellofemoral Research Retreats.

STATEMENT	NEW PAPERS	RECOMMENDATION
<b>Exercise therapy:</b>		
Exercise therapy (vs. control):		
1. Reduces pain the short-term	18	Appropriate
2. Improves function and symptoms in the short-term	18	Uncertain
3. Reduces pain the medium- to long-term		Appropriate
4. Improves function and symptoms in the medium- to long-term		Appropriate
Combined hip and knee targeted exercise therapy:		
5. Reduces pain compared to knee targeted exercise therapy in the short-term	17 19	Appropriate
6. Improves function compared to knee targeted exercise therapy in the short-term	19	Appropriate
7. Reduces pain compared to knee targeted exercise therapy in the medium- to long-term		Appropriate
8. Improves function compared to knee targeted exercise therapy in the medium- to long-term		Appropriate
Hip targeted exercise therapy:		
9. Reduces pain compared to knee targeted exercise therapy in the short-term		Uncertain
10. Improves function compared to knee targeted exercise therapy in the short-term		Uncertain
11. Reduces pain compared to knee targeted exercise therapy in the medium- to long-term		Uncertain
12. Improves function compared to knee targeted exercise therapy in the medium- to long-term		Uncertain
<b>Combined interventions:</b>		
1. Reduces pain in the short-term	21	Appropriate
2. Reduces pain in the medium-term		Appropriate
3. Reduces pain in the long-term among adolescents with PFP		Uncertain
<b>Prefabricated foot orthoses:</b>		
1. Reduces pain in the short-term		Appropriate
2. Improves function in the short-term		Uncertain
<b>Patellar taping and bracing:</b>		
1. Tailored or untailored patellar taping to reduce pain immediately during functional tasks	25	Uncertain
2. Combine untailored patellar taping with exercise to further reduce pain in the long-term		Uncertain
3. Patellar bracing to reduce pain in the short- and medium-term	22 26	Uncertain
<b>Other adjunctive interventions:</b>		
1. <i>Acupuncture or dry needling to trigger points reduces pain in the short/medium term*</i>	27	<i>Uncertain*</i>
2. Patellofemoral and knee mobilisation improves pain or functional outcomes in the short-term		Inappropriate
3. Lumbar mobilisation/manipulation improves pain or functional outcomes in the short-term	29	Inappropriate
4. Electrophysical agents (e.g. ultrasound, phonophoresis, laser therapy etc.) improves pain or functional outcomes in the short-term		Inappropriate

5. <i>Manual soft tissue techniques (e.g. ischemic compression to peri- and retro-patellar trigger points; myofascial techniques) are beneficial in the short-term*</i>	29 32	<i>Uncertain*</i>
6. <i>Blood flow restriction training is superior to knee-focused exercise therapy to reduce pain with activities of daily living in the short-term*</i>	28	<i>Uncertain*</i>
7. <i>Gait retraining reduces pain and improve function in the short-term*</i>	18 30	<i>Uncertain*</i>

Short-term, <6 months; medium-term, 6-12 months; long-term, >12 months.

*Statements in italics were voted on as new/revised statements in 2017.* \*Wording has been amended for consistency with 2015 statements.





**1. It is uncertain whether acupuncture or dry needling to trigger points reduce pain in the short or medium term**

**Evidence:** One moderate quality systematic review<sup>36</sup> reported acupuncture to reduce pain more than a control at 5 months. One high quality RCT<sup>27</sup> reported no difference in pain at 2 weeks and 3 months when dry needling to trigger points was added to knee & hip targeted exercises and manual therapy.

**5. It is uncertain whether manual soft tissue techniques (e.g. ischemic compression to peri- and retro-patellar trigger points; myofascial techniques) are beneficial in the short-term**

**Evidence:** One high quality RCT<sup>37</sup> reported no improvement in pain with ischaemic compression to trigger points at the knee *versus* hip. One moderate quality RCT<sup>29</sup> reported greater pain reduction and functional improvement at 3 months with ischaemic compression compared to lumbopelvic manipulation. One low quality RCT<sup>32</sup> reported no difference in pain reduction or functional improvement at 5 weeks when myofascial techniques (massage + stretching) were added to hip-focused exercise therapy.

**6. It is uncertain whether blood flow restriction training is superior to knee focused exercise therapy to reduce pain with activities of daily living in the short term.**

**Evidence:** One high quality RCT<sup>28</sup> reported reduced pain during activities of daily living at 8 weeks with blood flow restriction + (low load) knee focused exercise therapy, compared with (standard) knee focused exercise therapy.

**7. It is uncertain whether gait retraining is effective to reduce pain and improve function, in the short term**

**Evidence:** One moderate quality RCT<sup>18</sup> reported no difference in pain reduction, function or symptom improvement at 8 weeks and 20 weeks with gait retraining (step-rate  $\pm$  strike pattern) + education compared to education alone. One moderate quality RCT<sup>30</sup> reported significant pain at 1 month following gait retraining (forefoot strike pattern) compared to no intervention.

