



**Manchester
Metropolitan
University**

Romano-Smith, S, Wood, G, Wright, DJ and Wakefield, CJ (2018) Simultaneous and alternate action observation and motor imagery combinations improve aiming performance. *Psychology of Sport and Exercise*, 38. pp. 100-106. ISSN 1469-0292

Downloaded from: <https://e-space.mmu.ac.uk/621377/>

Version: Accepted Version

Publisher: Elsevier

DOI: <https://doi.org/10.1016/j.psychsport.2018.06.003>

Usage rights: Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Please cite the published version

<https://e-space.mmu.ac.uk>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22

Simultaneous and Alternate Action Observation and Motor Imagery Combinations Improve
Aiming Performance

S. Romano-Smith^a, G. Wood^b, D. J. Wright^b and C.J. Wakefield^a

- a. School of Health Sciences, Liverpool Hope University, Taggart Avenue, Liverpool,
L16 9JD, UK
- b. Department of Exercise and Sport Science, Manchester Metropolitan University,
Crewe Green Road, Crewe CW1 5DU, UK

23 **Abstract**

24 Motor imagery (MI) and action observation (AO) are techniques that have been
25 shown to enhance motor skill learning. While both techniques have been used independently,
26 recent research has demonstrated that combining action observation and motor imagery
27 (AOMI) promotes better outcomes. However, little is known about the most effective way to
28 combine these techniques. This study examined the effects of simultaneous (i.e., observing an
29 action whilst imagining carrying out the action concurrently) and alternate (i.e., observing an
30 action and then doing imagery related to that action consecutively) AOMI combinations on
31 the learning of a dart throwing task. Participants (n=50) were randomly allocated to one of
32 five training groups: action observation (AO), motor imagery (MI), simultaneous action
33 observation and motor imagery (S-AOMI), alternate action observation and motor imagery
34 (A-AOMI) and a control group. Interventions were conducted three times per week for six
35 weeks and pre- and post-measures of total score were collected. Results revealed that all
36 intervention groups, with the exception of the AO and control groups, significantly improved
37 performance following the intervention. Posthoc analyses showed that S-AOMI group
38 improved to a significantly greater degree than the MI and AO groups, and participants in the
39 A-AOMI group improved to a significantly greater degree than the AO group. Participants in
40 the A-AOMI group did not improve to a significantly greater degree than the S-AOMI group
41 ($p=1.00$). These findings suggest that combining AOMI, regardless of how it's combined,
42 may be the beneficial method for improving the learning and performance of aiming skills.

43 *Keywords:* Motor skill learning; Motor imagery; Aiming; Action observation

44

45 **Introduction**

46 Motor imagery (MI) is the process of mentally rehearsing actions, typically without
47 overt action or physical output (Jeannerod, 2001). It is well established that MI interventions
48 can contribute to improvements in performance and learning in a wide variety of motor skills
49 (Cumming & Williams, 2012; Wakefield, Smith, Moran, & Holmes, 2013). To explain such
50 benefits, researchers have posited several explanations to explain improvements in
51 performance. The psychoneuromuscular theory (Jacobson, 1931) suggest mental practice
52 facilitates performance and the learning of a movement by causing a similar pattern of
53 muscular activation as during movement execution, which sequentially aids subsequent
54 movement execution. In contrast, the symbolic learning theory (Sackett, 1934) proposes that
55 the sequence of a movement is coded through symbols. Thus, by mentally rehearsing a
56 movement sequence through the repetition of symbolic components of the movement
57 sequence results in an improved symbolic representation.

58 Neuroscientific research has also provided an indication of the mechanism by which
59 imagery interventions contribute to such improvements in motor skill performance and
60 learning. Specifically, there is evidence that motor imagery activates similar brain regions to
61 those involved in motor skill planning and execution (Filimon, Nelson, Hagler, & Sereno,
62 2007). As such, MI practice is thought to activate and strengthen the cortical pathways
63 involved in motor skill execution and thereby contribute to improvements in motor
64 performance (Wakefield et al., 2013).

65 Like imagery, action observation (AO) interventions also offer an effective method
66 for improving performance and learning in a variety of motor skills (Ste-Marie et al., 2012).
67 Action observation involves the deliberate and structured observation of successful motor
68 skill execution (Neuman & Gray, 2013). The facilitation effect of AO is thought to reflect

69 involuntary activation of motor codes that are consistent with observed actions (bottom up
70 mechanism; Gibson 1966). The bottom-up mechanism is referred to as influences driven by
71 the extrinsic properties of stimuli (Baluch & Itti, 2011). Supporting this postulation is
72 evidence that observers copy the movement kinematics (speed) exhibited by a human model
73 which are coded through biological motion through lower level mechanisms of the AO
74 network (AON; Wild, Poliakoff & Gowen, 2010). AO also evokes activity in the areas of the
75 brain responsible for movement execution (Caspers, Zilles, Laird & Eickhoff, 2010).

76 Traditionally, MI and AO have been viewed as separate intervention techniques, with
77 researchers often comparing the two methods against each other to establish the most
78 effective for improving performance (e.g., Ram et al., 2007; Neumann & Gray, 2013). More
79 recently, however, researchers have begun to investigate the effects of combining action
80 observation and motor imagery (i.e., AOMI) by instructing participants to observe an action
81 presented in a video whilst simultaneously focusing on imagining the physiological
82 sensations and behavioural responses associated with the observed scenario (Scott, Taylor,
83 Chesterton, Vogt, & Eaves, 2017; Taube, Lorch, Zeiter, & Keller, 2014; Sun et al., 2016).
84 There is now a convincing body of evidence indicating that such AOMI interventions
85 produce increased activity in the motor regions of the brain, compared to either AO or MI
86 alone (see Eaves, Riach, Holmes, & Wright, 2016 for a review). As such, combined AOMI
87 approaches may be more effective for improving motor skill performance and learning than
88 the more traditional use of either independent AO or MI (Holmes & Wright, 2017).

89 Despite evidence that AOMI may produce greater activity in the motor regions of the
90 brain than the independent use of AO or MI, to date, relatively few experiments have
91 explored the effects of AOMI on the performance and learning of sport-related tasks. Those
92 studies that have been conducted have shown consistently positive effects for AOMI
93 interventions, compared to AO or MI alone, in strength (Scott et al., 2017; Wright & Smith,

94 2009), balance (Taube, et al., 2014) and golf putting (Smith & Holmes, 2004) tasks.
95 However, one unexplored issue in this area is how best to combine AOMI. In a recent study,
96 Sun et al. (2016) manipulated the structure of AOMI interventions in patients recovering
97 from stroke by asking patients to either combine AOMI simultaneously (S-AOMI) or by
98 alternating AO and MI components (A-AOMI). Specifically, these authors employed a 4-
99 week AOMI intervention where one group was instructed to observe a limb movement and
100 then subsequently asked to produce a mental image of the movement (A-AOMI) whilst the
101 other group practiced AOMI simultaneously (S-AOMI). Results showed that larger
102 improvements in grip strength and dexterity were observed within the effected limb in the S-
103 AOMI group.

104 To explain this finding the authors outlined two possible explanation: (1) that systems
105 shared by observation and imagery may be executed simultaneously in the S-AOMI condition
106 which may enhance cortex excitation or (2) that the observed action may enhance the
107 effectiveness and quality of simultaneous MI by providing learners with more direct
108 perceptual cues for the imagination of the same movement (Grèzes & Decety, 2001). Indeed,
109 there is some neuroscientific evidence that could support this. For example, Filimon et al.
110 (2015) and Hardwick et al. (2017) have showed that whilst both AO and MI activate the
111 similar areas of the brain (e.g., the premotor cortex), AO activates some areas more (e.g.,
112 inferior frontal gyrus; ventral premotor areas) than MI and MI activates other areas more
113 strongly (e.g., angular gyrus; dorsal premotor area) than AO. Given this evidence, it is
114 possible that S-AOMI (i.e., combining both approaches concurrently) would produce
115 increased and more widespread, activity in the premotor cortex than A-AOMI does and this is
116 what produces beneficial motor learning effects.

117 The aim of this experiment was replicate and extend these findings, from a clinical
118 population to individuals learning an aiming skill, in an effort to explore how generalizable

119 these effects are to other, more complex, motor skills (i.e., dart throwing) that require higher
120 levels of coordination, are temporally constrained actions and require greater levels of
121 accuracy. It was hypothesised that AO, MI, A-AOMI and S-AOMI practice would all
122 produce performance improvements from pre-test to post-test, relative to a control group. The
123 extent of the performance improvements were predicted to be greater in both combined
124 AOMI groups, compared to the independent AO or MI intervention (Eaves, Riach, Holmes,
125 & Wright, 2016). Finally, it was predicted that the S-AOMI group would exhibit greater
126 performance improvements when compared to A-AOMI group (as Sun et al., 2016).

127

128 **Method**

129 Participants

130 Fifty university students (25 males, 25 females; *Mean age = 23.88 years, SD = 3.78*)
131 were recruited. The number of participants recruited was established to be comparable to that
132 of previous research of a similar nature (Taube et al., 2014; Wright and Smith, 2009). All
133 participants self-reported being right-handed using the Edinburgh Handedness Inventory
134 (Oldfield, 1971). Participants also self-reported normal or corrected to normal vision and
135 were novice performers who had limited darts throwing experience and had not participated
136 in any previous MI training. The experiment was approved by the faculty ethics board at the
137 first author's institution.

138 Measures

139 *Movement Imagery Questionnaire-Revised (MIQR; Hall & Martin, 1997)*. The MIQ-
140 R is an eight-item inventory that assesses an individual's ability to perform visual and
141 kinaesthetic imagery on four movements: a knee lift, jump, arm movement and toe touch. In
142 this study, the MIQ-R was used as a screening tool, used by previous research (Smith &
143 Holmes, 2004; Wright & Smith, 2009). Participants physically performed each of the

144 requested actions a single time. Following execution of the action, participants were
145 instructed to image the movement, using an internal visual or kinaesthetic modality.
146 Participants then rated the ease or difficulty with which they completed the imagery on a 7-
147 point Likert type scale ranging from 1 (*very hard to see/feel*) to 7 (*very easy to see/feel*). The
148 validity and consistency of the MIQ-R has been demonstrated by Gregg, Hall, & Butler
149 (2010) and has been used previously in imagery studies investigating aiming tasks (e.g.,
150 Smith, Wright, & Cantwell, 2008)

151 *Imagery Diary*

152 Participants were provided with an imagery diary which they could complete after
153 each MI session by the guidelines of Goginsky and Collins (1996). Participants were
154 instructed to record any difficulties or concerns they experienced when performing imagery
155 during the intervention period. Furthermore, engagement with the session was measured
156 using a frequency count of sessions completed, out of a possible eighteen. The vividness and
157 controllability of the imagery were also rated on a 7-point Likert scale (ranging from 1 being
158 *not at all controllable not at all vivid* / 7 being *very controllable and very vivid*). Thorough
159 use of manipulation checks to ensure the completion of and focus of the intervention have
160 also been employed in a number of recent studies examining the efficacy of MI on
161 performance (e.g., Frank, Land, Popp, & Schack, 2014; Guillot, Genevois, Desliens, Saieb, &
162 Rogowski, 2012)

163 *The Aiming Task*

164 Concentric circle dartboard was used to collect performance data (see Figure 1). The
165 dartboard was positioned at the centre fixed point, 1.73cm from the floor and 2.37 cm
166 horizontally from the throwing line, as per standard darts rules. Performance (throwing
167 accuracy score) was measured using a similar system employed by Williams, and Cumming

168 (2012) measured in 10 concentric circles (2cm wide). The throws were scored in relation to
169 where the dart landed within the 10 circles, the centre of the scoring 10 points and the outer
170 circle scoring 1 point. Darts that landed outside the circumference of the dartboard were
171 awarded a score of zero.

172 **Procedure**

173 Prior to commencing the study, all participants provided informed consent and
174 completed the MIQ-R. All participants were randomly allocated to one of four experimental
175 groups (n =10/ group): action observation (AO); motor imagery (MI); simultaneous imagery
176 and observation (S-AOMI); and alternate imagery and observation (A-AOMI). Each group
177 contained five male and five female participants. All participants were given identical brief
178 instructions of the correct dart throwing technique that they should attempt to use when
179 completing the experiment. For example, participants were asked to focus on the centre of the
180 board, ensuring their dart and target were in line. They were also informed about the scoring
181 system and were instructed to aim for the centre of the board. After five practice throws,
182 participants completed their pre-test. This enabled the participants to experience the physical
183 sensation associated with holding a dart and executing a dart throw. The number of practice
184 throws were comparable to that of research of a similar target based task (Williams and
185 Cumming, 2012).

186 Pre and post-tests consisted of 30 dart throws split into six blocks of five dart throws.
187 Total score was taken as the performance measure during both pre and post-tests. Based on
188 the recommendations of others (Wakefield & Smith, 2009; Wright, McCormick, Birks,
189 Loporto, & Holmes, 2015) participants were instructed to perform each intervention for three
190 times per week, for a 6-week period. As previously indicated, participants' imagery diaries
191 also served as manipulation checks, ensuring that participants had correctly performed their

192 imagery as well as discussing deviations from normal behaviours such as sleeping patterns
193 and physical exertion. Any further information of issues or difficulties encountered with the
194 following MI interventions were also noted.

195 **Interventions**

196 Following the pre-test, the interventions were introduced to the participants. All
197 participants, except those in the control group and AO group, received stimulus response
198 training (SRT; Lang, Kozak, Miller, Levin, & McLean, 1980). Based on the bio
199 informational theory proposed by Lang et al. (1980), participants were instructed to attend to
200 specific stimulus details of the scenario that he/she finds easy to image (e.g., specific details
201 about the environment) and response propositions such as physiological sensations (e.g.,
202 muscle tension in their muscles), visceral events (e.g., increased heart rate) and sense organs
203 adjustments (e.g., postural changes). It has been suggested that imagery containing response
204 propositioning can produce more vivid imagery and consequently, improves the execution of
205 motor skills (Williams, Cooley, & Cumming, 2013). Over the 6 weeks, participants were
206 instructed to perform imagery in the first person perspective, with their eyes open and build
207 the image up by including additional details and/or by making the details more vivid or life
208 like. It is important to note however, this process was participant generated and participants
209 were not directed to specific propositions by the researchers.

210 *Control group*

211 The control group watched a video interview with a professional darts player three
212 times per week, which took the same amount of time as the videos presented to the other
213 treatment groups. The video was a documentary about darts, but did not provide advice on
214 the technique to aid the execution of a dart throw performance. Control participants were
215 informed that the study was designed to investigate the perception of dart throwing amongst

216 university students over a 6-week period. This procedure similar to the placebo used by
217 Smith and Holmes, (2004) and Smith et al. (2008).

218 *Action observation intervention*

219 The AO group were provided with the short pre-recorded observational video
220 containing six blocks of five dart throws, equalling thirty throws. Participants in this
221 treatment group were instructed to watch one of the pre-recorded videos (female hand/male
222 hand) equivalent to their sex. Video recordings provided participants with a view of the
223 model's right hand and forearm from a first person perspective (see Figure 1). A first person
224 perspective was employed for two reasons. First, there is evidence that action observation
225 from a first person perspective produces greater activity in the motor system than when
226 viewed from a third person perspective (Alaerts, Heremans, Swinnen, & Wenderoth, 2009).
227 Second, this perspective provides a closer behavioural match with physical performance than
228 would a third person perspective (Wakefield et al., 2013) and also ensured consistency with
229 conditions involving motor imagery which utilized a first person perspective based on the
230 PETTLEP imagery guidelines (Holmes & Collins, 2001). The video recording consisted of
231 observing an intermediate player executing thirty throws while attempting to hit the bullseye,
232 with a total score of 222/300. The characteristics of the model were comparable to that of
233 previous research of a similar nature suggesting the observation of trials that contained
234 degrees of error facilitated rapid learning of a fine motor task than observing trials that
235 contained minimal error (LeBel, Haverstock, Cristancho, van Eimeren, & Buckingham,
236 2017). The observational video was recorded in the same laboratory and with the same
237 equipment as used by participants in the study, allowing the combined intervention groups to
238 emphasise the environment component of the PETTLEP model.

239 *Imagery intervention group*

240 Each participant started by generating a simple image of themselves holding a dart,
241 with attention being drawn to aspects of the imaged scenario that they found easy to image.
242 Additional details to the relevant scenario were then progressively added (e.g. sensory
243 modalities, physiological sensations and emotional response). The completed script was then
244 used by the participant to practice during each imagery session. All aspects of the PETTLEP
245 model imagery (Holmes and Collins, 2001) were addressed in the interventions. The MI
246 group, along with all groups that incorporated MI into the intervention (A-AOMI and S-
247 AOMI) completed all elements of the model (see Table 1 for details of the PETTLEP
248 intervention).

249 *Alternate imagery and action observation (A-AOMI) group*

250 The A-AOMI group were provided with the pre-recorded observational video
251 containing six blocks of five dart throws, equalling 30 throws. Participants were required to
252 observe a block of five dart throws and were then were instructed to engage in PETTLEP MI
253 for a further five dart throws in an alternate manner until 30 throws were competed. The
254 structure of the trials allowed the participant to become accustomed to the requirements of the
255 intervention and were comparable to the trial structure of the study by Sun et al. (2016). The
256 PETTLEP MI aspect of the video was regulated by real time, as the screen during this
257 intervention showed a static dartboard and incorporated audio cues of the darts hitting the
258 board to ensure participants were imaging with the same timing as the observational element
259 of their intervention.

260 *Simultaneous imagery and action observation (S-AOMI) group*

261 The S-AOMI group were provided with the pre-recorded video containing six blocks
262 of five dart throws, equalling 30 throws. The video content was equivalent; however,
263 participants were given additional imagery instructions. Participants were instructed to

264 observe the dart throws shown in the video whilst simultaneously imaging the physiological
265 feelings and sensations that they would experience when executing performing the dart
266 throw.

267 *Data Analysis*

268 The data obtained from the MIQ-R imagery ability questionnaire were analysed using
269 separate one-way analyses of variance (ANOVAs) for the visual and kinaesthetic sub-scales
270 to establish any differences in imagery ability prior to the start of any intervention. Dart
271 throwing performance was measured as the mean of total throwing accuracy score (out of 300
272 points) for each group. This data was analysed using a 5 (group) x 2 (time) mixed between
273 within analysis of variance (ANOVA). Significance was measured at the .05 level. Where the
274 ANOVAs revealed significant effects, post-hoc Tukey tests were used to establish where any
275 significant differences existed. Effect sizes were calculated using partial eta squared (η_p^2) for
276 omnibus comparisons and Cohen's *d* for pairwise comparisons (Lakens, 2013).

277 **Results**

278 *Self-report data*

279 Results from the one-way ANOVAs revealed a significant difference in MIQ-K
280 scores, $F(4, 49) = 6.225, p < .001, \eta_p^2 = .356$ and MIQ-V scores, $F(4, 49) = 9.92, p < .001,$
281 $\eta_p^2 = .469$. Post-hoc Tukey tests showed that participants in the control group scored
282 significantly lower than participants in the intervention groups (all $p < .05$) for both visual
283 and kinaesthetic imagery ability (see Table 2). This result was expected as, prior to the pre-
284 test, low scoring imagers were deliberately placed into the control group prior to testing to
285 reduce the likelihood of control group participants engaging in spontaneous imagery of the
286 task throughout the intervention period. Importantly, no significant differences between

287 imagery ability were apparent for intervention groups on MIQ-K scores and MIQ-V scores
288 (all $p > .05$).

289 *Self-report data: manipulation checks*

290 Inspection of the imagery diaries and manipulation checks conducted revealed that
291 participants reported performing their imagery as instructed by the researcher. Prior to the
292 completion of the testing, a minimum of 14 intervention sessions was set as the cut-off point,
293 and completion of less than 14 would result in the participant's data being removed from the
294 study. As all participants reported completing at minimum of 14 sessions, all data were
295 included in the study. Furthermore, there were no significant imagery content differences for
296 imaging, ease of visual or kinaesthetic imagery, or imagery vividness (p 's $> .05$). These data
297 are presented in Table 3.

298 *Performance*

299 Results revealed a significant main effect for time, $F(1, 9) = 20.37, p < .001, \eta_p^2 =$
300 $.694$, and a significant main effect of group, $F(4, 36) = 3.172, p = 0.03, \eta_p^2 = .261$. There was
301 also a significant time x group interaction, $F(4, 36) = 6.44, p < .001, \eta_p^2 = .417$. Within group
302 post hoc comparisons using the Tukey test revealed significant improvements from pre-test to
303 post-test in the A-AOMI ($p = .001, d = 1.57$), S-AOMI ($p = .001, d = 1.79$) and MI ($p = .020,$
304 $d = 1.14$) groups. Participants in both the AO group and control group did not significantly
305 improve performance from pre- to post-test. Between group post hoc tests showed that the S-
306 AOMI group improved to a significantly greater degree than the AO ($p = .03, d = 1.17$), MI (p
307 $= .05, d = 1.11$), and control ($p = .001, d = 1.74$) groups. Participants in the A-AOMI group
308 improved to a significantly greater degree than the AO ($p = .05, d = 0.95$) and control ($p = .002,$
309 $d = 1.61$) groups. Participants in the A-AOMI group did not improve to a significantly greater
310 degree than the S-AOMI group ($p = 1.00$; see Figure 2).

311

312 **Discussion**

313 The aim of this experiment was to explore the effects of differing combinations of
314 AOMI practice against independent AO or MI practice on performance in an aiming task.
315 The results indicate that both combinations of imagery and observation training (i.e., S-
316 AOMI and A-AOMI) can improve target performance over-and-above AO or MI
317 interventions alone. This corroborates the findings of previous research that has reported
318 similar improvements in motor performance after AOMI interventions (see Eaves et al.,
319 2016). The findings of this experiment indicate that combining imagery and observation may
320 provide the optimal method for producing performance improvements in target throwing
321 tasks.

322 Importantly, however, both S-AOMI and A-AOMI appear to provide equivalent
323 performance enhancements during for this type of skill, which is in direct contrast to the
324 findings of Sun et al. (2016). One possible explanations for the discrepancy could be due
325 differences between the participants in both studies. For example, Sun et al. (2016) recruited
326 patients recovering from stroke while our study used ‘non-affected’ adults. As patients
327 recovering from stroke usually have impairments in working memory (WM) (Constantinidis
328 & Klingberg, 2016) it could be the case that the S-AOMI condition reduced the demand on
329 WM resources by eliminating the need to remember the action observed in order to guide
330 their MI. We propose that the participants in our study, whom presumably had normal levels
331 of WM, had sufficient WM resources to cope with the demands of either AOMI combination.
332 Therefore the optimal structure for AOMI interventions may be an important consideration
333 for clinical populations who have impairments in WM such as the elderly (Schott, 2012),
334 children with developmental disorders (Alloway, 2011) or patients with Parkinson’s disease

335 (Lees & Smith, 1983). Future research is warranted to evaluate the merits of such AOMI
336 combinations in these populations.

337 One explanation for why the two AOMI interventions resulted in greater performance
338 improvements than the independent AO or MI interventions may relate to the manner in
339 which they produced activity in the motor regions of the brain. Although no measure of
340 neural activity was included in this experiment, it is well established that both AO and MI
341 evoke activity in the motor regions of the brain (e.g., Grezes & Decety, 2001), and that
342 AOMI interventions elicit greater activity in these brain regions than independent AO or MI
343 (Eaves et al., 2016). As such, by engaging in both AO and MI three times per week for six
344 weeks, either in a simultaneous or alternate manner, participants in the S-AOMI and A-
345 AOMI groups may have experienced increased activity in motor-related brain regions during
346 their intervention than either the independent AO or MI groups. Although the independent
347 AO or MI interventions would likely still have elicited activity in similar regions of the motor
348 system, this is likely to have occurred to a lesser extent than in the two AOMI groups, and
349 this may explain why their performance did not improve to the same level as either
350 combination groups. To substantiate this explanation, further research utilizing mobile
351 electroencephalography technology to record cortical activity during AOMI interventions
352 alongside performance measures would be welcome.

353 Another explanation for the greatest improvements being found in the two AOMI
354 intervention groups may be that AOMI helps to develop a common motor representation that
355 helps to prime top-down attentional processes (e.g., action intention, movement programming
356 and preparation) which are important for task execution (Jeannerod, 2001). Evidence to
357 support this explanation can be taken from studies that have shown similar eye-movement
358 patterns during physical practice and MI (Heremans et al., 2009), physical practice and AO
359 (Flanagan & Johansson, 2003) and MI and AO group (McCormick et al., 2012). This

360 suggests that eye-movement patterns observed in motor simulation interventions may reflect
361 the shared neural network used to plan and control visually guided actions during physical
362 practice. Therefore, it is possible that the improvement in darts throwing performance in this
363 study was attributable to the development of optimal eye-movement strategies important for
364 aiming. In fact, previous research by Frank, Land and Schack (2015) has shown that mental
365 simulation of a golf-putting task resulted in more elaborate motor representations which
366 facilitated more optimal eye-movement behaviours (quiet-eye (QE) durations; Vickers, 2007)
367 shown to be important in aiming skills. Future research should therefore explore the utility of
368 AOMI interventions for implicitly facilitating QE aiming durations in such tasks.

369 Our data showed no significant change in performance in the AO group, yet
370 significant improvements in the MI group. This is surprising, as previous studies that have
371 employed AO in isolation have showed this to be effective (e.g., Battaglia et al., 2014; Gatti
372 et al., 2013). One potential explanation for this finding may be that MI is more cognitively
373 demanding compared to AO. For example, MI depends on the individual's ability to rehearse
374 or recruit the relevant motor representation and to perform the action covertly while
375 generating visual and kinaesthetic imagery. On the other hand, AO interventions provide a
376 model of the action with minimal instruction and therefore imposes a lower cognitive
377 demand. This disparity in the mental resources employed during either intervention in
378 isolation may explain these differing effects on performance and learning.

379 A potential limitation of the study is our decision to place poor imagers into the
380 control group. However, this decision was taken to reduce the likelihood of spontaneous
381 imagery throughout the intervention period that has been suggested in similar research (i.e.,
382 Smith et al., 2008). Despite this justification, this decision will have an impact on how
383 generalizable these findings maybe be individuals with poor imagery ability. Another
384 limitation of our study relates to the nature of the performance measurement used. Criticism

385 of this method suggests that it lacks sensitivity and is inappropriate for the capture of the true
386 characteristics of performance such as direction and variability around the target (see
387 Fischman, 2015). Finally, the decision to ask participants to complete the intervention at
388 home may be a further limitation of the study design, as we cannot ensure subjects integrity
389 to engage in the intervention period. However, the improvements in performance suggest that
390 this was not the case.

391 In conclusion, in this study we have shown that two types of AOMI interventions
392 improved dart throwing performance over-and-above AO or MI interventions alone. This
393 offers further behavioural evidence to support the efficacy of AOMI for improving
394 performance in sport. As such, sport psychologists should consider adapting their practice to
395 include the delivery of combined AOMI interventions. Finally, further research should seek
396 to explore whether the two combinations AOMI provide similar benefits when employed in
397 other populations and with other, more complex motor skills.

398

399

400

401

402

403

404

405

406

407

408

409 **Funding**

410 This research did not receive any specific grant from funding agencies in the public,
411 commercial, or not-for-profit sectors.

412 **Conflict of interest**

413 None.

414

415

416

417

418

419

420

421

422

423

424

425

426

427 **References**

- 428 Alaerts, K., Heremans, E., Swinnen, S. P., & Wenderoth, N. (2009). How are observed
429 actions mapped to the observer's motor system? Influence of posture and perspective.
430 *Neuropsychologia*, *47*(2), 415–422.
431 <https://doi.org/10.1016/j.neuropsychologia.2008.09.012>
- 432 Alloway, T. P. (2011). A comparison of working memory profiles in children with ADHD
433 and DCD. *Child Neuropsychology*, *17*(5), 483-494.
- 434 Baluch, F., & Itti, L. (2011). Mechanisms of top-down attention. *Trends in Neurosciences*,
435 *34*(4), 210–224. <https://doi.org/10.1016/j.tins.2011.02.003>
- 436 Battaglia, C., D'Artibale, E., Fiorilli, G., Piazza, M., Tsopani, D., Giombini, A., di Cagno, A.
437 (2014). Use of video observation and motor imagery on jumping performance in
438 national rhythmic gymnastics athletes. *Human Movement Science*, *38*, 225–234.
439 <https://doi.org/10.1016/j.humov.2014.10.001>
- 440 Cumming, J., & Williams, S. E. (2012). Imagery: The role of imagery in performance. In S.
441 Murphy (Ed.), *Handbook of sport and performance psychology* (pp. 213-232). New
442 York, NY: Oxford University Press. doi:10.1093/oxfordhb/9780199731763.013.0011
- 443 Caspers, S., Zilles, K., Laird, A. R., & Eickhoff, S. B. (2010). ALE meta-analysis of action
444 observation and imitation in the human brain. *NeuroImage*, *50*(3), 1148–1167.
445 <https://doi.org/10.1016/j.neuroimage.2009.12.112>
- 446 Constantinidis, C., & Klingberg, T. (2016). The neuroscience of working memory capacitand
447 training. *Nature Reviews. Neuroscience*, *17*(7), 438–449.
448 <https://doi.org/10.1038/nrn.2016.43>

- 449 Eaves, D. L., Riach, M., Holmes, P. S., & Wright, D. J. (2016). Motor Imagery during Action
450 Observation: A Brief Review of Evidence, Theory and Future Research
451 Opportunities. *Frontiers in Neuroscience, 10*. <https://doi.org/10.3389/fnins.2016.0051>
- 452 Flanagan, J. R., & Johansson, R. S. (2003). Action plans used in action observation. *Nature,*
453 *424(6950), 769–771*. <https://doi.org/10.1038/nature01861>
- 454 Filimon, F., Nelson, J. D., Hagler, D. J., & Sereno, M. I. (2007). Human cortical
455 representations for reaching: mirror neurons for execution, observation, and imagery.
456 *NeuroImage, 37(4), 1315–1328*. <https://doi.org/10.1016/j.neuroimage.2007.06.008>
- 457 Filimon, F., Rieth, C. A., Sereno, M. I., & Cottrell, G. W. (2015). Observed, Executed, and
458 Imagined Action Representations can be Decoded From Ventral and Dorsal Areas.
459 *Cerebral Cortex (New York, N.Y.: 1991), 25(9), 3144–3158*.
460 <https://doi.org/10.1093/cercor/bhu110>
- 461 Fischman, M. G. (2015). On the continuing problem of inappropriate learning measures:
462 Comment on Wulf et al. (2014) and Wulf et al. (2015). *Human Movement Science,*
463 *42, 225–231*. <https://doi.org/10.1016/j.humov.2015.05.011>
- 464 Frank, C., Land, W. M., Popp, C., & Schack, T. (2014). Mental Representation and Mental
465 Practice: Experimental Investigation on the Functional Links between Motor Memory
466 and Motor Imagery. *PLOS ONE, 9(4), e95175*.
467 <https://doi.org/10.1371/journal.pone.0095175>
- 468 Frank, C., Land, W. M., & Schack, T. (2015). Perceptual-cognitive changes during motor
469 learning: The influence of mental and physical practice on mental representation, gaze
470 behavior, and performance of a complex action. *Frontiers in psychology, 6*.
- 471 Gatti, R., Tettamanti, A., Gough, P. M., Riboldi, E., Marinoni, L., & Buccino, G. (2013).
472 Action observation versus motor imagery in learning a complex motor task: a short

- 473 review of literature and a kinematics study. *Neuroscience Letters*, 540, 37–42.
474 <https://doi.org/10.1016/j.neulet.2012.11.039>
- 475 Goginsky, A. M., & Collins, D. (1996). Research design and mental practice. *Journal of*
476 *Sports Sciences*, 14(5), 381–392. <https://doi.org/10.1080/02640419608727725>
- 477 Gregg, M., Hall, C., & Butler, A. (2010). The MIQ-RS: A Suitable Option for Examining
478 Movement Imagery Ability. *Evidence-Based Complementary and Alternative*
479 *Medicine : eCAM*, 7(2), 249–257. <https://doi.org/10.1093/ecam/nem170>
- 480 Grèzes, J., & Decety, J. (2001). Functional anatomy of execution, mental simulation,
481 observation, and verb generation of actions: a meta-analysis. *Human Brain Mapping*,
482 12(1), 1–19.
- 483 Guillot, A., Genevois, C., Desliens, S., Saieb, S., & Rogowski, I. (2012). Motor imagery and
484 “placebo-racket effects” in tennis serve performance. *Psychology of Sport and*
485 *Exercise*, 13(5), 533–540. <https://doi.org/10.1016/j.psychsport.2012.03.002>
- 486 Hardwick, R. M., Caspers, S., Eickhoff, S. B., & Swinnen, S. P. (2017). Neural Correlates of
487 Motor Imagery, Action Observation, and Movement Execution: A Comparison
488 Across Quantitative Meta-Analyses. *BioRxiv*, 198432. <https://doi.org/10.1101/198432>
- 489 Heremans, E., Helsen, W. F., De Poel, H. J., Alaerts, K., Meyns, P., & Feys, P. (2009).
490 Facilitation of motor imagery through movement-related cueing. *Brain Research*,
491 1278, 50–58. <https://doi.org/10.1016/j.brainres.2009.04.041>
- 492 Holmes, P. S., & Collins, D. J. (2001). The PETTLEP Approach to Motor Imagery: A
493 Functional Equivalence Model for Sport Psychologists. *Journal of Applied Sport*
494 *Psychology*, 13(1), 60–83. <https://doi.org/10.1080/10413200109339004>
- 495

- 496 Holmes, P. S., & Wright, D. J. (2017). Motor cognition and neuroscience in sport
497 psychology. *Current Opinion in Psychology*, *16*, 43–47.
498 <https://doi.org/10.1016/j.copsyc.2017.03.009>
- 499 JR, S. Y. E. (1969a). James J. Gibson, The Senses Considered as Perceptual Systems. *The Art*
500 *Bulletin*, *51*(3), 310–311. <https://doi.org/10.1080/00043079.1969.10790296>
- 501 Jacobson, E. (1931). Electrical measures of neuromuscular states during mental activities. V.
502 *American Journal of Physiology*, *96*, 1 15-121.
- 503 Jeannerod, M. (2001). Neural simulation of action: a unifying mechanism for motor
504 cognition. *NeuroImage*, *14*(1 Pt 2), S103-109. <https://doi.org/10.1006/nimg.2001.0832>
- 505 Lakens, D. (2013). Calculating and reporting effect sizes to facilitate cumulative science: a
506 practical primer for t-tests and ANOVAs. *Frontiers in Psychology*, *4*, 863.
507 <https://doi.org/10.3389/fpsyg.2013.00863>
- 508 Lang, P. J., Kozak, M. J., Miller, G. A., Levin, D. N., & McLean, A. (1980). Emotional
509 imagery: conceptual structure and pattern of somato-visceral response.
510 *Psychophysiology*, *17*(2), 179–192.
- 511 Lees, A. J., & Smith, E. (1983). Cognitive deficits in the early stages of Parkinson's disease.
512 *Brain*, *106*(2), 257-270.
- 513 LeBel, M.-E., Haverstock, J., Cristancho, S., van Eimeren, L., & Buckingham, G. (2017).
514 Observational Learning During Simulation-Based Training in Arthroscopy: Is It
515 Useful to Novices? *Journal of Surgical Education*.
516 <https://doi.org/10.1016/j.jsurg.2017.06.005>
- 517 McCormick, S. A., Causer, J., & Holmes, P. S. (2012). Eye gaze metrics reflect a shared
518 motor representation for action observation and movement imagery. *Brain and*
519 *Cognition*, *80*(1), 83–88. <https://doi.org/10.1016/j.bandc.2012.04.010>

- 520 Neuman, B., & Gray, R. (2013). A direct comparison of the effects of imagery and action
521 observation on hitting performance, *Abstract. Movement & Sport Sciences*, (79), 11–
522 21.
- 523 Oldfield, R. C. (1971). The assessment and analysis of handedness: the Edinburgh inventory.
524 *Neuropsychologia*, 9(1), 97–113.
- 525 Ram, N., Riggs, S. M., Skaling, S., Landers, D. M., & McCullagh, P. (2007). A comparison
526 of modelling and imagery in the acquisition and retention of motor skills. *Journal of*
527 *Sports Sciences*, 25(5), 587–597. <https://doi.org/10.1080/02640410600947132>
- 528 Sackett, R. S. (1934). Influence of symbolic rehearsal upon retention of maze habit. *Journal*
529 *of General Psychology*, 10, 376-396.
- 530 Scott, M., Taylor, S., Chesterton, P., Vogt, S., & Eaves, D. L. (2017). Motor imagery during
531 action observation increases eccentric hamstring force: an acute non-physical
532 intervention. *Disability and Rehabilitation*, 1–9.
533 <https://doi.org/10.1080/09638288.2017.1300333>
- 534 Schott, N. (2012). Age-related differences in motor imagery: working memory as a mediator.
535 *Experimental Aging Research*, 38(5), 559-583.
- 536 Smith, D., & Holmes, P. (2004). The Effect of Imagery Modality on Golf Putting
537 Performance. *Journal of Sport and Exercise Psychology*, 26(3), 385–395.
538 <https://doi.org/10.1123/jsep.26.3.385>
- 539 Smith, D., Wright, C. J., & Cantwell, C. (2008). Beating the bunker: the effect of PETTLEP
540 imagery on golf bunker shot performance. *Research Quarterly for Exercise and Sport*,
541 79(3), 385–391. <https://doi.org/10.1080/02701367.2008.10599502>
- 542 Ste-Marie, D. M., Law, B., Rymal, A. M., Jenny, O., Hall, C., & McCullagh, P. (2012).
543 Observation interventions for motor skill learning and performance: an applied model

- 544 for the use of observation. *International Review of Sport and Exercise Psychology*,
545 5(2), 145–176. <https://doi.org/10.1080/1750984X.2012.665076>
- 546 Sun, Y., Wei, W., Luo, Z., Gan, H., & Hu, X. (2016). Improving motor imagery practice with
547 synchronous action observation in stroke patients. *Topics in Stroke Rehabilitation*,
548 23(4), 245–253. <https://doi.org/10.1080/10749357.2016.1141472>
- 549 Taube, W., Lorch, M., Zeiter, S., & Keller, M. (2014). Non-physical practice improves task
550 performance in an unstable, perturbed environment: motor imagery and observational
551 balance training. *Frontiers in Human Neuroscience*, 8.
552 <https://doi.org/10.3389/fnhum.2014.00972>
- 553 Vickers, J. N. (2007). *Perception, Cognition, and Decision Training: The Quiet Eye in*
554 *Action*. Human Kinetics.
- 555 Villiger, M., Estévez, N., Hepp-Reymond, M.-C., Kiper, D., Kollias, S. S., Eng, K., & Hotz-
556 Boendermaker, S. (2013). Enhanced activation of motor execution networks using
557 action observation combined with imagination of lower limb movements. *PloS One*,
558 8(8), e72403. <https://doi.org/10.1371/journal.pone.0072403>
- 559 Wakefield, C. J., & Smith, D. (2009). Impact of differing frequencies of PETTLEP imagery
560 on netball shooting performance. *Journal of Imagery Research in Sport and Physical*
561 *Activity*, 4(1), 1–12.
- 562 Wakefield, C., Smith, D., Moran, A. P., & Holmes, P. (2013). Functional equivalence or
563 behavioural matching? A critical reflection on 15 years of research using the
564 PETTLEP model of motor imagery. *International Review of Sport and Exercise*
565 *Psychology*, 6(1), 105–121. <https://doi.org/10.1080/1750984X.2012.724437>
- 566 Williams, S. E., & Cumming, J. (2012). Challenge vs. threat imagery: Investigating the effect
567 of using imagery to manipulate cognitive appraisal of a dart throwing task. *Sport and*
568 *Exercise Psychology Review*, 8, 4–21.

- 569 Williams, S. E., Cooley, S. J., & Cumming, J. (2013). Layered stimulus response training
570 improves motor imagery ability and movement execution. *Journal of Sport &*
571 *Exercise Psychology*, 35(1), 60–71.
- 572 Wild, K. S., Poliakoff, E., Jerrison, A., & Gowen, E. (2010). The influence of goals on
573 movement kinematics during imitation. *Experimental Brain Research*, 204(3), 353–
574 360. <https://doi.org/10.1007/s00221-009-2034-8>
- 575 Wood, J. N. (2007). Visual working memory for observed actions. *Journal of Experimental*
576 *Psychology: General*, 136(4), 639.
- 577 Wright, C. J., & Smith, D. (2009). The effect of PETTLEP imagery on strength performance.
578 *International Journal of Sport and Exercise Psychology*, 7(1), 18–31.
579 <https://doi.org/10.1080/1612197X.2009.9671890>
- 580 Wright, D. J., McCormick, S. A., Birks, S., Loporto, M., & Holmes, P. S. (2015). Action
581 Observation and Imagery Training Improve the Ease With Which Athletes Can
582 Generate Imagery. *Journal of Applied Sport Psychology*, 27(2), 156–170.
583 <https://doi.org/10.1080/10413200.2014.968294>
584
585
586
587
588
589
590
591

592 **Figure Captions**593 **Figure 1.** An example still shot from the Action Observation video594 **Figure 2.** Mean (\pm s.e.m) pre and post-test throwing accuracy scores for each experimental
595 group ($*p < .05$, $**p < .001$).

596

597

598

599

600

601

602

603

604

605

606

607

608

609

610

611

612 **Figure 1**



613

614

615

616

617


618

619

620

621 **Figure 2**

622



623

624

625

Table 1. Summary of the PETLEP motor imagery content for all imagery instructions

PETTLEP category	Description
Physical	Participants were instructed to stand while holding a cylindrical object similar to a dart or pen suggested by Holmes and Collins (2001). Participants were also instructed to adopt the stance recognised in dart throwing performance.
Environment	PETTLEP MI was performed at home. Participants were instructed to watch the video static dartboard within the video from their pre-test
Task	Participants performed a series of dart throws to emulate the performance measure as closely as possible. This included the intricacies associated with their specific skill level on the task.
Timing	Participants were instructed to perform MI in ‘real time’, rather than in slow motion or faster than normal. Auditory cues. For example, audio feedback of the darts making contact with the board during pre-test conditions.
Learning	Participant were instructed to revisit their imagery scripts after every two week period of the intervention and make any necessary adaptations depending on their perceived development of the skill.
Emotion	Scripts were created after the pre-test allowing familiarisation with the dart throwing action. This was based on the results of the stimulus and response training (Lang et al., 1980) that had been undertaken. Participants often identified associations with the physical sensations or of dart throwing.
Perspective	Participants were instructed to image in the first person perspective in order to best reflect the perspective from physical completion of the task.

626

627

628

629

630

631

632

Table 2. Mean MIQ-R scores and (SD) for each experimental group.

Group	MIQ-R Visual	MIQ-R Kinaesthetic
A-AOMI	6.7 (0.64)	5.9 (0.98)
S-AOMI	6.4 (0.52)	6.2 (0.64)
MI	6.3 (0.66)	6.3 (0.93)
AO	6.0 (0.62)	5.8 (0.61)
Control	4.8 (0.81)	4.5 (0.61)

639

640

Table 3. Manipulation check mean scores (SD) for number of sessions completed, ease of visual, kinaesthetic imagery, and imagery vividness for each experimental group.

	A-AOMI	S-AOMI	MI
Frequency of imaging	16.1 (0.54)	16.4 (0.47)	15.8 (0.53)
Ease of imagery (see)	6.7 (0.15)	6.5 (0.17)	6.7 (0.15)
Ease of imagery (feel)	6.5 (0.16)	6.5 (0.18)	6.7 (0.15)
Vividness of imagery	6.5 (0.16)	6.5 (0.16)	6.7(0.15)