

**Please cite the Published Version**

Dean, AP, Hartley, A, McIntosh, OA, Smith, A, Feord, HK, Holmberg, NH, King, T, Yardley, E, White, KN and Pittman, JK (2019) Metabolic adaptation of a *Chlamydomonas acidophila* strain isolated from acid mine drainage ponds with low eukaryotic diversity. *Science of the Total Environment*, 647. pp. 75-87. ISSN 0048-9697

**DOI:** <https://doi.org/10.1016/j.scitotenv.2018.07.445>

**Publisher:** Elsevier

**Version:** Accepted Version

**Downloaded from:** <https://e-space.mmu.ac.uk/621624/>

**Usage rights:**



[Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0](#)

**Additional Information:** Copyright Elsevier.

**Enquiries:**

If you have questions about this document, contact [openresearch@mmu.ac.uk](mailto:openresearch@mmu.ac.uk). Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

**Metabolic adaptation of a *Chlamydomonas acidophila* strain isolated from acid mine drainage ponds with low eukaryotic diversity**

Andrew P. Dean<sup>a</sup>, Antoni Hartley<sup>d</sup>, Owen A. McIntosh<sup>d</sup>, Alyssa Smith<sup>b,1</sup>, Helen Feord<sup>b,2</sup>,  
Nicolas H. Holmberg<sup>b</sup>, Thomas King<sup>b</sup>, Ellen Yardley<sup>c</sup>, Keith N. White<sup>b,d</sup>, and Jon K.  
Pittman<sup>b,d,\*</sup>

<sup>a</sup>School of Science and the Environment, Manchester Metropolitan University, Oxford Road,  
Manchester, M1 5GD, UK; <sup>b</sup>Faculty of Life Sciences, University of Manchester, Michael  
Smith Building, Oxford Road, Manchester M13 9PT, UK; <sup>c</sup>Department of Geography,  
University of Sheffield, Sheffield S10 2TN, UK; <sup>d</sup>School of Earth and Environmental  
Sciences, Faculty of Science and Engineering, University of Manchester, Michael Smith  
Building, Oxford Road, Manchester M13 9PT, UK

\*Corresponding author.

Email address: [jon.pittman@manchester.ac.uk](mailto:jon.pittman@manchester.ac.uk) (J.K. Pittman)

<sup>1</sup>Present address: Department of Chemistry, University of Cambridge, Cambridge, UK

<sup>2</sup>Present address: School of Biological Sciences, University of Edinburgh, Edinburgh, UK

## Abstract

The diversity and biological characteristics of eukaryotic communities within acid mine drainage (AMD) sites is less well studied than for prokaryotic communities. Furthermore, for many eukaryotic extremophiles the potential mechanisms of adaptation are unclear. This study describes an evaluation of eight highly acidic (pH 1.6 – 3.1) and one moderately acidic (pH 5.6) metal-rich acid mine drainage ponds at a disused copper mine. The severity of AMD pollution on eukaryote biodiversity was examined, and while the most species-rich site was less acidic, biodiversity did not only correlate with pH but also with the concentration of dissolved and particulate metals. Acid-tolerant microalgae were present in all ponds, including the species *Chlamydomonas acidophila*, abundance of which was high in one very metal-rich and highly acidic (pH 1.6) pond, which had a particularly high PO<sub>4</sub>-P concentration. The *C. acidophila* strain named PM01 had a broad-range pH tolerance and tolerance to high concentrations of Cd, Cu and Zn, with bioaccumulation of these metals within the cell. Comparison of metal tolerance between the isolated strain and other *C. acidophila* strains previously isolated from different acidic environments found that the new strain exhibited much higher Cu tolerance, suggesting adaptation by *C. acidophila* PM01 to excess Cu. An analysis of the metabolic profile of the strains in response to increasing concentrations of Cu suggests that this tolerance by PM01 is in part due to metabolic adaptation and changes in protein content and secondary structure.

**Keywords:** *Chlamydomonas acidophila*, acid tolerance, metal tolerance, acid mine drainage, bioremediation, copper, zinc, cadmium

## 1. Introduction

Many freshwater bodies worldwide are highly acidic either due to natural causes or anthropogenic activities such as mining (Schultze, 2013; Smucker et al., 2014). Acid mine drainage (AMD) due principally to pyrite oxidation, is the cause of significant acidity in lakes and ponds situated in areas impacted by mining, and in rivers receiving mine water

discharge (Johnson, 2003; Nordstrom, 2000). Because decreasing pH causes increased solubility of metals, AMD results in high concentrations of dissolved Fe, S and various trace metals such as Cu, Cd and Zn in the contaminated waters. Concentrations of nutrients, especially inorganic phosphate ( $\text{PO}_4\text{-P}$ ), are also frequently very low (Nixdorf et al., 1998). The combination of toxic metals and nutrient limitation limits biodiversity and can cause significant ecosystem damage (Deneke, 2000; Smucker et al., 2014). Evaluation of the biological impacts of AMD allows quantification of pollution damage, allows understanding of fundamental processes of adaptation and can identify AMD-tolerant species that have biotechnological applications, such as bioremediation (Ñancucheo and Johnson, 2011; Yun et al., 2014).

While prokaryotes in AMD environments has been extensively studied and reviewed (Johnson and Hallberg, 2003; Mendez-Garcia et al., 2015), there is still limited knowledge regarding the presence and roles of eukaryotes in these aquatic environments (Aguilera et al., 2006; Baker et al., 2004; Nixdorf et al., 1998). Photosynthetic microorganisms are found in many AMD ecosystems; however, the biodiversity of phytoplankton in such waters is severely limited and dominated by just a few acid-tolerant genera, such as *Chlamydomonas*, *Dunaliella*, *Euglena* and *Ochromonas* (Aguilera et al., 2006; Hargreaves et al., 1975; Ñancucheo and Johnson, 2012; Nixdorf et al., 1998; Pedrozo et al., 2001). Despite being able to tolerate the highly acidic and metal-rich conditions, productivity of these extremophile microalgae is often limited by low inorganic carbon and nutrient availability in acidic waters (Beamud et al., 2007; Spijkerman et al., 2007b). A fairly broad diversity of heterotrophic fungi and protists has also been observed in acidic waters (Baker et al., 2004; Das et al., 2009), while the diversity and abundance of zooplankton is typically very low as most species are unable to tolerate these environments (Deneke, 2000).

The high concentrations of dissolved metals in AMD can cause toxicity to microorganisms through a wide variety of mechanisms, some of which are shared between metals and across different organisms, such as competition with essential metals, direct interactions with proteins and other molecules within the cell, and induction of oxidative

stress (Sharma and Dietz, 2009). Metals such as Cu are particularly efficient at inducing the formation of reactive oxygen species (ROS) in contrast to non-redox active metals such as Zn and Cd (Valko et al., 2005). In most photosynthetic organisms, excess Cu has many detrimental effects with the photosynthetic apparatus, including direct inhibition of photosynthetic activity and degradation of chloroplast structures (Bernal et al., 2006; Küpper et al., 2003). Furthermore, non-extremophile microalgae exposed to high Cu conditions exhibit high concentrations of ROS and subsequent ROS-induced damage including lipid membrane peroxidation (Jamers et al., 2013; Jiang et al., 2016; Sabatini et al., 2009).

The adaptive mechanisms by which eukaryotic microorganisms including extremophile microalgae can survive in acid and metal rich conditions are still poorly researched but potential insights into these mechanisms are increasing. For example, proteomic approaches have indicated the importance of metal and acidity tolerance proteins, such as molecular chaperones of the Heat Shock Protein family (Cid et al., 2010; Gerloff-Elias et al., 2006). Likewise, genome sequencing and transcriptomics studies are beginning to identify the array of genes that might explain extremophile functional characteristics, some of which may have been obtained by horizontal gene transfer from bacteria. Genome sequences of the acidophiles *Chlamydomonas eustigma* (Hirooka et al., 2017) and *Galdieria sulphuraria* (Schönknecht et al., 2013) have recently been determined. Furthermore, transcriptomic approaches are beginning to provide insight into the molecular mechanisms of *Chlamydomonas acidophila* tolerance in response to Cd and Cu exposure (Olsson et al., 2015; Puente-Sánchez et al., 2018), and *Dunaliella acidophila* in response to Cd (Olsson et al., 2017; Puente-Sánchez et al., 2016), although further experimental analyses of these transcriptomic datasets are needed.

AMD tolerant biota might have potential for bioremediation, with biological-based processes potentially more cost effective and sustainable than chemical based methods such as anoxic limestone drains and chemical addition (Geller and Schultze, 2013; Hedin et al., 2010; Johnson and Hallberg, 2005). Bioremediation methods can include utilisation of bacterial SO<sub>4</sub> reduction and neutralisation (Neculita et al., 2007) or aerobic wetlands that can

oxidise and precipitate dissolved metals (Dean et al., 2013). However, eukaryotic algae that can tolerate AMD conditions may be an alternative bioremediation agent (Abinandan et al., 2018; Das et al., 2009). Novel extremophile algal strains that show high acid and metal tolerance, and metal bioaccumulation traits are therefore needed for such applications. In addition, extremophile algae may have other biotechnological applications, such as a source of novel high-value chemicals including nutritional vitamins and anti-oxidants, food additives, and biofuels (Varshney et al., 2015).

The aim of this study was to identify eukaryotes, especially extremophile microalgae, in a series of standing waters affected by AMD with the intention to characterise a strain of microalgae for evidence of AMD adaptation. Following a screen of eukaryotic biota within nine Cu-rich AMD ponds, an extremophile chlorophyte microalgal strain identified as *C. acidophila* was examined in detail due to its abundance and ubiquity across the site and its high tolerance to acidity and dissolved metal concentrations, especially to Cu.

## **2. Materials and Methods**

### **2.1. Study site**

The site for this study is Parys Mountain, a disused Cu mine, in Anglesey North Wales, UK. The site has been mined for Cu from the Bronze Age, until mining activities ceased in the early 1900s (Dean et al., 2013). The area consists of large amounts of exposed spoil, with large pits and depressions that have filled with rainwater, and now retain large amounts of metal-rich and acidic water (Fig. 1). In addition, precipitation ponds and lagoons were constructed at the base of Parys Mountain, which were built in order to extract metals from the water as part of the mining process, and also contain large volumes of AMD polluted water (Younger and Potter, 2012). All the ponds are situated at close proximity within a similar geology with the rocks naturally rich in Cu, Pb and Zn. It is the only known example of Kuroko type volcanogenic massive sulfides in the UK, though the geology has been

disturbed by many millennia of underground and surface mining activities (Younger and Potter, 2012).

Of the various mining ponds and lagoons at the Parys Mountain site, nine ponds were examined (Fig. 1). Ponds 1 – 4 are located at an elevated position on the spoil outcrop, with one of these (Pond 4) on the side of a steep incline of the now drained large opencast (Fig. 1). The remaining five larger ponds (Ponds 5 – 9) are the precipitation ponds and lagoons at the base of the Parys Mountain outcrop, each adjacent to agricultural land. Ponds 1 – 4 and 9 are shallow and less than 1 m depth and subject to rapid variation in depth due to seasonal evaporation and rainfall. Ponds 5 – 8 are deeper and are typically 2 m in depth. All ponds showed little variation in depth across each pond. Ponds 1, 2 and 4 had no vegetation in or surrounding them, whereas the other ponds had surrounding vegetation and marginal wetland plants.

## *2.2. Field site sampling*

Sampling at the nine AMD ponds was carried out in 2013 to 2015, including a spring (February and March), summer (June) and autumn (October) sampling regime in 2015. Water chemistry samples were taken in triplicate at each pond on each sampling occasion and were taken at approximately 15 cm depth 1 – 2 m from the edge of the pond. Water pH, conductivity, temperature and dissolved oxygen were measured using a YSI 556 probe (Xylem Analytics). For analysis of dissolved water chemistry metals 250 mL of pond water was filtered through a 0.45 µm cellulose acetate filter, as described previously (August et al., 2002; Boulton et al., 1994), and a 50 mL volume was retained for the analysis of dissolved nutrients ( $\text{PO}_3\text{-P}$ ,  $\text{NO}_3\text{-N}$ ,  $\text{NH}_4\text{-N}$ ). This dissolved fraction will also include some colloidal metals (Florence et al., 2016). A further 50 mL volume was acidified to 1% (v/v) nitric acid final concentration for the analysis of dissolved metals (Al, As, Cd, Cu, Fe, Mn, Pb, S, Zn). The filter paper was retained, and used for the determination of particulate metals.

For the analysis of algae 250 mL of unfiltered water was collected and aliquots were preserved with Lugol's iodine for enumeration of algae cells and further unfiltered water

sample was taken in a sterile container for the isolation and identification of algae. Algal samples were taken from all ponds, including samples taken from where algal biofilms were observed (Ponds 5 and 6). For chlorophyll-a measurement, 250 mL of pond water was filtered onto a GF/C filter paper.

Sediment samples were also taken to a depth of approximately 2 cm depth for determination of acid-extractable metals. For invertebrate sampling, sediment samples were taken from the littoral, approximately 1 m from the pond edge and 20 – 50 cm depth depending on the pond and sieved to remove organisms. This was followed by a 3 min sweep using a hand-held net with a mesh size of 1 mm, and a 3 min examination under large stones. Ethanol (70%) was added to preserve the invertebrate biota for identification to family level using standard keys (Greenhaigh and Ovenden, 2007; Quigley, 1977).

### *2.3 Nutrient, chlorophyll and metal analysis*

Dissolved  $\text{PO}_3\text{-P}$ ,  $\text{NO}_3\text{-N}$ , and  $\text{NH}_4\text{-N}$  were measured from the 0.45  $\mu\text{m}$  filtered water samples using a Skalar Sans Plus autoanalyser. Chlorophyll-a concentrations were determined by absorbance spectroscopy following extraction in 96% (v/v) ethanol, as described previously (Dean et al., 2010). Dissolved metals were determined by inductively coupled plasma atomic emission spectroscopy (ICP-AES) using a Perkin-Elmer Optima 5300, exactly as described previously (Dean et al., 2013). Acid extractable sediment metals and suspended particulate metals were determined by acid digestion of 0.1 g of 250  $\mu\text{m}$  sieved dried sediments and digestion of the pond water filter papers, respectively.

Sediments and filter papers were digested in 5 mL ultrapure-grade nitric acid at 70°C for 4 h, diluted to 2% (v/v) nitric acid, and the metals measured by ICP-AES. Certified Reference Standard TM25.5 was used for all ICP-AES analyses. All samples were calibrated using a matrix-matched serial dilution of Specpure multi-element plasma standard solution 4 (Alfa Aesar) set by linear regression. Only results with a relative standard deviation < 20% were considered.



## 2.4. Microalgae cultivation and analysis

Microalgae was visually identified to genus level and enumerated from the Lugol's iodine preserved water samples by light microscopy using a Sedgewick-Rafter cell counting slide and a morphological taxonomy key (John et al., 2002). Isolation of individual algae strains was carried out by incubating serial dilutions of water samples on modified acid medium (MAM) agar plates at 22 °C with a 16-h light:8-h dark light regime and a photon flux of approximately 150  $\mu\text{mol m}^{-2} \text{s}^{-1}$ . MAM is a defined inorganic medium developed previously (Olaveson and Stokes, 1989), and used here as described by the Canadian Phycological Culture Centre, University of Waterloo, Canada containing 0.5 g L<sup>-1</sup> (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 0.01 g L<sup>-1</sup> CaCl<sub>2</sub>·2H<sub>2</sub>O, 0.5 g L<sup>-1</sup> MgSO<sub>4</sub>·7H<sub>2</sub>O, 0.3 g L<sup>-1</sup> KH<sub>2</sub>PO<sub>4</sub>, 0.03 g L<sup>-1</sup> NaCl, 0.01 g L<sup>-1</sup> Na<sub>2</sub>EDTA·2H<sub>2</sub>O, 4.98 mg L<sup>-1</sup> FeSO<sub>4</sub>·7H<sub>2</sub>O, 2.86 mg L<sup>-1</sup> H<sub>3</sub>BO<sub>3</sub>, 1.81 mg L<sup>-1</sup> MnCl<sub>2</sub>·4H<sub>2</sub>O, 0.22 mg L<sup>-1</sup> ZnSO<sub>4</sub>·7H<sub>2</sub>O, 0.39 mg L<sup>-1</sup> NaMoO<sub>4</sub>·2H<sub>2</sub>O, 79  $\mu\text{g L}^{-1}$  CuSO<sub>4</sub>·5H<sub>2</sub>O, 49.4  $\mu\text{g L}^{-1}$  Co(NO<sub>3</sub>)<sub>2</sub>·6H<sub>2</sub>O, 1  $\mu\text{g L}^{-1}$  vitamin B12, 1  $\mu\text{g L}^{-1}$  biotin, and 0.2 mg L<sup>-1</sup> thiamine-HCl, adjusted to pH 3.0. Individual colonies were then extracted and grown in liquid MAM.

For identification of isolates using 18S rRNA gene amplification, DNA was extracted using an UltraClean Tissue & Cells DNA isolation kit (Mo Bio) and 18S rRNA gene amplicon sequences amplified using universal primers EukF and EukR (DeLong, 1992). DNA amplicons were purified (Qiagen PCR Purification kit) before sequencing (to give ~1 kb sequence reads) by GATC-Biotech, with subsequent sequence analysis performed by BLAST, using the NCBI GenBank database (Table S1). The species identification of putative *C. acidophila* strain was further confirmed by partial length 18S rDNA gene amplicon sequencing, using PCR primers (18SFOR 5'-WAC CTG GTT GAT CCT GCC AGT-3', and 18SREV 5'-GAT CCT TCY GCA GGT TCA CCT AC-3') and PCR conditions as described (Huss et al., 1999), and sequenced as described above but with sequence reads of ~1.7 kb size. Phylogenetic analysis was then performed essentially as described previously (Osundeko et al., 2013). 18S rRNA nucleotide sequences of selected unicellular Chlorophyta microalgae of the *Chlamydomonas moewusii* clade were obtained from GenBank and sequences were aligned using ClustalW. The phylogenetic tree was generated using the

maximum likelihood method using RAxML-GUI and the GTR-GAMMA model (Stamatakis, 2006). Confidence in the tree was assessed using the thorough bootstrap method by performing 10 runs of 100 replications.

An isolate of *C. acidophila* from Pond 1 (named PM01) was cultivated to quantify its tolerance to high metal concentrations and pH ranges alongside two strains of *C. acidophila* previously isolated from an acidic pond near Fratiskovy Lazne, Czech Republic (CCAP 11/136) (Fott and McCarthy, 1964), and an acidic mining Lake 111 in eastern Germany (CCAP 11/137) (Gerloff-Elias et al., 2005). All cultures were maintained in MAM pH 3.0. Cultures were serially inoculated in fresh media and were grown in batch culture conditions on an orbital shaker at 120 rpm at 22 °C with a 16-h light:8-h dark light regime and a photon flux of approximately 150  $\mu\text{mol m}^{-2} \text{s}^{-1}$ . For metal exposure treatments, various concentrations of metals as chloride salts were added to liquid MAM pH 3.0 or solid MAM pH 3.0 agar plates. For pH range treatments, MAM was adjusted and buffered to the desired pH (from pH 1.0 to 7.0) as described previously (Gerloff-Elias et al., 2006). Essentially the medium was buffered through the presence of Fe in the medium for pH 1.0 to 3.0, with 10 mM citric acid for pH 4.0 to 5.0, and with 10 mM HEPES for pH 6.0 to 7.0. The pH could be maintained to within ~0.5 pH unit during the growth period. Starting cell densities were normalised by optical density measurement at 680 nm ( $\text{OD}_{680\text{nm}}$ ). Algae growth was determined by cell number, and in some instances by  $\text{OD}_{680\text{nm}}$ , total chlorophyll or growth rate measurement, exactly as described previously (Osundeko et al., 2013). Algae samples were prepared for metal content measurement by ICP-AES, as described previously (Webster et al., 2011). Cells were EDTA washed to remove externally bound metals, as validated elsewhere for Cd, Cu, Pb and Zn (Hassler et al., 2004). Cells were centrifuged for 10 min at 3000 g, followed by resuspension of the cell pellet in 10 mL of 1 mM EDTA for 5 min, then re-centrifuged and washed with a 15 mL volume of Milli-Q water. After further centrifugation, cell pellets were oven-dried at 60 °C for 24 h and then digested in 0.5 mL of ultrapure concentrated nitric acid at 70 °C for 3 h. Samples were diluted in Milli-Q water to 2% (v/v) concentration of acid and analysed by ICP-AES as described above. Cells of *C.*

*acidophila* were measured (width and length) using an eyepiece graticule, and cell volume was calculated according the prolate spheroid formula (Hillebrand et al., 1999). The cell volume measurements were used to calculate the internal (cellular) concentration of metal (determined from EDTA-washed cells) on a volume basis. This value was then divided by the external (MAM or pond water) concentration of metal in order to calculate concentration factor ( $K_{conc}$ ) values.

## 2.5. Fourier transform-infrared (FT-IR) spectroscopy

Each algal strain was grown in liquid MAM (pH 3.0), with Cu concentrations of 0, 6.5, 13 and 130 mg L<sup>-1</sup> for 14 days, at which point cultures had an OD<sub>750nm</sub> of between 0.3 and 0.5. Cultures were normalised to an OD<sub>750nm</sub> of 0.3 and 10 mL was centrifuged at 1800 *g* for 5 min. The supernatant was removed and the cells washed in 1.5 mL of 0.9% (v/v) NaCl, then centrifuged and washed again before final resuspension of the cell pellet in 1 mL of 0.9% (v/v) NaCl before 20 µL of this suspension was deposited onto a 96-well silicon microplate. The samples were then oven dried at 40°C for 1 h, and an additional 20 µL sample was added and the samples once again dried at 40°C for 1 h. The plate was placed in a HTS-XT high-throughput microplate extension and FT-IR spectra collected using an Equinox 55 FT-IR spectrometer (Bruker Corporation), equipped with a deuterated triglycerine sulphate detector. Spectra were collected over the wavenumber range 4000-600 cm<sup>-1</sup>. Each sample was analysed as nine technical replicates. Spectra were pre-processed using extended multiplicative signal correction (Martens and Stark, 1991) prior to multivariate analysis. Band assignments were determined as described previously (Driver et al., 2015).

## 2.6. Modelling and statistical analysis

Metal speciation modelling was performed using Visual Minteq version 3.0 (Gustafsson, 2010). All data was statistically analyzed by one-way ANOVA using Tukey post-hoc test performed using Prism v.6.04 (GraphPad). Principal component analysis (PCA) of environmental data was performed using PRIMER v.6 (Primer-E) and plotted using XLSTAT

(Addinsoft), while PCA of FT-IR spectra was performed using MATLAB version R2016a. All environmental data (except for pH values) were natural log transformed for PCA and linear regression analysis.

### 3. Results

#### 3.1. Chemical characteristics of extremely acidic metal-rich mining ponds

The nine ponds were situated in close proximity (Fig. 1) but differed in water chemistry (Fig. 2). In all ponds, concentrations of suspended particulate metals were present at much lower concentrations compared to dissolved metals (Fig. S1). None of the metal or nutrient concentrations in the ponds differed significantly across the spring, summer and autumn samples. Ponds 1, 2 and 4 were situated on top of the Parys Mountain site in depressions of the spoil waste. These ponds were characterised by very low pH ( $< 2.0$ ) and high conductivity values (Fig. 2A). Pond 1 in particular was highly acidic (mean pH of 1.6) and had a very high conductivity (mean 10.03 mS), which is predominantly controlled by the high dissolved S concentration, as well as high concentrations of dissolved Fe, Al, Cd, Cu and Zn (Fig. 2B, C). In addition, Pond 1 also had a very high  $\text{PO}_4\text{-P}$  concentration (Fig. 2D) and the highest dissolved As concentration (Fig. S1). Ponds 2 and 4 also had high dissolved metal and S concentrations but differed from Pond 1 due to lower conductivity ( $\sim 3$  mS) and much lower  $\text{PO}_4\text{-P}$ . Pond 3 was also situated on the top of the mine site, however, this pond stood out as the least polluted, with a weakly acidic pH (mean pH 5.3), and low conductivity and dissolved metals and S (Fig. 2A; Fig. S1). The precipitation ponds and lagoons (Ponds 5 – 9) were also highly acidic but had lower conductivity due to much lower dissolved metal concentrations, with the exception of dissolved Mn (Fig. S1). In most ponds, ammonium ( $\text{NH}_4\text{-N}$ ) was the dominant form of N rather than nitrate ( $\text{NO}_3\text{-N}$ ), as is expected in acidic waters where nitrification rates are low (Baffico et al., 2004). Pond 6 had unusually high  $\text{NO}_3\text{-N}$  (Fig. 2D) possibly because the pond is directly adjacent to fertilised farmland. Chlorophyll-a was detectable in all ponds, indicating the presence of photosynthetic

microorganisms, but Pond 1 also differed from the other ponds with respect to having significantly higher concentration of chlorophyll-a, which increased substantially during the year as the water temperature and insolation increased (Fig. 2E).

The differences in water chemistry between the ponds situated within the mine spoil waste (Ponds 1, 2 and 4), the lagoons and precipitation ponds (Ponds 5 – 9), and the less polluted Pond 3, are reflected in the PCA plot (Fig. 3). In particular, Pond 1 and Pond 3 were distinguished from each other and from the other ponds through difference in pH. Furthermore, the high sediment Al and Mn concentrations explained the differentiation of Pond 3, likely a consequence of the high pH of this water resulting in the precipitation of dissolved Al. Pond 1 was situated within the PC space particularly on the basis of high chlorophyll-a, PO<sub>4</sub>-P and dissolved As concentration. It is therefore apparent that many of these ponds are highly toxic environments, particularly Pond 1 with its very high acidity and high concentration of toxic dissolved trace metals including As, Al, Cd, and Cu.

### 3.2. *Taxa diversity*

Overall taxa diversity, as determined by number of different taxonomic families, was lowest in the highly acidic, high conductivity Pond 1 (with two taxa) and also low in the Ponds 2 and 4, which are also situated among open mine spoil. The highest number of taxa were found in the low conductivity, mildly acidic (pH 5.3) Pond 3, with 15 taxa, while the lagoons/precipitation ponds had intermediate taxa diversity (7-10 families). Linear regression analyses of the relationships between taxa diversity and water chemistry (Fig. S2) showed a significant positive correlation between increasing pH values and increasing taxa number ( $R^2 = 0.59$ ;  $p = 0.03$ ) but also a very strong negative correlation between decreasing conductivity and increasing taxa number ( $R^2 = 0.85$ ;  $p = 0.001$ ), indicative of the dissolved metal concentration. In particular, there was significant correlation between dissolved As, Cd, Cu, Fe and S and taxa number, as well as between particulate Fe and S with taxa number, but no significant correlation on the basis of sediment metal concentration (Fig. S2).

Invertebrates were observed in all ponds, though only Chironomidae was identified in all, although there were only one or two individuals recorded from Pond 2 and 4 (Fig. 5A). In Ponds 1 and 4, this was the only invertebrate taxa present, indicating that the conditions in these ponds are not suited for a diverse invertebrate community, but are able to support a few species that have adapted to the extreme AMD conditions. Other invertebrate taxa that were abundant are Corixidae, present in five of the ponds, and Sialidae, present in four of the ponds, but both absent in the three most acidic ponds (Fig. 4). The observed biota in Pond 3 included 14 invertebrate families, including many that are pollution-sensitive. Ponds 5 – 9 contained between 5 and 7 invertebrate taxa. All ponds were devoid of macrophytes within the open water.

A number of distinct eukaryotic microorganisms were isolated from the open water including microalgal, fungal and protozoan species (Table S1). The highest eukaryotic diversity was in Ponds 2, 5 and 6 (Fig. 4). In Ponds 5 and 6 and their adjacent channels much of the microorganisms were observed to be associated in biofilms. These included a strain which showed 99% identity based on the 18S rRNA sequence to *Euglena mutabilis* (Table S1), a well-known acid-tolerant species that has been previously observed in AMD environments, including coal mine waste sites (Brake et al., 2001), and including at the adit draining from the Parys Mountain mine (Ñancucheo and Johnson, 2012). In addition, a non-motile chlorophyte (likely to be *Koliella corcontica*; 96% identity) that has been previously observed in mildly acidic lakes (Vrba et al., 2003), and a diatom (likely to be *Eunotia naegeli*; 98% identity) was found associated with biofilm adjacent to Pond 6. However, diatoms were not found in open water in any of the Parys Mountain ponds. Ponds 1, 3 and 7 had the lowest number of eukaryotic microorganisms, with just one chlorophyte algae taxa identified in each pond (Fig. 4). There was substantial heterogeneity between the ponds, with all microbial taxa present in just one pond sample with the exception of the *Euglena* sp. present in two ponds, and a *Chlamydomonas* sp. identified in all ponds (100% abundance). Sequencing of the ~1 kb 18S rRNA gene amplicon from the *Chlamydomonas* sp. showed highest sequence identity (99%) to a strain annotated as *C. acidophila* (CCAP 11/134)

(Table S1). Quantification of cell density of this strain found highest density in Pond 1 (Fig. 5A), which correlated significantly with the highest pond PO<sub>4</sub>-P concentration ( $R^2 = 0.60$ ;  $p = 0.02$ ), but cell density was also lowest in the most alkaline pond water where there was the lowest dissolved metal concentrations, likely explaining the positive correlation between cell density and conductivity or dissolved metal concentrations, such as Cu (Fig. 5B).

### 3.3. Phylogeny of a *C. acidophila* strain

*C. acidophila* was found across the Parys Mountain site in all nine ponds and was highly abundant in the most acidic and metal rich Pond 1. Therefore this strain was studied further in more detail. The isolate of *C. acidophila* from Pond 1 (named PM01) was examined by longer (1.7 kb) 18S rRNA gene read sequence and phylogenetic analysis (Fig. 6). PM01 18S rRNA sequence was identical apart from one nucleotide within this 1.7 kb region to sequences from three strains annotated as *C. acidophila* (Gerloff-Elias et al., 2005): CCAP 11/134 (an isolate from Argentina), CCAP 11/136 (an isolate from Czech Republic) and CCAP 11/137 (an isolate from Germany). PM01 was also identical within this 18S region to an unidentified strain (Rt1n1) originally isolated from the acidic Rio Tinto river in Spain (Amaral Zettler et al., 2002). The CCAP 11/136 strain (Fott and McCarthy, 1964), previously regarded as an authentic strain of *C. acidophila* (Gerloff-Elias et al., 2005), is also deposited in a different culture collection as strain UTCC 354 (Pollio et al., 2005), but this has distinct 18S rDNA sequence (Fig. 6), indicating that CCAP 11/136 and UTCC 354 are not in fact identical. It was previously argued that the UTCC 354 strain assigned as *C. acidophila* may be more appropriately assigned as *Chlamydomonas pitschmannii*, another highly acidotolerant species (Pollio et al., 2005). Other strains recorded as *C. acidophila* including UTCC 121 (an isolate from Canada) (Twiss, 1990) and OU 030/a (an isolate from Japan) (Nishikawa and Tominaga, 2001), were also distinct from PM01 and CCAP 11/136 but grouped more closely to the UTCC 354 strain and *C. pitschmannii* (Fig. 6). Although strain OU 030/a has also been previously regarded as an authentic strain of *C. acidophila* (Nishikawa and Tominaga, 2001; Pollio et al., 2005), here we refer to the CCAP 11/136 and

11/137 strains as *C. acidophila* species, in line with previous analysis (Gerloff-Elias et al., 2005), and thus the PM01 strain is further referred to as a strain of this species.

#### 3.4. AMD tolerance by *C. acidophila* PM01

To examine the relationship between pH and *C. acidophila* further, strain PM01 from Pond 1 was grown in MAM artificial pond water at a range of pH values from pH 1.0 to 7.0. PM01 exhibited a very broad pH tolerance range, with optimal growth at pH 3.0 – 5.0 (mean growth rate ranging between 0.158 – 0.175 d<sup>-1</sup>; no significant difference between treatments;  $p > 0.05$ ), and with the highest cell density after 25 d obtained in pH 3.0 conditions. Strong growth was still observed in pH 7.0 (mean growth rate 0.153 d<sup>-1</sup>) and pH 2.0 (mean growth rate 0.157 d<sup>-1</sup>) conditions, but with a significant reduction ( $p < 0.05$ ) in growth after 25 d in pH 2.0 and pH 7.0 by 25% and 37%, respectively compared to pH 3.0 MAM. and could grow at pH 1.0 after an extended lag phase of 13 d (growth rate 0.018 d<sup>-1</sup>), but with a significant reduction in growth by 73% after 25 days compared to pH 3.0 MAM. The pH characteristics of the ponds may explain in part the microalgae biodiversity and cell density profiles, and as described above, there was a significant negative correlation ( $R^2 = 0.57$ ;  $p = 0.02$ ) between pH and *C. acidophila* cell density (Fig. 5B).

Samples of PM01 taken directly from Pond 1 showed high concentration of absorbed and internalised metals, as determined by measurement of EDTA-washed cells, to remove externally cell wall-bound metals. There was a high concentration of Cu and Zn accumulated almost entirely within the cell (no significant difference between EDTA washed versus unwashed cells) (Table 1). Relative to mean Pond 1 dissolved Cu concentration of 58.6 mg L<sup>-1</sup>, the internal cellular Cu concentration in pond 1 cells was 84.3 fg cell<sup>-1</sup> (equivalent to 185.5 mg L<sup>-1</sup>). The mean concentration of dissolved Zn in Pond 1 was 37.9 mg L<sup>-1</sup>, while the internal cellular Zn concentration was 115.0 fg cell<sup>-1</sup> (253.0 mg L<sup>-1</sup>). This gives concentration factor ( $K_{conc}$ ) values of 3.2 for Cu and 6.7 for Zn. Substantial bioconcentration was also observed for Pb and Cd. Relative to a mean Pb concentration in Pond 1 of 0.8 mg L<sup>-1</sup>, the accumulation and bioconcentration of Pb by the strain was particularly high, with a cellular



concentration of 146.3 fg cell<sup>-1</sup> (321.8 mg L<sup>-1</sup>) giving a  $K_{conc}$  value of 421.8. Cd was also accumulated almost entirely within the cell to a concentration of 2.8 fg cell<sup>-1</sup> (6.2 mg L<sup>-1</sup>), and relative to pond water concentration of 0.3 mg L<sup>-1</sup> gives a  $K_{conc}$  value of 24.5. In contrast, only 56% of accumulated Fe was taken up into the cell (Fig. S3).

Metal tolerance and accumulation by *C. acidophila* PM01 was further examined in an artificial growth medium to assess the tolerance range of three of the trace metals found within the Parys Mountain ponds. PM01 was grown in increasing concentrations of Cd, Cu and Zn in MAM at pH 3.0. For cells grown under controlled conditions, the maximum cell density achieved and the total chlorophyll concentration per cell (as a measure of cell physiological status) was higher than that observed for the cells analysed *in situ* (in Pond 1) (Table 1). The PM01 strain displayed tolerance to high concentrations of these three metals. To allow comparison with Pond 1 water concentrations, the free ionic Cd<sup>2+</sup>, Cu<sup>2+</sup> and Zn<sup>2+</sup> concentrations were calculated using the Visual Minteq speciation model. PM01 could tolerate up to 2.6 mg L<sup>-1</sup> Cd<sup>2+</sup>, which was 16-times higher than present in Pond 1 water, with no significant inhibition of growth rate, maximum cell density or chlorophyll content when compared to no added Cd<sup>2+</sup> (Table 1). However, at a concentration of 6.0 mg L<sup>-1</sup> Cd<sup>2+</sup> all three parameters (growth rate, cell density, chlorophyll content) were significantly reduced ( $p < 0.05$ ). PM01 was highly tolerant to Cu, and none of the concentrations up to 78.3 mg L<sup>-1</sup> Cu<sup>2+</sup> treatment significantly inhibited growth or chlorophyll-a concentration, the apparent reduction in cell density was not significant. Substantial Zn tolerance was also observed, with no significant inhibition to growth rate at 855.4 mg L<sup>-1</sup> Zn<sup>2+</sup>, which was nearly 30-times higher than the Zn<sup>2+</sup> concentration in Pond 1, but at 1760.8 mg L<sup>-1</sup> Zn<sup>2+</sup> there was a significant ( $p < 0.05$ ) reduction in maximum cell density, although growth rate and chlorophyll concentration was not significantly inhibited. The strain was still growing in 3002.8 mg L<sup>-1</sup> Zn<sup>2+</sup> despite cell density being inhibited by 94%, with also a significant ( $p < 0.05$ ) reduction in growth rate and chlorophyll concentration (Table 1).

Accumulation of Cd, Cu and Zn was also quantified in the metal-treated PM01 strain after 25 d growth in MAM pH 3.0 (Table 1). There was a concentration-dependent increase

in cellular accumulation of Cu and Cd with no significant difference between the values with or without EDTA washing, indicating that almost all of the Cu and Cd was taken up within the cell. In contrast, there were significant ( $p < 0.05$ ) differences in Zn concentration following EDTA washing, indicating that a smaller proportion of Zn was internalised (Table 1). As the Zn concentration in the medium increased, the relative concentration within the cell decreased, suggesting that Zn transport into the cell saturated at higher concentrations. Overall, the characteristics of metal accumulation in artificial media were broadly similar to those observed in the pond.

### 3.5. Metabolic adaptation to Cu tolerance by *C. acidophila* PM01

It was unknown whether the metal tolerance properties of *C. acidophila* differ between strains isolated from AMD sites with differing water chemistry. To begin to assess this, the tolerance of PM01 to a range of metals (Al, Cd, Cu, Fe, Mn and Zn) was compared to two other strains of *C. acidophila* (CCAP 11/136 and CCAP 11/137) that had previously been isolated from different field sites. Strains CCAP 11/136 and CCAP 11/137 were validated by 18S rDNA sequencing and phylogenetic analysis as *C. acidophila* species (Fig. 6). The CCAP 11/136 strain was originally isolated from a highly acidic (~pH 1.0) humic acid-rich peat water environment but the metal characteristics of the site were not described (Fott and McCarthy, 1964). In contrast, strain CCAP 11/137 originated from German mining Lake 111 at pH 2.6, very low total P ( $8 \mu\text{g L}^{-1}$ ) and fairly high levels of Zn ( $0.75 \text{ mg L}^{-1}$ ) (Spijkerman et al., 2007a).

For Al, Cd and Mn there was no growth difference between the three strains. However, Fe treatment substantially inhibited growth of CCAP 11/136, while Zn treatment slightly inhibited growth of PM01 and CCAP 11/137 (Fig. S4). However, there was a very clear-cut difference in growth between the three strains following Cu exposure (Fig. 7). On solid media containing  $130 \text{ mg L}^{-1}$  Cu, the CCAP 11/137 strain was unable to grow, and the CCAP 11/136 strain grew very weakly in contrast to strong growth by the PM01 strain (Fig. 7A). In liquid media with addition of  $0.0 \text{ mg L}^{-1}$ ,  $6.5 \text{ mg L}^{-1}$  and  $3 \text{ mg L}^{-1}$  growth rate and cell

density after 14 days was identical between all three strains. However, with 130 mg L<sup>-1</sup> Cu addition, cell growth was unchanged for PM01 but both CCAP 11/136 and 11/137 strains were barely able to grow (Fig. 7B).

To examine whether there were any macromolecular changes within the strains in response to the increasing copper treatment, and to examine whether the different strains could be distinguished on the basis of their metabolic 'fingerprint', the FT-IR spectroscopy technique was used. FT-IR spectra were collected for replicates of each strain cultivated in the absence of added Cu or with the addition of 6.5 and 13 mg L<sup>-1</sup> Cu (Fig. S5). In addition, PM01 was tested at the 130 mg L<sup>-1</sup> concentration that inhibited growth of 11/136 and 11/137, thereby preventing FT-IR spectroscopy analysis of these strains at the higher Cu concentration. PCA of all FT-IR spectra showed that the CCAP 11/137 strain samples cluster separately from CCAP 11/136 and PM01 in all treatments, with the close clustering of spectra indicating that Cu addition did not significantly alter the metabolic fingerprint of CCAP 11/137 (Fig. 8A). Likewise Cu addition did not significantly alter the metabolic fingerprint of the CCAP 11/136 strain, with all samples clustering with the PM01 control samples. However, there was a clear difference in the FT-IR spectra-derived metabolic profile of PM01 samples following addition of increasing Cu concentrations, with the sample position within the PCA plot changing on the basis of both PC1 and PC2. At the highest (130 mg L<sup>-1</sup>) Cu concentration all replicate PM01 samples are distinct from the control samples (Fig. 8A). The PC loading plots (Fig. 8B) indicate that along PC1 the spectral changes are based predominantly on increased abundance of amide peaks at 1655 and 1545 cm<sup>-1</sup>, as well as a decrease in one of the carbohydrate peaks at approximately 1036 cm<sup>-1</sup>.

#### 4. Discussion

This study has demonstrated that while AMD substantially impacts biodiversity in aquatic environments, there is still substantial taxa abundance in these extreme locations. The scarcity of invertebrates across the ponds clearly indicates the severity of pollution at this

abandoned mine site. With the exception of the near-neutral pH Pond 3, invertebrate diversity was very poor in all ponds. Taxa diversity overall, including invertebrate diversity in particular, was strongly correlated with pH and with dissolved metal concentration, which was in line with expectations and previous studies (Courtney and Clements, 2002; Malmqvist and Hoffsten, 1999). Nevertheless, some invertebrate species can adapt to extreme AMD conditions (De Bisthoven et al., 2005; Deneke, 2000), and can be used as indicators of mine-waste pollution (Gray and Delaney, 2008). Here Chironomidae were observed in all ponds, including the very acidic Pond 1 and 4, and highly abundant in Pond 9, probably because this pond is fairly shallow and has fine sediment in contrast to the other ponds. Certain Chironomidae species are able to survive in highly acidic waters, such as the acid-tolerant *Chironomus acidophilus* that has been found nearby in the highly acidic (pH 2.4) and metal-rich Afon Goch river draining Parys Mountain (Michailova et al., 2009).

While Chironomidae were present in all ponds so there was also the consistent presence of chlorophyte microalgae, and in particular a strain confirmed as *C. acidophila*. The widespread occurrence of *C. acidophila* at Parys Mountain reflects the extremely acidic and metal-rich pond water and is consistent with other acidic and metal-rich sites (Fott and McCarthy, 1964; Gerloff-Elias et al., 2005; Hargreaves et al., 1975; Twiss, 1990). The widely differing *C. acidophila* abundance between ponds may reflect nutrient availability. AMD environments typically have very low productivity, due in part to low nutrients such as PO<sub>4</sub>-P (Spijkerman et al., 2007a; Spijkerman et al., 2007b), and in this study nutrients, and in particular PO<sub>4</sub>-P, were at low concentration in all but Pond 1. The low cell density of this species in Ponds 2 – 9, where PO<sub>4</sub>-P concentrations were low suggests growth limitation due to low PO<sub>4</sub>-P availability. In contrast, Pond 1 had very high PO<sub>4</sub>-P and a high concentration of *C. acidophila* cells. Other studies have shown high PO<sub>4</sub>-P levels associated with acidic lakes in areas with abundant PO<sub>4</sub>-P-containing FeS minerals (Spijkerman, 2008); however, as Pond 2 is within 5 m of Pond 1 and has the same mineralogy but significantly lower PO<sub>4</sub>-P levels, the high concentration of PO<sub>4</sub>-P in Pond 1 is unlikely to be due to minerology of the surrounding area. It may be that the high PO<sub>4</sub>-P concentration is due to a

eutrophication event, as there is evidence of construction waste deposition that is likely to explain the PO<sub>4</sub>-P entry into the pond.

The high microalgae productivity of Pond 1 is due almost exclusively to *C. acidophila* abundance. Acidophilic microalgae present in the other ponds, such as *K. corcontica*, *E. naegelii* and *E. mutabilis*, were absent from Pond 1. *E. naegelii* and other acid-tolerant diatoms are increasingly used as indicator species for acid polluted environments (Zalack et al., 2010) and have been previously found in surface sediment from one of the rivers draining from the Parys Mountain site (Dean et al., 2013), as well as in acid pit lakes with equivalent water chemistry (Geller, 2013). However, their apparent low abundance in these ponds might partly be due to the sampling regime used here from the near-surface open water rather than from the bottom of the ponds. *E. mutabilis* was observed in two of the nine ponds (Ponds 5 and 6), which are both ~pH 3.0, a pH range that has been shown to be preferential for this organism (Brake et al., 2001). Furthermore, other acidophilic microalgae such as *Ochromonas* sp. and *Dunaliella* sp., that are widely abundant in many acidic sites (Aguilera et al., 2006; Nixdorf et al., 1998), were not identified in Pond 1, or in any other Parys Mountain ponds. That *C. acidophila* was the only microalgal species identified in Pond 1 is likely to be due to the extreme conditions that restricts microalgal diversity. One of these factors appears to be water pH, and the *C. acidophila* strain studied here was particularly acid tolerant. In waters with pH > 3.0, it has been previously observed that algal biodiversity is generally higher (Smucker et al., 2014), yet the least acidic pond studied here, Pond 3 (pH 5.3), still had very low algal biodiversity and very low *C. acidophila* abundance, suggesting that other factors are also important, such as nutrient availability or water chemistry.

Although the dissolved Fe, Zn, Cu and Al concentrations in Pond 1 did not exceed the very high concentrations seen in some AMD lakes and rivers such as those of the Iberian Pyrite Belt (Sánchez España et al., 2008), they did exceed those seen in mine pit lakes in Germany, Poland, Australia and USA (Geller, 2013). The metal concentrations of the shallow Pond 1 can thus be considered high. The strain of *C. acidophila* isolated from Pond 1 is therefore not just extremely acid-tolerant, but can also tolerate high metal

concentrations. The PM01 strain showed substantial tolerance to Cd, Cu, and Zn. A comparison of PM01 to other genera of acid tolerant algae shows the Zn tolerance of PM01 was higher than that reported for *Chlorella protothecoides* var. *acidicola* isolated from AMD sites in a Spanish mine and *E. mutabilis* isolated from the adit flowing from Parys Mountain. However, both of these strains showed a higher Cu tolerance than PM01 (Ñancucheo and Johnson, 2012).

Other studies looking at metal tolerance in *C. acidophila* are fairly scarce, though a study looking at a putative strain of *C. acidophila* OU 030/a isolated from a volcanic acid lake also showed high tolerance to Cd, Cu and Zn in metal-rich minimal media (at pH 4.0) when compared to other algal species (Nishikawa and Tominaga, 2001). Furthermore, a strain of *C. acidophila* RT46 isolated from Río Tinto in Spain, showed unaffected photosynthetic activity in response to 0.5 mM Cu (~32 mg L<sup>-1</sup> Cu) exposure (Olsson et al., 2015). A putative *C. acidophila* strain (UTCC 121) isolated from Cu contaminated soil was previously shown to tolerate up to 100 mg L<sup>-1</sup> Cu. However, in contrast, a laboratory strain of *C. acidophila* (CCAP 11/96) was Cu sensitive, as was the non-acidophilic freshwater alga *Chlamydomonas reinhardtii* (Twiss, 1990), indicating that the *Chlamydomonas* genera is not intrinsically Cu tolerant. In this study we demonstrate that the *C. acidophila* strain isolated in this study (PM01) has higher Cu tolerance than strains (CCAP 11/136 and 11/137) from AMD field sites with less Cu pollution. This shows that the PM01 strain has adapted to the high dissolved Cu concentrations of Pond 1, rather than this species having innate Cu tolerance properties.

FT-IR spectroscopy analysis demonstrates that the tolerance of PM01 to copper is partly due to its ability to modulate its metabolism in response to increasing Cu exposure, as indicated by a dose-dependent change in spectra characteristics, which does not occur with the other *C. acidophila* strains. Examination of the FT-IR spectra indicates that this metabolic adaptation is predominantly due to protein increase and potential modification of protein secondary structure, as shown by significant increase in amide I peak height associated with C=O stretching, and in amide II peak height associated with N-H bending and C-N stretching

(Giordano et al., 2001). A previous study using FT-IR spectroscopy to examine sensitivity and subsequent acclimation of microalgae to wastewater treatment, also found that the acclimation process was coincident with a relative increase in amide I and amide II peak height (Osundeko et al., 2014). Moreover, this same study showed that particularly sensitive strains including *Chlamydomonas debaryana* and *Desmodesmus intermedius* exhibited accumulation of carbon storage products including glycerolipids and starch, while acclimated strains that could tolerate the wastewater conditions did not show this response. Likewise, other metabolic indicators of stress such as increased carbohydrate and lipid peaks within the FT-IR spectra were not observed in response to high Cu treatment in PM01.

Because Cu exerts toxicity in part through inhibition of key cellular processes including photosynthesis, either directly due to Cu binding or indirectly via accumulation of ROS (Jamers et al., 2013; Küpper et al., 2003; Sabatini et al., 2009), it would be expected that adaptive mechanisms would counteract these processes in some ways. Proteomic responses to Cu stress linked to Cu tolerance have previously been observed in other organisms. Cu exposure experiments in plants and fungi have observed increases in soluble protein that has been linked to induction of anti-oxidant enzymes (Cavalcanti Luna et al., 2015; Gao et al., 2008; Rout et al., 2013) while induction of Cu-binding proteins has been demonstrated in a Cu-tolerant variety of rice (Chen et al., 2015). Mechanisms of Cu tolerance in algae are not well understood, and may involve differential Cu uptake and internalisation in some cases (Levy et al., 2008). Although the molecular mechanisms of stress tolerance by *C. acidophila* are also poorly understood but there have been some recent insights. Heat shock proteins, which are a family of evolutionarily conserved stress tolerance molecular chaperones, have been previously found to increase in abundance in *C. acidophila* CCAP 11/137 in response to very low pH and metal-rich lake water treatment, partly in response to high Fe concentration (Gerloff-Elias et al., 2006; Spijkerman et al., 2007a). To date, no proteomic or enzymatic analysis has been performed in *C. acidophila* in response to Cu stress, but a transcriptomics approach observed differences in mRNA transcript profiles following Cu treatment in *C. acidophila* RT46. A range of gene transcripts

were up-regulated in response to Cu treatment including those involved in photosynthesis, signaling and stress-response (Olsson et al., 2015). Future experiments will aim to examine the proteomic adaptive response of *C. acidophila* PM01 in more detail in order to enhance our fundamental understanding of metal tolerance in microalgae, also to appreciate the effects that environmental pollution has on adaptive evolution and the ecological consequences of such adaptation.

A potential application of highly metal tolerant microalgae such as *C. acidophila* PM01 is the potential use of such organisms for metal bioremediation. Indeed PM01 was confirmed to bioconcentrate metals through a combination of cell wall binding and internalisation. Algae strains, such as those isolated in this study, may be used for *in situ* lake bioremediation in surface water mesocosms by controlled eutrophication and harvesting (Dessouki et al., 2005), or for *ex situ* bioremediation, such as immobilised algae in bioreactors (Mehta and Gaur, 2005). The accumulated metals may then be harvested and processed to allow metal recovery (Minoda et al., 2015; Raikova et al., 2016). Alternatively, highly acid and metal tolerant microalgae such as the PM01 strain, may have an important role in sustaining SO<sub>4</sub>-reducing bacteria by providing organic carbon and thus increasing the efficiency of AMD remediation microbial bioreactors (Diez-Ercilla et al., 2014; Nancucheo and Johnson, 2012; Totsche et al., 2006). For example, it was demonstrated that microalgal addition to mine tailing mesocosms containing pyrite-oxidizing bacteria caused higher production of alkalinity, higher concentrations of ferrous Fe, and increased immobilization of Cu and Zn (Nancucheo and Johnson, 2011).

The main aim of this study was to identify microorganisms from the open, standing waters of the AMD ponds. Some of the ponds were surrounded by vegetation including wetland plant species, which will harbor associated microorganisms (Aguinaga et al., 2018). Although not the scope of this study, future research can examine the role of the plants on microbial communities within isolated AMD pond environments, as well an examination of biota along spatial transects of the ponds including within the sediment. Moreover, future studies will be needed to examine how other microorganisms found in these environments



have adapted to AMD stresses, and whether there are common mechanisms between different extremophile species.

## 5. Conclusions

AMD is a major source of freshwater pollution worldwide for which restoration is very important. However, as found in this study, while extremely acidic and metal rich AMD substantially impacts biota, there is still substantial biodiversity, with tolerance derived through natural adaptation. In particular, a strain of *C. acidophila* is abundant in all ponds at this Cu mine site, especially in waters with high acidity and coupled with high PO<sub>4</sub>-P concentration. Although Cu toxicity is a significant challenge to most photosynthetic organisms, this strain of *C. acidophila* has specifically adapted to the high Cu status of the ponds in contrast to other strains of the same species isolated from field sites elsewhere. Moreover, the *C. acidophila* strain displays evidence of Cu-dependent metabolic plasticity. The marked metal tolerance and metal accumulation characteristics of *C. acidophila* PM01 indicates that organisms from these environments have biotechnological potential, such as bioremediation.

## Acknowledgments

We thank Oscar Aguinaga and Rachel Webster for assistance with sampling work, Lorna Slater and Shunzheng Wang for performing preliminary laboratory analysis of microalgae, Claude-Eric Souquieres for assistance with sequence data analysis, and Paul Lythgoe for ICP-AES analysis. This work was supported in part by BBSRC DTP PhD studentship funding to O.A.M. (grant number BB/M011208/1).

## References

Abinandan S, Subashchandrabose SR, Venkateswarlu K, Megharaj M. Microalgae–bacteria biofilms: a sustainable synergistic approach in remediation of acid mine drainage. *Applied Microbiology and Biotechnology* 2018; 102: 1131-1144.

663 Aguilera A, Manrubia SC, Gómez F, Rodríguez N, Amils R. Eukaryotic community  
 664 distribution and its relationship to water physicochemical parameters in an extreme  
 665 acidic environment, Río Tinto (Southwestern Spain). *Applied and Environmental*  
 666 *Microbiology* 2006; 72: 5325-5330.

667 Aguinaga OE, McMahon A, White KN, Dean AP, Pittman JK. Microbial community shifts in  
 668 response to acid mine drainage pollution within a natural wetland ecosystem.  
 669 *Frontiers in Microbiology* 2018; 9: 1445.

670 Amaral Zettler LA, Gomez F, Zettler E, Keenan BG, Amils R, Sogin ML. Microbiology:  
 671 eukaryotic diversity in Spain's River of Fire. *Nature* 2002; 417: 137-137.

672 August EE, McKnight DM, Hrnčíř DC, Garhart KS. Seasonal variability of metals transport  
 673 through a wetland impacted by mine drainage in the Rocky Mountains.  
 674 *Environmental Science & Technology* 2002; 36: 3779-3786.

675 Baffico GD, Diaz MM, Wenzel MT, Koschorreck M, Schimmele M, Neu TR, et al. Community  
 676 structure and photosynthetic activity of epilithon from a highly acidic ( $\text{pH} \leq 2$ ) mountain  
 677 stream in Patagonia, Argentina. *Extremophiles* 2004; 8: 463-473.

678 Baker BJ, Lutz MA, Dawson SC, Bond PL, Banfield JF. Metabolically active eukaryotic  
 679 communities in extremely acidic mine drainage. *Applied and Environmental*  
 680 *Microbiology* 2004; 70: 6264-6271.

681 Beamud SG, Diaz MM, Pedrozo FL. Summer phytoplankton composition and nitrogen  
 682 limitation of the deep, naturally-acidic ( $\text{pH} \sim 2.2$ ) Lake Caviahue, Patagonia,  
 683 Argentina. *Limnologia* 2007; 37: 37-48.

684 Bernal M, Ramiro MV, Cases R, Picorel R, Yruela I. Excess copper effect on growth,  
 685 chloroplast ultrastructure, oxygen-evolution activity and chlorophyll fluorescence in  
 686 *Glycine max* cell suspensions. *Physiologia Plantarum* 2006; 127: 312-325.

687 Boulton S, Collins DN, White KN, Curtis CD. Metal transport in a stream polluted by acid mine  
 688 drainage - the Afon Goch, Anglesey, UK. *Environmental Pollution* 1994; 84: 279-284.

689 Brake SS, Dannelly HK, Connors KA. Controls on the nature and distribution of an alga in  
690 coal mine-waste environments and its potential impact on water quality.  
691 Environmental Geology 2001; 40: 458-469.

692 Cavalcanti Luna MA, Vieira ER, Okada K, Campos-Takaki GM, do Nascimento AE. Copper-  
693 induced adaptation, oxidative stress and its tolerance in *Aspergillus niger* UCP1261.  
694 Electronic Journal of Biotechnology 2015; 18: 418-427.

695 Chen C, Song Y, Zhuang K, Li L, Xia Y, Shen Z. Proteomic analysis of copper-binding  
696 proteins in excess copper-stressed roots of two rice (*Oryza sativa* L.) varieties with  
697 different Cu tolerances. PLoS ONE 2015; 10: e0125367.

698 Cid C, Garcia-Descalzo L, Casado-Lafuente V, Amils R, Aguilera A. Proteomic analysis of  
699 the response of an acidophilic strain of *Chlamydomonas* sp. (Chlorophyta) to natural  
700 metal-rich water. Proteomics 2010; 10: 2026-2036.

701 Courtney LA, Clements WH. Assessing the influence of water and substratum quality on  
702 benthic macroinvertebrate communities in a metal-polluted stream: an experimental  
703 approach. Freshwater Biology 2002; 47: 1766-1778.

704 Das BK, Roy A, Koschorreck M, Mandal SM, Wendt-Potthoff K, Bhattacharya J. Occurrence  
705 and role of algae and fungi in acid mine drainage environment with special reference  
706 to metals and sulfate immobilization. Water Research 2009; 43: 883-894.

707 De Bisthoven LJ, Gerhardt A, Soares AMVM. Chironomidae larvae as bioindicators of an  
708 acid mine drainage in Portugal. Hydrobiologia 2005; 532: 181-191.

709 Dean AP, Lynch S, Rowland P, Toft BD, Pittman JK, White KN. Natural wetlands are  
710 efficient at providing long-term metal remediation of freshwater systems polluted by  
711 acid mine drainage. Environmental Science & Technology 2013; 47: 12029-12036.

712 Dean AP, Sigee DC, Estrada B, Pittman JK. Using FTIR spectroscopy for rapid  
713 determination of lipid accumulation in response to nitrogen limitation in freshwater  
714 microalgae. Bioresource Technology 2010; 101: 4499-4507.

715 DeLong EF. Archaea in coastal marine environments. Proceedings of the National Academy  
716 of Sciences, USA 1992; 89: 5685-5689.

717 Deneke R. Review of rotifers and crustaceans in highly acidic environments of pH values  $\leq 3$ .  
718 Hydrobiologia 2000; 433: 167-172.

719 Dessouki TCE, Hudson JJ, Neal BR, Bogard MJ. The effects of phosphorus additions on the  
720 sedimentation of contaminants in a uranium mine pit-lake. Water Research 2005; 39:  
721 3055-3061.

722 Diez-Ercilla M, Sánchez-España J, Yusta I, Wendt-Potthoff K, Koschorreck M. Formation of  
723 biogenic sulphides in the water column of an acidic pit lake: biogeochemical controls  
724 and effects on trace metal dynamics. Biogeochemistry 2014; 121: 519-536.

725 Driver T, Bajhaiya AK, Allwood JW, Goodacre R, Pittman JK, Dean AP. Metabolic responses  
726 of eukaryotic microalgae to environmental stress limit the ability of FT-IR  
727 spectroscopy for species identification. Algal Research 2015; 11: 148-155.

728 Florence K, Sapsford DJ, Johnson DB, Kay CM, Wolkersdorfer C. Iron-mineral accretion  
729 from acid mine drainage and its application in passive treatment. Environmental  
730 Technology 2016; 37: 1428-1440.

731 Fott B, McCarthy AJ. Three acidophilic Volvocine flagellates in pure culture. The Journal of  
732 Protozoology 1964; 11: 116-120.

733 Gao S, Yan R, Cao M, Yang W, Wang S, Chen F. Effects of copper on growth, antioxidant  
734 enzymes and phenylalanine ammonia-lyase activities in *Jatropha curcas* L. seedling.  
735 Plant Soil Environment 2008; 54: 117-122.

736 Geller W. Case Studies and Regional Surveys. In: Geller W, Schultze M, Kleinmann R,  
737 Wolkersdorfer C, editors. Acidic Pit Lakes: The Legacy of Coal and Metal Surface  
738 Mines. Springer Berlin Heidelberg, Berlin, Heidelberg, 2013, pp. 265-435.

739 Geller W, Schultze M. Remediation and Management of Acidified Pit Lakes and Outflowing  
740 Waters. In: Geller W, Schultze M, Kleinmann R, Wolkersdorfer C, editors. Acidic Pit  
741 Lakes: The Legacy of Coal and Metal Surface Mines. Springer Berlin Heidelberg,  
742 Berlin, Heidelberg, 2013, pp. 225-264.

743 Gerloff-Elias A, Barua D, Mölich A, Spijkerman E. Temperature- and pH-dependent  
 744 accumulation of heat-shock proteins in the acidophilic green alga *Chlamydomonas*  
 745 *acidophila*. FEMS Microbiology Ecology 2006; 56: 345-354.

746 Gerloff-Elias A, Spijkerman E, Proschold T. Effect of external pH on the growth,  
 747 photosynthesis and photosynthetic electron transport of *Chlamydomonas acidophila*  
 748 Negoro, isolated from an extremely acidic lake (pH 2.6). Plant, Cell & Environment  
 749 2005; 28: 1218-1229.

750 Giordano M, Kansiz M, Heraud P, Beardall J, Wood B, McNaughton D. Fourier transform  
 751 infrared spectroscopy as a novel tool to investigate changes in intracellular  
 752 macromolecular pools in the marine microalga *Chaetoceros muellerii*  
 753 (Bacillariophyceae). Journal of Phycology 2001; 37: 271-279.

754 Gray NF, Delaney E. Comparison of benthic macroinvertebrate indices for the assessment  
 755 of the impact of acid mine drainage on an Irish river below an abandoned Cu–S  
 756 mine. Environmental Pollution 2008; 155: 31-40.

757 Greenhaigh M, Ovenden D. Freshwater Life: Britain and Northern Europe. London, UK:  
 758 Collins, 2007.

759 Gustafsson JP. Visual MINTEQ ver. 3.0. Department of Land and Water Resources  
 760 Engineering, The Royal Institute of Technology, Sweden., 2010.

761 Hargreaves JW, Lloyd EJH, Whitton BA. Chemistry and vegetation of highly acidic streams.  
 762 Freshwater Biology 1975; 5: 563-576.

763 Hassler CS, Slaveykova VI, Wilkinson KJ. Discriminating between intra- and extracellular  
 764 metals using chemical extractions. Limnology and Oceanography-Methods 2004; 2:  
 765 237-247.

766 Hedin R, Weaver T, Wolfe N, Weaver K. Passive treatment of acidic coal mine drainage:  
 767 The Anna S mine passive treatment complex. Mine Water and the Environment  
 768 2010; 29: 165-175.

769 Hillebrand H, Durselen CD, Kirschtel D, Pollinger U, Zohary T. Biovolume calculation for  
 770 pelagic and benthic microalgae. Journal of Phycology 1999; 35: 403-424.

771 Hirooka S, Hirose Y, Kanesaki Y, Higuchi S, Fujiwara T, Onuma R, et al. Acidophilic green  
 772 algal genome provides insights into adaptation to an acidic environment.  
 773 Proceedings of the National Academy of Sciences, USA 2017; 114: E8304-E8313.  
 774 Huss VAR, Frank C, Hartmann EC, Hirmer M, Kloboucek A, Seidel BM, et al. Biochemical  
 775 taxonomy and molecular phylogeny of the genus *Chlorella* sensu lato (Chlorophyta).  
 776 Journal of Phycology 1999; 35: 587-598.  
 777 Jamers A, Blust R, De Coen W, Griffin JL, Jones OAH. Copper toxicity in the microalga  
 778 *Chlamydomonas reinhardtii*: an integrated approach. BioMetals 2013; 26: 731-740.  
 779 Jiang Y, Zhu Y, Hu Z, Lei A, Wang J. Towards elucidation of the toxic mechanism of copper  
 780 on the model green alga *Chlamydomonas reinhardtii*. Ecotoxicology 2016; 25: 1417-  
 781 1425.  
 782 John DM, Whitton BA, Brook AJ. The Freshwater Algae of the British Isles. An Identification  
 783 Guide to Freshwater and Terrestrial Algae. Cambridge: Cambridge University Press,  
 784 2002.  
 785 Johnson DB. Chemical and microbiological characteristics of mineral spoils and drainage  
 786 waters at abandoned coal and metal mines. Water, Air, & Soil Pollution: Focus 2003;  
 787 3: 47-66.  
 788 Johnson DB, Hallberg KB. The microbiology of acidic mine waters. Research in Microbiology  
 789 2003; 154: 466-473.  
 790 Johnson DB, Hallberg KB. Acid mine drainage remediation options: a review. Science of the  
 791 Total Environment 2005; 338: 3-14.  
 792 Küpper H, Šetlík I, Šetliková E, Ferimazova N, Spiller M, Küpper FC. Copper-induced  
 793 inhibition of photosynthesis: limiting steps of *in vivo* copper chlorophyll formation in  
 794 *Scenedesmus quadricauda*. Functional Plant Biology 2003; 30: 1187-1196.  
 795 Levy JL, Angel BM, Stauber JL, Poon WL, Simpson SL, Cheng SH, et al. Uptake and  
 796 internalisation of copper by three marine microalgae: Comparison of copper-sensitive  
 797 and copper-tolerant species. Aquatic Toxicology 2008; 89: 82-93.

798 Malmqvist B, Hoffsten P-O. Influence of drainage from old mine deposits on benthic  
799 macroinvertebrate communities in central Swedish streams. Water Research 1999;  
800 33: 2415-2423.

801 Martens H, Stark E. Extended multiplicative signal correction and spectral interference  
802 subtraction: new preprocessing methods for near infrared spectroscopy. Journal of  
803 Pharmaceutical and Biomedical Analysis 1991; 9: 625-635.

804 Mehta SK, Gaur JP. Use of algae for removing heavy metal ions from wastewater: progress  
805 and prospects. Critical Reviews in Biotechnology 2005; 25: 113-152.

806 Mendez-Garcia C, Pelaez AI, Mesa V, Sánchez J, Golyshina OV, Ferrer M. Microbial  
807 diversity and metabolic networks in acid mine drainage habitats. Frontiers in  
808 Microbiology 2015; 6: 475.

809 Michailova P, Ilkova J, Kerr R, White KN. Chromosome variability in *Chironomus acidophilus*  
810 Keyl, 1960 from the Afon Goch, UK – a river subject to long-term trace metal  
811 pollution. Aquatic Insects 2009; 31: 213-225.

812 Minoda A, Sawada H, Suzuki S, Miyashita S-i, Inagaki K, Yamamoto T, et al. Recovery of  
813 rare earth elements from the sulfothermophilic red alga *Galdieria sulphuraria* using  
814 aqueous acid. Applied Microbiology and Biotechnology 2015; 99: 1513-1519.

815 Ñancucheo I, Johnson DB. Acidophilic algae isolated from mine-impacted environments and  
816 their roles in sustaining heterotrophic acidophiles. Frontiers in Microbiology 2012; 3:  
817 325.

818 Ñancucheo I, Johnson DB. Significance of microbial communities and interactions in  
819 safeguarding reactive mine tailings by ecological engineering. Applied and  
820 Environmental Microbiology 2011; 77: 8201-8208.

821 Neculita C-M, Zagury GJ, Bussière B. Passive treatment of acid mine drainage in  
822 bioreactors using sulfate-reducing bacteria. Journal of Environmental Quality 2007;  
823 36: 1-16.

824 Nishikawa K, Tominaga N. Isolation, growth, ultrastructure, and metal tolerance of the green  
825 alga, *Chlamydomonas acidophila* (Chlorophyta). Bioscience, Biotechnology, and  
826 Biochemistry 2001; 65: 2650-2656.

827 Nixdorf B, Mischke U, Leßmann D. Chrysophytes and chlamydomonads: pioneer colonists in  
828 extremely acidic mining lakes (pH <3) in Lusatia (Germany). Hydrobiologia 1998;  
829 369: 315-327.

830 Nordstrom DK. Advances in the hydrogeochemistry and microbiology of acid mine waters.  
831 International Geology Review 2000; 42: 499-515.

832 Olaveson MM, Stokes PM. Responses of the acidophilic alga *Euglena mutabilis*  
833 (Euglenophyceae) to carbon enrichment at pH 3. Journal of Phycology 1989; 25:  
834 529-539.

835 Olsson S, Penacho V, Puente-Sánchez F, Díaz S, Gonzalez-Pastor JE, Aguilera A.  
836 Horizontal gene transfer of phytochelatin synthases from bacteria to extremophilic  
837 green algae. Microbial Ecology 2017; 73: 50-60.

838 Olsson S, Puente-Sánchez F, Gómez MJ, Aguilera A. Transcriptional response to copper  
839 excess and identification of genes involved in heavy metal tolerance in the  
840 extremophilic microalga *Chlamydomonas acidophila*. Extremophiles 2015; 19: 657-  
841 672.

842 Osundeko O, Davies H, Pittman JK. Oxidative stress-tolerant microalgae strains are highly  
843 efficient for biofuel feedstock production on wastewater. Biomass and Bioenergy  
844 2013; 56: 284-294.

845 Osundeko O, Dean AP, Davies H, Pittman JK. Acclimation of microalgae to wastewater  
846 environments involves increased oxidative stress tolerance activity. Plant and Cell  
847 Physiology 2014; 55: 1848-1857.

848 Pedrozo F, Kelly L, Diaz M, Temporetti P, Baffico G, Kringel R, et al. First results on the  
849 water chemistry, algae and trophic status of an Andean acidic lake system of  
850 volcanic origin in Patagonia (Lake Caviahue). Hydrobiologia 2001; 452: 129-137.



851 Pollio A, Cennamo P, Ciniglia C, De Stefano M, Pinto G, A.R. Huss V. *Chlamydomonas*  
852 *pitschmannii* Ettl, a little known species from thermoacidic environments. Protist  
853 2005; 156: 287-302.

854 Puente-Sánchez F, Díaz S, Penacho V, Aguilera A, Olsson S. Basis of genetic adaptation to  
855 heavy metal stress in the acidophilic green alga *Chlamydomonas acidophila*. Aquatic  
856 Toxicology 2018; 200: 62-72.

857 Puente-Sánchez F, Olsson S, Aguilera A. Comparative transcriptomic analysis of the  
858 response of *Dunaliella acidophila* (Chlorophyta) to short-term cadmium and chronic  
859 natural metal-rich water exposures. Microbial Ecology 2016; 72: 595-607.

860 Quigley M. Invertebrates of Streams and Rivers: A Key to Identification. London, UK:  
861 Edward Arnold, 1977.

862 Raikova S, Smith-Baedorf H, Bransgrove R, Barlow O, Santomauro F, Wagner JL, et al.  
863 Assessing hydrothermal liquefaction for the production of bio-oil and enhanced metal  
864 recovery from microalgae cultivated on acid mine drainage. Fuel Processing  
865 Technology 2016; 142: 219-227.

866 Rout JR, Ram SS, Das R, Chakraborty A, Sudarshan M, Sahoo SL. Copper-stress induced  
867 alterations in protein profile and antioxidant enzymes activities in the in vitro grown  
868 *Withania somnifera* L. Physiology and Molecular Biology of Plants 2013; 19: 353-361.

869 Sabatini SE, Juárez ÁB, Eppis MR, Bianchi L, Luquet CM, Ríos de Molina MdC. Oxidative  
870 stress and antioxidant defenses in two green microalgae exposed to copper.  
871 Ecotoxicology and Environmental Safety 2009; 72: 1200-1206.

872 Sánchez España J, Pamo EL, Pastor ES, Ercilla MD. The acidic mine pit lakes of the Iberian  
873 Pyrite Belt: An approach to their physical limnology and hydrogeochemistry. Applied  
874 Geochemistry 2008; 23: 1260-1287.

875 Schönknecht G, Chen W-H, Ternes CM, Barbier GG, Shrestha RP, Stanke M, et al. Gene  
876 transfer from bacteria and archaea facilitated evolution of an extremophilic eukaryote.  
877 Science 2013; 339: 1207-1210.

878 Schultze M. Limnology of Pit Lakes. In: Geller W, Schultze M, Kleinmann R, Wolkersdorfer  
879 C, editors. Acidic Pit Lakes: The Legacy of Coal and Metal Surface Mines. Springer  
880 Berlin Heidelberg, Berlin, Heidelberg, 2013, pp. 23-224.

881 Sharma SS, Dietz K-J. The relationship between metal toxicity and cellular redox imbalance.  
882 Trends in Plant Science 2009; 14: 43-50.

883 Smucker NJ, Drerup SA, Vis ML. Roles of benthic algae in the structure, function, and  
884 assessment of stream ecosystems affected by acid mine drainage. Journal of  
885 Phycology 2014; 50: 425-436.

886 Spijkerman E. Phosphorus limitation of algae living in iron-rich, acidic lakes. Aquatic  
887 Microbial Ecology 2008; 53: 201-210.

888 Spijkerman E, Barua D, Gerloff-Elias A, Kern J, Gaedke U, Heckathorn SA. Stress  
889 responses and metal tolerance of *Chlamydomonas acidophila* in metal-enriched lake  
890 water and artificial medium. Extremophiles 2007a; 11: 551-562.

891 Spijkerman E, Bissinger V, Meister A, Gaedke U. Low potassium and inorganic carbon  
892 concentrations influence a possible phosphorus limitation in *Chlamydomonas*  
893 *acidophila* (Chlorophyceae). European Journal of Phycology 2007b; 42: 327-339.

894 Stamatakis A. RAxML-VI-HPC: Maximum likelihood-based phylogenetic analyses with  
895 thousands of taxa and mixed models. Bioinformatics 2006; 22: 2688-2690.

896 Totsche O, Fyson A, Steinberg CEW. Microbial alkalinity production to prevent reacidification  
897 of neutralized mining lakes. Mine Water and the Environment 2006; 25: 204-213.

898 Twiss MR. Copper tolerance of *Chlamydomonas acidophila* (Chlorophyceae) isolated from  
899 acidic, copper-contaminated soils. Journal of Phycology 1990; 26: 655-659.

900 Valko M, Morris H, Cronin MTD. Metals, toxicity and oxidative stress. Current Medicinal  
901 Chemistry 2005; 12: 1161-1208.

902 Varshney P, Mikulic P, Vonshak A, Beardall J, Wangikar PP. Extremophilic micro-algae and  
903 their potential contribution in biotechnology. Bioresource Technology 2015; 184: 363-  
904 372.

905 Vrba J, Kopáček J, Fott J, Kohout L, Nedbalová L, Pražáková M, et al. Long-term studies  
 906 (1871–2000) on acidification and recovery of lakes in the Bohemian Forest (central  
 907 Europe). *Science of The Total Environment* 2003; 310: 73-85.  
 908 Webster RE, Dean AP, Pittman JK. Cadmium exposure and phosphorus limitation increases  
 909 metal content in the freshwater alga *Chlamydomonas reinhardtii*. *Environmental*  
 910 *Science & Technology* 2011; 45: 7489-7496.  
 911 Younger PL, Potter HAB. Parys in Springtime: Hazard Management and Steps Towards  
 912 Remediation of the UKs Most Polluted Acidic Mine Discharge. 9th International  
 913 Conference on Acid Rock Drainage (ICARD), Ottawa, Canada, 2012.  
 914 Yun H-S, Lee H, Park Y-T, Ji M-K, Kabra AN, Jeon C, et al. Isolation of novel microalgae  
 915 from acid mine drainage and its potential application for biodiesel production. *Applied*  
 916 *Biochemistry and Biotechnology* 2014; 173: 2054-2064.  
 917 Zalack JT, Smucker NJ, Vis ML. Development of a diatom index of biotic integrity for acid  
 918 mine drainage impacted streams. *Ecological Indicators* 2010; 10: 287-295.

919

920

## Figures




Fig. 1. Map and satellite image of Parys Mountain copper mine in Anglesey, North Wales, UK. The sampled AMD ponds are labelled 1–9 and the AMD-polluted Southern Afon Goch river is indicated. The location coordinates of each sampled pond are also shown (A). (B) Satellite image of the mine and the sampled ponds. Source, Google Earth, 2016.




Fig. 2. Water chemistry of the AMD ponds. (A - C) pH and conductivity values (A), and concentration of dissolved Fe, Cd (B), Cu and Zn (C) for each pond during 2015 (spring, summer, autumn samples). Boxes show the 25<sup>th</sup> and 75<sup>th</sup> percentiles, the line within the boxes shows the median values, and the whisker bars show minimum and maximum values ( $n = 3 - 18$ ). (D) PO<sub>4</sub>-P, NO<sub>3</sub>-N and NH<sub>4</sub>-N concentration in each pond (June sample). Pond 7 data is not available. Values are means ( $n = 3 - 12$ ) and error bars correspond to the standard error of the mean. (E) Mean chlorophyll-a (Chl a) concentration in each pond during 2015 (spring, summer, autumn samples). Chl a change in Pond 1 during the year and pond water temperature is shown (inset). Each pond was sampled in triplicate on three separate occasions (spring, summer, autumn). Values are means ( $n = 9$ ) and error bars correspond to the standard error of the mean. For all data, bars that do not share a lower case letter show significant difference ( $p < 0.05$ ) between pond sites.




Fig. 3. PCA of water chemistry data from each of the AMD ponds. (A) Observation plot for each pond. The upper site small ponds are shown as blue triangles, the near-neutral pH pond as a red circle, and the lower site large lagoon ponds as green squares. (B) Variables plot with selected environmental variables labelled.




Fig. 4. Taxa diversity of eukaryotic organisms in the AMD ponds as determined by numbers of distinct taxonomic families of microalgae, moss/liverwort, fungi, protozoa, as identified by 18S rRNA gene amplicon sequencing, and families of invertebrates, as identified by visual observation and use of taxonomic identification keys. Data for Pond 8 is not available.




Fig. 5. (A) Abundance of *C. acidophila* microalgae and Chironomidae insects in each pond from summer sampling. Values of *C. acidophila* are means ( $n = 3$ ) and error bars correspond to the standard error of the mean, while value of Chironomidae are total counts. For *C. acidophila* data, bars that do not share a lower case letter show significant difference ( $p < 0.05$ ) between pond sites. Chironomidae data for Pond 8 is not available. (B) Linear regression analyses for dissolved phosphate, pH, conductivity and dissolved Cu in relation to *C. acidophila* cell counts in AMD ponds. Apart from pH values, all data were natural log transformed.






Fig. 6. 18S rDNA sequence analysis of *C. acidophila* strains. Phylogenetic tree based on 18S nucleotide sequence obtained from the PM01 strain isolated from Pond 1, sequences of *C. acidophila* CCAP 11/136 and CCAP 11/137 strains, other known or putative *C. acidophila* strains, and selected unicellular Chlorophyta microalgae of the *Chlamydomonas moewusii* clade, including three strains originally classified as *C. acidophila*. The sequence lengths were between 1528 – 1792 nucleotides. *Chlamydomonas reinhardtii* is included as an out-group. Accession numbers are shown for each sequence. Bootstrap percentage values are indicated at the tree nodes of branches for 100 replications and indicate confidence in tree node positions. The branch length scale bar indicates evolutionary distance.




Fig. 7. Cu tolerance of *C. acidophila* PM01 in comparison to *C. acidophila* strains CCAP 11/136 and 11/137. (A) Growth of algae spot dilutions on MAM pH 3.0 plates with or without added Cu and photographed after 12 d. Image is representative of 3 independent experiments. (B) Growth of strains in liquid MAM pH 3.0 determined after 14 d cultivation in response to a range of Cu concentrations. Values are means (n = 3 - 4) and error bars correspond to the standard error of the mean. Bars that do not share a lower case letter show significant difference (p < 0.05) between strains.




Fig. 8. PCA clustering of FT-IR spectra from *C. acidophila* PM01 in comparison to *C. acidophila* strains CCAP 11/136 and 11/137 in response to a range of Cu concentrations. (A) PCA scores plot of replicate ( $n = 4$ ) FT-IR spectra obtained from cells grown after 14 d cultivation in liquid MAM pH 3.0 with increasing concentrations of added Cu or no added Cu. Only PM01 cells could grow in 130 mg L<sup>-1</sup> Cu. (B) PC1 and PC2 loading plot. Band peaks which explain most of the variation for each PC are  $\nu$ C=O of amide I (1655 cm<sup>-1</sup>) and  $\delta$ N-H of amide II (1545 cm<sup>-1</sup>).