

Please cite the Published Version

Symeonakis, Ilias, Petroulaki, K, Higginbottom, T and Marques-Mateu, A (2017) Optimisation of regional scale woody vegetation cover mapping with optical, thermal and radar data. In: European Space Agency WorldCover 2017, 14 March 2017 - 16 March 2107, Frascati, Italy.

Publisher: ESA

Downloaded from: <https://e-space.mmu.ac.uk/622287/>


Usage rights: © In Copyright

Additional Information: Abstract and poster presented at European Space Agency WorldCover 2017.

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

Optimisation of Woody Vegetation Cover Mapping with Optical, Thermal and Radar data



Elías Symeonakis, K. Petroulaki, T. Higginbottom, Á. Marqués-Mateu²

School of Science and the Environment, Manchester Metropolitan University, UK

²School of Engineering in Geodesy, Cartography and Surveying, Universitat Politècnica de València, Spain

E.Symeonakis@mmu.ac.uk, Kyriaki.Petroulaki@stu.mmu.ac.uk, T.Higginbottom@mmu.ac.uk

www.land-degradation.org



Rationale

- Woody perennial vegetation is an integral part of savannah ecosystems
- Accurately mapping its presence can provide useful input to global carbon emissions models as well regional policy decision making efforts regarding bush control or the overexploitation of fuelwood
- Recent attempts to map the extent of savannah woody cover over the regional scale have employed Earth observation data either from optical or radar sensors, and most commonly from the dry season when the spectral difference from the 'background' grasses is maximised
- By far the most common practice has been the use of Landsat optical bands, but some studies have also used vegetation indices or L-band or C-band SAR data
- However, conflicting reports with regards to the effectiveness of the different approaches have emerged leaving the respective land cover mapping community with unclear methodological pathways to follow
- We address this issue by employing Landsat and ALOS PALSAR data, together with colour aerial photography for training and validation of random forest regressions, to assess the accuracy of mapping woody vegetation when:
 - (a) data from either or both (wet and dry) seasons are considered
 - (b) PALSAR data are used on their own or together with the optical data
 - (c) vegetation indices are calculated and are used either on their own or together with the Landsat bands
 - (iv) thermal infrared information is not discarded but included in the parameterisation

Study area (Figure 1)

Falls within the Northwest Province (NWP), South Africa:

- Covers an area >53,000 km²
- 6 Landsat scenes required for mosaic
- Temperatures:
 - 17° to 31 °C summer
 - 3° to 21 °C winter
- Annual rainfall:
 - ~360 mm, ~all in summer months, (October to April)

Methods

Sampling (training & validation)

~ 10,000 point samples of three land cover types were selected:

- Woody vegetation cover (WVC)
- Non-woody vegetation cover (NWVC), including grasses and crops
- No vegetation (NVC, urban areas and bare areas)

Regressions

Random forest (RF) regressions carried out using R

Accuracy statistics reviewed and training samples modified to achieve optimum predictive models.




Figure 2. Flowchart of the methodological framework

Results & Discussion

- Table 2 is a summary of the overall statistics (accuracy and k) and the per-class balanced accuracy figures for the 16 models tested
- The inclusion of the wet season data, the thermal band and vegetation indices and, most importantly, the radar data, improves the overall accuracy of the classifications by 13% and the balanced accuracy for the mapping of the woody vegetation cover by 9%.
- Simply adding the HV polarized SAR data to the dry season optical bands, improves the overall accuracy by 10% and the woody cover balanced accuracy by 7%.
- The accuracies achieved are in agreement with a number of research studies comparing radar and optical data, e.g. Armston et al. (2009), Laurin et al. (2012), Lehmann et al. (2015), Higginbottom et al. (subm), Symeonakis and Higginbottom (2014)

ID	Model	~5000 samples (=50% of total, other 50% for validation)										
		Per Class Balanced Accuracy			Prod. Acc. (om.)			User Acc. (comm.)			Overall Acc (%)	K
		WVC	NWVC	NVC	WVC	NWVC	NVC	WVC	NWVC	NVC		
1	Lan dry	81.33	77.48	81.81	66.97	77.52	77.36	75.91	75.31	76.09	75.69	0.60
2	Lan wet	80.21	78.73	83.53	65.40	78.90	79.22	72.88	76.55	78.56	76.72	0.62
3	Lan (dry+wet)	86.53	83.91	86.44	76.52	85.46	81.24	81.93	81.01	84.63	82.41	0.71
4	(Lan+SARHH) _{dry}	85.35	82.93	85.32	74.65	83.09	81.03	79.27	80.72	81.89	80.92	0.69
5	(Lan+SARHV) _{dry}	88.67	88.11	87.68	80.10	88.91	83.88	85.41	85.93	85.15	85.57	0.77
6	(Lan+SAR(HH,HV)) _{dry}	88.40	88.08	86.64	79.42	91.20	80.42	85.97	84.07	86.72	85.27	0.76
7	(Lan+SAR(HH,HV,diff,div)) _{dry}	88.37	87.70	87.10	79.80	89.48	81.99	84.07	84.84	85.93	85.10	0.76
8	SAR(HH,HV,diff,div) _{dry}	60.87	82.65	65.15	31.78	86.28	54.10	39.07	78.22	56.67	65.34	0.43
9	3VIs (NDVI,MSAVI,TNDVI) _{dry}	82.31	69.15	71.46	69.31	69.48	62.55	75.39	65.80	64.68	66.92	0.46
10	4VIs (NDVI,MSAVI,TNDVI,GDVI ²) _{dry}	81.15	67.86	71.43	67.01	66.19	64.82	74.64	65.35	62.83	65.83	0.45
11	(Lan+2VIs) _{dry}	83.72	78.76	82.02	71.55	79.70	76.17	78.18	75.99	77.99	77.04	0.63
12	(Lan+3VIs) _{dry}	83.80	78.84	82.80	71.52	78.25	78.84	78.87	76.74	77.43	77.32	0.63
13	(Lan+TIR) _{dry}	84.11	81.42	85.84	72.12	81.58	82.12	79.12	79.43	81.58	80.16	0.68
14	Dry+3VIs+TIR+SAR(HH,HV)	88.78	90.09	88.87	79.71	92.56	84.38	88.26	86.42	88.26	87.36	0.79
15	Dry+wet+3VIs+3VIs _w +TIR _w +TIR _{dry}	87.20	85.13	87.78	77.14	87.46	82.73	85.22	81.57	86.89	83.98	0.74
16	Dry+wet+3VIs+3VIs _w +TIR _w +TIR _{dry} +SAR(HH,HV) _{dry}	90.59	90.75	90.25	83.06	93.50	85.55	90.11	86.78	90.94	88.76	0.82

Table 2. Summary table of overall accuracies for the 16 models and the four different training sample sizes tested for the year 2007. 'dry': dry season Landsat TM or ETM+ bands; 'wet': same for wet season; 'TIR': thermal Landsat band; 'NDVI': Normalised Difference Vegetation Index; 'MSAVI': Modified Soil Adjusted Vegetation Index; 'HH' and 'HV': HH- or HV-polarised ALOS PALSAR data

Datasets

Aerial photos

- 0.5m-pixel colour aerial photography (free for 2008 onwards by South African National Geospatial Information (NGI) mapping agency, <http://www.ngi.gov.za/index.php>)

Landsat

- The Landsat imagery employed for the mosaics are shown in Table 1
- Where ETM+ SLC-off data had to be used, gaps were filled in using the Gapfill plug-in for ENVI 5.2

SAR data

We used ALOS PALSAR data from the Alaska Satellite Facility (<https://www.asf.alaska.edu/>) level 1.5 high resolution terrain corrected data in dual HH and HV polarisation and a spatial resolution of 12.5m

Season	Sensor	Path	Row	DOY	Year
Dry	5 TM	171	79	228	2007
	5 TM	172	77	200	2006
	5 TM	172	78	184	2006
	5 TM	172	79	187	2007
	5 TM	173	78	191	2006
	5 TM	173	79	223	2006
Wet	5 TM	171	79	4	2007
	7 ETM+	172	77	35	2007
	7 ETM+	172	77	19	2007
	7 ETM+	172	78	35	2007
	7 ETM+	172	78	51	2007
	7 ETM+	172	79	35	2007
	7 ETM+	172	79	19	2007
	5 TM	173	78	114	2007
5 TM	173	79	114	2007	

Table 1. Landsat data used for the dry and wet seasons of 2007. When data for 2007 were not available, data for 2006 were used. Two scenes per SLC-off Landsat 7 scene were used to deal with the stripping issue

Conclusions

- Remote sensing methods that accurately map woody vegetation cover in southern African savannahs are important as an initial stage in the attempt to map bush encroachment, a process repeatedly acknowledged as a form of land degradation in these areas
- Our research has identified the combination of Landsat TM multi-seasonal optical and thermal data, together with ALOS PALSAR1 HH and HV polarized data, as the most accurate in mapping woody vegetation cover in the Northwest Province of South Africa. We also concluded, however, that in the absence of other data, employing dry season Landsat data is able to provide highly accurate estimates of woody cover
- Further research is currently underway in South Africa aiming to employ the presented methodological framework to assess woody cover change through time, on the one hand, and the mapping and monitoring of fractional woody cover, on the other

References:

- Armston, J.D., Denham, R.J., Danaher, T.J., Scarth, P.F., Moffiet, T.N. 2009. Prediction and validation of foliage projective cover from Landsat-5 TM and Landsat-7 ETM+ imagery. *Journal of Applied Remote Sensing*, 3 (1), art. no. doi: 10.1117/1.3216031
- Higginbottom, T.P., Symeonakis, E., Meyer, H., van der Linden, S. subm. Mapping Woody Coverage in Semi-arid Savannahs: the role of scale, seasonality, and sensor fusion. *International Journal of Applied Earth Observation and Geoinformation*.
- Laurin, G.V., Liesenber, V., Chen, Q., Guerriero, L., Del Frate, F., Bartolini, A., Coomes, D., (...), Valentini, R. 2012. Optical and SAR sensor synergies for forest and land cover mapping in a tropical site in West Africa. *International Journal of Applied Earth Observation and Geoinformation*, 21 (1), pp. 7-16.
- Lehmann, E.A., Caccetta, P., Lowell, K., Mitchell, A., Zhou, Z.-S., Held, A., Milne, T., (...), Tapley, I. 2015. SAR and optical remote sensing: Assessment of complementarity and interoperability in the context of a large scale operational forest monitoring system. *Remote Sensing of Environment*, 156, pp. 335-348
- Symeonakis, E., Higginbottom, T. P. 2014. Bush encroachment monitoring using multi-temporal Landsat data & random forests. *Int. Arch. Photogramm. Remote Sens. Spatial Inf. Sci.*, XL-2, 29-35. doi:10.5194/isprsarchives-XL-2-29-2014.

Acknowledgements

Elías Symeonakis is supported by an EU FP7 Marie Curie Career Integration Grant (PCIG12-GA-2012-3374327) Thomas Higginbottom is supported by an MMU PhD Fellowship