



**Manchester  
Metropolitan  
University**

---

Kougkoulos, I, Cook, S, Clarke, L, Symeonakis, Ilias, Edwards, L and Dortch, J (2018) Volumetric changes of glaciers in the Bolivian Andes between 1986 and 2017. In: EGU General Assembly, 08 April 2018 - 13 April 2018, Vienna, Austria.

---

**Downloaded from:** <https://e-space.mmu.ac.uk/622298/>

**Publisher:** EGU

Please cite the published version

<https://e-space.mmu.ac.uk>



## **Volumetric changes of glaciers in the Bolivian Andes between 1986 and 2017**

Ioannis Kougkoulos (1), Simon J. Cook (2), Leon J. Clarke (1), Elias Symeonakis (1), Laura A. Edwards (3), and Jason M. Dortch (4)

(1) School of Science and the Environment, Manchester Metropolitan University, Chester Street, Manchester, M1 5GD, UK, (2) Geography, School of Social Sciences, University of Dundee, Nethergate, Dundee DD1 4HN, UK, (3) School of Natural Sciences and Psychology, Liverpool John Moores University, Liverpool, L3 3AF, UK, (4) Department of Geography, University of Manchester, Oxford Road, Manchester, M13 9PL, UK

Glaciers represent an important water resource for Andean cities and mountain communities. However, a recent study has shown that Bolivian glaciers have shrunk by  $\sim 43\%$  in area over the last  $\sim 30$  years. If current rates of glacier recession are sustained then there could be potentially important consequences for downstream water supply, especially during the dry season. A first step in assessing the severity of this problem is to estimate the current volume of glacier ice in Bolivia, and how this has changed over recent decades. Here, we use VOLTA (created by James and Carrivick, 2016 – Computers & Geosciences), an ArcGIS tool requiring only a Digital Elevation Model (DEM) and glacier outlines to give a first-order ice thickness estimate and therefore derive volume changes for the entire Bolivian Andes between 1986 and 2017. The VOLTA tool also models bed topography, which we use to make a preliminary assessment of the locations of subglacial overdeepenings, which will become loci for future proglacial lakes capable of generating glacial lake outburst floods (GLOFs) and storing meltwater.