


Please cite the Published Version

Burnell, OW, Connell, SD, Irving, AD, Watling, JR  and Russell, BD (2014) Contemporary reliance on bicarbonate acquisition predicts increased growth of seagrass *Amphibolis antarctica* in a high-CO₂ world. *Conservation Physiology*, 2 (1).

DOI: <https://doi.org/10.1093/conphys/cou052>

Publisher: Oxford University Press (OUP)

Version: Published Version

Downloaded from: <https://e-space.mmu.ac.uk/623066/>

Usage rights:  [Creative Commons: Attribution 4.0](https://creativecommons.org/licenses/by/4.0/)

Additional Information: This is an Open Access article in *Conservation Physiology* published by Oxford University Press.

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

Contemporary reliance on bicarbonate acquisition predicts increased growth of seagrass *Amphibolis antarctica* in a high-CO₂ world

Owen W. Burnell^{1*}, Sean D. Connell¹, Andrew D. Irving², Jennifer R. Watling¹ and Bayden D. Russell¹

¹School of Earth & Environmental Sciences, University of Adelaide, Adelaide, SA 5005, Australia

²School of Medical and Applied Sciences, Central Queensland University, Bruce Highway, Rockhampton, QLD 4702, Australia

*Corresponding author: School of Earth & Environmental Sciences, University of Adelaide, Adelaide, SA 5005, Australia. Tel: +61 8 8313 6125. Email: owen.burnell@adelaide.edu.au

Rising atmospheric CO₂ is increasing the availability of dissolved CO₂ in the ocean relative to HCO₃⁻. Currently, many marine primary producers use HCO₃⁻ for photosynthesis, but this is energetically costly. Increasing passive CO₂ uptake relative to HCO₃⁻ pathways could provide energy savings, leading to increased productivity and growth of marine plants. Inorganic carbon-uptake mechanisms in the seagrass *Amphibolis antarctica* were determined using the carbonic anhydrase inhibitor acetazolamide (AZ) and the buffer tris(hydroxymethyl)aminomethane (TRIS). *Amphibolis antarctica* seedlings were also maintained in current and forecasted CO₂ concentrations to measure their physiology and growth. Photosynthesis of *A. antarctica* was significantly reduced by AZ and TRIS, indicating utilization of HCO₃⁻-uptake mechanisms. When acclimated plants were switched between CO₂ treatments, the photosynthetic rate was dependent on measurement conditions but not growth conditions, indicating a dynamic response to changes in dissolved CO₂ concentration, rather than lasting effects of acclimation. At forecast CO₂ concentrations, seedlings had a greater maximum electron transport rate (1.4-fold), photosynthesis (2.1-fold), below-ground biomass (1.7-fold) and increase in leaf number (2-fold) relative to plants in the current CO₂ concentration. The greater increase in photosynthesis (measured as O₂ production) compared with the electron transport rate at forecasted CO₂ concentration suggests that photosynthetic efficiency increased, possibly due to a decrease in photorespiration. Thus, it appears that the photosynthesis and growth of seagrasses reliant on energetically costly HCO₃⁻ acquisition, such as *A. antarctica*, might increase at forecasted CO₂ concentrations. Greater growth might enhance the future prosperity and rehabilitation of these important habitat-forming plants, which have experienced declines of global significance.

Key words: *Amphibolis antarctica*, carbon dioxide, carbonic anhydrase, electron transport rate, oxygen evolution, photosynthesis

Editor: Lauren Sack

Received 9 January 2014; Revised 30 July 2014; accepted 16 October 2014

Cite as: Burnell OW, Connell SD, Irving AD, Watling JR, Russell BD (2014) Contemporary reliance on bicarbonate acquisition predicts increased growth of seagrass *Amphibolis antarctica* in a high-CO₂ world. *Conserv Physiol* 2: doi:10.1093/conphys/cou052.

Introduction

Seagrasses are habitat-forming marine plants that provide a number of critical ecological services to coastal zones, such as

stabilization of sediments, support of trophic food webs, nutrient cycling and carbon sequestration (Costanza *et al.*, 1997; Duarte, 2002; Waycott *et al.*, 2009; Irving *et al.*, 2011). When angiosperms first entered the aquatic realm nearly

90 million years ago, atmospheric CO₂ levels were much greater (approximately three to seven times) than today (Beer and Koch, 1996; Beardall *et al.*, 1998; Berner and Kothavala, 2001). Since that time, atmospheric CO₂ concentrations have generally declined (Berner and Kothavala, 2001), a trend which has resulted in potential carbon limitation for many marine plants. This low atmospheric CO₂ has reduced the availability of inorganic carbon (C_i) for photosynthesis, which is compounded in marine systems by the slow diffusion of CO₂ in seawater and the slow rate of conversion of HCO₃⁻ to CO₂ when uncatalysed (Beer, 1989; Larkum *et al.*, 1989; Schwarz *et al.*, 2000). The recent spike in atmospheric CO₂ linked to anthropogenic activities is changing C_i availability in marine systems and is thus likely to affect carbon acquisition and growth in primary producers, such as seagrasses (Koch *et al.*, 2013).

While HCO₃⁻ is more readily available than dissolved CO₂ in marine systems, it cannot diffuse passively across the cell plasma membrane, and therefore, extracellular mechanisms have evolved to aid aquatic plants in the acquisition of CO₂ from HCO₃⁻ for photosynthesis, thereby reducing C_i limitation (Larkum, *et al.*, 1989; Invers *et al.*, 1999; Hellblom *et al.*, 2001). Three primary extracellular HCO₃⁻ acquisition systems have been described for seagrasses (Beer and Koch, 1996; James and Larkum, 1996; Hellblom *et al.*, 2001; Beer *et al.*, 2002). First, the enzyme carbonic anhydrase (CA) can catalyse the rapid conversion of HCO₃⁻ to CO₂ to restore CO₂/HCO₃⁻ equilibrium at the plasma membrane or concentrate CO₂ at the chloroplast level (System A, *sensu* Beer *et al.*, 2002). Second, the outward pumping of protons (H⁺) from cells can create H⁺ gradients to aid the cotransport of H⁺ and HCO₃⁻ back across the plasma membrane (System B). Finally, the combination of extracellular CA-catalysed dehydration of HCO₃⁻ to CO₂ within acidified zones created by the extrusion of H⁺ across the plasma membrane can concentrate CO₂ and encourage diffusion into cells (System C). For a full review of these mechanisms, see Beer *et al.* (2002).

All seagrasses tested to date appear to be reliant, to some extent, on the extracellular activity of CA for carbon acquisition, suggesting that they could experience some degree of carbon limitation at current atmospheric CO₂ concentrations (Larkum *et al.*, 2006; Koch *et al.*, 2013). Also, there is an energetic cost associated with both the production of extracellular CA and the more active mode of carbon acquisition using H⁺ extrusion (Spalding and Ogren, 1982; Raven and Lucas, 1985; Falk and Palmqvist, 1992; Kübler and Raven, 1995; Fridlyand *et al.*, 1996). As atmospheric, and thus oceanic, CO₂ continues to accumulate, this energetic cost could diminish if reliance on these carbon-acquisition mechanisms decreases, relative to direct CO₂ usage (Beardall and Giordano, 2002; Raven *et al.*, 2011; Koch *et al.*, 2013). For example, some lower order marine producers (i.e. cyanobacteria and eukaryotic algae) can adjust their HCO₃⁻-acquisition strategies within a number of hours of exposure to different CO₂ conditions (Falk and Palmqvist, 1992; Matsuda and Colman, 1995; Brueggeman *et al.*, 2012). Seagrasses may

have a similar ability to regulate energetically costly C_i acquisition, but our understanding of this has advanced slowly relative to lower order producers (i.e. cyanobacteria and eukaryotic algae) that have shorter generation times and can therefore be manipulated more easily for genetic expression studies (Larkum, *et al.*, 2006).

Increasing CO₂ availability and any down-regulation of HCO₃⁻ acquisition could result in improved photosynthetic efficiency as the energy required to acquire carbon decreases, which may also translate to greater photosynthetic rates (Badger and Andrews, 1982; Raven *et al.*, 2011; Koch *et al.*, 2013). Likewise, if CA-mediated HCO₃⁻ mechanisms are maintained, they may become more efficient at lower ambient pH levels (Koch *et al.*, 2013). However, this potential for more efficient and greater photosynthesis may be accompanied by a net gain in leaf growth or energy storage only when other resources, such as light or nitrogen, are not limiting (Zimmerman *et al.*, 1997; Palacios and Zimmerman, 2007; Alexandre *et al.*, 2012). In many seagrass species with a heavy reliance on HCO₃⁻ for C_i acquisition, it is unknown whether they will undergo subsequent changes in growth as dissolved CO₂ increases.

In the present study, the reliance of *Amphibolis antarctica* (Labill.) Sonder et Ascherson on HCO₃⁻ pathways of C_i acquisition was investigated by using an inhibitor of the enzyme CA (i.e. acetazolamide, AZ) and the biological buffer (i.e. tris(hydroxymethyl)aminomethane, TRIS). Having established that *A. antarctica* has a significant reliance on energetically costly HCO₃⁻ acquisition, a second experiment was conducted in which juvenile *A. antarctica* were grown in the presence of ambient (~390 ppm) and forecasted CO₂ concentrations (~900 ppm). It was hypothesized that photosynthesis and growth would increase for *A. antarctica* when CO₂ was enriched, because the greater availability of CO₂ relative to HCO₃⁻ might increase the photosynthetic efficiency of plants, as they may partition relatively fewer resources to energetically costly processes, such as HCO₃⁻-uptake mechanisms or photorespiration.

Materials and methods

Plant material

Mature seagrasses were collected from a depth of 4 m at Marino Rocks in the Gulf St Vincent, South Australia (35°02.806 S, 138°30.350 E). Seagrasses were transported to The University of Adelaide and kept in recirculating aerated aquaria with lighting conditions similar to the collection site (~60 μmol m⁻² s⁻¹) in a 12 h–12 h light–dark cycle for 1 week, during which time experiments to determine C_i-uptake mechanisms took place.

Inorganic carbon-acquisition mechanisms

The C_i-uptake mechanisms of seagrasses were investigated by inhibiting HCO₃⁻ pathways to carbon acquisition. Seagrasses were exposed to the inhibitor AZ or the biological buffer

TRIS, either separately or in combination. The primary seagrass of interest was *A. antarctica*, but to identify the generality of the mechanism within and across genera, C₁-uptake mechanisms were also investigated for two other co-occurring seagrass species, the congener *Amphibolis griffithii* (Blackden Hartog) and the species *Posidonia sinuosa* (Cambridge and Kuo), using the same methodology.

Oxygen evolution rates were determined using a Clark-type oxygen electrode and the logging program Biograph (Axword Software, Adelaide, South Australia). An entire leaf of *A. antarctica* (~20 mm long) that was free of epiphyte growth was placed in the electrode chamber in 4 ml of seawater filtered to 0.45 µm. The fourth or fifth youngest leaf on each leaf head was chosen because these leaves were mature and consistently free of epiphytes and other biota, as well as being the correct length to fit the photosynthetic chamber. The chamber was illuminated using a fibre-optic light, which delivered a photon flux density (PFD) of ~500 µmol m⁻² s⁻¹ at the leaf surface that was sufficient to saturate photosynthesis in *A. antarctica*. This light level was chosen to represent peak irradiances previously recorded in *A. antarctica* meadows (A. D. Irving, unpublished data; Bryars *et al.*, 2011), but below levels shown to have a photoinhibitory effect on *A. antarctica* and other co-occurring species (Masini *et al.*, 1995; Ralph *et al.*, 1998). Light between 380 and 710 nm was measured by positioning the fibre quantum sensor of a diving pulse amplitude modulated (PAM) fluorometer (Walz, Effeltrich, Germany) in the electrode chamber. The PAM fluorometer fibre quantum sensor was calibrated against a LI-COR quantum sensor (Li-192SA; Lincoln, NE, USA). The electrode chamber was maintained at a constant temperature of 20°C using a transparent recirculating water jacket and was constantly mixed using a magnetic stirrer.

Seagrass leaves were sealed in the chamber and covered with a black cloth to measure dark respiration. Respiration rates were allowed to stabilize for 3 min before recording. The chamber was then illuminated, and where necessary, the appropriate stock solutions (see below) were injected into the chamber using micro-syringes. Photosynthetic rates were allowed to stabilize for 2 min before recording. Photosynthetic and respiration rates were averaged over 3 min and are expressed on a chlorophyll basis (as micromoles of oxygen per gram of chlorophyll per minute).

A 20 mM stock solution of the CA inhibitor AZ (Sigma Aldrich) was prepared in 50 mM sodium hydroxide (NaOH). A 1 M stock solution of TRIS (Sigma Aldrich) was prepared and the pH adjusted to ambient seawater pH (8.05). The buffer yielded a pH of 8.06 when injected into the electrode chamber. For AZ, 20 µl of stock solution was injected to achieve a final chamber concentration of 100 µM. For TRIS, 200 µl of stock solution was injected to achieve a final chamber concentration of 50 mM. Chamber conditions, including carbonate chemistry, are presented in Table 1a.

Effects of CO₂ on photosynthesis and growth

Juvenile *A. antarctica* seedlings were collected at the same depth and location as described above for adult plants and transported to an outdoor glasshouse at The University of Adelaide. Four individual seedlings were planted in sediment from the collection site in each of 12 transparent 2 litre microcosms (25 cm depth). Seagrasses were maintained at two CO₂ concentrations, representing current atmospheric CO₂ levels (~390 ppm) and forecasted future CO₂ levels (~900 ppm)

Table 1: Sea water chemistry in the oxygen electrode chamber during buffer/inhibitor experiments (a; see Figs 1 and S1), during 12 week growth experiments (b; see Figs 2a and 3) and reciprocal switch measurements (c; Fig. 2b)

	pH	A _T [µmol (kg sea water ⁻¹)]	Salinity (‰)	Temperature (°C)	Total CO ₂ [µmol (kg sea water ⁻¹)]	Partial pressure of CO ₂ (µatm)	HCO ₃ ⁻ [µmol (kg sea water ⁻¹)]	CO ₃ [µmol (kg sea water ⁻¹)]	CO ₂ [µmol (kg sea water ⁻¹)]
(a)									
Control	8.05 ± 0.01	2533 ± 49	38	20	2223	440	1981	228	14.0
TRIS	8.06 ± 0.01	2533 ± 49	38	20	2215	424	1968	234	13.5
AZ	8.08 ± 0.02	2533 ± 49	38	20	2204	404	1950	241	12.9
TRIS + AZ	8.06 ± 0.00	2533 ± 49	38	20	2211	418	1962	236	13.3
(b)									
L[CO ₂]	8.12 ± 0.002	2701 ± 27	40.3 ± 0.06	19.8 ± 0.01	2309	376	2011	285	11.8
H[CO ₂]	7.82 ± 0.003	2697 ± 35	40.0 ± 0.05	19.9 ± 0.01	2498	872	2309	161	27.5
(c)									
L[CO ₂]	8.15 ± 0.003	2648 ± 43	40.6 ± 1.49	20	2236	338	1930	295	10.6
H[CO ₂]	7.82 ± 0.007	2656 ± 48	40.3 ± 1.35	20	2454	849	2266	162	26.7

Abbreviations: AZ, acetazolamide; TRIS, tris(hydroxymethyl)aminomethane. L[CO₂] = low CO₂, H[CO₂] = high CO₂. Measurements of pH, total alkalinity (A_T), salinity and temperature (fixed at 20°C during photosynthetic trials) were used to calculate carbonate chemistry. Salinity was fixed at 38‰ during inhibitor experiments.

under emission scenario A1FI for the year 2100 (Meehl *et al.*, 2007), hereafter referred to as low CO₂ (L[CO₂]) and high CO₂ (H[CO₂]), respectively. Carbon dioxide was enriched by aerating the microcosms with a combination of ambient air and pure CO₂ using a two-channel gas mixer (Columbus Instruments, Columbus, OH, USA), which is equivalent to maintaining treatments within an enriched CO₂ atmosphere. Aeration helps to accelerate the diffusion of CO₂ into seawater, also minimizing any variation driven by plant photosynthesis or respiration. The accuracy of the mixed gas aerating treatments can vary slightly dependent on ambient atmospheric conditions, but any such changes are minor and occur naturally whenever ambient air is used for aeration; ambient CO₂ treatments are also subject to such natural fluctuations. Experimental conditions, including carbonate chemistry, in experimental microcosms were monitored throughout the experiment (Table 1b). Seventy per cent of seawater was replaced twice weekly to maintain salinity and alkalinity.

Microcosms were in an outdoor glass house and they were shaded from full surface irradiance using a combination of 50 and 70% shade cloth. HOBO® waterproof light loggers (Onset, Wareham, MA, USA) were used to record light in lux, which was then converted to give an approximation of photosynthetically active radiation using the constant for natural sunlight (1 lux = 54 $\mu\text{mol m}^{-2} \text{s}^{-1}$) recommended by Thimijan and Heins (1983). Average daily PFD was 42.5 $\mu\text{mol m}^{-2} \text{s}^{-1}$ over a 12 h photoperiod, with an average daily maximum of 111.3 \pm 11.7 $\mu\text{mol m}^{-2} \text{s}^{-1}$ recorded at 13.00 h. Shading was designed to replicate closely the light conditions recorded during a monitoring study on the Adelaide metropolitan coast, which found average daily PFD of ~44 $\mu\text{mol m}^{-2} \text{s}^{-1}$ over a 12 h photoperiod (Irving, 2009). One specific 40 day light monitoring deployment at ~2 m (low tide) on the Adelaide metropolitan coast found averagedaily maximal PFD of 155.95 \pm 20.54 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (A. D. Irving, unpublished data), which is comparable to the peak levels recorded in our acclimation study. The daily maximum in this monitoring study ranged from 7 to 453 $\mu\text{mol m}^{-2} \text{s}^{-1}$, highlighting the dynamic nature of the light climate in the shallow sub-tidal near-shore zone, where the swell direction, wind direction, rainfall, runoff and topography can all influence turbidity and thus light attenuation.

Chlorophyll fluorescence

A submersible diving PAM fluorometer (Walz, Effeltrich, Germany) was used to record rapid light curves (RLCs) and maximal quantum yield (QY_{max}). Seagrasses were measured *in situ* in microcosms at L[CO₂] and H[CO₂] after 12 weeks. The photon flux densities used during the RLCs were 0, 18, 37, 62, 92, 125, 186, 256 and 400 $\mu\text{mol m}^{-2} \text{s}^{-1}$, each of 10 s duration, followed by a saturating pulse of light, to record the effective quantum yield (Φ_{PSII}). All RLC measurements were taken between 11.00 and 12.30 h. The light absorbance of every sample leaf was measured by placing a quantum sensor directly behind the leaf and recording the percentage of ambient light that was absorbed. The electron transport rate (ETR) was calculated as follows: $\text{ETR} = \Phi_{\text{PSII}} \times \text{PFD} \times \text{leaf}$

absorbance \times 0.5. For quantitative comparisons of RLCs [i.e. initial slope (α), maximal electron transport rate (ETR_{max}) and light saturation (I_k)], the ETR data were fitted with a least-squares non-linear regression curve based on the exponential difference equation from Platt *et al.* (1980) using the Microsoft Excel (Microsoft, Redmond, WA, USA) solvers provided by Ritchie (2008). Light saturation was calculated as $\text{ETR}_{\text{max}} / \alpha$, where α is the initial slope of the non-inhibited section of the fitted curve. The value of QY_{max} was determined from pre-dawn fluorescence measurements.

Photosynthesis and respiration

Following 12 weeks at experimental CO₂ levels, photosynthesis and respiration rates were measured as described above, but without the addition of buffers and inhibitors. Reciprocal measurements were made, where individuals from each CO₂ treatment were measured at both low and high CO₂, to determine whether physiological acclimation had occurred over the course of the experiment (see Table 1c for a description of chamber conditions). Electron transport rates at 400 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and maximal gross photosynthetic rates were then used to calculate ETR-to-O₂ ratios (i.e. moles of electrons per mole of O₂ evolved). This was done for L[CO₂] and H[CO₂] plants tested in their respective growth treatments.

Growth

Growth parameters were measured after 12 weeks. Leaves and shoots (above ground) were separated from roots and rhizomes (below ground) and dried for 48 h at 60°C to measure dry mass (DM). Above-ground components were then washed in 5% HCl to remove any calcified epiphytes, redried for 48 h at 60°C and reweighed.

Leaf initiation was recorded by trimming the corner of the second youngest leaf on the highest leaf head of each seedling at the commencement of the growth experiment. On juveniles, this method of monitoring leaf growth is less obtrusive than sheath marking or using leaf ties, which can impact meristem growth and plant buoyancy (O. W. Burnell, personal observation). The total number of leaves on each leaf head was recorded before planting and again after 12 weeks to calculate the change in total leaf number, which accounted for different rates of leaf shedding and production from any new meristems.

Chlorophyll determination

Chlorophyll content was measured to enable expression of photosynthesis and respiration per unit of chlorophyll. Following oxygen electrode measurements, seagrass material was removed from the chamber, rinsed in milliQ water, and immediately frozen in liquid nitrogen. Samples were then stored at -80°C until extractions were conducted. Chlorophyll concentrations were determined using the methodology of Granger and Lizumi (2001). Briefly, leaves were soaked in 1 ml of 100% acetone in dark, refrigerated conditions for 1 h. Seagrass and acetone were then transferred to a chilled

mortar and pestle and ground using an additional 1 ml of 86.7% acetone and acid washed sand until they were reduced to a flocculent slurry. Two drops of 1% MgCO_3 were added before diluting to 4 ml with 86.7% acetone to achieve a final sample concentration of 90% acetone. Samples were then centrifuged for 5 min at 670.8 g of relative centrifugal force. Absorbance of the supernatant was measured in a spectrophotometer (Jenway 6405, Stone, Staffordshire, UK) at 647, 664 and 725 nm. Chlorophyll concentrations were calculated using equations from Jeffrey and Humphrey (1975).

Results

It was evident that *A. antarctica* uses HCO_3^- pathways for C_i acquisition, because light-saturated photosynthesis declined by ~54% when AZ and TRIS were added in combination (Fig. 1). The sensitivity of *A. antarctica* to both AZ and the TRIS buffer was consistent with a System A + B mode of C_i acquisition (Beer *et al.*, 2002). The lack of an interactive effect in the two-factor ANOVA provided no support for a System C mode of acquisition (Table 2a). Inorganic carbon uptake in the congener *A. griffithii* followed a similar pattern to *A. antarctica*, in that photosynthesis declined by ~57% when AZ and TRIS were added in combination; however, an interaction between AZ and TRIS in the two-factor ANOVA provided better support for a System C mode of acquisition ($F_{1,28} = 7.88$, $P = 0.008$; Supplementary material Fig. S1a and Supplementary material Table S1). In contrast, the species *P. sinuosa* was insensitive to the addition of the TRIS buffer and only moderately affected by the inhibitor AZ, highlighting a lesser dependency on HCO_3^- pathways for photosynthesis (Supplementary material Fig. S1b and Supplementary material Table S1). Dark respiration did not differ between experimental treatments for any of the three species, and therefore, analyses are not presented (Fig. 1 and Supplementary material Fig. S1). For rates of photosynthesis and respiration per gram of fresh mass, see Supplementary material Table S2.

Forecasted CO_2 increased both the ETR and photosynthesis of *A. antarctica* when measured in growth conditions (Fig. 2 and Table 2b). The mean ETR_{max} from the fitted regression curves was $7.39 \pm 0.80 \mu\text{mol electrons m}^{-2} \text{s}^{-1}$ for $\text{L}[\text{CO}_2]$ plants, compared with $10.04 \pm 0.55 \mu\text{mol electrons m}^{-2} \text{s}^{-1}$ for $\text{H}[\text{CO}_2]$ plants ($F_{1,10} = 7.561$, $P = 0.028$). The onset of light saturation (I_k) for ETR occurred at $27.12 \pm 2.60 \mu\text{mol m}^{-2} \text{s}^{-1}$ for $\text{L}[\text{CO}_2]$, compared with $35.79 \pm 2.82 \mu\text{mol m}^{-2} \text{s}^{-1}$ for $\text{H}[\text{CO}_2]$ plants ($F_{1,10} = 5.13$, $P = 0.049$). In contrast, there was no difference in the initial slope (α) of ETR (0.28 ± 0.03 and 0.29 ± 0.02) or QY_{max} (0.656 ± 0.033 and 0.683 ± 0.017) for $\text{L}[\text{CO}_2]$ and $\text{H}[\text{CO}_2]$, respectively, thus analyses are not presented.

When plants were reciprocally switched between CO_2 treatments, measurement CO_2 had a significant effect on photosynthesis of both $\text{L}[\text{CO}_2]$ and $\text{H}[\text{CO}_2]$ grown plants (Fig. 2b and Table 2b). In contrast, there was no effect of growth CO_2 ; that is, seagrasses photosynthesis was responding primarily to measurement conditions (i.e. dissolved CO_2), rather than exhibit-




Figure 1: The effects of inhibitors [tris(hydroxymethyl)aminomethane (TRIS), acetazolamide (AZ) and TRIS + AZ] on rates of photosynthesis and respiration in *Amphibolis antarctica* measured in seawater. Bars are means \pm SEM ($n = 8$).

Table 2: ANOVA for the effects on photosynthesis in *A. antarctica* of buffer/inhibitor (a; in seawater/control, TRIS, AZ and TRIS + AZ; see Fig. 1) and reciprocal switch (b; i.e. plants were grown at either $\text{L}[\text{CO}_2]$ or $\text{H}[\text{CO}_2]$ and then measured at both $\text{L}[\text{CO}_2]$ and $\text{H}[\text{CO}_2]$; see Fig. 2b)

Source	d.f.	MSE	F	P-Value
(a) Buffer/inhibitor, <i>A. antarctica</i>				
TRIS	1	24 540	6.36	0.016
AZ	1	17 517	4.54	0.035
TRIS \times AZ	1	2157	0.56	0.467
Residual	28	3858		
(b) Reciprocal switch, <i>A. antarctica</i>				
Measurement	1	578 810	9.66	0.005
Growth	1	23 996	0.40	0.574
Measurement \times growth	1	90 060	1.50	0.252
Residual	16	59 916		

Abbreviations: d.f., degrees of freedom; F, f-statistic; MSE, mean squared error. No significant differences were found between respiration rates; therefore, analyses are not presented.

ing significant physiological acclimation to growth in the different CO_2 treatments. Molar ratios between ETR and gross oxygen evolution based on leaf area were $9.51 \text{ mol electrons (mol O}_2\text{)}^{-1}$ for $\text{L}[\text{CO}_2]$ plants and $8.08 \text{ mol electrons (mol O}_2\text{)}^{-1}$ for $\text{H}[\text{CO}_2]$ plants.

Growth of juvenile seagrass was greater at $\text{H}[\text{CO}_2]$, with total biomass ($F_{1,10} = 4.26$, $P = 0.050$), below-ground biomass ($F_{1,10} = 5.42$, $P = 0.043$) and change in leaf number




Figure 2: (a) Rapid light curves of electron transport rate for *A. antarctica* grown and measured at 390 (L[CO₂]) or 900 ppm (H[CO₂]). Filled circles indicate L[CO₂] and filled triangles H[CO₂]. Data points are means \pm SEM ($n = 6$). (b) Rates of photosynthesis and respiration in *A. antarctica* measured at either low (shaded bars) or high CO₂ (open bars) and grown at either low (G L[CO₂]) or high CO₂ (G H[CO₂]). Bars are means \pm SEM ($n = 5$).

($F_{1,10} = 5.40$, $P = 0.047$) significantly higher than for L[CO₂] grown plants. However, there was no significant difference in above-ground biomass ($F_{1,10} = 2.79$, $P = 0.139$) or leaf initiation from each individual meristem ($F_{1,10} = 3.96$, $P = 0.087$; Fig. 3a–e). Significant correlations between below-ground biomass and each of the above-ground growth parameters suggested that interdependent positive relationships existed between above- and below-ground growth (Fig. 4a–c).

Discussion

The three seagrass species tested were sensitive to the inhibition of HCO₃⁻ uptake mechanisms, indicating that at or under contemporary CO₂ concentrations they are reliant on energetically costly C_i acquisition. Seagrasses are commonly known to use these HCO₃⁻ pathways for photosynthesis (Beer *et al.*,

1977; Millhouse and Strother, 1986; James and Larkum, 1996; Beardall, *et al.*, 1998; Hellblom *et al.*, 2001); however, we have a limited understanding of how they regulate these mechanisms when faced with environmental variation (Larkum, *et al.*, 2006) and any implications of this for long-term growth. Any increase in the proportion of direct CO₂ uptake relative to energetically costly HCO₃⁻ acquisition could benefit their carbon balance and thus growth rate (Beardall and Giordano, 2002; Raven *et al.*, 2011; Koch *et al.*, 2013). In accordance with such predictions, we found that increased photosynthesis and growth of *A. antarctica* was accompanied by lower ETR-to-O₂ ratios, indicating that not only growth but also photosynthetic efficiency could increase at forecasted CO₂ concentrations.

Greater availability of CO₂ increased the ETR_{max} in *A. antarctica*, as has been reported in other seagrass species (Jiang *et al.*, 2010; Alexandre *et al.*, 2012). However, changes in ETR at forecasted CO₂ concentrations were small relative to increases in light-saturated photosynthesis measured by O₂ evolution (i.e. 1.4-fold for ETR vs. 2.1-fold for O₂ evolution). This difference translated to lower ETR-to-O₂ ratios calculated for H[CO₂] plants when compared with L[CO₂] plants (i.e. 8.08 vs. 9.51 mol electrons (mol O₂)⁻¹, respectively). This molar ratio is also commonly referred to in its inverse form, which would equate to 0.105 and 0.124 mol O₂ (mol electrons)⁻¹ for L[CO₂] and H[CO₂] plants, respectively. While these values differ from the theoretical maximum of 0.25 mol O₂ (mol electrons)⁻¹ that is based on the minimal number of electrons needed to produce a given amount of O₂ during photosynthesis (i.e. 4 mol electrons (mol O₂)⁻¹; Walker, 1987), recorded values for different seagrass species are known to vary widely (Beer *et al.*, 1998; Silva *et al.*, 2009). Given that ETR-to-O₂ ratios were measured at relatively high light intensities, it is likely that processes such as photorespiration contributed to these high ETR-to-O₂ ratios. Photorespiration increases in conditions of low C_i availability, where plants fix O₂ rather than CO₂ (Black *et al.* 1976; Buapet *et al.* 2013). It is well known that the usually linear relationship between ETR and O₂ evolution can deteriorate at high light intensities, at which photorespiration or other processes that can act as sinks for electrons increase (Beer *et al.*, 1998; Carr and Bjork, 2003; Silva and Santos 2004).

Photorespiration may also have contributed to the treatment differences that we observed between ETR-to-O₂ ratios for L[CO₂] and H[CO₂] plants. Importantly, the finding that L[CO₂] plants transported more electrons per mole of O₂ evolved than H[CO₂] plants could indicate greater photorespiration where plants were CO₂ limited. In such circumstances, photorespiration is likely to have acted as a greater sink for electrons measured using fluorescence in L[CO₂] plants. Thus, O₂ evolution measurements may provide a better estimate of changing photosynthetic efficiency than ETR at higher irradiances, because photorespiration in CO₂-limited individuals remains undetected by fluorescence measurements. However, photorespiration is generally low in seagrasses with effective carbon concentrating mechanisms that maintain high concentrations of CO₂ around rubisco (Beer, 1989; Touchette




Figure 3: Effect of growth [CO₂] on total dry mass (DM; **a**), above-ground DM (**b**), below-ground DM (**c**), leaf initiation (**d**) and change in leaves per plant (**e**) for *A. antarctica* grown in L[CO₂] and H[CO₂] conditions for 12 weeks. Bars are means ± SEM ($n = 6$).

and Burkholder, 2000). It is also possible that greater ETR-to-O₂ ratios at low CO₂ might be consistent with a greater proportion of electron flow supporting C_i-uptake mechanisms. However, this may be of greater consequence at lower irradiances, where plants are forced to partition limited energetic resources between photosynthesis and CO₂ acquisition.

Despite greater photosynthesis and growth at forecasted CO₂ concentrations, there was limited evidence to suggest any down-regulation of HCO₃⁻-uptake mechanisms in *A. antarctica* grown at H[CO₂], because when seagrasses were switched to L[CO₂] to measure photosynthesis, there was no effect of CO₂ growth conditions. This suggests that, on transition to a different CO₂ concentration, the photosynthetic rate was determined by changing dissolved CO₂, rather than by different affinities for HCO₃⁻ acquired during the 3 month growth period. If H[CO₂]-grown plants were no longer able to acquire CO₂ from HCO₃⁻, much lower O₂ production would have

been expected when they were switched back to L[CO₂]. Shorter-term studies have likewise found that the relative level of HCO₃⁻ uptake by seagrass is often maintained across pH gradients (James and Larkum, 1996; Campbell and Fourqurean, 2013b). Thus, it appears that changes in growth and photosynthesis may have occurred based primarily on greater passive uptake of CO₂, not on down-regulation of HCO₃⁻-uptake mechanisms. Alternatively, down-regulation may have been masked if plants rapidly modified HCO₃⁻-acquisition rates during the brief reciprocal CO₂ transitions, although this appears unlikely. While rapid re-establishment (2–5 min) of HCO₃⁻ acquisition can occur following dark-to-light transitions (Carr and Axelsson, 2008), the time for activation and deactivation for HCO₃⁻ mechanisms with CO₂ transitions would appear to be much greater. Studies of lower order producers (i.e. cyanobacteria and eukaryotic algae) and freshwater angiosperms show hours to days for full re-activation and down-regulation following CO₂ transitions




Figure 4: Correlations between below-ground biomass (expressed as DM per seagrass) and above-ground growth parameters (expressed as DM biomass per seagrass, leaf initiation per meristem and change in leaves per seagrass) for *A. antarctica*. Data points are the average in each microcosm ($n = 6$).

(Spalding and Ogren, 1982; Falk and Palmqvist, 1992; Matsuda and Colman, 1995; Maberly and Madsen, 2002).

The greater increase in below-ground biomass at $H[CO_2]$ suggests that *A. antarctica* may preferentially allocate resources to roots and rhizomes for growth and energy storage. These findings are consistent with studies of natural and *in situ* CO_2 enrichment that indicate disproportionate accumulation of below-ground biomass in other seagrasses,

relative to above-ground tissue (Hall-Spencer *et al.*, 2008; Russell *et al.*, 2013; Campbell and Fourqurean, 2013a). However, the significant correlation between below-ground biomass and the three above-ground growth parameters could suggest that the former simply responds more rapidly to CO_2 enrichment, and this could later translate to changes in above-ground parameters. This translation to greater above-ground biomass would appear possible given that seagrass at $H[CO_2]$ also showed a greater increase in leaf number; nonetheless, with potentially greater carbon fixation per leaf area at enriched CO_2 levels, there may be limited need for large increases in above-ground biomass.

The increase in below-ground biomass could offer seagrasses greater resistance to environmental perturbations in an era when anthropogenic influences have been reported to be almost exclusively negative (Orth *et al.*, 2006). Greater below-ground energetic resources could sustain juveniles during periods of physiological stress, including reduced light availability (Burke *et al.*, 1996; Mackey *et al.*, 2007; Ooi *et al.*, 2011), high temperature (Walker and Cambridge, 1995; Niu *et al.*, 2012; Koch *et al.*, 2013) and grazing pressure (Hughes *et al.*, 2004). Preferential allocation to below-ground resources that are protected from herbivores might be particularly beneficial given that enriched CO_2 conditions can result in seagrass that is more palatable to grazers owing to a reduction of phenolic deterrents (Arnold *et al.*, 2012). Likewise, faster establishment of below-ground structures could lower the vulnerability of seagrass to physical disturbances, such as wave action and bioturbation/erosion (Townsend and Fonseca, 1998; Bastyan and Cambridge, 2008). An increase in recruitment success resulting from reproduction could be particularly beneficial, given that most seagrasses are heavily reliant on clonal growth for meadow expansion, with low rates of seed or seedling survival and natural recolonization (Kirkman and Kuo, 1990; Irving, 2013). Any prolonged increase in seagrass growth, however, will require other favourable conditions, such as adequate light, nutrients and facilitative interactions with other biota (i.e. grazer activity and epiphyte abundance), all of which are known to limit seagrass biomass where CO_2 is enriched (O. W. Burnell, unpublished data; Palacios and Zimmerman, 2007; Alexandre, *et al.*, 2012; Burnell *et al.*, 2013).

As historically high CO_2 levels potentially aided the initial transition of angiosperms into the aquatic realm (Beer and Koch, 1996; Beardall, *et al.*, 1998), forecasted increases in CO_2 could return such benefits, by increasing the availability of dissolved CO_2 . We found that *A. antarctica* has a contemporary reliance on HCO_3^- -uptake mechanisms, as well as greater photosynthesis and growth with prolonged acclimation to forecasted CO_2 conditions. Importantly, any lasting increase in productivity and growth could enhance the host of ecosystem benefits provided by seagrass meadows.

Supplementary material

Supplementary material is available at *Conservation Physiology* online.

Acknowledgements

We would like to thank members of the Plant Ecophysiology Laboratory at The University of Adelaide for welcoming us into their laboratory, as well as Julie Francis for assistance with oxygen electrodes.

Funding

S.D.C. and B.D.R. were funded by an ARC grant and S.D.C. by an ARC Future Fellowship.

References

- Alexandre A, Silva J, Buapet P, Björk M, Santos R (2012) Effects of CO₂ enrichment on photosynthesis, growth, and nitrogen metabolism of the seagrass *Zostera noltii*. *Ecol Evol* 2: 2620–2630.
- Arnold T, Mealey C, Leahey H, Miller AW, Hall-Spencer JM, Milazzo M, Maers K (2012) Ocean acidification and the loss of phenolic substances in marine plants. *PLoS ONE* 7: e35107.
- Badger MR, Andrews TJ (1982) Photosynthesis and inorganic carbon usage by the marine cyanobacterium, *Synechococcus* sp. *Plant Physiol* 70: 517–523.
- Bastyan GR, Cambridge ML (2008) Transplantation as a method for restoring the seagrass *Posidonia australis*. *Estuar Coast Shelf Sci* 79: 289–299.
- Beardall J, Giordano M (2002) Ecological implications of microalgal and cyanobacterial CO₂ concentrating mechanisms, and their regulation. *Funct Plant Biol* 29: 335–347.
- Beardall J, Beer S, Raven JA (1998) Biodiversity of marine plants in an era of climate change: some predictions based on physiological performance. *Bot Marina* 41: 113–123.
- Beer S (1989) Photosynthesis and photorespiration of marine angiosperms. *Aquat Bot* 34: 153–166.
- Beer S, Koch E (1996) Photosynthesis of marine macroalgae and seagrasses in globally changing CO₂ environments. *Mar Ecol Prog Ser* 141: 199–204.
- Beer S, Eshel A, Waisel Y (1977) Carbon metabolism in seagrasses. I. Utilization of exogenous inorganic carbon species in photosynthesis. *J Exp Bot* 28: 1180–1189.
- Beer S, Vilenkin B, Weil A, Veste M, Susel L, Eshel A (1998) Measuring photosynthetic rates in seagrasses by pulse amplitude modulated (PAM) fluorometry. *Mar Ecol Prog Ser* 174: 293–300.
- Beer S, Björk M, Hellblom F, Axelsson L (2002) Inorganic carbon utilization in marine angiosperms (seagrasses). *Funct Plant Biol* 29: 349–354.
- Berner RA, Kothavala Z (2001) Geocarb III: a revised model of atmospheric CO₂ over phanerozoic time. *Am J Sci* 301: 182–204.
- Black CC, Burris JE, Everson RG (1976) The influence of oxygen concentration on photosynthesis in marine plants. *Aust J Plant Physiol* 3: 81–86.
- Brueggeman AJ, Gangadharaiah DS, Cserhati MF, Casero D, Weeks DP, Ladunga I (2012) Activation of the carbon concentrating mechanism by CO₂ deprivation coincides with massive transcriptional restructuring in *Chlamydomonas reinhardtii*. *Plant Cell* 24: 1860–1875.
- Bryars S, Collings G, Miller D (2011) Nutrient exposure causes epiphytic changes and coincident declines in two temperate Australian seagrasses. *Mar Ecol Prog Ser* 441: 89–103.
- Buapet P, Rasmusson LM, Gullström M, Björk M (2013) Photorespiration and carbon limitation determine productivity in temperate seagrasses. *PLoS ONE* 8: e83804.
- Burke MK, Dennison WC, Moore KA (1996) Non-structural carbohydrate reserves of eelgrass *Zostera marina*. *Mar Ecol Prog Ser* 137: 195–201.
- Burnell OW, Russell BD, Irving AD, Connell SD (2013) Eutrophication off-sets increased sea urchin grazing on seagrass caused by ocean warming and acidification. *Mar Ecol Prog Ser* 485: 37–46.
- Campbell JE, Fourqurean JW (2013a) Effects of in situ CO₂ enrichment on the structural and chemical characteristics of the seagrass *Thalassia testudinum*. *Mar Biol* 160: 1465–1475.
- Campbell JE, Fourqurean JW (2013b) Mechanisms of bicarbonate use influence the photosynthetic carbon dioxide sensitivity of tropical seagrasses. *Limnol Oceanogr* 58: 839–848.
- Carr H, Björk M (2003) A methodological comparison of photosynthetic oxygen evolution and estimated electron transport rate in tropical *Ulva* (Chlorophyceae) species under different light and inorganic carbon conditions. *J Phycol* 39: 1125–1131.
- Carr H, Axelsson L (2008) Photosynthetic utilization of bicarbonate in *Zostera marina* is reduced by inhibitors of mitochondrial ATPase and electron transport. *Plant Physiol* 147: 879–885.
- Costanza R, D'Arge R, de Groot R, Farber S, Grasso M, Hannon B, Limburg K, Naeem S, O'Neill RV, Paruelo J *et al.* (1997) The value of the world's ecosystem services and natural capital. *Nature* 387: 253–260.
- Duarte CM (2002) The future of seagrass meadows. *Environ Conserv* 29: 192–206.
- Falk S, Palmqvist K (1992) Photosynthetic light utilization efficiency, photosystem II heterogeneity, and fluorescence quenching in *Chlamydomonas reinhardtii* during the induction of the CO₂-concentrating mechanism. *Plant Physiol* 100: 685–691.
- Fridlyand L, Kaplan A, Reinhold L (1996) Quantitative evaluation of the role of a putative CO₂-scavenging entity in the cyanobacterial CO₂-concentrating mechanism. *Biosystems* 37: 229–238.
- Granger S, Lizumi H (2001) Water quality measurement methods for seagrass habitat. In Short FT, Coles RG, eds, *Global Seagrass Research Methods*. Elsevier, Amsterdam, The Netherlands, pp 393–406.
- Hall-Spencer JM, Rodolfo-Metalpa R, Martin S, Ransome E, Fine M, Turner SM, Rowley SJ, Tedesco D, Buia MC (2008) Volcanic carbon dioxide vents show ecosystem effects of ocean acidification. *Nature* 454: 96–99.
- Hellblom F, Beer S, Björk M, Axelsson L (2001) A buffer sensitive inorganic carbon utilisation system in *Zostera marina*. *Aquat Bot* 69: 55–62.

- Hughes AR, Bando KJ, Rodriguez LF, Williams SL (2004) Relative effects of grazers and nutrients on seagrasses: a meta-analysis approach. *Mar Ecol Prog Ser* 282: 87–99.
- Invers O, Perez M, Romero J (1999) Bicarbonate utilization in seagrass photosynthesis: role of carbonic anhydrase in *Posidonia oceanica* (L) delile and *Cymodocea nodosa* (Ucria) Ascherson. *J Exp Mar Biol Ecol* 235: 125–133.
- Irving AD (2009) Seagrass rehabilitation in Adelaide's coastal waters VI. Refining techniques for the rehabilitation of *Amphibolis* spp. Final report prepared for the coastal management branch of the Department for Environment and Heritage SA, South Australian Research and Development Institute (Aquatic Sciences), Adelaide, South Australia.
- Irving AD (2013) A century of failure for habitat recovery. *Ecography* 36: 414–416.
- Irving AD, Connell SD, Russell BD (2011) Restoring coastal plants to improve global carbon storage: reaping what we sow. *PLoS ONE* 6: e18311.
- James PL, Larkum AWD (1996) Photosynthetic inorganic carbon acquisition of *Posidonia australis*. *Aquat Bot* 55: 149–157.
- Jeffrey SW, Humphrey GF (1975) New spectrophotometric equations for determining chlorophylls A, B, C1 and C2 in higher plants, algae and natural phytoplankton. *Biochemie und Physiologie der Pflanzen* 167: 191–194.
- Jiang ZJ, Huang XP, Zhang JP (2010) Effects of CO₂ enrichment on photosynthesis, growth, and biochemical composition of seagrass *Thalassia hemprichii* (Ehrenb.) Aschers. *J Integr Plant Biol* 52: 904–913.
- Kirkman H, Kuo J (1990) Pattern and process in southern Western Australian seagrasses. *Aquat Bot* 37: 367–382.
- Koch M, Bowes G, Ross C, Zhang XH (2013) Climate change and ocean acidification effects on seagrasses and marine macroalgae. *Glob Change Biol* 19: 103–132.
- Kübler JE, Raven JA (1995) The interaction between inorganic carbon acquisition and light supply in *Palmaria palmata* (Rhodophyta). *J Phycol* 31: 369–375.
- Larkum AWD, Roberts G, Kuo J, Strother S (1989) Gaseous movement in seagrasses. In Larkum AWD, McComb AJ, Shepherd SA, eds, *Biology of Seagrasses: A Treatise of the Biology of Seagrasses with a Special Reference to the Australian Region*. Elsevier, Amsterdam, The Netherlands, pp 686–722.
- Larkum AWD, Drew EA, Ralph PJ (2006) Photosynthesis and metabolism in seagrasses at the cellular level. In Larkum AWD, Orth RJ, Duarte CM, eds, *Seagrasses: Biology, Ecology and Conservation*. Springer, Dordrecht, The Netherlands, pp 323–345.
- Maberly SC, Madsen TV (2002) Freshwater angiosperm carbon concentrating mechanisms: processes and patterns. *Funct Plant Biol* 29: 393–405.
- Mackey P, Collier CJ, Lavery PS (2007) Effects of experimental reduction of light availability on the seagrass *Amphibolis griffithii*. *Mar Ecol Prog Ser* 342: 117–126.
- Masini RJ, Cary JL, Simpson CJ, McComb AJ (1995) Effects of light and temperature on the photosynthesis of temperate meadow-forming seagrasses in Western Australia. *Aquat Bot* 49: 239–254.
- Matsuda Y, Colman B (1995) Induction of CO₂ and bicarbonate transport in the green alga *Chlorella ellipsoidea*: 1. Time course of induction of the two systems. *Plant Physiol* 108: 247–252.
- Meehl GA, Stocker TF, Collins WD, Friedlingstein P, Gaye AT, Gregory JM, Kitoh A, Knutti R, Murphy JM, Noda A *et al.* (2007) Global climate projections. In Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL, eds, *Climate Change 2007: The Physical Science Basis. Contribution of the Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK and New York, NY, USA.
- Millhouse J, Strother S (1986) The effect of pH on the inorganic carbon source for photosynthesis in the seagrass *Zostera muelleri* Irmisch ex Aschers. *Aquat Bot* 24: 199–209.
- Niu S, Zhang P, Liu J, Guo D, Zhang X (2012) The effect of temperature on the survival, growth, photosynthesis, and respiration of young seedlings of eelgrass *Zostera marina* L. *Aquaculture* 350: 98–108.
- Ooi JLS, Kendrick GA, Van Niel KP (2011) Effects of sediment burial on tropical ruderal seagrasses are moderated by clonal integration. *Cont Shelf Res* 31: 1945–1954.
- Orth RJ, Carruthers TJB, Dennison WC, Duarte CM, Fourqurean JW, Heck KL, Hughes AR, Kendrick GA, Kenworthy WJ, Olyarnik S *et al.* (2006) A global crisis for seagrass ecosystems. *Bioscience* 56: 987–996.
- Palacios SL, Zimmerman RC (2007) Response of eelgrass *Zostera marina* to CO₂ enrichment: possible impacts of climate change and potential for remediation of coastal habitats. *Mar Ecol Prog Ser* 344: 1–13.
- Platt T, Gallegos CL, Harrison WG (1980) Photoinhibition of photosynthesis in natural assemblages of marine phytoplankton. *J Mar Res* 38: 687–701.
- Ralph PJ, Gademann R, Dennison WC (1998) In situ seagrass photosynthesis measured using a submersible, pulse-amplitude modulated fluorometer. *Mar Biol* 132: 367–373.
- Raven JA, Lucas WJ (1985) Energy costs of carbon acquisition. In Lucas WJ, Berry JA, eds, *Inorganic Carbon Uptake by Aquatic Photosynthetic Organisms*. American Society of Plant Physiologists, Rockville, MD, USA, pp 305–324.
- Raven JA, Giordano M, Beardall J, Maberly SC (2011) Algal and aquatic plant carbon concentrating mechanisms in relation to environmental change. *Photosynth Res* 109: 281–296.
- Ritchie RJ (2008) Fitting light saturation curves measured using modulated fluorometry. *Photosynth Res* 96: 201–215.
- Russell BD, Connell SD, Uthicke S, Muehllehner N, Fabricius KE, Hall-Spencer JM (2013) Future seagrass beds: can increased productivity lead to increased carbon storage? *Mar Pollut Bull* 73: 463–469.

- Schwarz AM, Björk M, Buluda T, Mtolera H, Beer S (2000) Photosynthetic utilisation of carbon and light by two tropical seagrass species as measured *in situ*. *Mar Biol* 137: 755–761.
- Silva J, Santos R (2004) Can chlorophyll fluorescence be used to estimate photosynthetic production in the seagrass *Zostera noltii*? *J Exp Mar Biol Ecol* 307: 207–216.
- Silva J, Sharon Y, Santos R, Beer S (2009) Measuring seagrass photosynthesis: methods and applications. *Aquat Biol* 7: 127–141.
- Spalding MH, Ogren WL (1982) Photosynthesis is required for induction of the CO₂ concentrating system in *Chlamydomonas reinhardtii*. *FEBS Lett* 145: 41–44.
- Thimijan RW, Heins RD (1983) Photometric, radiometric, and quantum light units of measure: a review of procedures for interconversion. *Hortscience* 18: 818–822.
- Touchette BW, Burkholder JM (2000) Overview of the physiological ecology of carbon metabolism in seagrasses. *J Exp Mar Biol Ecol* 250: 169–205.
- Townsend EC, Fonseca MS (1998) Bioturbation as a potential mechanism influencing spatial heterogeneity of North Carolina seagrass beds. *Mar Ecol Prog Ser* 169: 123–132.
- Walker D (1987) *The Use of the Oxygen Electrode and Fluorescence Probes in Simple Measurements of Photosynthesis*. Oxygraphics, Sheffield, UK.
- Walker DI, Cambridge ML (1995) An experimental assessment of the temperature responses of two sympatric seagrasses, *Amphibolis antarctica* and *Amphibolis griffithii*, in relation to their biogeography. *Hydrobiologia* 302: 63–70.
- Waycott M, Duarte CM, Carruthers TJB, Orth RJ, Dennison WC, Olyarnik S, Calladine A, Fourqurean JW, Heck KL, Hughes AR *et al.* (2009) Accelerating loss of seagrasses across the globe threatens coastal ecosystems. *Proc Natl Acad Sci USA* 106: 12377–12381.
- Zimmerman RC, Kohrs DG, Steller DL, Alberte RS (1997) Impacts of CO₂ enrichment on productivity and light requirements of eelgrass. *Plant Physiol* 115: 599–607.