




Please cite the Published Version

Fatoye, Francis , Yeowell, G , Wright, JM and Gebrye, Tadesse  (2019) Clinical and cost effectiveness of physiotherapy interventions following total knee replacement - A systematic review and meta-analysis. In: ISPOR 2019, 18 May 2019 - 22 May 2019, New Orleans, USA.

DOI: <https://doi.org/10.1016/j.jval.2019.04.1132>

Publisher: Elsevier

Downloaded from: <https://e-space.mmu.ac.uk/625682/>

Usage rights:  [Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0](https://creativecommons.org/licenses/by-nc-nd/4.0/)

Additional Information: Abstract only. Published in Value in Health, Vol. 21, Iss. S1

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

PMS22

CLINICAL AND COST EFFECTIVENESS OF PHYSIOTHERAPY INTERVENTIONS FOLLOWING TOTAL KNEE REPLACEMENT- A SYSTEMATIC REVIEW AND META-ANALYSIS

Fatoye F,¹ Yeowell G,² Wright JM,² Gebrye T²

¹Manchester Metropolitan University, Manchester, LIN, United Kingdom,

²Manchester Metropolitan University, Manchester, United Kingdom

Objectives: Osteoarthritis (OA) of the knee is the leading cause of pain and disability. When pharmacological treatments fail, total knee replacement (TKR) is recommended. Physiotherapy interventions help to reduce pain, restore mobility and function, improve quality of life (QoL), and reduce healthcare costs following TKR. This review examined the clinical and cost effectiveness of physiotherapy interventions following TKR. **Methods:** A systematic review was conducted according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA). MEDLINE, CINAHL, AMED, DARE, HTA and NHS EED databases were searched up to September 2018. Search terms related to the clinical and cost effectiveness of physiotherapy interventions was used. Studies meeting the inclusion criteria were identified and key data were extracted. Quality assessment of the studies was performed by one reviewer and checked by a second. Random effect meta-analysis was conducted for pain, physical function and range of motion (ROM). **Results:** 467 studies were identified. Of these, seventeen studies were included; methodological quality of most studies was moderate. Physiotherapy interventions were more effective than control for physical function (-0.655 [95% CI $-1.262, -0.050$ $p = 0.034$]) and ROM (-0.354 [95% CI $-0.640, -0.112$ $p = 0.015$]) for a follow up of 2 or 3 months and six months, respectively. It was also found that physiotherapy interventions were not statistically significant compared to the control, for pain. Physiotherapy interventions were not more effective compared to control for pain, physical function and ROM for a 1 year follow up. No evidence on the pooled estimate of cost-effectiveness of physiotherapy interventions for people with TKR was found. **Conclusions:** The findings of this review suggest that physiotherapy interventions were effective for improving physical function and ROM for 2 or 3 months and six months follow up, respectively. Further research should examine the pooled cost-effectiveness estimate of physiotherapy interventions.

