


**Please cite the Published Version**

D'Cruze, Neil, Bates, Jodie, Assou, Délagnon, Ronfot, Delphine, Coulthard, Emma, Segniagbeto, Gabriel Hoinsoudé, Auliya, Mark, Megson, David  and Rowntree, Jennifer (2020) A preliminary assessment of bacteria in "ranché" ball pythons (*Python regius*), Togo, West Africa. *Nature Conservation*, 39. pp. 73-86. ISSN 1314-6947

**DOI:** <https://doi.org/10.3897/natureconservation.39.48599>

**Publisher:** Pensoft Publishers

**Version:** Published Version

**Downloaded from:** <https://e-space.mmu.ac.uk/625727/>

**Usage rights:**  [Creative Commons: Attribution 4.0](https://creativecommons.org/licenses/by/4.0/)

**Additional Information:** This is an open access article published in *Nature Conservation*, published by Pensoft Publishers, copyright The Author(s).

**Enquiries:**

If you have questions about this document, contact [openresearch@mmu.ac.uk](mailto:openresearch@mmu.ac.uk). Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

# A preliminary assessment of bacteria in “ranched” ball pythons (*Python regius*), Togo, West Africa

Neil D’Cruze<sup>1,2,3</sup>, Jodie Bates<sup>3</sup>, Délagnon Assou<sup>4,5</sup>, Delphine Ronfort<sup>6</sup>,  
Emma Coulthard<sup>3</sup>, Gabriel Hoinsoudé Segniagbeto<sup>4,5</sup>, Mark Auliya<sup>6,7</sup>,  
David Megson<sup>3</sup>, Jennifer Rowntree<sup>3</sup>

**1** World Animal Protection, 222 Gray’s Inn Rd., London WC1X 8HB, UK **2** Wildlife Conservation Research Unit, Department of Zoology, University of Oxford, Recanati-Kaplan Centre, Tubney House, Abingdon Road, Tubney, Abingdon OX13 5QL, UK **3** Ecology and Environment Research Centre, Department of Natural Sciences, Manchester Metropolitan University, All Saints Building, All Saints, Manchester M15 6BH, UK **4** Laboratory of Ecology and Ecotoxicology, Department of Zoology and Animal Biology, Faculty of Sciences, University of Lomé, BP 1515 Lomé-Togo **5** Togolese Society for Nature Conservation (AGBO-ZEGUE NGO). 06 BP 6057 Lomé-Togo **6** Zoological Research Museum Alexander Koenig, Department Herpetology, Adenauerallee 160, 53113 Bonn, Germany **7** Department of Conservation Biology, Helmholtz Centre for Environmental Research GmbH – UFZ, 04318 Leipzig, Germany

Corresponding author: Neil D’Cruze ([neil.dcruze@zoo.ox.ac.uk](mailto:neil.dcruze@zoo.ox.ac.uk))

Academic editor: F. Andreone | Received 19 November 2019 | Accepted 26 March 2020 | Published 13 May 2020

<http://zoobank.org/35CA7D69-9AAD-419B-A6A7-40E814A5FF77>

**Citation:** D’Cruze N, Bates J, Assou D, Ronfort D, Coulthard E, Segniagbeto GH, Auliya M, Megson D, Rowntree J (2020) A preliminary assessment of bacteria in “ranched” ball pythons (*Python regius*), Togo, West Africa. Nature Conservation 39: 73–86. <https://doi.org/10.3897/natureconservation.39.48599>

## Abstract

Captive reptiles are routinely identified as reservoirs of pathogenic bacteria and reports of reptile-associated infections relating to some species are well documented (e.g., salmonellosis). Currently, relatively little is known about the epidemiology and bacteria of ball pythons. We carried out a survey of ball python farms in Togo, West Africa to assess the presence of any potentially pathogenic bacterial taxa that have been identified in recent scientific literature relating to this species. The presence of bacteria belonging to the genera *Acinetobacter*, *Bacteroides*, *Citrobacter*, *Enterobacter*, *Lysobacter*, *Proteus*, *Pseudomonas*, *Staphylococcus*, and *Tsukamurella* in oral and cloacal samples taken from five individual ball pythons is of potential concern for horizontal transmission given that pathogenic species belonging to these genera have been previously documented. The presence of bacteria belonging to the genera *Clostridium*, *Escherichia*, *Moraxella*, and *Stenotrophomonas* in the oral and rectal samples taken from five mice used to feed ball

pythons suggests that they represent a potential reservoir of infection for wild caught ball pythons and their progeny. Furthermore, possible sources of environmental contamination include other captive amphibians, birds, reptiles and mammals, as well as free ranging birds and small mammals. Additional surveillance of ball pythons in the wild and in captivity at python farms in West Africa will shed light on whether or not this type of commercial activity is increasing pathogen exposure and lowering barriers to transmission. Meanwhile, as a precautionary measure, it is recommended that python farms should immediately establish biosecurity and disease surveillance practices to minimize potential horizontal and vertical bacterial transfer.

### Keywords

ball python, *Python regius*, reptile, wildlife trade, zoonosis

## Introduction

Global demand for reptiles as exotic pets is a relatively recent phenomenon (Mitchell 2009). Their popularity, however, has risen to the extent that they are now thought to represent the second most species-rich vertebrate class (after birds) in the international exotic pet trade (Bush et al. 2014). Reptiles are particularly rife in European and North American markets (Auliya 2003; Jensen et al. 2018), with conservative estimates of c. 0.7 million individuals being kept in the UK and 9.4 million in the USA, respectively (PFMA 2017; APPA 2019).

Global trade in wildlife (whether it legal or illegal) has also been cited as a disease transmission mechanism of growing concern in recent decades (Smith et al. 2009; Can et al. 2019). Specifically, these concerns relate to how pathogens are spread when humans capture wild animals from their natural habitats, transport them by land, sea and air and trade them dead or alive in different parts of a country or the world (e.g., Morens et al. 2004; Karesh et al. 2005; OIE 2017).

Captive reptiles are routinely identified as reservoirs of pathogenic bacteria and reports of reptile-associated infections for some species are well documented, such as salmonellosis (Arena et al. 2012; Bošnjak et al. 2016; Green et al. 2020). Several studies have investigated and highlighted the potential for horizontal and vertical transfer of disease at commercial captive breeding operations [e.g., Green iguanas (*Iguana iguana*) (Mitchell et al. 1999; Mitchell and Shane 2000) and Green sea turtles (*Chelonia mydas*) (Warwick et al. 2013)]. In some scenarios reptile-associated infections can spread to humans who have had direct or indirect contact with pet snakes and feeder rodents (used as reptile food) before their illnesses occurred [e.g., Canada in late 2019; (Government of Canada 2019)].

The ball python (*Python regius*), a species native to western and central Africa, is being exported in relatively large numbers [1,657,814 live individuals since 1978 (Convention on International Trade in Endangered Species of Wild Fauna and Flora [CITES] Trade Database; <https://trade.cites.org>)]. In fact, it is the single most traded CITES listed species (currently under CITES Appendix II) that is legally exported

alive from Africa (D'Cruze et al. 2020). Much of this international trade can be traced back to a number of python “farms” that are in operation across West Africa, most notably in Benin, Ghana and Togo (Robinson et al. 2015).

Since c. 1996, these python farms have been engaged in “ranching” (UNEP 2019), which refers to *rearing, in a controlled environment, snakes taken as eggs or juveniles from the wild, where they would otherwise have had a low probability of surviving to adulthood* (CITES 2019), and releasing a proportion back into the wild (Ineich 2006). Additionally, gravid females are also collected, and after laying their eggs in captivity are released back into the wild (Ineich 2006).

Recent studies have confirmed that the wild capture of ball pythons (for the export of specimens as “ranching” individuals the CITES source code “R” is used) often involves the removal of snakes from rodent burrows and live transport in sacks filled with other reptiles (D'Cruze et al. 2020). Once at farms, the snakes are reportedly housed separately, but they can also be housed at times in overcrowded enclosures in rooms that are filled with many other reptile species (D'Cruze et al. 2020). Mature ball pythons are typically fed live mice that are sourced from breeders or housed, or even bred, on site at farms.

Despite the international scope, large scale, and national wild release component of ball python “ranching” in Togo, there has been no current research focused on the epidemiology of this commercial trade activity. Therefore, we aimed to carry out an initial review, using amplicon sequence variants (ASVs) methods, to determine the presence of any potentially pathogenic genera of bacteria present in ball pythons and the live mice used as their food. We hope our findings will inform biosecurity surveillance practices to minimize potential horizontal and vertical transfer of zoonotic diseases.

## Methods

### Literature review

We conducted a systematic review of the scientific literature featured in PubMed, Scopus and Web of Science, from 2009–2019 to identify bacteria that are known to have affected the well-being of ball pythons. The following search terms were used (disease, pathogen, bacteria, bacterial). Each search term was employed with the Boolean operator “AND”, with three additional terms (ball python, royal python, *Python regius*).

### Laboratory analysis

A total of 20 dry swab samples were taken from five snakes and five mice at a python farm in Togo in September 2019 (Fig. 1, Suppl. material 1). Two swab samples were taken from each animal, one each from oral (both snake and mice), cloacal (snakes only) and rectal (mice only) orifices. Swabs were immediately transferred into 2 ml

screwcap microcentrifuge tubes containing approximately 600 µl of DNA/RNA Shield and stored at -20 °C until transport to the UK.

Prior to DNA extraction, samples (swab and reagent) were transferred into a fresh 2 ml screwcap microcentrifuge bead-beating tube, containing approximately 0.06 g of 0.1 mm glass-silica beads (Thistle Scientific, Glasgow), and vortexed twice for 30 seconds. The swabs were then discarded and the supernatant/liquid portion of the sample transferred to a 1.5 ml tube containing 274 µl polyethylene glycol (6000) and 141 µl 5M sodium chloride and incubated at 5 °C for 15–45 minutes. DNA was extracted using a modified phenol-chloroform method (Rogers et al. 2013) with reduced reagent volumes (200 µl each of molecular biology grade water, phenol, phenol-chloroform and ammonium-acetate: isopropanol compared to the 500 µl used previously). DNA pellets were suspended in 10 µl of molecular biology grade water and stored at -20 °C.

For PCR, 16S rRNA gene amplicons were generated following the Illumina two-step protocol (Illumina 2019) using primers designed by Caporaso et al. (2012) for the first step PCR and Nextera XT Index primers (Illumina, USA) for the second step. PCR conditions are shown in Appendix I. Amplicon clean-up was performed using AMPure XP beads (Beckman Coulter, UK), and normalised using a 96-well Sequal-Prep Normalisation Plate (Thermo Fisher Scientific, USA). Amplicons were sequenced using a 300-cycle MiSeq Reagent Micro Kit v2 (Illumina, USA) with a read length of  $2 \times 150$  bp – on the MiSeq platform.

Sequence quality of the top six samples was visually assessed within R (R Core Team 2019) using the DADA2 R package (Callahan et al. 2016), which was used for taxonomic assignment in combination with the Genome Taxonomy Database (GTDB) (Parks et al. 2018; Chaumeil et al. 2019) using  $1 \times 150$  bp forward read sequences. Raw reads were processed in accordance with the DADA2 Pipeline Tutorial (1.12) (Callahan 2019). In addition to using the standard filtering parameters for trimming, an extra step was added to ensure only amplicons of the expected length were included (i.e., with a minimum length of 148 and a maximum length of 151 bp). The resultant ASV and taxonomy csv files were combined to make a single database, which was further trimmed in Excel (Microsoft, USA). Prior to the assignment of presumptive genus-level classification for the ASVs, NA values for taxonomic levels at family level and above were removed. NA values at the genus level were kept and included as part of the 'other' genus category. Species level assignment was not possible due to the short length of the targeted 16S rRNA region, which may be indistinguishable among species and / or strains (Bulman et al. 2018; Osawa et al. 2015).

Bacterial genera that had been reported in the published scientific literature were identified using the search function within Excel. The assigned identity was confirmed using nucleotide BLAST searches against the 16S rRNA sequences (Bacteria and Archaea) coupled with megablast (highly similar sequences). Sequence read values for each genus were combined to create a stacked bar chart. No cut-off values for reads per sample were applied. The overall relative abundance for some genera were calculated by converting the number of reads for ASVs that have been assigned to a particular genus to a percentage relative to the total number of reads (derived using ASVs assigned genera in addition to the 'other' category).

## Results

The literature review identified 29 different species of bacteria across 26 genera that have negatively impacted the health of ball pythons [according to 15 scientific papers published between 2017 and 2019 (Table 1, Fig. 1, Suppl. material 1)]. Sequencing the microbiota of snake and mouse samples collected from the python farm in Togo provided presumptive identification of 13 (50%) of these 26 genera. One of the samples (Mouse Oral 1) did not contain amplifiable DNA and hence was not successfully sequenced. (Table 1, Fig. 1). Searches with BLAST resulted in a query cover range of 94 – 100% and percentage identity range of 89 – 100% (Suppl. material 1).


In terms of overall abundance, 85% of ASVs were assigned to genera of bacteria that were not identified as being of zoonotic concern by the literature review (Table 1, Fig. 1, Suppl. material 1). However, all but one of the samples (95%) contained at least one assigned genus of potential zoonotic concern (Table 1, Fig. 1). Between zero to six of the literature-identified genera were assigned within each sample (mean of two) (Suppl. Mat. I). The relative abundance of the literature-identified genera ranged between 0–35% of isolates per sample (mean of 13%) (Suppl. material 1).

Of the literature-identified genera, *Lysobacter* was the most prevalent among the genera-assigned ASVs, although it was only associated with snake samples. *Lysobacter* assigned ASVs also accounted for just over 10% with regards to the overall relative abundance of ASVs. Furthermore, these ASVs were present within eight out of the 19 samples (Fig. 1, Suppl. material 1). The second most abundant genus was *Bacteroides*, which accounted for 2% of the overall relative abundance of ASVs. *Bacteroides* assigned ASVs were present in each of the four sample types (snake oral, snake cloacal, mouse oral and mouse rectal), although relative abundance was greatest in the mouse rectal sample set (Fig. 1; Suppl. material 1).

## Discussion

The purpose of this study was to evaluate the presence of any potentially pathogenic bacterial taxa in ball pythons and the live mice used as their food at a commercial python farm that could impact negatively on the health of these snakes and/or those keeping them. The target facility reportedly releases all previously gravid females, and approximately 20% of their hatchlings, back into the wild and exports the remainder internationally for use as exotic pets (primarily to the USA). Of particular interest was relating the epidemiology of infection to potential vertical and horizontal transmission.

This study reported 13 different genera of bacteria, which include species that are known pathogens of ball pythons. The assignment of ASVs to *Acinetobacter*, *Bacteroides*, *Citrobacter*, *Enterobacter*, *Lysobacter*, *Proteus*, *Pseudomonas*, *Staphylococcus*, and *Tsukamurella* in the oral and cloacal samples taken from ball pythons is of potential concern for vertical and horizontal transmission, given that recent scientific literature reports pathogenic species belonging to these genera (Fig. 1, Table 1).



**Figure 1.** **A** Relative abundance of bacteria genera identified in samples taken from mice and ball pythons (*Python regius*) at a python farm in Togo. Genera containing potential pathogens of known zoonotic concern to ball pythons (as reported in the scientific literature) are highlighted in colour. The majority of reads (shown in grey) were assigned to “other” least concern groups, which consisted of ASVs that were either assigned to non-target genera or no genus. Samples were from swabs of: MR – mouse rectal; MO – mouse oral, SC – snake cloacal; and SO – snake oral. **B** Relative abundance of bacteria genera of known zoonotic concern to ball pythons (*Python regius*) (as reported in the scientific literature) identified in samples taken from mice and snakes at a python farm in Togo. Samples were from swabs of: MR – mouse rectal; MO – mouse oral, SC – snake cloacal; and SO – snake oral. No bacterial genera of zoonotic concern were identified from sample MO2.



**Table 1.** List of potentially pathogenic bacteria from ball pythons (*Python regius*) (see Methods).

| Genera                  | Species                                                                                                                         | References                                                                                                                |
|-------------------------|---------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------|
| <i>Acinetobacter</i>    | <i>Acinetobacter calcoaceticus</i> ,<br><i>Acinetobacter lawsonii</i>                                                           | Dipineto et al. 2014; Zancoli et al. 2015                                                                                 |
| <i>Aeromonas</i>        | <i>Aeromonas hydrophila</i> ,<br><i>Aeromonas veronii</i>                                                                       | Dipineto et al. 2014; Zancoli et al. 2015                                                                                 |
| <i>Anaplasma</i>        | <i>Anaplasma phagocytophilum</i>                                                                                                | Nowak et al. 2010                                                                                                         |
| <i>Bacteroides</i>      | <i>Bacteroides</i> spp.                                                                                                         | Dipineto et al. 2014                                                                                                      |
| <i>Bordetella</i>       | <i>Bordetella hinzii</i>                                                                                                        | Schmidt et al. 2013                                                                                                       |
| <i>Chlamydomphila</i>   | <i>Chlamydomphila</i> spp.                                                                                                      | Hoon-Hanks et al. 2018                                                                                                    |
| <i>Citrobacter</i>      | <i>Citrobacter freundii</i>                                                                                                     | Dipineto et al. 2014; Zancoli et al. 2015; Schmidt et al. 2013                                                            |
| <i>Clostridium</i>      | <i>Clostridium</i> spp.                                                                                                         | Dipineto et al. 2014                                                                                                      |
| <i>Elizabethkingia</i>  | <i>Elizabethkingia meningoseptica</i>                                                                                           | Schmidt et al. 2013                                                                                                       |
| <i>Enterobacter</i>     | <i>Enterobacter cloacae</i>                                                                                                     | Dipineto et al. 2014; Schmidt et al. 2013                                                                                 |
| <i>Enterococcus</i>     | <i>Enterococcus pallens</i>                                                                                                     | Zancoli et al. 2015                                                                                                       |
| <i>Escherichia</i>      | <i>Escherichia coli</i>                                                                                                         | Moss et al. 2007; Dipineto et al. 2014;<br>Larsen et al. 2011                                                             |
| <i>Klebsiella</i>       | <i>Klebsiella</i> spp., <i>Klebsiella oxytoca</i> ,<br><i>Klebsiella pneumoniae</i>                                             | Bardi et al. 2019; Schmidt et al. 2013; White et al. 2011                                                                 |
| <i>Leptospira</i>       | <i>Leptospira grippotyphosa</i>                                                                                                 | Ajayi et al. 2017                                                                                                         |
| <i>Lysobacter</i>       | <i>Lysobacter pythonis</i>                                                                                                      | Busse et al. 2019                                                                                                         |
| <i>Moraxella</i>        | <i>Moraxella osloensis</i>                                                                                                      | Zancoli et al. 2015                                                                                                       |
| <i>Morganella</i>       | <i>Morganella morganii</i>                                                                                                      | Dipineto et al. 2014; White et al. 2011                                                                                   |
| <i>Mycoplasma</i>       | <i>Mycoplasma</i> spp.                                                                                                          | Hoon-Hanks et al. 2018                                                                                                    |
| <i>Proteus</i>          | <i>Proteus vulgaris</i> , <i>Proteus</i> spp.                                                                                   | Dipineto et al. 2014; Schmidt et al. 2013                                                                                 |
| <i>Providencia</i>      | <i>Providencia rettgeri</i>                                                                                                     | Myers et al. 2009                                                                                                         |
| <i>Pseudomonas</i>      | <i>Pseudomonas aeruginosa</i> ,<br><i>Pseudomonas fluorescens</i> ,<br><i>Pseudomonas japonica</i> ,<br><i>Pseudomonas</i> spp. | Bardi et al. 2019; Zancoli et al. 2015; Sala et al. 2019;<br>Dipineto et al. 2014; Schmidt et al. 2013; White et al. 2011 |
| <i>Salmonella</i>       | <i>Salmonella enterica</i> , <i>Salmonella</i><br><i>paratyphi</i> B, <i>Salmonella</i> spp.,<br><i>Salmonella</i> Muenchen     | Moss et al. 2007; Krishnasamy et al. 2018; Dipineto et al.<br>2014; Schmidt et al. 2013; White et al. 2011                |
| <i>Serratia</i>         | <i>Serratia plymuthica</i>                                                                                                      | Zancoli et al. 2015                                                                                                       |
| <i>Staphylococcus</i>   | <i>Staphylococcus</i> spp.,<br><i>Staphylococcus warneri</i>                                                                    | Dipineto et al. 2014; Zancoli et al. 2015; White et al. 2011;<br>Schmidt et al. 2013                                      |
| <i>Stenotrophomonas</i> | <i>Stenotrophomonas maltophilia</i>                                                                                             | Zancoli et al. 2015; Schmidt et al. 2013; Klinger et al. 2018                                                             |
| <i>Tsukamurella</i>     | <i>Tsukamurella paurometabola</i>                                                                                               | Zancoli et al. 2015                                                                                                       |

The relatively high frequency of ASVs assigned to the genus *Lysobacter* – 80% ( $n = 8$ ) of oral and cloacal samples from ball pythons – is consistent with a recent report of isolates from the trachea of a ball python suffering from respiratory tract infection [*Lysobacter pythonis* sp. nov. (Busse et al. 2019)]. *Lysobacter* spp. have been described as “ubiquitous inhabitants of soil and water” (Christensen and Cook 1978) making the soil from the dens of wild caught pythons a potential source of infection.

The absence of ASVs assigned to the genera *Citrobacter*, *Enterobacter*, *Lysobacter*, *Proteus*, and *Tsukamurella* in the oral and rectal samples taken from mice used to feed ball pythons suggests that their diet was not a source of infection in this commercial operation, at least on the days of sampling, for these particular genera. Nonetheless, a much larger sample of mice is needed to determine the true bacterial status of the rodents that are typically used to feed “ranch” ball pythons when in captivity.

The presence of ASVs assigned to the genera *Clostridium*, *Escherichia*, *Moraxella*, and *Stenotrophomonas* in the oral and rectal samples taken from mice used to feed ball



pythons suggests that these mice represent a potential reservoir of infection, for these particular genera, that could impact negatively on the health of wild caught ball pythons (i.e., gravid females) and their progeny. Other potential sources of environmental contamination include other captive amphibians, birds, reptiles and mammals that are traded by python farms (cf. Bell et al. 2004), as well as free ranging birds and small mammals (including rodents), which were observed on the farm.

Ball python production systems in West Africa have the potential to encourage disease transmission and the evolution of increased pathogen virulence. Python farms that practice the “ranching” of ball pythons operate at high stocking densities and with poor hygiene measures, where animals are sourced from geographically and ecologically diverse areas with minimal quarantine. These practices can increase pathogen exposure and lower barriers to transmission (Stenglein et al. 2014).

Furthermore, the ball python is the most traded CITES-listed live wild animal currently being exported from Africa, with more than 963,334 snakes exported from Togo alone between 1978 and 2017 (D'Cruze et al. 2020). Commercial breeders in importing countries (predominantly the USA and countries of the EU) also operate at high stocking densities and commonly attend trade shows, where animals from different sources are juxtaposed (Stenglein et al. 2014).

## Limitations

The present study was restricted to 20 samples taken from five snakes and five mice at one of the seven python farms currently operating in Togo. Furthermore, it reports only on assigned bacterial genera identified as possessing pathogenic species that are known to have affected ball pythons (as reported by recent scientific literature); thus, this study is not a comprehensive or exhaustive list of genera that may contain zoonotic pathogens. Only 15% of the ASVs in our samples were assigned to genera of concern (as reported in the literature), while other potentially pathogenic genera may be present and could be identified by further analysis.

Species level identification could not be achieved with the samples in this preliminary assessment due to the short length of the targeted 16S rRNA region, which may be indistinguishable among species and/or strains (a low level taxonomic rank used at the intraspecific level) (Bulman et al. 2018; Osawa et al. 2015). Future studies could overcome this limitation and improve taxonomic resolution by using more sensitive techniques, such as quantitative polymerase chain reaction (qPCR) with species-specific primers (Osawa et al. 2015).

Similarly, the present study did not distinguish between pathogenic and non-pathogenic ASVs. This is an important distinction, since the same species of bacteria can act as a harmless commensal, as well as a dangerous pathogen (e.g., *Escherichia coli*) (Proença et al. 2017). To overcome this limitation, future studies should adopt a highly targeted and individual approach per species [e.g., as taken by Delannoy et al. (2017) who detected pathogenic strains of *Escherichia coli* using a qPCR assay that target the

K1 capsule]. Similar analyses should also look to target other pathogen types (e.g., viruses) in “ranching” ball pythons and other wild animal species held in captivity at python farms.

We recognize that the present study represents a preliminary evaluation that should be treated as an initial indicator of both the bacteria present in commercial python farms in West Africa and their potential involvement in zoonotic disease. However, given the international scope, large scale, and national wild release component of the “ranching” process that currently underpins commercial trade of live ball pythons, we believe that these initial findings provide an important insight into the potential for vertical and horizontal bacterial transmission and highlight the need for further research.

## **Recommendations**

Additional surveillance of ball pythons, both in the wild and in captivity at python farms in West Africa, will shed light on whether this type of commercial activity increases pathogen exposure and lowers barriers to transmission. However, in light of other management concerns (Auliya et al. 2020; D’Cruze et al. 2020), and as a precautionary measure, it is strongly advised that farms maintaining reptiles and other wildlife adjust to standard hygiene and quarantine measures, (e.g., biosecurity and disease surveillance practices [cf. Woodford 2000]) to minimize horizontal and vertical transfer.

Biosecurity measures should also be applied to snakes that are being released back into the wild as part of the “ranching” system in Togo. Theoretically, when they are properly released within an area of its indigenous range, this type of wild population “reinforcement” can improve the conservation status of the focal species (IUCN/SSC 2013). However, the IUCN/SSC (2013) recommends effective monitoring as an essential activity, and that reinforcement efforts should include the assessment of disease, welfare conditions, and mortality to maximise positive conservation outcomes.

Biosecurity surveillance practices should extend to importing countries. Such initiatives should also aim to inform those who trade and own ball python of the potential risks associated with zoonotic infection. Providing an appropriate environment and adequate nutrition for ball pythons is also important for maintaining their health. Washing of hands after handling ball pythons is strongly recommended (Centers for Disease Control and Prevention 2018) and they are inappropriate pets for immunocompromised owners and in households with young children (Centers for Disease Control and Prevention 2018).

## **Conclusion**

This survey represents the first investigation into the epidemiology of bacterial genera at a commercial ball python farm in West Africa. This study was developed through the opportunity to collect samples during a broader official scientific review. It is recommended that further research should be carried out at python farms in Benin, Ghana

and Togo. These studies should look to fully assess the species diversity, relative abundance, and pathogenic status of any bacteria (and other types of pathogen such as viruses) present in “ranching” ball pythons and the rodents that are used to feed them.

## Acknowledgements

We wish to thank the CITES Management Authorities of Togo (Mr. Okoumassou Kotchikpa) who facilitated the access to python farms. Thanks especially to Kinam Kombiagnou (Directeur de l'Élevage, Ministère de l'Agriculture, de l'Élevage et de la Pêche) for issuing the relevant permit. We further thank all farm owners who accepted the examination of the specimens and the sampling of swabs. Furthermore, the help of the Master students in Ecology and Wildlife Management was indispensable for us during fieldwork. We sincerely thank Agbo-Zegue NGO for providing logistical support. Jodie Bates, Emma Coulthard, Mark Auliya, David Megson and Jennifer Rowntree received a grant from World Animal Protection to carry out this research. We also sincerely thank Becky Dharmapaul, Jennah Green, John Norrey, and Laura Norrey for their assistance in reviewing the existing scientific literature and Damian Rivett for invaluable laboratory supervision.

## References

- APPA [American Pet Products Association] (2019) National Pet Owners Survey 2017–2018. [http://www.americanpetproducts.org/press\\_industrytrends.asp](http://www.americanpetproducts.org/press_industrytrends.asp) [Accessed on 11 November 2019]
- Arena PC, Steedman C, Warwick C (2012) Amphibian and reptile pet markets in the EU: an investigation and assessment. Animal Protection Agency, Animal Public, International Animal Rescue, Eurogroup for Wildlife and Laboratory Animals, Fundación para la Adopción, el Apadrinamiento y la Defensa de los Animales, 52.
- Auliya M (2003) Hot trade in cool creatures: A review of the live reptile trade in the European Union in the 1990s with a focus on Germany. TRAFFIC Europe, Brussels, Belgium.
- Auliya M, Hofmann S, Segniagbeto GH, Assou D, Ronfot D, Astrin JJ, Forat S, Ketoh GKK, D'Cruze N (2020) The first genetic assessment of wild and farmed ball pythons (Reptilia, Serpentes, Pythonidae) in southern Togo. *Nature Conservation* 38: 37–59. <https://doi.org/10.3897/natureconservation.38.49478>
- Bell D, Robertson S, Hunter PR (2004) Animal origins of SARS coronavirus: Possible links with the international trade in small carnivores. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* 359(1447): 1107–1114. <https://doi.org/10.1098/rstb.2004.1492>
- Bošnjak I, Zdravković N, Čolović S, Randelović S, Galić N, Radojičić M, Šekler M, Aleksić-Kovačević S, Krnjaić D (2016) Neglected zoonosis – The Prevalence of *Salmonella* spp. in pet reptiles in Serbia. *Vojnosanit Pregl* 73: 980–982. <https://doi.org/10.2298/VSP160809222B>

- Bulman SR, McDougal RL, Hill K, Lear G (2018) Opportunities and limitations for DNA metabarcoding in Australasian plant-pathogen biosecurity. *Australasian Plant Pathology* 47(5): 467–474. <https://doi.org/10.1007/s13313-018-0579-3>
- Bush ER, Baker SE, Macdonald DW (2014) Global trade in exotic pets 2006–2012. *Conservation Biology* 28(3): 663–676. <https://doi.org/10.1111/cobi.12240>
- Busse HJ, Huptas C, Baumgardt S, Loncaric I, Spargser J, Scherer S, Wenning M, Kämpfer P (2019) Proposal of *Lysobacter pythonis* sp. nov. isolated from royal pythons (*Python regius*). *Systematic and Applied Microbiology* 42: 326–333. <https://doi.org/10.1016/j.syapm.2019.02.002>
- Callahan B (2019) DADA2 Pipeline Tutorial (1.12) <https://benjjneb.github.io/dada2/tutorial.html> [Accessed on 01 September 2019]
- Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP (2016) DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods* 13(7): 581–583. <https://doi.org/10.1038/nmeth.3869>
- Can OE, D’Cruze N, Macdonald DW (2019) Dealing in deadly pathogens taking stock of the legal trade in live wildlife and potential risks to human health. *Global Ecology and Conservation* 17: e00515. <https://doi.org/10.1016/j.gecco.2018.e00515>
- Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, Owens SM, Betley J, Fraser L, Bauer M, Gormley N, Gilbert JA, Smith G, Knight R (2012) Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. *The ISME Journal* 6(8): 1621–1624. <https://doi.org/10.1038/ismej.2012.8>
- Centers for Disease Control and Prevention (2018) Multistate Outbreak of *Salmonella agbeni* Infections Linked to Pet Turtles, 2017. <https://www.cdc.gov/salmonella/agbeni-08-17/index.html> [Accessed 19/12/2018]
- Chaumeil PA, Mussig A, Hugenholtz P, Parks DH (2019) GTDB-Tk: A toolkit to classify genomes with the Genome Taxonomy Database. *Bioinformatics (Oxford, England)* <https://doi.org/10.1093/bioinformatics/btz848>
- Christensen P, Cook FD (1978) *Lysobacter*, a new genus of nonfruiting, gliding bacteria with a high base ratio. *International Journal of Systematic Bacteriology* 28: 367–393. <https://doi.org/10.1099/00207713-28-3-367>
- CITES [Convention on International Trade in Endangered Species of Wild Fauna and Flora] (2019) CITES glossary <https://www.cites.org/eng/resources/terms/glossary.php#r> [Accessed on 11 November 2019]
- D’Cruze N, Harrington LA, Assou D, Ronfot D, Macdonald DW, Segniagbeto GH, Auliya M (2020) Searching for snakes: Ball python hunting in southern Togo, West Africa. *Nature Conservation* 38: 13–36. <https://doi.org/10.3897/natureconservation.38.47864>
- Delannoy S, Beutin L, Mariani-Kurkdjian P, Fleiss A, Bonacorsi S, Fach P (2017) The *Escherichia coli* serogroup O1 and O2 lipopolysaccharides are Encoded by Multiple O-antigen gene clusters. *Frontiers in Cellular and Infection Microbiology* 7: 30. <https://doi.org/10.3389/fcimb.2017.00030>
- Government of Canada (2019) <https://www.canada.ca/en/public-health/services/public-health-notices/2019/outbreak-salmonella-infections-snakes-rodents.html>

- Green J, Coulthard E, Megson D, Norrey J, Norrey L, Rowntree JK, Bates J, Dharmapaul B, Auliya M, D'Cruze N (2020) Blind trading: A literature review of research addressing the welfare of ball pythons in the exotic pet trade. *Animals (Basel)* 10(2): 193. <https://doi.org/10.3390/ani10020193>
- Illumina (2019) <https://support.illumina.com> [Accessed on 11 November 2019]
- Ineich I (2006) Les élevages de reptiles et de scorpions au Bénin, Togo et Ghana, plus particulièrement la gestion des quotas d'exportation et la définition des codes 'source' des spécimens exportés. Rapport d'étude réalisée pour le Secrétariat de la CITES. Projet CITES A-251: 1–113.
- IUCN/SSC (2013) Guidelines for Reintroductions and Other Conservation Translocations. Version 1.0. IUCN Species Survival Commission, Gland, Switzerland, 57 pp.
- Jensen TJ, Auliya M, Burgess ND, Aust PW, Pertoldi C, Strand J (2018) Exploring the international trade in African snakes not listed on CITES: highlighting the role of the internet and social media. *Biodiversity and conservation*, 28: 1–19. <https://doi.org/10.1007/s10531-018-1632-9>
- Karesh WB, Cook RA, Bennett EL, Newcomb J (2005) Wildlife trade and global disease emergence. *Emerging Infectious Diseases* 11(7): 1000–1002. <https://doi.org/10.3201/eid1107.050194>
- Mitchell MA (2009) History of exotic pets. *Manual of exotic pet practice*, 1–3. <https://doi.org/10.1016/B978-141600119-5.50004-4>
- Mitchell MA, Shane SM, Roy A (1999) Detection of *Salmonellae* in the green iguana using the polymerase chain reaction technique. *Proc ARAV*, Columbus, OH, 115–117.
- Mitchell MA, Shane SM (2000) Preliminary findings of *Salmonella* spp. in captive green iguanas (*Iguana iguana*) and their environment. *Preventive Veterinary Medicine* 45: 297–304. [https://doi.org/10.1016/S0167-5877\(00\)00124-0](https://doi.org/10.1016/S0167-5877(00)00124-0)
- Morens DM, Folkers GK, Fauci A (2004) The challenge of emerging and re-emerging infectious diseases. *Nature* 430(6996): 242–249. <https://doi.org/10.1038/nature02759>
- OIE (2017) The OIE Worldwide Monitoring System for Wild Animal Diseases [http://www.oie.int/wahis\\_2/public/wahidwild.php#/](http://www.oie.int/wahis_2/public/wahidwild.php#/) (Accessed 1 November 2019).
- Osawa K, Shigemura K, Shirai H, Kato A, Okuya Y, Jikimoto T, Arakawa S, Fujisawa M, Shirakawa T (2015) Bacterial identification using *ssrA* encoding transfer-messenger RNA. *The Southeast Asian Journal of Tropical Medicine and Public Health* 46: 720–727.
- Parks DH, Chuvochina M, Waite DW, Rinke C, Skarszewski A, Chaumeil PA, Hugenholtz P (2018) A standardized bacterial taxonomy based on genome phylogeny substantially revises the tree of life. *Nature Biotechnology* 36(10): 996–1004. <https://doi.org/10.1038/nbt.4229>
- PFMA (2017) PFMAUK pet population statistics. Pet Food Manufacturers' Association. <http://www.pfma.org.uk/pet-population-2017> [Accessed 14/02/2018]
- Proença JT, Barral DC, Gordo I (2017) Commensal-to-pathogen transition: One-single transposon insertion results in two pathoadaptive traits in *Escherichia coli* -macrophage interaction. *Scientific Reports* 7(1): 4504. <https://doi.org/10.1038/s41598-017-04081-1>
- R Core Team (2019) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>

- Robinson JE, Griffiths RA, St. John FAV, Roberts DL (2015) Dynamics of the global trade in live reptiles: Shifting trends in production and consequences for sustainability. *Biological Conservation* 184: 42–50. <https://doi.org/10.1016/j.biocon.2014.12.019>
- Rogers GB, Cuthbertson L, Hoffman LR, Wing PA, Pope C, Hooftman DAP, Lilley AK, Oliver A, Carroll MP, Bruce KD, van der Gast CJ (2013) Reducing bias in bacterial community analysis of lower respiratory infections. *The ISME Journal* 7(4): 697–706. <https://doi.org/10.1038/ismej.2012.145>
- Smith KF, Behrens M, Schloegel LM, Marano N, Burgiel S, Daszak P (2009) Reducing the Risks of the Wildlife Trade. *Science* 324(5927): 594–595. <https://doi.org/10.1126/science.1174460>
- Stenglein MD, Jacobson ER, Wozniak EJ, Wellehan JFX, Kincaid A, Gordon M, Porter BF, Baumgartner W, Stahl S, Kelley K, Towner JS, Derisi JL (2014) Ball python nidovirus: A candidate etiologic agent for severe respiratory disease in *Python regius*. *mBio* 5(5): 1–16. <http://derisilab.ucsf.edu/pdfs/mBio-2014-Stenglein-.pdf> <https://doi.org/10.1128/mBio.01484-14>
- UNEP (2019) The Species+ Website. Nairobi, Kenya. Compiled by UNEP-WCMC, Cambridge, UK. [www.speciesplus.net](http://www.speciesplus.net) [Accessed 07/01/2019]
- Warwick C, Arena P, Lindley S, Jessop M, Steedman C (2013) Assessing reptile welfare using behavioural criteria. *In Practice* 35(3): 123–131. <https://doi.org/10.1136/inp.f1197>
- Woodford MH (2000) Quarantine and health screening protocols for wildlife prior to translocation and release in to the wild. Published jointly by the IUCN Species Survival Commission's Veterinary Specialist Group, Gland, Switzerland, the Office International des Epizooties (OIE), Paris, France, Care for the Wild, U.K., and the European Association of Zoo and Wildlife Veterinarians, Switzerland. <http://www.iucnwhsg.org/sites/default/files/Quarantine%20and%20Health%20Screening%20Protocol.pdf>

## Appendix I

PCR conditions for amplicon preparation using the Illumina two-step protocol.

| Step 1 Thermocycler conditions |                  |          |        |
|--------------------------------|------------------|----------|--------|
| Stage                          | Temperature (°C) | Duration | Cycles |
| Initial Denaturation           | 98               | 3 min    | 1      |
| Denaturation                   | 95               | 30 s     | ×25    |
| Annealing                      | 59               | 30 s     |        |
| Extension                      | 72               | 30 s     |        |
| Final Extension                | 72               | 5 min    | 1      |
| Step 2 Thermocycler conditions |                  |          |        |
| Stage                          | Temperature (°C) | Duration | Cycles |
| Initial Denaturation           | 98               | 30 s     | 1      |
| Denaturation                   | 98               | 10 s     | ×10    |
| Annealing                      | 62               | 20 s     |        |
| Extension                      | 72               | 30 s     |        |
| Final Extension                | 72               | 2 min    | 1      |

## Supplementary material I

### **Genera of bacteria identified in swab samples of mice and ball pythons (*Python regius*)**

Authors: Neil D'Cruze, Jodie Bates, Délagnon Assou, Delphine Ronfot, Emma Coulthard, Gabriel Hoinsoudé Segniagbeto, Mark Auliya, David Megson, Jennifer Rowntree

Data type: Excel sheet

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/natureconservation.39.48599.suppl1>