

Please cite the Published Version

Read, D , Jones, B and Till, K (2016) The use of relative speed thresholds in team sports: Applications for GPS analysis. In: Annual British Association of Sport and Exercise Science (BASES) Conference, 29 November 2016 - 30 November 2016, East Midlands Conference Centre, UK.

DOI: <https://doi.org/10.1080/02640414.2016.1260807>

Downloaded from: <https://e-space.mmu.ac.uk/625801/>

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

The use of relative speed thresholds in team sports: Applications for GPS analysis

*Dale Read^{1,2}, Ben Jones^{1,2} and Kevin Till^{1,2}

¹Leeds Beckett University, Leeds, LS6 3QS, United Kingdom

²Yorkshire Carnegie Rugby Union Football Club, Leeds, LS6 3BR, United Kingdom

*Corresponding author: d.read@leedsbeckett.ac.uk
@DaleRead4

Data from global positioning system (GPS) technology are typically presented as the distances covered in specific locomotor categories (e.g., walking, jogging, striding, sprinting). Differences are found when categorisations are made using either pre-defined absolute thresholds or thresholds relative to maximum speed. However, there are two distinct methods of using relative speed thresholds currently employed in the literature, although no study has attempted to compare them. Therefore, the purpose of this study was to compare the differences in data when analysing the same GPS files relative to speed, using either a maximum velocity sprint (V_{max}) or a maximum velocity achieved during match-play (V_{peak}). Following institutional ethics approval, 99 GPS files were analysed from rugby union match-play and split between forwards ($n=59$) and backs ($n=40$). The male participants involved were part of a Regional Academy and had the following characteristics (age: 17.5 ± 0.7 years; stature: 183.6 ± 6.6 cm; body mass: 90.6 ± 10.6 kg). V_{max} was established by players performing a maximum 40 m sprint, whilst V_{peak} was defined as the maximum velocity achieved during each match. The locomotor categories were defined as walking 0-20%, jogging 20-50%, striding 50-80% and sprinting 80-100% (Duthie et al., 2006) of either V_{max} or V_{peak} . Data were analysed using magnitude based inferences. The mean V_{max} and V_{peak} for all players were 8.7 ± 0.6 m.s⁻¹ and 7.2 ± 0.9 m.s⁻¹, respectively. There were *almost certain* differences in walking (2088 ± 298 vs. 1611 ± 435 m), striding (670 ± 244 vs. 1197 ± 375 m) and sprinting (28 ± 29 vs. 145 ± 73 m) between V_{max} and V_{peak} , for forwards. There was also a *likely* difference in jogging (2674 ± 313 vs. 2502 ± 301 m). *Very likely* differences were found for walking (2414 ± 288 vs. 2177 ± 347 m) and striding (708 ± 159 vs. 927 ± 347 m) for backs. There was also an *almost certain* difference in sprinting (66 ± 41 vs. 151 ± 49 m) whilst an *unclear* difference was found for striding (2409 ± 433 vs. 2338 ± 352 m) for backs. The use of relative thresholds using V_{peak} seems to overestimate the distance covered in striding and sprinting whilst underestimating walking and jogging. Practitioners should look to use V_{max} for relative speed thresholds as V_{peak} from match-play is likely to change from match-to-match and consequently misrepresent the movement demands that players are exposed to.