

# Please cite the Published Version

Ibuot, A, Dean, AP and Pittman, JK (2020) Multi-genomic analysis of the cation diffusion facilitator transporters from algae. Metallomics : integrated biometal science, 12 (4). pp. 617-630. ISSN 1756-5901

**DOI:** https://doi.org/10.1039/d0mt00009d

Publisher: Royal Society of Chemistry

Version: Accepted Version

Downloaded from: https://e-space.mmu.ac.uk/625861/

Usage rights: O In Copyright

**Additional Information:** This is an Author Accepted Manuscript of a paper accepted for publication in Metallomics, published by and copyright Royal Society of Chemistry.

# Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines)

## Multi-genomic analysis of the Cation Diffusion Facilitator transporters from algae

Aniefon Ibuot, <sup>a</sup> Andrew P. Dean, <sup>b</sup> and Jon K. Pittman <sup>c,\*</sup>

<sup>a</sup> Department of Science Technology, Akwa Ibom State Polytechnic, Ikot Osurua, P.M.B
 1200.Ikot Ekpene, Akwa Ibom State, Nigeria
 <sup>b</sup> Department of Natural Sciences, Faculty of Science and Engineering, Manchester
 Metropolitan University, Oxford Road, Manchester, M1 5GD, UK
 <sup>c</sup> Department of Earth and Environmental Sciences, School of Natural Sciences, Faculty of
 Science and Engineering, The University of Manchester, Michael Smith Building, Oxford
 Road, Manchester M13 9PT, UK

\*Corresponding Author:

Dr Jon Pittman, Department of Earth and Environmental Sciences, The University of Manchester, Michael Smith Building, Oxford Road, Manchester M13 9PT, UK; Tel: +44 (0)161 275 5235; Fax: +44 (0)161 275 5082; Email: <u>jon.pittman@manchester.ac.uk</u>

Orchid ID: 0000-0001-7197-1494

#### Abstract

Metal transport processes are relatively poorly understood in algae in comparison to higher plants and other eukaryotes. A screen of genomes from 33 taxonomically diverse algal species was conducted to identify members of the Cation Diffusion Facilitator (CDF) family of metal ion transporter. All algal genomes contained at least one CDF gene with four species having >10 CDF genes (median of 5 genes per genome), further confirming that this is a ubiquitous gene family. Phylogenetic analysis suggested a CDF gene organisation of five groups, which includes Zn-CDF, Fe/Zn-CDF and Mn-CDF groups, consistent with previous phylogenetic analyses, and two functionally undefined groups. One of these undefined groups was algal specific although excluded chlorophyte and rhodophyte sequences. The majority of sequences (22 out of 26 sequences) from this group had a putative ion binding site motif within transmembrane domain 2 and 5 that was distinct from other CDF proteins, such that alanine or serine replaced the conserved histidine residue. The phylogenetic grouping was supported by sequence cluster analysis. Yeast heterologous expression of CDF proteins from Chlamydomonas reinhardtii indicated Zn2+ and Co2+ transport function by CrMTP1, and Mn<sup>2+</sup> transport function by CrMTP2, CrMTP3 and CrMTP4, which validated the phylogenetic prediction. However, the Mn-CDF protein CrMTP3 was also able to provide zinc and cobalt tolerance to the Zn- and Co-sensitive zrc1cot1 yeast strain. There is wide diversity of CDF transporters within the algae lineage, and some of these genes may be attractive targets for future applications of metal content engineering in plants or microorganisms.

### Introduction

Knowledge of the transport pathways for metal influx, efflux and subcellular metal partitioning in plant and algal cells is critical in order to understand the processes of essential metal homeostasis within the cell. These will include pathways for essential trace metals such as zinc, copper and manganese, which are required for many biochemical functions including as enzyme co-factors or structural components of proteins. <sup>1</sup> Furthermore, both essential and non-essential metals, such as cadmium, are toxic at elevated concentrations and so transporters play a key role in protecting the cell against metal toxicity. <sup>2</sup> This protection can be mediated by metal efflux from the cell or intracellular metal sequestration into compartments such as the vacuole. <sup>3</sup> Manipulation of expression of specific transporter proteins, such as by genetic engineering, can therefore be used to enhance metal tolerance of an organism. <sup>4</sup> Moreover, manipulation of metal sequestration may also provide increased total metal content within a cell whilst still providing metal tolerance, and would be useful for metal bioremediation applications. <sup>5, 6</sup>

Algae are one such group of organisms that have been proposed as potentially suitable for metal bioremediation, <sup>7</sup> but if genetic engineering of metal transporters is to be tested in algae then molecular targets that have been functionally characterised are needed. Furthermore, knowledge of the gene families that encode metal transporters across taxonomically diverse algal species provides understanding of the evolution of metal homeostasis within these important photosynthetic organisms. There is a long history of the study of metal homeostasis and transport processes in algae during the second half of the last century (particularly for green microalgae species and brown seaweeds), including physiological investigation of metal micronutrient requirements and metal toxicity responses, and biochemical characterisation of metal transport (Fig. S1, ESI). However, the molecular identification of individual metal transporters and genomic scale analysis of whole gene families has only begun in the last fifteen years since the availability of algal genome sequences. For example, genomic analyses of green microalgae, such as *Chlamydomonas* 

*reinhardtii*, and red microalgae, such as *Cyanidioschyzon merolae* indicate that there are a wide variety of metal transporter pathways that are conserved between microalgae and higher plants alongside other eukaryotes such as yeast. <sup>8, 9</sup> The identification of putative algal metal transporter genes has been complemented by other genome-scale analyses such as by proteomic and transcriptomic methods, <sup>10-14</sup> and subsequent functional characterisation of individual metal transporters. <sup>6, 15, 16</sup>

One important metal transporter family is the Cation Diffusion Facilitator (CDF) family, which are referred to as Metal Transport Proteins (MTP) in higher plants. <sup>17, 18</sup> CDFtype transporters have been reported to facilitate the transport of metals including cadmium (Cd), cobalt (Co), copper (Cu), iron (Fe), manganese (Mn), nickel (Ni), and zinc (Zn), and are found virtually in all organisms covering the Eukaryote, Eubacteria and Archaea kingdoms. <sup>19, 20</sup> In yeast and plants, CDF/MTP proteins localise to membranes of various intracellular compartments, in particular the vacuolar membrane but also the mitochondria, endoplasmic reticulum (ER), Golgi/trans-Golgi and pre-vacuolar compartment, where they are involved in the efflux of metals from the cytoplasm into these organelles. <sup>18, 21</sup> For example, in *Saccharomyces cerevisiae*, CDF proteins mediate Zn<sup>2+</sup> and Co<sup>2+</sup> transfer into the vacuole (ScZrc1 and ScCot1), Zn<sup>2+</sup> transfer into the ER (ScZrg17 and ScMsc2) and Fe<sup>2+</sup> transfer from the mitochondria (ScMmt1 and ScMmt2). <sup>22-24</sup> In *Arabidopsis thaliana*, AtMTP1 and related orthologues including AtMTP3 play a major role in the storage and sequestration of Zn<sup>2+</sup> in the vacuole, <sup>25, 26</sup> AtMTP5 and AtMTP12 transport Zn<sup>2+</sup> into the Golgi, <sup>27</sup> while AtMTP8 and AtMTP11 mediate Mn<sup>2+</sup> transport into the vacuole and Golgi. <sup>28-30</sup>

These main metal substrates ( $Zn^{2+}$ ,  $Co^{2+}$ ,  $Mn^{2+}$ ,  $Fe^{2+}$ ) of the higher plant and fungal CDF family proteins are all essential micronutrients but can cause major toxicity if they accumulate at excess concentration. Zn is an important co-factor for many enzymes, such as carbonic anhydrase that is required for photosynthetic carbon fixation, and is associated with many Zn-finger containing transcription factors, and has a structural role with some proteins. <sup>31, 32</sup> Co is an essential metal in a few organisms, such as where it is required for vitamin B<sub>12</sub> and used by coenzyme B<sub>12</sub>-dependent enzymes. <sup>33</sup> Mn also has many essential

enzyme co-factor requirements, such as Mn-dependent superoxide dismutase and the water splitting oxygen evolution process of Photosystem II. <sup>34, 35</sup> Finally, Fe also plays many co-factor roles and protein structural roles, and is essential for electron transport processes. <sup>32</sup>

The current understanding of CDF/MTP proteins in algae is limited. The CDF gene family appears to be present in all previously examined algal species but specific details about CDF family structures within individual species are unknown.<sup>9, 36, 37</sup> Phylogenetic analysis will be able to reveal some useful details since substrate specificity seems to be related to phylogeny. <sup>19, 38</sup> Within the well-studied model microalga *C. reinhardtii* there are five putative MTP genes (CrMTP1 to CrMTP5). Some of these genes are transcriptionally induced by Zn or Mn deficiency conditions, <sup>10, 39</sup> but to date only CrMTP4 has been functionally characterised as a Mn<sup>2+</sup> and Cd<sup>2+</sup> transporter. <sup>6</sup> This was determined by performing over-expression of CrMTP4 in C. reinhardtii yielding enhanced Cd tolerance, and by using yeast (S. cerevisiae) heterologous expression where CrMTP4 was able to rescue the Mn sensitive phenotype of the *pmr1* yeast mutant. In this study, we aimed to perform a large-scale phylogenetic analysis of putative MTP genes from available algal genome sequences in order to evaluate the potential diversity within this broad taxonomic lineage. Furthermore, a characterisation of additional members of the C. reinhardtii MTP gene family was carried out by using yeast heterologous expression to infer the metal substrates and determine metal tolerance functions.

## Experimental

### **Bioinformatic analysis**

Thirty three algal species genomes were screened from one Charophyta: *Klebsormidium nitens*; 13 Chlorophyta: *Bathycoccus prasinos* RCC1105, *Botryococcus braunii*, *Chlamydomonas reinhardtii*, *Chlorella* sp. NC64A, *Chromochloris zofingiensis*, *Coccomyxa subellipsoidea* C-169, *Dunaliella salina*, *Micromonas pusilla* CCMP1545, *Micromonas* sp. RCC299, *Ostreococcus lucimarinus*, *Ostreococcus* sp. RCC809, *Ostreococcus tauri* 

RCC4221, *Volvox carteri*; two Pelagophyta: *Aureococcus anophagefferens* clone 1984, *Pelagophyceae* sp. CCMP2097; two Cryptophyta: *Cryptophyceae* sp. CCMP2293, *Guillardia theta* CCMP2712; three Rhodophyta: *Cyanidioschyzon merolae*, *Galdieria sulphuraria* 074W, *Porphyra umbilicalis*; four Haptophyta: *Emiliania huxleyi* CCMP1516, *Pavlovales* sp. CCMP2436, *Phaeocystis antarctica* CCMP1374, *Phaeocystis globosa* Pg-G; four Bacillariophyta: *Fragilariopsis cylindrus* CCMP 1102, *Phaeodactylum tricornutum* CCAP 1055/1, *Pseudo-nitzschia* multiseries CLN-47, *Thalassiosira pseudonana* CCMP 1335; three Ochrophyta: *Ectocarpus siliculosus* Ec 32, *Nannochloropsis oceanica* CCMP1779, *Ochromonadaceae* sp. CCMP2298; and one Chlorarachniophyta: *Bigelowiella natans* CCMP2755. In addition, other species genomes were screened for comparison from the cyanobacterium *Nostoc punctiforme* PCC 73102, human *Homo sapiens*, land plant *Arabidopsis thaliana*, yeast *Saccharomyces cerevisiae*, and the thraustochytrid *Schizochytrium aggregatum* ATCC 28209. Finally sequences from ten bacteria and one archeae were used. Genome details and individual ID numbers or accession numbers for the sequences used are provided in Supplementary Information and Table S1 (ESI).

Genomes were screened by BLAST using *C. reinhardtii* and *S. cerevisiae* CDF protein sequences. In addition, genomes were screened for the presence of the Pfam domain PF01545. All sequences were then manually examined by sequence alignment to remove any false positive sequences that had been mis-annotated. Amino acid multiple sequence alignments were performed using Clustal Omega. <sup>40</sup> For visual analysis sequence alignments were shaded using BoxShade v.3.21 and manually annotated. Prediction of transmembrane domains (TMD) was performed by hydropathy analysis using TMHMM v.2.0, <sup>41</sup> and was performed by consensus topology prediction using TOPCONS. <sup>42</sup> His-rich loop presence was determined by manual analysis by identifying the presence of three or more histidine residues within the putative cytosolic loop region between TMD 4 and 5. Conserved residues within TMD2 and TMD5 of Group 1 (Zn-CDF) and Group 5 sequences were identified using WebLogo v.2.8.2. <sup>43</sup>

Phylogenetic relationships at the amino acid level were performed using full length sequences, essentially as described previously. <sup>44, 45</sup> Sequence alignment outputs from Clustal Omega were used to generate maximum likelihood phylogenies under the WAG+F model of amino acid substitution with Γ-distributed rates across sites, as implemented in RAxML v.7.1, with tree confidence determined from 1000 replications using the fast bootstrap method. <sup>46, 47</sup> The trees were viewed using FigTree v.1.3 (http://tree.bio.ed.ac.uk/software/figtree).

The relationship between sequences was further quantified using Cluster Analysis of Sequences (CLANS), <sup>48</sup> which was used to conduct an all-against-all BLAST of unaligned sequences. Pairwise similarities between protein sequences were displayed in a 2-dimensional cluster plot using CLANS Jar from the MPI Bioinformatics Toolkit. <sup>49</sup> The P value was set to 1 x 10<sup>-5</sup> and the analysis was run for 50,000 iterations.

### Microalgae cultivation and RNA isolation

*C. reinhardtii* (CC125) was obtained from the Chlamydomonas Resource Center, USA, and was grown photo-heterotrophically in batch culture in Tris–acetate–phosphate (TAP) medium buffered at pH 7.0, as described previously, <sup>50</sup> in 200 mL glass flasks on an orbital shaker rotating at 2 Hz, at 25°C under cool-white fluorescent lights (150  $\mu$ mol m<sup>-2</sup> s<sup>-1</sup>) with a 16-h:8-h light:dark regime. Cultures were inoculated with the same starting cell density as determined by cell counting to give an initial cell count of ~65 x 10<sup>3</sup> cells mL<sup>-1</sup>. RNA was isolated from exponential growing *C. reinhardtii* CC125 cells using Trizol reagent (Life Technologies) and further purified by phenol/chloroform extraction and precipitation with isopropanol.

## MTP full length cDNA cloning

*C. reinhardtii* genomic sequence and gene model information was obtained from Phytozome v.12 using v.5.5 of the *C. reinhardtii* genome annotations and gene-specific oligonucleotide primers incorporating restriction enzyme sites (Table S2, ESI) were designed to amplify the

full-length *CrMTP1*, *CrMTP2*, and *CrMTP3* cDNA sequences by RT-PCR from *C. reinhardtii* CC125 RNA. First-strand cDNA was generated from 1 µg of DNase-treated RNA using Superscript III reverse transcriptase (Life Technologies) and an oligo(dT) primer, then KAPA HiFi DNA polymerase (Kapa Biosystems) and gene-specific primers were used for each PCR amplification, with an annealing temperature of 60°C and 35 amplification cycles. Following amplification, the PCR products were cloned into pGEM-T Easy plasmid (Promega) for propagation and sequencing (GATC Biotech) to confirm sequence fidelity. The yeast expression plasmid piUGpd was used to allow expression of each cDNA under the control of the constitutive yeast GAPDH promoter and selection of the *URA3* gene. <sup>51</sup> *CrMTP1* cDNA was sub-cloned into the Xbal and SacI sites of piUGpd, while *CrMTP2* and *CrMTP3* cDNA were sub-cloned into the BamHI and Xbal sites of piUGpd. *CrMTP4* in piUGpd was generated previously. <sup>6</sup>

## Quantitative gene expression analysis

*C. reinhardtii* CC125 was cultured in TAP medium for three days then cells were harvested and frozen in liquid N<sub>2</sub>. RNA was isolated and cDNA was produced as described above. Expression for each MTP gene was determined by quantitative real-time PCR (qPCR) using short amplicon primers that were designed to be specific to each gene (Table S3, ESI). For qPCR a SYBR Green core qPCR kit (Eurogentec) and a StepOnePlus machine (ThermoFisher) using the SYBR Green detection program was used. Transcript abundance for each *MTP* was normalised to *CBLP* transcript abundance. Reactions were run in triplicate and qPCR analysis was performed as described previously. <sup>52</sup>

### Yeast heterologous expression and metal tolerance assays

Yeast (*S. cerevisiae*) strains *pmr1* (*MATa; his3* $\Delta$ 1; *leu2* $\Delta$ 0; *met15* $\Delta$ 0; *ura3* $\Delta$ 0; *pmr1::kanMX4*) (Euroscarf, Frankfurt, Germany), *zrc1 cot1* (*MATa; his3* $\Delta$ 1; *leu2* $\Delta$ 0; *met15* $\Delta$ 0; *ura3* $\Delta$ 0; *zrc1::natMX*; *cot1::kanMX4*) <sup>53</sup> (kindly provided by Ute Krämer) and the corresponding wild type strain BY4741 (*MATa; his3* $\Delta$ 1; *leu2* $\Delta$ 0; *met15* $\Delta$ 0; *ura3* $\Delta$ 0) (Euroscarf) were each transformed with each *MTP*-piUGpd plasmid or empty piUGpd plasmid using the lithium acetate-polyethylene glycol method. Colonies were selected by growth at 30°C in synthetic defined medium minus uracil (SD –Ura) as described previously. <sup>54</sup> Expression of each *MTP* cDNA in yeast was confirmed by RT-PCR using the appropriate *MTP* qPCR primer pair (Table S3, ESI) and RNA extracted from yeast using Trizol reagent, then RT-PCR was performed as described above. PCR products were examined on a 1% agarose gel stained with SafeView (NBS Biologicals).

Metal tolerance assays were performed essentially as described previously, <sup>55</sup> on solid SD –Ura medium with or without 2 mM ZnCl<sub>2</sub>, 5 mM MnCl<sub>2</sub> or 0.8 mM CoCl<sub>2</sub> metal salts. These concentrations were chosen as the most appropriate to restrict growth of the Zn- and Co-sensitive zrt1 cot1 strain and the Mn-sensitive pmr1 strain but still allow the wild type BY4741 to grow. For growth in liquid SD –Ura medium, yeast strains were grown in medium with or without 10 mM or 20 mM CoCl<sub>2</sub>, MnCl<sub>2</sub>, or ZnCl<sub>2</sub> metal salts. Cell density was determined by absorbance measurement at 600 nm while yeast biomass was determined by weighing a cell pellet from a set volume of culture after it was dried at 60°C for 24 h. Internal metal content in yeast grown in liquid SD –Ura medium containing 20 mM ZnCl<sub>2</sub> or 20 mM MnCl<sub>2</sub>, respectively, was determined by inductively coupled plasma atomic emission spectroscopy (ICP-AES) essentially as described previously using a Perkin-Elmer Optima 5300. <sup>56</sup> It was estimated using Visual Minteq v.3.0 metal speciation modelling that the free ion concentrations available to the cells was 10.4 mM Zn<sup>2+</sup> and 12.6 mM Mn<sup>2+</sup>. Cells were centrifuged for 10 min at 3000 g then resuspended in 10 mL of 10 mM EDTA for 5 min, then re-centrifuged and washed with 15 mL Milli-Q water. Cell pellets were dried at 60 °C for 24 h and then digested in 0.5 mL of ultrapure concentrated nitric acid at 70 °C for 2 h. Samples were diluted in Milli-Q water to 2% (v/v) concentration of acid and analysed by ICP-AES. All samples were calibrated using a matrix-matched serial dilution of Specpure multielement plasma standard solution 4 (Alfa Aesar) set by linear regression.

### Results

### Phylogenetic analysis of algal CDF protein sequences

Previous phylogenetic analyses of CDF sequences are inconsistent in terms of clade structure and nomenclature of groups, with proposed numbers of CDF sub-groups ranging from 3 to 18 groups. <sup>19, 37, 38, 57</sup> Montanini et al. (2007) divided the CDF family into three major groups based on substrate specificity, referred to as Zn-CDF, Fe/Zn-CDF and Mn-CDF, <sup>26</sup> while Gustin et al. (2011) further sub-divided the higher plant CDF members into seven groups.<sup>25</sup> Cubillas et al. (2013) identified 18 CDF groups that were differentiated mostly depending on whether they contained eukaryote, bacteria or archaea derived sequences.<sup>12</sup> Here we performed a phylogenetic analysis that focussed on available algal genomes, including from one charophyte, 13 chlorophytes, two pelagophytes, two cryptophytes, three rhodophytes, four haptophytes, four bacillariophytes, three ochrophytes and one chlorarachniophyte (Table 1). A representative 'algal-like' thraustochytrid (*S. aggregatum*) was also included. First, each genome was analysed by a combination of BLAST and Pfam signature searching to collate CDF sequences. Each algal genome examined had at least one CDF sequence and the numbers of CDF transporter genes ranged from one in the red alga G. sulphuraria to 14 genes in the haptophyte bloom microalga P. antarctica (Table 1). The exact numbers of CDF genes within the more recently annotated genomes should be considered with caution since incorrect annotation and genome assembly errors can lead to genes being missed.

To generate the phylogenetic tree, the algal sequences were aligned with representative sequences from bacteria (*Escherichia coli*, *Bacillus subtilis*, *Bradyrhizobium diazoefficiens*, *Cupriavidus metallidurans*, *Rhizobium etli*, *Sinorhizobium meliloti*, *Streptococcus pneumoniae*, *Thermus thermophiles*), archaea (*Methanosarcina acetivorans*), cyanobacteria (*N. punctiforme*), animals (*H. sapiens*), fungi (*S. cerevisiae*), and land plants (*A. thaliana*), which had all been used in previous phylogenetic analyses and in most cases have known substrate specificity (Table S1, ESI). The constructed phylogenetic tree

contained 226 sequences, including 15 prokaryotic, 10 human, 12 plant, 6 yeast, 7 thraustochytrid and 176 algal sequences, of which 42 were chlorophyte sequences. Despite the dominance of many new algal sequences, the tree structure was consistent with previous CDF trees. <sup>37, 38</sup> It possessed distinct groups containing many of the known Zn<sup>2+</sup>, Fe<sup>2+</sup> and Mn<sup>2+</sup> transporting CDF proteins characterised in human, yeast, *A. thaliana* and bacteria, which were recognisable as the previously named Zn-CDF, Fe/Zn-CDF and Mn-CDF groups (Fig. 1). Group 1 (Zn-CDF) could have been considered as two or three separate groups, including a bacterial-specific clade (shown in dark blue in Fig. 1) and two eukaryotic groups (an AtMTP1 – AtMTP4 containing group, and an AtMTP5 and AtMTP12 containing group), which all include CDF proteins with experimentally confirmed Zn<sup>2+</sup> transport activity, e.g. <sup>24, 25, 26, 27, 58, 59</sup> indicating functional conservation across the whole Group 1. All taxonomic classes of algae possessed a Group 1 Zn-CDF, apart from the Chlorarachniophyta *B. natans*, although 5 of the 13 chlorophytes, including all three *Ostreococcus* spp. appear to lack a Zn-CDF (Table 1).

Group 2 (Fe/Zn-CDF), which includes Fe<sup>2+</sup> and/or Zn<sup>2+</sup> transporting proteins such as yeast ScMMT1 and ScMMT2, and *C. metallidurans* CmFieF, is rich in algal CDF sequences, including a large number of haptophyte, bacillariophyte, and ochrophyte sequences, but includes only two chlorophyte and no rhodophyte sequences (Fig. 1). A predominantly bacterial sub-group also containing five algal sequences (shown in light green in Fig. 1) might be considered as a separate group since it contains CDF proteins with known Mn<sup>2+</sup>, Ni<sup>2+</sup> and Co<sup>2+</sup> transport activities, <sup>19</sup> rather than Fe<sup>2+</sup>/Zn<sup>2+</sup>. Group 3 and Group 4 (Mn-CDF) are considered as two separate groups rather than a single group. Group 4 contains many known plant Mn<sup>2+</sup> transporting CDFs, such as AtMTP11, <sup>28, 29</sup> and is also enriched in chlorophyte sequences. Group 3 is also rich in sequences from all chlorophyte species, and includes CDF proteins such as AtMTP7 and HsZnt9 that have no currently confirmed substrate (Fig. 1). Although indirect evidence indicates Zn<sup>2+</sup> transport activity by HsZnt9, <sup>60</sup> Group 3 is referred to as a separate undefined CDF group. Finally, the tree displays a fifth algal-specific group, which is also referred to as undefined. Group 5

sequences are missing in all of the examined higher plant, charophyte and chlorophyte genomes (Fig. S2, ESI).

In order to provide validation for the group structure proposed for the CDF tree shown in Fig. 1, a phenetic sequence clustering method was performed using CLANS. <sup>48</sup> Nearly all of the sequences fell into five clusters, which completely matched the Groups 1 to 5 determined by the phylogenetic tree (Fig. 2). In particular, the analysis fully validated the separation of Group 3 and 4 into independent groups, which were both very tightly clustered. The Group 2 sequences were fairly tightly clustered and indicated partly that this group should not be further divided (shown by light green and dark green symbols overlapping in Fig. 2), although six of the sequences within this sub-group, including five algal proteins from *Pelagophyceae* sp., *T. pseudonana*, *E. siliculosus* and *Ochromonadaceae* sp., and the bacterial Sm\_0683 protein from *S. meliloti* did not cluster with the other Group 2 sequences, and are instead classified as 'unassigned' (Table 1). Group 1 and 5 sequences also formed clearly distinct clusters although these were not as tightly clustered as the other groups, and there was some overlap between four of the Group 5 sequences with Group 1. Overall, this CLANS analysis validated the five-group organisation for these CDF sequences.

## Structural features of Zn-CDF related sequences from algae

Conserved amino acid residues within TMD2 and 5 have been demonstrated to be critical for metal ion binding, in particular, two residues from TMD2 and two from TMD5 form a metal binding site referred to as 'binding site A'. <sup>17, 61</sup> The presence of residues HxxxD-HxxxD within the TMD2 and TMD5 regions, respectively, was conserved for the algal-dominated Group 1 proteins examined here (Fig. 3), which is consistent with previous analysis of Zn-CDF proteins. <sup>38</sup> In contrast, the Group 5 proteins displayed a binding site A motif that was predominantly A/SxxxD-HxxxD, such that Ala or Ser replaced the conserved His (Fig. 3). Only 4 of the 26 Group 5 proteins had the HxxxD-HxxxD motif that was highly conserved within Group 1 proteins. Indeed, these were the four Group 5 proteins that partially overlapped with the Group 1 cluster (Fig. 2).

Most of the Zn<sup>2+</sup>-transporting CDF proteins within Group 1 possess a His-rich cytosolic loop between TMD4 and 5, and this loop may play a role in metal selectivity, as some Zn-CDF transporters are specific for Zn<sup>2+</sup> while others can also transport metals including Co<sup>2+</sup>. <sup>62, 63</sup> However, the presence of a His-rich sequence is not universal, and while most the Zn-CDF proteins examined here possess an obvious His-rich loop region, including most of the chlorophyte and bacillariophyte sequences, a few do not, including some of the human Znt proteins, *A. thaliana* AtMTP5, and AtMTP5-related algal sequences (Fig. 4a). Another notable feature of many of the algal Zn-CDF sequences is that the His-rich cytosolic loop is often significantly longer than the equivalent loop region present within CDF proteins such as from *Arabidopsis*, human and yeast (Fig. 4b). For example, the CrMTP1 His-rich loop is over 300 amino acids in length while the AtMTP1 loop just over 50 amino acids in length (Fig. S3, ESI). However, some algal CDF proteins, such as from *Micromonas* sp. and *G. theta* have very short His-stretch regions of less than 10 amino acids (Fig. 4b).

## Functional characterisation of C. reinhardtii CDF proteins

*C. reinhardtii* has five CDF proteins, CrMTP1 within Group 1 (Zn-CDF), CrMTP2, CrMTP3 and CrMTP4, within Group 4 (Mn-CDF), and CrMTP5 within the undefined Group 3 (Fig. S2, ESI). To determine whether all genes are expressed within the organism, RT-PCR was performed using RNA extracted from exponential growth stage cells grown under standard conditions. Transcripts were detected for all five genes with *CrMTP1*, *CrMTP2* and *CrMTP5* showing an equivalent level of expression (between approximately 0.2 - 0.3% of *CBLP* expression), while *CrMTP3* and *CrMTP4* had low but detectable levels of expression that was significantly reduced (P < 0.05) by approximately 10 - 20-fold compared to that of *CrMTP1*, *CrMTP2* and *CrMTP5* (Fig. 5). Growth of *C. reinhardtii* in excess concentrations of metals such as Cd, Co, Fe, Mn and Zn had no significant effect on the expression level of any of the genes (data not shown) indicating that none of these CrMTP genes are transcriptionally regulated by excess metal status.

CrMTP4 has previously been confirmed to have Mn<sup>2+</sup> transport activity and the ability to transport Cd<sup>2+</sup>, <sup>6</sup> but the substrate specificity of the other *C. reinhardtii* CDF proteins is unknown. The full-length cDNA for each gene was functionally characterised by heterologous expression in yeast (Fig. 6), however, the expression of CrMTP5 was unsuccessful and therefore no functional information could be obtained for this protein. In contrast, CrMTP1, CrMTP2, CrMTP3 and CrMTP4 were all successfully expressed in a Znand Co-sensitive yeast strain zrc1 cot1 (lacking the yeast vacuolar Zn<sup>2+</sup> and Co<sup>2+</sup> transporters ScZRC1 and ScCOT1, respectively), and a Mn-sensitive yeast strain pmr1 (lacking the Mn<sup>2+</sup> ATPase transporter ScPMR1) (Fig. 6b). All cDNAs were functional as indicated by their ability to suppress metal sensitive phenotypes (Fig. 6a). CrMTP4 was used as a positive control as a confirmed Mn-CDF, and as seen with other higher plant Mn-CDF proteins, it was unable to rescue the growth of *zrc1 cot1* yeast on Zn or Co media. In contrast, CrMTP1 could strongly suppress both the Zn and Co sensitivity of zrc1 cot1 but was unable to suppress the Mn sensitivity of *pmr1*, indicating that CrMTP1 is a  $Zn^{2+}$  and Co<sup>2+</sup> transporter. CrMTP2 showed the identical characteristics of CrMTP4, indicating that it is a specific Mn<sup>2+</sup> transporter (Fig. 6a). CrMTP3 was interesting as it could strongly suppress the Mn sensitivity of *pmr1* but could also partially suppress the Zn and Co sensitivity of *zrc1* cot1, although not as strongly as CrMTP1. This suggested that CrMTP3 has broad substrate specificity for all three divalent cations tested. To date, none of the Mn-CDF proteins have been found to have Zn<sup>2+</sup> or Co<sup>2+</sup> transport activity, and appear to be largely Mn<sup>2+</sup>-specific. <sup>28,</sup> 29, 38

To further confirm the Zn, Co and Mn tolerance abilities of these proteins all cDNAs were expressed in a wild type yeast strain (BY4741) (Fig. 6b). Growth of the yeast in liquid medium was quantified in the presence of external metal addition, at concentrations that induced toxicity by inhibiting growth of wild type yeast and were relevant to metal contaminated environments. <sup>64</sup> CrMTP1 but not CrMTP3 was able to moderately but significantly increase tolerance to Zn in the yeast strain as determined by cell density measurement (Fig. 7b) and by yeast biomass (Fig. 8a). However, there was no significant

increase in yeast growth for any of the MTP-expressing strains in response to Co addition (Fig. 7c and Fig. 8b). All three CDF proteins (CrMTP2, CrMTP3, CrMTP4) that were able to strongly suppress the Mn sensitivity of *pmr1* yeast could also provide strong growth tolerance to BY4741 yeast in the presence of 20 mM Mn (Fig. 7d and Fig. 8c).

### Metal accumulation into yeast expressing C. reinhardtii CDF proteins

To determine whether the expression of algal CDF proteins in yeast can provide increased metal transport into the cell, Zn and Mn concentration in yeast cell biomass was quantified by ICP-AES following growth in liquid media with added Zn or Mn. Prior to measurement, cells were washed in an EDTA solution to remove cell wall bound metals so that only internalised metals were measured. There was no significant increase in Zn content for any of the strains including for the CrMTP1 expressing yeast (Fig. 9a), whereas the yeast expressing CrMTP2, CrMTP3 and CrMTP4 showed an approximately 2-fold increase in Mn content within the cells (Fig. 9b), indicating that the Mn tolerance of these strains was potentially due to internal sequestration of Mn.

## Discussion

## Conserved phylogenetic structure of algal CDF transporters

CDF metal transporters are a ubiquitous family of proteins across life, and this analysis confirms their presence across the wide range of taxonomically and physiologically diverse photosynthetic organisms that are collectively referred to as algae. CDF transporters appear to be principally involved in the transfer of various essential metal ions including Zn<sup>2+</sup>, Mn<sup>2+</sup>, Fe<sup>2+</sup> and Co<sup>2+</sup> into internal organelles either for delivery to specific subcellular-localised proteins that require metal co-factors or for providing metal tolerance through internal sequestration. <sup>17, 18</sup> Although other classes of primary or secondary energised metal transporters can play similar roles, <sup>9</sup> CDF transporters are clearly indispensable in algae since each algal genome that has been analysed here has at least one CDF isoform.

Furthermore, the algal-dominated phylogenetic analysis performed in this study was consistent with a previously determined CDF phylogenetic structure, whereby a small number of major groups can be differentiated on the basis of  $Zn^{2+}$ ,  $Fe^{2+}/Zn^{2+}$  and  $Mn^{2+}$  substrate specificity, plus undefined substrate specificity. <sup>36</sup> The fact that this phylogenetic structure is conserved in many algal species suggests that it was determined early in the evolution of photosynthetic organisms. In particular, the Zn-CDF and Mn-CDF groups are the most conserved across the algal genomes examined here. Nevertheless, some algal species have lost particular CDF types; for example, all three *Ostreococcus* spp. lack Group 1 (Zn-CDF) and Group 2 (Fe/Zn-CDF) genes, and in fact most chlorophytes lack Group 2 (Fe/Zn-CDF) genes, while all three red algae examined (*C. merolae*, *G. sulphuraria* and *P. umbilicalis*) lack Group 2 (Fe/Zn-CDF) and the undefined Group 3 genes. The functional relevance of this is unclear but may indicate that functional redundancy with other transporter classes (e.g.  $Zn^{2+}$  or  $Mn^{2+}$  P-type ATPases;  $Fe^{2+}$  or  $Mn^{2+}$  transporting VIT-like proteins) is not essential and such redundancy has therefore not been selected in many of these organisms.

#### A novel algal-specific phylogenetic group with distinct ion binding site motif

We have also proposed the presence of a fifth phylogenetic group (Group 5) that is most closely related to Group 1 (Zn-CDF) but appears to be distinct. This group is almost exclusively algal-specific apart from a *S. aggregatum* isoform, but includes no chlorophyte or rhodophyte sequences. The difference in sequence for the TMD2 half of the metal binding site A motif of the majority of the Group 5 proteins further demonstrates a divergence with the other groups. This TMD2 motif is typically HxxxD or DxxxD in most CDF proteins, such as in the Zn<sup>2+</sup>-transporting AtMTP1 and CrMTP1 (HxxxD), and in the Mn<sup>2+</sup>-transporting AtMTP11 and CrMTP4 (DxxxD), but the Group 5 TMD2 motif was A/SxxxD. Mutation of the His residue within this motif to Ala in the AtMTP1 Zn-CDF protein, the PtdMTP1 Zn-CDF protein from *Populus trichocarpa x deltoides* and the RmCzcD Zn-CDF protein from *Ralstonia metallidurans* all caused loss of function, <sup>62, 65, 66</sup> perhaps indicating that these

Group 5 proteins would not be functional for Zn<sup>2+</sup> transport. However, there are exceptions to these motifs where His or Asp in the first position is not essential, such as NxxxD within TMD2 of the Mn<sup>2+</sup> transporting SpMntA from *S. pneumoniae* or FxxxD within TMD2 of the Zn<sup>2+</sup> transporting ScZRG17 from *S. cerevisiae*. <sup>61</sup> Furthermore, the recently identified Na<sup>+-</sup> transporting MceT protein from the halophilic bacterium *Planococcus dechangensis*, which has been proposed to be a member of a novel Na-CDF clade, has a distinct YxxxS motif. <sup>67</sup> Currently we are unable to predict the substrate specificity of these Group 5 algae proteins from sequence comparisons therefore future experimental analysis is needed.

### C. reinhardtii MTP transporters have Zn-CDF and Mn-CDF functions

This study has identified *C. reinhardtii* members of the CDF family of metal efflux transporters as Zn<sup>2+</sup>, Co<sup>2+</sup> and Mn<sup>2+</sup> transporters, and validated the function of an algal Zn-CDF protein for the first time. CrMTP1 clearly groups in the Group 1 (Zn-CDF) clade while CrMTP2, CrMTP3 and CrMTP4 are clearly within the Group 4 (Mn-CDF) clade, and the yeast expression experiments validated these predicted activities, as we previously observed for CrMTP4. <sup>6</sup> An intriguing exception was the apparent Mn<sup>2+</sup> and Zn<sup>2+</sup> transport activity by CrMTP3, the first example we are aware of for a Mn<sup>2+</sup>-transporting CDF also being linked to Zn<sup>2+</sup> transport. A Mn-CDF from *A. thaliana* (AtMTP8) may transport both Mn<sup>2+</sup> and Fe<sup>2+</sup>, <sup>68</sup> while the Group 1 (Zn-CDF) member HsZnt10 can transport Mn<sup>2+</sup>. <sup>69</sup> While there is some sequence variation between CrMTP3 and CrMTP2/CrMTP4 that could explain this difference, future biochemical and mutagenesis studies are required to experimental validate this further.

CrMTP1 was able to efficiently provide Zn tolerance to yeast, indicating Zn<sup>2+</sup> transport activity, despite the structure of CrMTP1 being distinct from previously characterised Zn-CDF proteins such as AtMTP1 and ScZRC1 due to the significantly longer His-rich loop domain between TMD4 and 5. This seems to be a structural feature for many of the algal Zn-CDF proteins. The His-rich loop has been proposed to play a role in metal ion specificity and affinity, such that shortening of the AtMTP1 loop, which also reduces the

number of His residues, alters the kinetics of  $Zn^{2+}$  transport, while the shortened loop or specific point-mutations within the loop can allow  $Co^{2+}$  transport, which is otherwise poor in native AtMTP1. <sup>63, 70</sup> Therefore we might predict that the longer His-loop of CrMTP1 may provide different  $Zn^{2+}$  and  $Co^{2+}$  transport kinetics compared to some of the higher plant isoforms.

CrMTP2 and CrMTP4 but not the other C. reinhardtii CDF family genes have been shown to be transcriptionally induced by Mn deficiency, thus linking them further to Mn homeostasis roles, <sup>39</sup> while *CrMTP1* is induced by Zn deficiency, <sup>10</sup> although we could find no evidence by RT-PCR that any of these genes are transcriptionally regulated by metal excess. CrMTP1 was able to efficiently rescue the Zn sensitive phenotype of the yeast vacuolar Zn<sup>2+</sup>/H<sup>+</sup> CDF transporter ScZRC1 and provide Zn tolerance in wild type yeast, thus it is proposed to provide a pathway of vacuolar Zn<sup>2+</sup> sequestration equivalent to AtMTP1 and ScZRC1. <sup>24, 26</sup> Likewise, the significant Mn tolerance gained by CrMTP2, CrMTP3 or CrMTP4 that was coincident with increased Mn content within the cell is consistent with sequestration of Mn<sup>2+</sup> into an internal compartment, possibly a vacuolar structure. C. reinhardtii lacks a large central vacuole as in typical plant cells but possess multiple lysosomal/vacuolar-like structures that are also referred to as acidocalcisomes. <sup>71, 72</sup> These are acidic structures that accumulate polyphosphate and metals including Ca, Mg and Zn, and possess proton pumps that would be able to energise H<sup>+</sup>-coupled Zn<sup>2+</sup> or Mn<sup>2+</sup> transport by an MTP protein. <sup>73</sup> The identification of the red algal MTP1 orthologue from C. merolae (Cmer CMF058C) from an acidocalcisome proteomic study, <sup>74</sup> further strengthens the likely function of algal Zn-CDF proteins in vacuolar Zn<sup>2+</sup> sequestration.

## Conclusions

Knowledge of sub-cellular metal transport processes, such as by members of the CDF transporter family, not only provides improved understanding of fundamental mechanisms of metal homeostasis and metal tolerance within a cell for a given organism, but may provide tools for biotechnological approaches to manipulate metal content within a whole organism,

tissue, or sub-cellular compartment. Previous studies have proposed applications of CDF transporters for enhancing mineral biofortification of plants or for providing toxic metal bioremediation in plants or algae. <sup>6, 18, 75</sup> This present study has indicated the wide diversity of CDF transporters that exist within the algae lineage. Although the vast majority of these proteins are still uncharacterized and the potential metal substrates are unknown, it is quite possible that some of these proteins will have characteristics that are distinct to those CDF transporters already known in plants and other eukaryotes. Therefore algal CDF genes may be attractive gene targets for future genetic engineering approaches in plants or microorganisms for metal transport engineering.

### Acknowledgements

This work was supported by a Government of Nigeria TETFUND PhD studentship awarded to AI. We thank Paul Lythgoe (Manchester Analytical Geochemical Unit, Department of Earth and Environmental Sciences, University of Manchester) for ICP-AES analysis. We thank Ute Krämer for providing the *zrc1 cot1* yeast strain.

### **Conflict of interest**

There are no conflicts to declare.

### References

- C. Andreini, I. Bertini, G. Cavallaro, G. L. Holliday and J. M. Thornton, Metal ions in biological catalysis: from enzyme databases to general principles, *J. Biol. Inorg. Chem.*, 2008, **13**, 1205-1218.
- U. Kramer, I. N. Talke and M. Hanikenne, Transition metal transport, *FEBS Lett.*, 2007, 581, 2263-2272.
- S. S. Sharma, K.-J. Dietz and T. Mimura, Vacuolar compartmentalization as indispensable component of heavy metal detoxification in plants, *Plant, Cell Environ.*, 2016, **39**, 1112-1126.

- D. M. Antosiewicz, A. Barabasz and O. Siemianowski, Phenotypic and molecular consequences of overexpression of metal-homeostasis genes, *Front. Plant Sci.*, 2014, 5, 80.
- 5. S. Y. Cheng, P.-L. Show, B. F. Lau, J.-S. Chang and T. C. Ling, New prospects for modified algae in heavy metal adsorption, *Trends Biotechnol.*, 2019, **37**, 1255-1268.
- A. Ibuot, A. P. Dean, O. A. McIntosh and J. K. Pittman, Metal bioremediation by *CrMTP4* over-expressing *Chlamydomonas reinhardtii* in comparison to natural wastewater-tolerant microalgae strains, *Algal Res.*, 2017, 24, 89-96.
- K. Suresh Kumar, H.-U. Dahms, E.-J. Won, J.-S. Lee and K.-H. Shin, Microalgae A promising tool for heavy metal remediation, *Ecotoxicol. Environ. Saf.*, 2015, **113**, 329-352.
- M. Hanikenne, U. Kramer, V. Demoulin and D. Baurain, A comparative inventory of metal transporters in the green alga *Chlamydomonas reinhardtii* and the red alga *Cyanidioschizon merolae*, *Plant Physiol.*, 2005, **137**, 428-446.
- C. E. Blaby-Haas and S. S. Merchant, The ins and outs of algal metal transport, *Biochim. Biophys. Acta*, 2012, **1823**, 1531-1552.
- D. Malasarn, J. Kropat, S. I. Hsieh, G. Finazzi, D. Casero, J. A. Loo, M. Pellegrini, F.-A. Wollman and S. S. Merchant, Zinc deficiency impacts CO<sub>2</sub> assimilation and disrupts copper homeostasis in *Chlamydomonas reinhardtii*, *J. Biol. Chem.*, 2013, **288**, 10672-10683.
- 11. R. Beauvais-Flück, V. I. Slaveykova and C. Cosio, Comparative study of Cu uptake and early transcriptome responses in the green microalga *Chlamydomonas reinhardtii* and the macrophyte *Elodea nuttallii*, *Environ. Pollut.*, 2019, **250**, 331-337.
- F. Puente-Sánchez, S. Díaz, V. Penacho, A. Aguilera and S. Olsson, Basis of genetic adaptation to heavy metal stress in the acidophilic green alga *Chlamydomonas acidophila*, *Aquat. Toxicol.*, 2018, **200**, 62-72.
- S. I. Hsieh, M. Castruita, D. Malasarn, E. Urzica, J. Erde, M. D. Page, H. Yamasaki,
   D. Casero, M. Pellegrini, S. S. Merchant and J. A. Loo, The proteome of copper, iron,

zinc, and manganese micronutrient deficiency in *Chlamydomonas reinhardtii*, *Mol. Cell. Proteomics*, 2013, **12**, 65-86.

- B. Khatiwada, M. T. Hasan, A. Sun, K. S. Kamath, M. Mirzaei, A. Sunna and H. Nevalainen, Proteomic response of *Euglena gracilis* to heavy metal exposure Identification of key proteins involved in heavy metal tolerance and accumulation, *Algal Res.*, 2020, **45**, 101764.
- M. Tsednee, M. Castruita, P. A. Salomé, A. Sharma, B. E. Lewis, S. R. Schmollinger, D. Strenkert, K. Holbrook, M. S. Otegui, K. Khatua, S. Das, A. Datta, S. Chen, C. Ramon, M. Ralle, P. K. Weber, T. L. Stemmler, J. Pett-Ridge, B. M. Hoffman and S. S. Merchant, Manganese co-localizes with calcium and phosphorus in *Chlamydomonas* acidocalcisomes and is mobilized in manganese-deficient conditions, *J. Biol. Chem.*, 2019, **294**, 17626-17641.
- P. Chang, H. Yin, T. Imanaka, Y. Igarashi, N. Li and F. Luo, The metal transporter CrNRAMP1 is involved in zinc and cobalt transports in *Chlamydomonas reinhardtii*, *Biochem. Biophys. Res. Commun.*, 2020, **523**, 880-886.
- 17. O. Kolaj-Robin, D. Russell, K. A. Hayes, J. T. Pembroke and T. Soulimane, Cation Diffusion Facilitator family: Structure and function, *FEBS Lett.*, 2015, **589**, 1283-1295.
- F. Ricachenevsky, P. Menguer, R. Sperotto, L. Williams and J. Fett, Roles of plant metal tolerance proteins (MTP) in metal storage and potential use in biofortification strategies, *Front. Plant Sci.*, 2013, **4**, 144.
- C. Cubillas, P. Vinuesa, M. L. Tabche and A. García-de los Santos, Phylogenomic analysis of Cation Diffusion Facilitator proteins uncovers Ni<sup>2+</sup>/Co<sup>2+</sup> transporters, *Metallomics*, 2013, **5**, 1634-1643.
- D. H. Nies and S. Silver, Ion efflux systems involved in bacterial metal resistances, *J. Ind. Microbiol.*, 1995, **14**, 186-199.
- C. J. Haney, G. Grass, S. Franke and C. Rensing, New developments in the understanding of the cation diffusion facilitator family, *J. Ind. Microbiol. Biotechnol.*, 2005, 32, 215-226.

- C. D. Ellis, C. W. MacDiarmid and D. J. Eide, Heteromeric protein complexes mediate zinc transport into the secretory pathway of eukaryotic cells, *J. Biol. Chem.*, 2005, **280**, 28811-28818.
- L. Li, R. Miao, X. Jia, D. M. Ward and J. Kaplan, Expression of the yeast Cation Diffusion Facilitators Mmt1 and Mmt2 affects mitochondrial and cellular iron homeostasis: evidence for mitochondrial iron export, *J. Biol. Chem.*, 2014, **289**, 17132-17141.
- C. W. MacDiarmid, M. A. Milanick and D. J. Eide, Biochemical properties of vacuolar zinc transport systems of *Saccharomyces cerevisiae*, *J. Biol. Chem.*, 2002, **277**, 39187-39194.
- S. Arrivault, T. Senger and U. Kramer, The Arabidopsis metal tolerance protein AtMTP3 maintains metal homeostasis by mediating Zn exclusion from the shoot under Fe deficiency and Zn oversupply, *Plant J.*, 2006, **46**, 861-879.
- Y. Kobae, T. Uemura, M. H. Sato, M. Ohnishi, T. Mimura, T. Nakagawa and M. Maeshima, Zinc transporter of *Arabidopsis thaliana* AtMTP1 is localized to vacuolar membranes and implicated in zinc homeostasis, *Plant Cell Physiol.*, 2004, **45**, 1749-1758.
- T. Fujiwara, M. Kawachi, Y. Sato, H. Mori, N. Kutsuna, S. Hasezawa and M.
   Maeshima, A high molecular mass zinc transporter MTP12 forms a functional heteromeric complex with MTP5 in the Golgi in *Arabidopsis thaliana*, *FEBS J.*, 2015, **282**, 1965-1979.
- E. Delhaize, B. D. Gruber, J. K. Pittman, R. G. White, H. Leung, Y. S. Miao, L. W. Jiang, P. R. Ryan and A. E. Richardson, A role for the AtMTP11 gene of *Arabidopsis* in manganese transport and tolerance, *Plant J.*, 2007, **51**, 198-210.
- E. Peiter, B. Montanini, A. Gobert, P. Pedas, S. Husted, F. J. M. Maathuis, D. Blaudez, M. Chalot and D. Sanders, A secretory pathway-localized cation diffusion facilitator confers plant manganese tolerance, *Proc. Natl. Acad. Sci. U.S.A.*, 2007, **104**, 8532-8537.
- S. Eroglu, B. Meier, N. von Wirén and E. Peiter, The vacuolar manganese transporter MTP8 determines tolerance to iron deficiency-induced chlorosis in Arabidopsis, *Plant Physiol.*, 2016, **170**, 1030-1045.

- L. Decaria, I. Bertini and R. J. P. Williams, Zinc proteomes, phylogenetics and evolution, *Metallomics*, 2010, 2, 706-709.
- R. Hänsch and R. R. Mendel, Physiological functions of mineral micronutrients (Cu, Zn, Mn, Fe, Ni, Mo, B, Cl), *Curr. Opin. Plant Biol.*, 2009, **12**, 259-266.
- A. K. Renfrew, E. S. O'Neill, T. W. Hambley and E. J. New, Harnessing the properties of cobalt coordination complexes for biological application, *Coord. Chem. Rev.*, 2018, **375**, 221-233.
- 34. J. K. Pittman, Managing the manganese: molecular mechanisms of manganese transport and homeostasis, *New Phytol.*, 2005, **167**, 733-742.
- A. L. Socha and M. L. Guerinot, Mn-euvering manganese: the role of transporter gene family members in manganese uptake and mobilization in plants, *Front. Plant Sci.*, 2014, **5**, 106.
- C. E. Blaby-Haas and S. S. Merchant, Regulating cellular trace metal economy in algae, *Curr. Opin. Plant Biol.*, 2017, **39**, 88-96.
- 37. J. L. Gustin, M. J. Zanis and D. E. Salt, Structure and evolution of the plant cation diffusion facilitator family of ion transporters, *BMC Evol. Biol.*, 2011, **11**, 76.
- B. Montanini, D. Blaudez, S. Jeandroz, D. Sanders and M. Chalot, Phylogenetic and functional analysis of the Cation Diffusion Facilitator (CDF) family: improved signature and prediction of substrate specificity, *BMC Genomics*, 2007, 8, 107.
- M. D. Allen, J. Kropat, S. Tottey, J. A. Del Campo and S. S. Merchant, Manganese deficiency in Chlamydomonas results in loss of photosystem II and MnSOD function, sensitivity to peroxides, and secondary phosphorus and iron deficiency, *Plant Physiol.*, 2007, **143**, 263-277.
- F. Madeira, Y. m. Park, J. Lee, N. Buso, T. Gur, N. Madhusoodanan, P. Basutkar, A. R. N. Tivey, S. C. Potter, R. D. Finn and R. Lopez, The EMBL-EBI search and sequence analysis tools APIs in 2019, *Nucleic Acids Res.*, 2019, **47**, W636-W641.

- A. Krogh, B. Larsson, G. von Heijne and E. L. L. Sonnhammer, Predicting transmembrane protein topology with a hidden Markov model: Application to complete genomes, *J. Mol. Biol.*, 2001, **305**, 567-580.
- K. D. Tsirigos, C. Peters, N. Shu, L. Käll and A. Elofsson, The TOPCONS web server for consensus prediction of membrane protein topology and signal peptides, *Nucleic Acids Res.*, 2015, **43**, W401-W407.
- G. E. Crooks, G. Hon, J.-M. Chandonia and S. E. Brenner, WebLogo: a sequence logo generator, *Genome Res.*, 2004, **14**, 1188-1190.
- L. Emery, S. Whelan, K. D. Hirschi and J. K. Pittman, Protein phylogenetic analysis of Ca<sup>2+</sup>/cation antiporters and insights into their evolution in plants, *Front. Plant Sci.*, 2012, **3**, 1.
- 45. J. K. Pittman and K. D. Hirschi, Phylogenetic analysis and protein structure modelling identifies distinct Ca<sup>2+</sup>/Cation antiporters and conservation of gene family structure within Arabidopsis and rice species, *Rice*, 2016, **9**, 3.
- 46. A. Stamatakis, RAxML-VI-HPC: Maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models, *Bioinformatics*, 2006, **22**, 2688-2690.
- 47. A. Stamatakis, P. Hoover and J. Rougemont, A rapid bootstrap algorithm for the RAxML web servers, *Syst. Biol.*, 2008, **57**, 758-771.
- 48. T. Frickey and A. Lupas, CLANS: a Java application for visualizing protein families based on pairwise similarity, *Bioinformatics*, 2004, **20**, 3702-3704.
- 49. L. Zimmermann, A. Stephens, S.-Z. Nam, D. Rau, J. Kübler, M. Lozajic, F. Gabler, J. Söding, A. N. Lupas and V. Alva, A completely reimplemented MPI Bioinformatics Toolkit with a new HHpred server at its core, *J. Mol. Biol.*, 2018, **430**, 2237-2243.
- 50. E. H. Harris, *The Chlamydomonas Sourcebook*, Academic Press, San Diego, 1989.
- D. F. Nathan, M. H. Vos and S. Lindquist, Identification of SSF1, CNS1, and HCH1 as multicopy suppressors of a *Saccharomyces cerevisiae* Hsp90 loss-of-function mutation, *Proc. Natl. Acad. Sci. U.S.A.*, 1999, **96**, 1409-1414.

- 52. A. K. Bajhaiya, A. P. Dean, L. A. H. Zeef, R. E. Webster and J. K. Pittman, PSR1 is a global transcriptional regulator of phosphorus deficiency responses and carbon storage metabolism in *Chlamydomonas reinhardtii*, *Plant Physiol.*, 2016, **170**, 1216-1234.
- D. B. Drager, A. G. Desbrosses-Fonrouge, C. Krach, A. N. Chardonnens, R. C. Meyer, P. Saumitou-Laprade and U. Kramer, Two genes encoding *Arabidopsis halleri* MTP1 metal transport proteins co-segregate with zinc tolerance and account for high *MTP1* transcript levels, *Plant J.*, 2004, **39**, 425-439.
- 54. J. K. Pittman, N. H. Cheng, T. Shigaki, M. Kunta and K. D. Hirschi, Functional dependence on calcineurin by variants of the *Saccharomyces cerevisiae* vacuolar Ca<sup>2+</sup>/H<sup>+</sup> exchanger Vcx1p, *Mol. Microbiol.*, 2004, **54**, 1104-1116.
- 55. J. K. Pittman, C. Edmond, P. A. Sunderland and C. M. Bray, A cation-regulated and proton gradient-dependent cation transporter from *Chlamydomonas reinhardtii* has a role in calcium and sodium homeostasis, *J. Biol. Chem.*, 2009, **284**, 525-533.
- R. E. Webster, A. P. Dean and J. K. Pittman, Cadmium exposure and phosphorus limitation increases metal content in the freshwater alga *Chlamydomonas reinhardtii*, *Environ. Sci. Technol.*, 2011, **45**, 7489-7496.
- A. Migeon, D. Blaudez, O. Wilkins, B. Montanini, M. M. Campbell, P. Richaud, S. Thomine and M. Chalot, Genome-wide analysis of plant metal transporters, with an emphasis on poplar, *Cell. Mol. Life Sci.*, 2010, **67**, 3763-3784.
- 58. Y. Chao and D. Fu, Kinetic study of the antiport mechanism of an *Escherichia coli* zinc transporter, ZitB, *J. Biol. Chem.*, 2004, **279**, 12043-12050.
- J. P. Liuzzi and R. J. Cousins, Mammalian zinc transporters, *Annu. Rev. Nutr.*, 2004,
   24, 151-172.
- Y. Perez, Z. Shorer, K. Liani-Leibson, P. Chabosseau, R. Kadir, M. Volodarsky, D. Halperin, S. Barber-Zucker, H. Shalev, R. Schreiber, L. Gradstein, E. Gurevich, R. Zarivach, G. A. Rutter, D. Landau and O. S. Birk, SLC30A9 mutation affecting intracellular zinc homeostasis causes a novel cerebro-renal syndrome, *Brain*, 2017, **140**, 928-939.

- S. Barber-Zucker, B. Shaanan and R. Zarivach, Transition metal binding selectivity in proteins and its correlation with the phylogenomic classification of the cation diffusion facilitator protein family, *Sci. Rep.*, 2017, **7**, 16381.
- 62. M. Kawachi, Y. Kobae, S. Kogawa, T. Mimura, U. Kraemer and M. Maeshima, Amino acid screening based on structural modeling identifies critical residues for the function, ion selectivity and structure of Arabidopsis MTP1, *FEBS J.*, 2012, **279**, 2339-2356.
- D. Podar, J. Scherer, Z. Noordally, P. Herzyk, D. Nies and D. Sanders, Metal selectivity determinants in a family of transition metal transporters, *J. Biol. Chem.*, 2012, 287, 3185-3196.
- R. A. Wuana and F. E. Okieimen, Heavy metals in contaminated soils: a review of sources, chemistry, risks and best available strategies for remediation, *ISRN Ecology*, 2011, 2011, 402647.
- 65. D. Blaudez, A. Kohler, F. Martin, D. Sanders and M. Chalot, Poplar metal tolerance protein 1 confers zinc tolerance and is an oligomeric vacuolar zinc transporter with an essential leucine zipper motif, *Plant Cell*, 2003, **15**, 2911-2928.
- A. Anton, A. Weltrowski, C. J. Haney, S. Franke, G. Grass, C. Rensing and D. H.
   Nies, Characteristics of zinc transport by two bacterial Cation Diffusion Facilitators from *Ralstonia metallidurans* CH34 and *Escherichia coli*, *J. Bacteriol.*, 2004, **186**, 7499-7507.
- T. Xu, H. Chen, J. Li, S. Hong, L. Shao, X. Zheng, Q. Zou, Y. Wang, S. Guo and J. Jiang, Implications for cation selectivity and evolution by a novel Cation Diffusion Facilitator family member from the moderate halophile *Planococcus dechangensis*, *Front. Microbiol.*, 2019, **10**, 607.
- H.-H. Chu, S. Car, A. L. Socha, M. N. Hindt, T. Punshon and M. L. Guerinot, The Arabidopsis MTP8 transporter determines the localization of manganese and iron in seeds, *Scientific Reports*, 2017, 7, 11024.
- 69. Y. Nishito, N. Tsuji, H. Fujishiro, T.-a. Takeda, T. Yamazaki, F. Teranishi, F. Okazaki,
  A. Matsunaga, K. Tuschl, R. Rao, S. Kono, H. Miyajima, H. Narita, S. Himeno and T.
  Kambe, Direct comparison of manganese detoxification/efflux proteins and molecular

characterization of ZnT10 protein as a manganese transporter, *J. Biol. Chem.*, 2016, **291**, 14773-14787.

- M. Kawachi, Y. Kobae, T. Mimura and M. Maeshima, Deletion of a histidine-rich loop of AtMTP1, a vacuolar Zn2+/H+ antiporter of Arabidopsis thaliana, stimulates the transport activity, *J. Biol. Chem.*, 2008, **283**, 8374-8383.
- C. E. Blaby-Haas and S. S. Merchant, Lysosome-related organelles as mediators of metal homeostasis, *J. Biol. Chem.*, 2014, 289, 28129-28136.
- Y. Komine, L. L. Eggink, H. S. Park and J. K. Hoober, Vacuolar granules in Chlamydomonas reinhardtii: polyphosphate and a 70-kDa polypeptide as major components, *Planta*, 2000, **210**, 897-905.
- 73. F. A. Ruiz, N. Marchesini, M. Seufferheld, Govindjee and R. Docampo, The polyphosphate bodies of *Chlamydomonas reinhardtii* possess a proton-pumping pyrophosphatase and are similar to acidocalcisomes, *J. Biol. Chem.*, 2001, **276**, 46196-46203.
- F. Yagisawa, K. Nishida, M. Yoshida, M. Ohnuma, T. Shimada, T. Fujiwara, Y.
  Yoshida, O. Misumi, H. Kuroiwa and T. Kuroiwa, Identification of novel proteins in isolated polyphosphate vacuoles in the primitive red alga *Cyanidioschyzon merolae*, *Plant J.*, 2009, **60**, 882-893.
- N. Das, S. Bhattacharya and M. K. Maiti, Enhanced cadmium accumulation and tolerance in transgenic tobacco overexpressing rice metal tolerance protein gene
   *OsMTP1* is promising for phytoremediation, *Plant Physiol. Biochem.*, 2016, **105**, 297-309.

**Table 1.** Total number and distribution of CDF transporter genes from each analysed genome into five phylogenetic groups: Group 1 (Zn-CDF), Group 2 (Fe/Zn-CDF), Group 3 (undefined CDF), Group 4 (Mn-CDF), Group 5 (undefined CDF). Gene numbers were determined from the phylogenetic analysis (Fig. 1) and supported by the cluster analysis (Fig. 2). Six sequences were classed as unassigned (UA) including five sequences where the evidence for Group 2 assignment was not supported by cluster analysis.

Species		G.1 (Zn)	G.2 (Fe/Zn)	G.3 (n.d.)	G.4 (Mn)	G.5 (n.d.)	UA	Total
Cyanobacteria	Nostoc punctiforme	0	2	2	0	0	0	4
Metazoa	Homo sapiens	9	0	1	0	0	0	10
Ascomycota	Saccharomyces cerevisiae	3	2	0	0	0	1	6
Embryophyta	Arabidopsis thaliana	6	1	1	4	0	0	12
Thraustochytrid	Schizochytrium	3	1	1	1	1	0	7
	aggregatum							
Charophyta	Klebsormidium nitens	4	1	1	2	0	0	8
Chlorophyta	Bathycoccus prasinos	0	1	1	1	0	0	3
	Botryococcus braunii	1	0	1	0	0	0	2
	Chlamydomonas	1	0	1	3	0	0	5
	reinhardtii							
	Chlorella sp. NC64A	1	0	1	1	0	0	3
	Chromochloris	1	0	1	2	0	0	4
	zofingiensis							
	Coccomyxa	2	0	1	2	0	0	5
	subellipsoidea							
	Dunaliella salina	2	0	1	1	0	0	4
	Micromonas pusilla	0	0	1	1	0	0	2
	Micromonas sp.	2	1	1	1	0	0	5
	Ostreococcus	0	0	1	1	0	0	2
	lucimarinus							
	Ostreococcus sp.	0	0	1	1	0	0	2
	Ostreococcus tauri	0	0	1	1	0	0	2
	Volvox carteri	1	0	1	1	0	0	3
Pelagophyta	Aureococcus anophagefferens	1	2	0	1	0	0	4
	Pelagophyceae sp.	1	2	1	2	4	1	11
Cryptophyta	Cryptophyceae sp.	1	1	0	0	1	0	3
	Guillardia theta	6	1	1	1	1	0	10
Rhodophyta	Cyanidioschyzon	2	0	0	1	0	0	3
	merolae							
	Galdieria sulphuraria	1	0	0	0	0	0	1
	Porphyra umbilicalis	3	0	0	1	0	0	4
Haptophyta	Emiliania huxleyi	3	4	0	1	3	0	11
	Pavlovales sp.	2	2	0	2	3	0	9
	Phaeocystis antarctica	5	7	1	1	0	0	14
	Phaeocystis globosa	2	4	0	1	2	0	9
Bacillariophyta	Fragilariopsis cylindrus	1	1	1	0	3	0	6
	Phaeodactylum tricornutum	1	1	1	1	1	0	5
	Pseudo-nitzschia	2	1	1	1	1	0	6
	Thelessiosire	1	1	1	1	0	1	5
	nseudonana	1	I	I	I	0	I	5
Ochrophyta	Ectocarnus siliculosus	1	1	1	Ω	1	1	5
Ochiophyta	Nannochloronsis	3	1	1	1	0	0	6
	oceanica			ı 		0	0	
	Ochromonadaceae sp.	1	2	0	0	2	2	7
Chlorarachniophyta	Bigelowiella natans	0	2	1	1	3	0	7

**Figures** 



**Fig. 1.** Phylogenetic analysis of CDF transporter family proteins from various selected algal taxa in comparison to prokaryotic, animal (*H. sapiens*), land plant (*A. thaliana*) and yeast (*S. cerevisiae*) proteins. The genome identifier or accession numbers of all sequences used are provided in Table S1 (ESI). Coloured symbols indicate the major taxonomic domains of the species where the protein sequence is derived. Five major groups are highlighted. The tree was constructed using the maximum likelihood method and derived from alignments of full length amino acid sequence. A consensus tree following 1000 bootstrap replications is shown, with bootstrap percentage values indicated at the nodes of major branches. The branch length scale bar indicates the evolutionary distance of one amino acid substitution per site.



**Fig. 2.** A two-dimensional cluster analysis of sequences representing the phenetic sequence relationship of CDF transporter family proteins from various selected algal taxa in comparison to prokaryotic, animal (*H. sapiens*), land plant (*A. thaliana*) and yeast (*S. cerevisiae*) proteins. Each symbol represents a full length amino acid sequence of an individual CDF protein and is coloured on the basis of the phylogenetic groups shown in Fig. 1. Grey lines represent protein connections with reciprocal BLAST hits at P < 10<sup>-5</sup>. Six proteins are not shown (five Group 2 members: Psp\_584138, Tp\_22171, Es\_18100, Osp\_198761, Osp\_422041, and unassigned ScZRG17) as they did not cluster with any of the other sequences and are on the periphery of the plot off the scale.



**Fig. 3.** Comparison of conserved CDF residues within transmembrane (TM) domain 2 (**a**) and TM domain 5 (**b**) for Group 1 (Zn-CDF) and Group 5 proteins. Sequence logo representations of 18 residues within each TM domain determined for all of the Group 1 sequences and all of the Group 5 sequences. Asterisks indicate the positions of conserved active site 'A' residues.



**Fig. 4.** Analysis of Group 1 Zn-CDF proteins. (**a**) Phylogenetic analysis of Zn-CDF family proteins from various selected species. The coloured symbols indicating the major taxonomic domains of each species are as defined in Fig. 1. Proteins highlighted in red possess an obvious histidine-rich domain. (**b**) Topology models of selected Group 1 Zn-CDF proteins indicating the predicted transmembrane (TM) spans and the histidine-rich domain within the major cytosolic loop between TM4 and TM5.



**Fig. 5.** Expression the CrMTP family genes in *C. reinhardtii* cells grown in standard TAP medium at exponential growth stage. Expression of the selected mRNA transcripts as determined by real-time PCR was calculated as a percentage of *CBLP* expression. Data points are means ( $\pm$ SEM) calculated from biological triplicates. Bars sharing the same lowercase letter indicate no significant difference between treatments (P > 0.05) as determined by one-way ANOVA and Tukey post-hoc test.



**Fig. 6.** Heterologous expression of *CrMTP1*, *CrMTP2*, *CrMTP3* and *CrMTP4* in yeast strains. (a) Suppression of Zn, Co and Mn hypersensitivity of the Zn- and Co-sensitive *zrc1cot1* yeast mutant and the Mn-sensitive *pmr1* mutant by heterologous expression of *CrMTP* cDNA in comparison with empty vector (piUGpd) and WT (BY4741). Saturated liquid cultures of yeast strains were serially diluted to the indicated cell densities then spotted onto SD –Ura selection medium with or without added Zn, Co or Mn. Yeast growth is shown after 3 d. A representative experiment is shown. (b) RT-PCR of *MTP1*, *MTP2*, *MTP3* and *MTP4* expression in each yeast strain.



**Fig. 7.** Heterologous expression of *CrMTP1*, *CrMTP2*, *CrMTP3* and *CrMTP4* in wild type (BY4741) yeast. Cell density measurement of strains expressing each MTP cDNA in comparison with empty vector (piUGpd) grown after 3 d in liquid SD –Ura medium either without metal exposure (**a**) or with 10 mM Zn (**b**), 10 mM Co (**c**), or 20 mM Mn (**d**). Data points are means (±SEM) from triplicate samples. Bars with an asterisk indicate significant difference between the empty vector control (piUGpd) strain (P < 0.05) as determined by one-way ANOVA and Tukey post-hoc test.



**Fig. 8.** Heterologous expression of *CrMTP1*, *CrMTP2*, *CrMTP3* and *CrMTP4* in wild type (BY4741) yeast. Dry weight biomass measurement of strains expressing each *CrMTP* cDNA in comparison with empty vector (piUGpd) grown after 3 d in liquid SD –Ura medium either without metal exposure or with added Zn (**a**), Co (**b**), or Mn (**c**). Data points are means (±SEM) from triplicate samples. Bars with an asterisk indicate significant difference between the empty vector control (piUGpd) strain (P < 0.05) as determined by one-way ANOVA and Tukey post-hoc test.



**Fig. 9.** Zn and Mn uptake into wild type (BY4741) yeast expressing *CrMTP1*, *CrMTP2*, *CrMTP3* and *CrMTP4*. Metal content measurement of strains expressing each *CrMTP* cDNA in comparison with empty vector (piUGpd) grown after 3 d in liquid SD –Ura medium with added 20 mM Zn (**a**) or 20 mM Mn (**b**). Data points are means (±SEM) from triplicate samples. Bars with an asterisk indicate significant difference between the empty vector control (piUGpd) strain (P < 0.05) as determined by one-way ANOVA and Tukey post-hoc test.