



**Manchester
Metropolitan
University**

Dos'Santos, T and Bishop, C and Thomas, C and Comfort, P and Jones, PA (2019) The effect of limb dominance on change of direction biomechanics: A systematic review of its importance for injury risk. *Physical Therapy in Sport*, 37. pp. 179-189. ISSN 1466-853X

Downloaded from: <https://e-space.mmu.ac.uk/626015/>

Version: Accepted Version

Publisher: Elsevier

DOI: <https://doi.org/10.1016/j.ptsp.2019.04.005>

Please cite the published version

<https://e-space.mmu.ac.uk>

THE EFFECT OF LIMB DOMINANCE ON CHANGE OF DIRECTION BIOMECHANICS: A SYSTEMATIC REVIEW OF ITS IMPORTANCE FOR INJURY RISK



ABSTRACT

Objective: To critically evaluate the effect of limb dominance on change of direction (COD) biomechanics associated with increased ACL injury-risk.

Methods: A systematic review of the literature was conducted using Medline and Sport DISCUS databases. Studies that compared COD biomechanics (lower-limb/whole-body kinetics/kinematics) between limbs, contained an approach run, and included physically active participants were included.

Results: Of the 456 articles identified, six were included. All studies investigated a cutting action, while the majority defined limb dominance as the preferred kicking limb, whereas one study defined limb dominance as preferred push-off cutting limb. Conflicting observations were found, with one study indicating the non-dominant and one study indicating the dominant limb displayed biomechanical deficits associated with increased non-contact ACL injury-risk during COD. Conversely, the remaining studies demonstrated no significant or substantial differences in COD biomechanics between limbs.

Conclusions: Female soccer players, male rugby players, and female handball players exhibit subtle side-to-side differences when performing cutting manoeuvres. However, the limb displaying high-risk mechanics is inconsistent within and between studies and populations. Thus, it remains inconclusive for COD that limb dominance is an ACL injury-risk factor and whether a particular limb is of heightened injury-risk.

Level of evidence: Level 2, Systematic review

Keywords: asymmetries; anterior cruciate ligament; cutting; limb preference

Highlights:

- Female soccer players, male rugby players, and female handball players exhibit subtle side-to-side differences when performing cutting manoeuvres.
- The limb displaying “high-risk” mechanics is inconsistent within and between studies and populations.
- It is inconclusive that limb dominance is an ACL injury-risk factor and whether a particular limb is of heightened injury-risk during cutting.

- Practitioners should screen change of direction biomechanics in both push-off limbs to identify biomechanical deficits associated with non-contact ACL injury risk, so informed individualised preventative training interventions can be created.

1. INTRODUCTION

Central to the success of many multidirectional sports, the ability for athletes to change direction quickly and safely is of great importance.^{12, 45, 71, 105, 123, 132, 142, 149} Change of direction (COD) actions such as side-steps, crossover cuts, and pivots are regularly performed in multidirectional sports,^{12, 50, 128, 142, 143, 148, 151} and are often linked to decisive moments such as, evading an opponent to penetrate the defensive line in rugby (tackle-break success in rugby),^{102, 148, 151} getting into space to receive a pass in netball,⁴⁸ or creating goals in soccer.⁴⁵ However, of concern, directional changes are inciting events associated with non-contact anterior-cruciate ligament (ACL) injury.^{13, 16, 23, 43, 76, 99, 108, 147} This occurrence could be explained by the fact that COD actions have the propensity to create hazardous multiplanar knee joint loading when the foot is planted, such as high knee abduction moments (KAMs) and internal rotation moments (IRMs);^{7, 34, 35, 67} both of which can increase ACL strain.^{3, 73, 85, 107, 133} Importantly, knee joint loading during directional changes is exacerbated when biomechanical deficits and ‘high-risk’ mechanics such as lateral trunk flexion,^{35, 49, 64, 66} knee valgus,^{66, 68, 77, 92, 134} extended knee positions,^{32, 76} wide foot plants,^{35, 59, 66, 77} and high GRFs^{67, 131, 134, 136} during weight acceptance of COD tasks are exhibited. As such, ensuring athletes have the capability to change direction safely (i.e. optimal mechanics/frontal plane alignment) from both limbs by avoiding these ‘high-risk’ postures is a viable strategy to reduce ACL injury-risk.^{40, 47, 61, 109}

Anterior cruciate ligament injury is a serious, potentially career-threatening injury with negative health,^{61, 81} psychological,^{61, 79} and economic^{30, 61, 119} implications for athletes and the general population. Although the mechanisms of non-contact ACL injury are multifactorial,^{61, 120} strength, neuromechanical, and dynamic control between-limb differences (side to side differences/asymmetries), and overall lower-limb dominance has been suggested to be a potential ACL injury risk factor.^{14, 62, 63, 89, 110, 111, 116} Limb dominance, also known as lateral preference or laterality,^{22, 82} refers to the concept that humans will preferentially use one side of the body when performing a motor task, typically resulting in a more skilful and therefore dominant side.^{22, 82} A preferred leg to kick a ball is typically used to indicate limb/skill dominance and as such, practitioners and researchers are interested whether a particular leg is at a heightened injury risk. For example, seminal work from Hewett et al⁶² found significantly greater asymmetries (dominant [D] vs non-dominant [ND]) in landing peak KAM in nine youth

females athletes (soccer, basketball, volleyball) who injured their ACL compared to uninjured. Notably, six of the nine athletes injured their D limb, defined as their preferred kicking leg. Additionally, prospective research by Paterno et al¹¹⁶ reported athletes with a previous ACL injury that sustained a second ACL injury exhibited greater asymmetries (4.1 times) in landing knee extensor moments between limbs. Moreover, Kyritsis et al⁷⁸ showed previously injured athletes that did not meet return to play criteria in 6 tests (three of which required >90% limb symmetry indexes in hopping tasks) were four times more likely to sustain a second ACL rupture. Therefore, reducing between-limb biomechanical deficits could be a potential training strategy to reduce the relative risk of non-contact ACL injury.

Retrospective analysis of ACL injuries (i.e., questionnaires and interviews) report conflicting findings regarding whether the D or ND limb is at a greater risk of ACL injury. For example, previous studies have refuted the connection between limb dominance and ACL injury in athletes from multiple sports,^{88, 104} whereas limb dominance may serve as an aetiological risk factor regarding non-contact ACL injuries in soccer players and skiers.^{14, 126} Previously, researchers reported a greater occurrence of ACL injuries in the D limb compared to the ND limb (18 vs 8) in female soccer players;⁴⁴ however, they did not delineate between non-contact and contact ACL injuries. In contrast, Brophy et al¹⁴ demonstrated 74% (20/27) of males sustained a greater proportion of non-contact ACL injuries to the D limb, compared to 32% (10/31) in females. Thus, female soccer players were more likely to injure their ACL in the ND (supporting/stance) limb, whereas males demonstrated the opposite and a greater trend in the kicking limb. Corroborating the aforementioned findings, Boden et al¹³ found 68% of female athletes from multiple sports sustained a non-contact ACL injury in their ND limb, with a similar trend documented for female skiers compared to males (63% vs. 45%, $p = 0.020$).¹²⁶ As such, arguably, limb dominance may play a gender-based role in non-contact ACL injury in soccer players and female skiers; however, it is important to note that the retrospective analysis ACL injuries do not imply cause and effect; thus, further prospective research is required to confirm the role of limb dominance as an ACL injury risk factor.

While biomechanical investigations into COD biomechanics have considered the effect of sex,^{4, 29, 46, 92-94, 134, 135} approach velocity,^{31, 75, 77, 103, 146} anticipation,^{6, 24, 27, 34, 72, 74, 80, 112} and COD angle^{28, 52, 53, 58, 60, 129, 130, 134}, an emerging area of research is the effect of limb dominance on COD biomechanics.^{5, 21, 51, 86, 98, 117} Research into between-limb differences during COD provides further insight into the potential mechanisms of non-contact ACL injury. From both performance and risk of injury perspectives, it would be advantageous for athletes to have the

capacity to change direction safely and quickly from both limbs, given the unpredictable nature of multidirectional sports.^{36,37} However, as athletes can display strength, neuromechanical, and dynamic control deficits between limbs,^{10, 19, 33, 62, 63, 70, 90, 96, 116, 144} these deficits, hypothetically, could lead side to side asymmetries in COD biomechanics, whereby a particular limb could display ‘higher-risk’ mechanics, thus increased injury risk. Several studies have examined the effect of limb dominance on COD biomechanics^{5, 21, 51, 86, 98, 117} in an attempt to establish whether a particular limb displays greater biomechanical deficits associated with increased ACL injury risk. To the best of our knowledge, however, a systematic review and critical evaluation of the literature that has examined the effect of limb dominance on COD biomechanics does not exist.

The purpose of this systematic review, therefore, was to critically evaluate the literature to date, which has examined the effect of limb dominance on COD biomechanics associated with increased risk of injury, and to highlight the limitations, considerations, and future directions for research to improve our understanding regarding the effect of limb dominance and biomechanical asymmetries during COD. For the purpose of the review, limb dominance is defined as the leg which an athlete would prefer to kick a ball with (unless stated otherwise) and is synonymous with previous studies that used the terms limb preference,^{18, 145} leg preference,^{19, 21, 25} and leg dominance.⁵¹ Moreover, in this review, KAM and knee abduction angle (KAA) are synonymous with knee valgus moment and knee valgus.

2. METHODS

2.1 Literature search strategy

A literature search was performed using Medline and Sport DISCUS databases. Figure 1 provides a schematic of the search methodology in accordance to Prisma guidelines.⁹⁷ Search terms were as follows:

- 1; “limb dominance”, or “leg dominance”, or “leg preference”, or “limb preference”, or “asymmetries” or “side to side differences”, or “symmetry” AND
- 2; “change of direction”, or “cutting”, or “cut”, or “sidestep”, or “turning”, or “side-step”, or “agility”, “or multidirectional speed”

Bibliographies of potentially eligible studies were hand searched to identify any additional studies and citation tracking on Google Scholar was used to identify any additional material. Following the search, two authors from the current review independently screened each article

for inclusion. The screening process consisted of: 1, screening for duplicates; 2, screening the title; 3, screening the abstract; and 4, screening the full paper using the inclusion and exclusion criteria. If the two authors were not in agreement with the inclusion/exclusion criterion of the study, a third author independently reviewed the study and a discussion occurred until consensus was reached. The final search date was January 10th, 2019.

2.2 Study selection

Studies were included if they met the following criteria:

1. Investigated preplanned or unplanned cutting or turning tasks that contained an approach run
2. Compared COD biomechanics between D and ND limbs
3. Investigated lower-limb and/or whole-body kinetics/kinematics
4. Included participants who participated in sport or physical activity
5. Full-text in a peer reviewed journal, in English

Studies that failed to meet the abovementioned criteria were subsequently removed.




Figure 1. Flow diagram illustrating the different phases of the systematic review; based on PRISMA recommendations. 3D: Three-dimensional; GRF: Ground reaction force

2.3 Assessment of study quality

Following the article search and examination, a methodological quality assessment was performed based on the scale (Table 1) created by Brown et al²⁰ which conducted a similar systematic review regarding the effect of anticipation on knee mechanics during side-stepping. The scale by Brown et al²⁰ is argued to be more specific for evaluating COD biomechanical studies in contrast to the Delphi, Physiotherapy Evidence Database, or Cochrane scales, because a large proportion of studies would fail to achieve many of the criteria of the aforementioned scales such as random allocation, assessor blinding, and subject blinding. As such, each article was assessed against a nine-item scale (Table 1) comprising of an 18 point-scoring system (ranging from 0 to 18) where 0 = clearly no; 1 = maybe or inadequate information; and 2 = clearly yes.

Table 1. Methodological quality assessment of limb dominance COD studies

Question	Criteria	Bencke et al. ⁵	Brown et al. ²¹	Marshall et al. ⁸⁶	Greska et al. ⁵¹	Pollard et al. ⁵⁷	Mok et al. ⁹⁸
1	Power analysis was performed and justification of study sample size	0	2	0	2	0	0
2	Athlete demographics were clearly defined: gender, age, body height, and body mass at time of test	2	2	2	2	2	2
3	Athlete characteristics were clearly defined: sport, experience or activity level and level of play	1	2	2	2	2	2
4	Inclusion and exclusion criteria were clearly stated for athletes	0	2	1	2	2	0
5	Proper training and practice trials of the test were given to the athletes allowing for adequate familiarisation	1	2	1	1	1	2
6	Methods were described in great detail to allow replication of the test. Testing devices, n of trials, n and duration of rest, speed, angle of COD	1	2	2	2	2	1
7	Test-retest reliability of measurement device reported	0	0	2	0	0	2
8	Outcome variables clearly defined	1	2	2	2	2	1
9	Statistical analyses were appropriate	1	2	2	2	1	1
Total score (maximum 18)		7 (39%)	16 (89%)	14 (78%)	15 (83%)	12 (67%)	11 (61%)

Key: n: number; COD: Change of direction; 0: clearly no; 1: maybe or inadequate information; 2: clearly yes

3. RESULTS

Initial database searches resulted in the identification of 451 articles, with an additional 5 articles through bibliographies, citation tracking, and hand searching (Figure 1). After removing duplicates, 419 articles were retained for initial screening. Title and abstract screening resulted in 354 articles excluded. The remaining 65 articles were further examined using the inclusion/ exclusion criteria, and 59 studies were excluded, resulting in six studies included to examine the effect of limb dominance on COD biomechanics^{5, 21, 51, 86, 98, 117} (Figure 1 and Table 1). Methodology quality scores ranged from seven (39%) to sixteen (89%) (Table 1).

Four studies examined female athletes,^{5, 21, 51, 98} one study examined male athletes,⁸⁶ and one study used a mixed cohort.¹¹⁷ In addition, sporting populations varied with two studies investigating female collegiate soccer players,^{21, 51} one study investigating male international rugby players,⁸⁶ one study investigating (mixed) recreationally active,¹¹⁷ one study in female handball,⁵ while one study examined a mixture of elite female soccer and handball players.⁹⁸ The most common angle of COD was 45°,^{21, 51, 117} followed by 75°,⁸⁶ while two studies used a sports-specific cut^{5, 98} but did not specify COD angle. The majority defined limb dominance as the preferred kicking limb, whereas one study⁵ defined limb dominance as preferred push-off cutting limb (Table 2).

In terms of the effect of limb dominance on COD biomechanics associated with increased injury-risk, conflicting observations were found (Table 2). One study indicated that the ND²¹ limb and one study indicated the D⁸⁶ limb displayed biomechanical deficits associated with increased non-contact ACL injury risk during COD. Conversely, the remaining studies^{5, 51, 98, 117} demonstrated no significant or substantial differences in COD biomechanics between limbs (Table 2).

Table 2. Summary of research that has examined the effect of limb dominance on COD biomechanics in multidirectional athletes

Study	Subjects	COD task	Results
Bencke et al. ⁵	24 young female handball athletes	Handball specific cut – 5 step run up (COD angle not provided) D limb – based on preferred push- off limb	D vs ND <ul style="list-style-type: none"> ↔ ($p > 0.05$) hip and knee joint angles at IC; but small differences in D GCT (ES = 0.22), hip adduction angle (ES = -0.33), hip internal rotation angle (ES = 0.30), and knee flexion (ES = -0.33) ↔ ($p > 0.05$) hip and knee joint moments 100 ms after IC; but small differences in hip flexion moment (ES = 0.20), hip extension moment (ES = 0.27), hip adduction moment (ES = 0.54), and knee flexion moment (ES = -0.20) Trivial differences (ES = 0.00) for hip internal rotation, KAM, and knee internal rotation moment (only 1 d.p provided)
Brown et al. ²¹	16 female collegiate soccer NCAA Division I	Pre-planned 45° cut 5-m approach distance 4.5±0.5m.s ⁻¹ approach velocity D limb – kicking	ND vs D <ul style="list-style-type: none"> Knee flexion angles: slightly smaller at IC (ES = 0.19, -5.3%), WA (ES = 0.28, -4.5%), PPO (ES = 0.36, -3.5%) KAA: slightly greater WA (ES = 0.10, +9.8%) Knee internal rotation angle: greater WA (ES = 0.64, + 32%), slightly greater peak PO (ES = 0.58, + 25%), and final PO (ES = 0.22, + 20%) Knee extensor moments: slightly lower during peak PO (ES = 0.31, -5.8%) and final push-off (ES = 0.30, -22%) KAMs slightly greater WA (ES = 0.22, + 19%) Knee IRM slightly lower during PPO (ES = 0.42, -19%) and final PO (ES = 0.18, -11%) Peak power absorption and peak knee flexion velocity slightly greater (ES = 0.14, + 5.1% and ES = 0.09, + 2.7%) Peak power production and peak knee extension velocity were slightly lower (ES = 0.34, -8.3% and ES = 0.21, -5.2%)
Marshall et al. ⁸⁶	Twenty elite injury free international male rugby union players (11 forwards and 9 backs)	Pre-planned 75° cut (approach distance not provided)	<ul style="list-style-type: none"> ↔ 27/28 variables, but AI values ranged from 1–49 % (ES = 0.02-0.60) ND ↑ ankle IRM ($p = 0.04$, 67%, ES = 0.75) D ↑ KAA (ES = 0.23, AI = 21%), ↑ KAM (ES = 0.23, AI = 8%), ↑ knee IRM (ES = 0.43, AI = 29%), ↓ knee flexion angle (ES = 0.35, AI = 5%), and ↓ GCT (ES = 0.60, AI = 9%) ND ↑ vGRF (ES = 0.48, AI = 11%), ↑ mlGRF (ES = 0.25, AI = 14%), ↑ longitudinal GRF (ES = 0.31, AI = 7%) <i>ACP – across the whole waveform:</i> ND ↑ ankle IRM ($p = 0.02 - 0.04$, ES = 0.52) from 23-38% of the movement. ND ↑ dorsi-flexion during the latter stages (78–94 %) of the cut PO ($p = 0.011$, ES = 0.57)
Greska et al. ⁵¹	20 Female collegiate soccer players	Unanticipated 45° cut – 6-m approach distance Minimum 3m.s ⁻¹ approach velocity D limb – kicking	<ul style="list-style-type: none"> D ↑ peak knee flexion angle ($p = 0.034$, ES = 0.69) ↔ ($p > 0.05$) in hip and knee kinetics or kinematics, GRF, or EMG, but: D ↑ peak vGRF (ES = 0.39), but ND ↑vGRF at peak KAM (ES = 0.60) ND ↑ KAA at IC and peak (ES = 0.21-0.23) D ↑ peak KAM (ES = 0.26) GCT between limbs ($p > 0.004$), D ↑ (ES = 0.45) ↔ Approach velocities (ES = 0.06) Trivial to small differences to EMG activation (%MVIC) pre-contact (ES = 0.02-0.55) and IC (ES = 0.04-0.40) Trivial to small differences to EMG activation time course pre-contact (ES = 0.13-0.30) and peak stance (ES = 0.10-0.56)
Pollard et al. ¹¹⁷	31 healthy participants (15 males and 16 females)	Pre-planned 45° cut (approach distance and velocity not provided) D limb – kicking	D vs ND <ul style="list-style-type: none"> D ↓ knee internal rotation angle from IC to 40% ($p < 0.05$, ES = 0.74) and IC to pk knee flexion ($p < 0.05$, ES = 0.61) ↔ hip and knee kinematics (sagittal, transverse, and frontal) ($p > 0.05$, ES ≤ 0.18) ↔ hip and knee kinematics (sagittal, transverse, and frontal) ($p > 0.05$, ES ≤ 0.39) (note some variables are presented to only 1 d.p)

Mok et al. ⁹⁸	19 elite Female handball and 22 female soccer	Sports- specific cut 6-m approach distance (COD angle not provided)	<ul style="list-style-type: none">• 4/33 variables significantly different ($p < 0.05$)• Significant difference for peak hip abduction angle, peak knee internal rotation angle, peak knee valgus moment and peak knee flexion moment (Does not provide descriptive data for ND limb, thus direction of asymmetry cannot be established)
--------------------------	---	---	---

Limb D - kicking

Key: ↔: no significant differences; ↑: greater; ↓: lower; D: Dominant; ND: Non-dominant; KAM: Knee abduction moment; KAA: Knee abduction angle; GRF: Ground reaction force; vGRF: Vertical GRF; hGRF: Horizontal GRF; mGRF: Medio-lateral GRF; IC: Initial contact; PFC: Penultimate foot contact; FFC: Final foot contact; SS: Sidestep; ES: Effect size; WA: Weight acceptance; PO: Push-off; GCT: Ground contact time; EMG: Electromyography; MVIC: Maximum voluntary isometric contraction; COD: Change of direction; AI: Asymmetry index; ACP: Analysis of characterising phases; IC: Initial contact; d.p: decimal place; IRM: Internal rotation moment

4. DISCUSSION

The aim of this systematic review was to critically evaluate the literature to date, which has examined the effect of limb dominance on COD biomechanics associated with increased risk of injury. A secondary aim was and to highlight the limitations, considerations, and future directions for research to improve our understanding regarding the effect of limb dominance and biomechanical asymmetries during COD. As six studies were included in the final analysis, the effect of limb dominance on COD biomechanics will be discussed in sport-specific sections, relative to the sample in the included studies.

4.1 Soccer players

Given the findings from Brophy et al¹⁴ that limb dominance may play a gender-based role in non-contact ACL injury risk in soccer players, two studies have compared COD biomechanics between limbs for a greater understanding into the potential mechanisms of ACL injury which met the criteria for this review (Table 2). Brown et al²¹ conducted a comprehensive comparison of knee kinetics and kinematics between limbs during a pre-planned 45° cut in female collegiate soccer players. The authors observed subtle differences in knee kinematics and kinetics (Table 2) between limbs, reporting small differences in knee flexion angle, and slightly greater KAAs, knee internal rotation angle, and KAMs during weight acceptance in the ND limb. In addition, slightly lower knee extensor moments and knee IRMs were reported in the ND limb, while peak power absorption and peak knee flexion velocity in the ND leg were slightly greater than the D limb (Table 2). Moreover, peak power production and peak knee extension velocity in the ND limb were slightly lower than those in the D limb (Table 2), indicating that the push-off was executed faster in the D limb. Unfortunately, the authors did not examine centre of mass velocity at toe-off; therefore, it is uncertain whether the athletes displayed superior exit velocity, thus performance from the D limb. Collectively, these results indicate that the ND limb exhibits biomechanical deficits associated with greater risk of non-contact ACL injury, which may partially support the greater ND limb ACL incidence rates in female soccer players reported by Brophy et al.¹⁴

Investigating an unanticipated 45° cut in female collegiate soccer players, Greska et al⁵¹ demonstrated no significant differences between limbs in hip and knee moments, GRFs, and peak electromyography activation (Table 2). Interestingly, based on the effect sizes (Table 2), differences in neuromechanical characteristics were inconsistent between limbs. For example, the D limb displayed greater peak knee flexion angles, greater peak KAMs, and

greater peak vertical GRFs, while the ND limb exhibited greater KAAs (IC and peak) and greater vertical GRF at peak KAM (Table 2). Moreover, the authors found the time course of muscle activation was also different between limbs (Table 2). These findings contrast to Brown et al²¹ that found the ND to display potentially more hazardous knee kinetics and kinematics. However, it is worth noting that the conflicting and inconsistent findings between studies could be attributed to task differences because Brown et al²¹ examined a pre-planned 45° cut, compared to an unanticipated 45° cut investigated by Greska et al.⁵¹ Furthermore, greater approach velocities were reported by Brown et al²¹ compared to Greska et al⁵¹ ($4.5 \pm 0.5 \text{ m}\cdot\text{s}^{-1}$ vs. $3.26 \pm 0.18 \text{ m}\cdot\text{s}^{-1}$), whereby velocities $\geq 4 \text{ m}\cdot\text{s}^{-1}$ have been recommended to screening and evaluating COD biomechanics in female athletes.¹⁴⁶ As such, further research is required examining between-limb differences in COD biomechanics during unanticipated cuts $\geq 4 \text{ m}\cdot\text{s}^{-1}$, in line with recommendations from Vanrenterghem et al.¹⁴⁶

4.2 Rugby players

Five of the six studies (Tables 1 & 2) have compared between-limb COD biomechanics based on discrete point analysis (DPA) (i.e. peak value during weight acceptance or push-off) (Table 2). This approach, however, leads to regional focus bias (focusing only on one aspect of the waveform),^{101, 113-115, 124} does not provide information regarding temporal differences (timing differences),^{121, 122, 138} and a large proportion of potentially valuable and meaningful information of the waveform (i.e. moment, GRF, angle waveform) is left unexamined.^{17, 118}

In light of the issues with DPA, Marshall et al⁸⁶ compared between-limb biomechanics in male international rugby players during a 75° cut using not only DPA, but also compared the continuous data between limbs using a method known as analysis of characterising phases (ACP: a continuous data analysis techniques that detects and examines phases of variance within a sample a sample of curves utilising the time, magnitude, and magnitude-time domains).^{121, 122} Interestingly, based on DPA, ankle IRM was the only variable (1/28) that showed a significant difference between limbs (Table 2). Notably, however, when the full waveform for variables were compared between D and ND limbs, ACP revealed ankle IRM was significantly greater for the ND limb from 23-38% of the movement, while significantly greater ankle dorsi-flexion angles during the latter stages of push off (78-94%) were also observed for the ND limb. Although not significantly different, based on DPA, slightly greater KAAs (asymmetry index (AI = 8%), knee IRMs (AI = 29%), and lower knee flexion angles (AI = 5%), and greater vertical GRF (AI = 11%) were observed for the D limb (Table 2). These findings are concerning because multiplanar joint loading can increase ACL strain,^{3, 73, 85, 107,}

¹³³ while model-based image-matching has revealed knee valgus, extended knee postures, and high vertical GRFs as characteristics of non-contact ACL injury during COD.⁷⁶ Furthermore, Marshall et al⁸⁶ observed a slightly longer ground contact time (GCT) for the ND limb (ES = 0.60, AI = 9%), potentially indicating slower performance from that push-off limb.^{38, 84, 87, 127} Therefore, rugby players may benefit from improving their unilateral reactive strength to improve COD speed performance from their ND limb,^{11, 83} though it is imperative that athletes have a solid foundation of strength to fully reap the benefits of unilateral reactive strength training.^{40, 140, 141}

4.3 Handball players and mixed cohorts

Only two studies have examined asymmetries in COD biomechanics in handball players (Table 2). Bencke et al⁵ reported no significant asymmetries in knee and hip joint angles, and no significant differences in hip and knee moments (flexion, extension, abduction, and internal rotation) during a sports-specific cut in female handball players (Table 2). However, it should be noted that Bencke et al⁵ failed to calculate AI values or effect sizes for the between-limb comparisons. Based on the descriptive data provided, though some data were presented only to one decimal place, small effect sizes would have been observed (Table 2), with the D limb displaying a slightly longer GCT, smaller knee and hip adduction angles, and a greater hip internal rotation angle. Furthermore, the D limb displayed slightly greater hip flexion, hip extension, hip adduction, and lower knee flexor moments; however, notably, trivial differences in KAM and knee internal rotation moments were observed. It is important to note, however, that Bencke et al⁵ defined limb dominance as the preferred push-off leg which may not necessarily correspond to the D kicking limb⁴¹ as used by the other limb dominance studies (Tables 2).

Similarly, Mok et al⁹⁸ also compared between-limb biomechanics during a sports-specific cut in elite female handball and soccer players, finding four of 33 variables displayed a significant difference (peak hip abduction angle, peak knee internal rotation angle, peak KAM, and peak knee flexion moment) between limbs. Unfortunately, the authors failed to provide the descriptive data for the ND limb; thus, it is unclear which limb displayed the greater biomechanical deficits, and the magnitude of the differences could not be established. Recently, Pollard et al¹¹⁷ reported knee internal rotation angle at 40% of stance and internal rotation angle displayed significant difference between limbs during a 45° cut in a mixed cohort (Table 2). Although effect sizes were not calculated, the ND limb displayed greater knee internal rotation

angles (ES = 0.61-0.74), while non-significant and trivial differences between limbs were observed for all other knee and hip kinematics (Table 2). In addition, the authors observed no significant differences in knee kinetics between limbs, though effect sizes ranged from trivial to small (Table 2); however, it is worth noting that values were only presented to one decimal place which may slightly alter effect size calculations.

5. CONSIDERATIONS, LIMITATIONS, AND FUTURE DIRECTIONS FOR RESEARCH

Collectively, it remains inconclusive whether limb dominance has a direct association with COD biomechanics connected with non-contact ACL injuries and whether a particular limb displays mechanics associated with greater ACL injury risk, with conflicted findings reported in female soccer, male rugby, and female handball athletes (Table 2). It is worth noting that the published literature to date are limited to female soccer, male rugby, female handball, and physically active populations (Tables 2); thus, these findings can only be extrapolated within this context and cannot be generalised to other athletic populations. Further research is required exploring the effect of limb dominance on COD biomechanics from different athletic populations, such as netball, American football, and Australian rules football given the importance of COD actions from both performance and risk of non-contact ACL injury perspectives.^{65, 99, 139, 142, 148} In addition, the published studies (Table 1) have only examined one COD task; thus, further research is required, investigating a greater range of COD tasks of different angles (i.e. 45° vs 90° vs 135 vs 180°) because the biomechanical demands of directional changes are angle-dependent.^{28, 40, 52, 53, 58, 60, 129, 130, 134}

Although failing to achieve the eligibility criteria for this systematic review, two studies,^{18, 145} presented as posters, observed greater KAAs in the ND limb during a 180° turn in female athletes (ES = 0.22-0.63), though trivial differences in peak KAMs (ES = 0.13) were observed. Furthermore, Brown et al¹⁹ compared sidestepping (COD angle not provided) biomechanics between D and ND limbs in thirty male academy rugby players. In contrast to Marshall et al,⁸⁶ the ND limb displayed mechanics associated with increased risk of injury with 8% less knee flexion (ES = -0.26), 17 % greater lateral trunk flexion (ES = 0.42), 11% greater COM distance relative to the joint centre (ES = 0.97), and 25% greater peak KAMs (ES = 0.43) observed.¹⁹ These results further highlight the inconsistency in ‘high-risk’ mechanics displayed

by the D and ND limb, and demonstrate the individual variation regarding the relationship between limb dominance and ACL injury risk.

Specifically, all eligible studies investigated a cutting action $\leq 75^\circ$ (Table 2), while no published peer-reviewed study has examined the braking characteristics and braking strategy differences between limbs over the penultimate foot contact (PFC). The PFC has been recently highlighted as playing a pivotal role in deceleration prior to changing direction and is also considered a preparatory step in facilitating effective directional changes.^{38, 39, 67, 69} Although published in poster format only, Thomas et al¹⁴⁵ found female soccer players displayed different braking strategies during a 180° turn between directions. Greater magnitudes of PFC horizontal braking forces (i.e., earlier braking with the ND limb) when changing direction from the D limb were demonstrated by the female soccer players and this also resulted in faster turning performance. Conversely, slower performance and greater emphasis and magnitudes of braking forces were displayed during the final foot contact when changing direction from the ND limb. The better performance and PFC dominant braking strategy demonstrated by the soccer players may be attributed to the similarities of the kicking action with their preferred leg, whereby the ND limb (i.e., stance) would experience greater eccentric and braking demands,^{1, 2, 15, 56, 91} and may therefore be a more skilful and efficient limb for braking. Consequently, further research is required exploring the between-limb differences in braking strategies during sharper directional changes, considering the PFC.

It should also be acknowledged that in order for an asymmetry to be deemed ‘real’, the between-limb difference must be greater than the variability for that variable.^{8, 42} A fundamental shortcoming of the COD limb dominance studies is the failure to report their variability statistics (i.e., coefficient of variation/ typical error) (Table 2). Moreover, only one study has directly calculated AI values,⁸⁶ though the equation used was in contrast to the recent recommendations of Bishop et al^{8, 9} for quantifying asymmetries, and practitioners and researchers are encouraged to use these recommendations to correctly calculate between-limb percentage imbalances. Briefly, when calculating AI values, if imbalances are not calculated in respect to the maximum value, then the percentage is mathematically incorrect.^{8, 9} Practitioners run the risk of incorrectly calculating AIs when defining the D limb as the kicking limb because the kicking limb may not necessarily be the limb that displays the greater value. Therefore, limb dominance should be defined as the limb that displays the highest value (i.e., highest peak KAM, KAA, vGRF, etc.) and subsequent AI and comparison between D and ND

limbs should be based on this approach to nullify the inconsistencies based on kicking limb preference.

A problematic issue in the COD limb dominance literature (Table 1 & 2), is no study has accounted for the effect of variability when interpreting between-limb differences. This absence has large implications because it is uncertain the between-limb differences presented in the aforementioned literature can be interpreted as ‘real’^{8, 42} (Table 2). Additionally, no study, to the best of our knowledge, has comprehensively examined the between-session reliability of the between-limb differences in COD biomechanics. It is not yet understood whether the magnitudes and directions of asymmetries between limbs are consistent between sessions (i.e. left limb consistently displays 15% AI for both session 1 and 2). This is important because if the magnitudes and directions of asymmetries are inconsistent between sessions, this could lead to different clinical diagnoses which could influence the future training for that athlete. Going forward, future research that investigates between-limb differences and asymmetries in COD biomechanics should account for the variability to establish ‘significant’ and ‘real’ differences between limbs.

While five of the six published studies have compared between-limb differences in COD biomechanics based on limb dominance (i.e., preferred leg to kick a ball), the assumption that the preferred kicking leg is a more coordinated and potentially stronger limb is flawed because previous studies have shown that the kicking limb may not necessarily be the stronger limb,^{41, 70, 90} or lead to superior functional performance from that limb,^{10, 41, 90} with research indicating that the preferred kicking limb does not necessarily correspond to faster COD speed performance from the same push-off limb.⁴¹ The published COD limb dominance studies have pooled their data with respect to limb dominance which can mask and conceal potentially meaningful between-limb differences because some athletes may display greater biomechanical deficits in the kicking limb, while the ND limb will display biomechanical deficits for other athletes. Going forward, it could be more suitable to compare COD biomechanics between limbs between preferred and non-preferred turning (push-off) limbs to identify if a particular limb is a greater risk of injury, as done by Bencke et al.⁵ This recommendation is extremely pertinent when wanting to examine the effect of limb dominance in athletes from sports where kicking is not a regularly performed action such as handball, basketball, netball, and rugby.

To the best of our knowledge, no study has examined whether between-limb force asymmetries (muscle strength asymmetries) affect COD biomechanics. Moreover, it has been documented that athletes display faster performance, based on completion times, from a push-off limb during cutting and turning tasks,^{36, 37, 41, 57, 84, 106, 125, 150} yet the underpinning kinetic and kinematic mechanisms which explain the differences between faster cutting or turning performance to a side are yet to be established and warrant further investigation. As such, comparing COD biomechanics between stronger and weaker limbs, or faster and slower sides, may provide different between-limb differences and subsequent evaluations regarding an athlete's risk of injury and performance. Performing such investigations would improve our understanding regarding the effect of between-limb force asymmetries on COD biomechanics and mechanical differences that explain differences in COD performance between limbs.

Finally, the majority of studies have conducted DPA (Tables 2); however, this method fails to account for the whole waveform for angle, moments, and GRF data whereby valuable information is left unexamined.^{17, 118} Only one study⁸⁶ has conducted statistical analysis across the whole waveform for kinetic and kinematic variables to provide greater insight into the temporal differences between-limbs. Thus, further research is warranted using temporal phase analysis,^{26, 95} ACP,^{121, 122} or one-dimensional statistical parametric mapping^{113, 124, 146} to explore differences between limbs across the whole waveform. Furthermore, no study to our knowledge, has examined the joint-joint coordination differences (angle-angle plots) between-limbs during COD, which may provide insight into the coupling behaviours between multiple segments,^{54, 55, 100, 101, 118, 137} and thus, is a recommendation for future research.

6. CONCLUSIONS

In conclusion, female soccer players, male rugby players, and female handball players, in general, exhibit subtle side to side differences when performing cutting manoeuvres, though the magnitude and direction of the differences are inconsistent within and between studies and populations. Based on the published literature to date, it remains inconclusive whether limb dominance is a risk factor associated with non-contact ACL biomechanical risk factors during COD. Studies to date have demonstrated conflicting results, indicating that the ND²¹ or D⁸⁶ limb may display biomechanical deficits associated with increased non-contact ACL injury risk, whereas previous research have demonstrated no significant or substantial differences between limbs during COD,^{5, 51, 117} refuting the notion of limb dominance as a risk factor of non-contact ACL injury. However, a fundamental flaw of the majority of the studies is that

limb dominance was defined as the preferred kicking limb, and thus the assumption that the kicking limb will be the more skilful and coordinated limb when changing direction is inherently incorrect. Nevertheless, practitioners are encouraged to screen COD biomechanics in both push-off limbs to identify biomechanical deficits associated with non-contact ACL injury risk, so informed individualised preventative training interventions can be created. Furthermore, as the aim of the sports medicine, sports science, and strength and conditioning is to improve athletic performance and minimise risk of injury, it would be advantageous that athletes have the capacity and are equally proficient in changing direction safely and quickly from both limbs, due to the unpredictable nature of multidirectional sport.

REFERENCES

1. Ball K. Loading and performance of the support leg in kicking. *Journal of science and medicine in sport*. 2013;16(5):455-459.
2. Ball KA. Kinematic comparison of the preferred and non-preferred foot punt kick. *Journal of Sports Sciences*. 2011;29(14):1545-1552.
3. Bates NA, Myer GD, Shearn JT, Hewett TE. Anterior cruciate ligament biomechanics during robotic and mechanical simulations of physiologic and clinical motion tasks: a systematic review and meta-analysis. *Clinical biomechanics*. 2015;30(1):1-13.
4. Beaulieu MLL, Lamontagne M, Xu L. Lower limb muscle activity and kinematics of an unanticipated cutting manoeuvre: a gender comparison. *Knee Surgery, Sports Traumatology, Arthroscopy*. 2009;17(8):968-976.
5. Bencke J, Curtis D, Krogshede C, Jensen LK, Bandholm T, Zebis MK. Biomechanical evaluation of the side-cutting manoeuvre associated with ACL injury in young female handball players. *Knee Surgery, Sports Traumatology, Arthroscopy*. 2013;21(8):1876-1881.
6. Besier TF, Lloyd DG, Ackland TR, Cochrane JL. Anticipatory effects on knee joint loading during running and cutting maneuvers. *Medicine and science in sports and exercise*. 2001;33(7):1176-1181.
7. Besier TF, Lloyd DG, Cochrane JL, Ackland TR. External loading of the knee joint during running and cutting maneuvers. *Medicine & Science in Sports & Exercise*. 2001;33(7):1168-1175.
8. Bishop C, Read P, Chavda S, Turner A. Asymmetries of the Lower Limb: The Calculation Conundrum in Strength Training and Conditioning. *Strength & Conditioning Journal*. 2016;38(6):27-32.
9. Bishop C, Read P, Lake J, Chavda S, Turner A. Inter-limb asymmetries: understanding how to calculate differences from bilateral and unilateral tests. *Strength & Conditioning Journal*. 2018.
10. Bishop C, Turner A, Read P. Effects of inter-limb asymmetries on physical and sports performance: a systematic review. *Journal of sports sciences*. 2017;36(10):1135-1144.
11. Bishop C, Turner A, Read P. Training Methods and Considerations for Practitioners to Reduce Interlimb Asymmetries. *Strength & Conditioning Journal*. 2018;40(2):40-46.
12. Bloomfield J, Polman R, Donoghue P. Physical demands of different positions in FA Premier League soccer. *Journal of Sports Science and Medicine*. 2007;6(1):63-70.
13. Boden BP, Dean GS, Feagin JA, Garrett WE. Mechanisms of anterior cruciate ligament injury. *Orthopedics*. 2000;23(6):573-578.
14. Brophy R, Silvers HJ, Gonzales T, Mandelbaum BR. Gender influences: the role of leg dominance in ACL injury among soccer players. *British Journal of Sports Medicine*. 2010;44:694-697.

15. Brophy RH, Backus SI, Pansy BS, Lyman S, Williams RJ. Lower extremity muscle activation and alignment during the soccer instep and side-foot kicks. *Journal of Orthopaedic & Sports Physical Therapy*. 2007;37(5):260-268.
16. Brophy RH, Stepan JG, Silvers HJ, Mandelbaum BR. Defending puts the anterior cruciate ligament at risk during soccer: a gender-based analysis. *Sports health*. 2015;7(3):244-249.
17. Brown C, Bowser B, Simpson KJ. Movement variability during single leg jump landings in individuals with and without chronic ankle instability. *Clinical biomechanics*. 2012;27(1):52-63.
18. Brown N, Donelon T, Smith LC, Jones PA. An investigation into the effect of limb preference on knee mechanics and braking strategy during a 180° change of direction task: An exploratory study. *J Sports Sci*. 2016;34:s57.
19. Brown SR. Lower-extremity symmetry influences knee abduction moments during sidestepping in rugby. *Br J Sports Med*. 2017:bjsports-2017-097868.
20. Brown SR, Brughelli M, Hume PA. Knee mechanics during planned and unplanned sidestepping: a systematic review and meta-analysis. *Sports Medicine*. 2014;44(11):1573-1588.
21. Brown SR, Wang H, Dickin DC, Weiss KJ. The relationship between leg preference and knee mechanics during sidestepping in collegiate female footballers. *Sports Biomechanics*. 2014;13(4):351-361.
22. Carpes FP, Mota CB, Faria IE. On the bilateral asymmetry during running and cycling-A review considering leg preference. *Physical therapy in Sport*. 2010;11(4):136-142.
23. Cochrane JL, Lloyd DG, Buttfield A, Seward H, McGivern J. Characteristics of anterior cruciate ligament injuries in Australian football. *Journal of Science and Medicine in Sport*. 2007;10(2):96-104.
24. Collins JD, Almonroeder TG, Ebersole KT, O'connor KM. The effects of fatigue and anticipation on the mechanics of the knee during cutting in female athletes. *Clinical Biomechanics*. 2016;35:62-67.
25. Condello G, Kernozek TW, Tessitore A, Foster C. Biomechanical Analysis of a Change-of-Direction Task in College Soccer Players. *International journal of sports physiology and performance*. 2016;11(1):96-101.
26. Cormie P, McBride JM, McCaulley GO. Power-time, force-time, and velocity-time curve analysis of the countermovement jump: impact of training. *The Journal of Strength & Conditioning Research*. 2009;23(1):177-186.
27. Cortes N, Blount E, Ringleb S, Onate JA. Soccer-specific video simulation for improving movement assessment. *Sports Biomechanics*. 2011;10(01):22-34.
28. Cortes N, Onate J, Van Lunen B. Pivot task increases knee frontal plane loading compared with sidestep and drop-jump. *Journal of sports sciences*. 2011;29(1):83-92.
29. Cronstrom A, Creaby MW, Nae J, Ageberg E. Gender differences in knee abduction during weight-bearing activities: A systematic review and meta-analysis. *Gait & posture*. 2016;49:315-328.
30. Cumps E, Verhagen E, Annemans L, Meeusen R. Injury rate and socioeconomic costs resulting from sports injuries in Flanders: data derived from sports insurance statistics 2003. *British journal of sports medicine*. 2008;42(9):767-772.
31. Dai B, Garrett WE, Gross MT, Padua DA, Queen RM, Yu B. The effect of performance demands on lower extremity biomechanics during landing and cutting tasks. *Journal of Sport and Health Science*. 2016;43(9):1680-1608.
32. Dai B, William EG, Michael TG, Darin AP, Robin MQ, Bing Y. The Effects of 2 Landing Techniques on Knee Kinematics, Kinetics, and Performance During Stop-Jump and Side-Cutting Tasks. *The American journal of sports medicine*. 2014;43(2):466-474.

33. Del Bel MJ, Fairfax AK, Jones ML, Steele K, Landry SC. Effect of limb dominance and sex on neuromuscular activation patterns in athletes under 12 performing unanticipated side-cuts. *Journal of Electromyography and Kinesiology*. 2017;36:65-72.
34. Dempsey AR, Lloyd DG, Elliott BC, Steele JR, Munro BJ. Changing sidestep cutting technique reduces knee valgus loading. *The American journal of sports medicine*. 2009;37(11):2194-2200.
35. Dempsey AR, Lloyd DG, Elliott BC, Steele JR, Munro BJ, Russo KA. The effect of technique change on knee loads during sidestep cutting. *Medicine & Science in Sports & Exercise*. 2007;39(10):1765-1773.
36. Dos' Santos T, Thomas C, Jones PA, Comfort P. Assessing Asymmetries in Change of Direction Speed Performance; Application of Change of Direction Deficit. *Journal of strength and conditioning research*. 2018:Published ahead of print.
37. Dos' Santos T, Thomas C, Jones PA, Comfort P. Asymmetries in isometric force-time characteristics are not detrimental to change of direction speed. *The Journal of Strength & Conditioning Research*. 2018;32(2):520-527.
38. Dos'Santos T, Thomas C, Jones AP, Comfort P. Mechanical determinants of faster change of direction speed performance in male athletes. *The Journal of Strength & Conditioning Research*. 2017;31:696-705.
39. Dos'Santos T, Thomas C, Comfort P, Jones P. The Role of the Penultimate Foot Contact During Change of Direction: Implications on Performance and Risk of Injury. *Strength & Conditioning Journal*. 2018:Published ahead of print.
40. Dos'Santos T, Thomas C, Comfort P, Jones PA. The effect of angle and velocity on change of direction biomechanics: an angle-velocity trade-off. *Sports Medicine*. 2018;48(10):2235-2253.
41. Dos'Santos T, Thomas C, Jones PA, Comfort P. Asymmetries in single and triple hop are not detrimental to change of direction speed. *Journal of Trainology*. 2017;6(2):35-41.
42. Exell TA, Irwin G, Gittoes MJ, Kerwin DG. Implications of intra-limb variability on asymmetry analyses. *Journal of Sports Sciences*. 2012;30(4):403-409.
43. Faude O, Junge A, Kindermann W, Dvorak J. Injuries in female soccer players a prospective study in the german national league. *The American journal of sports medicine*. 2005;33(11):1694-1700.
44. Faude O, Junge A, Kindermann W, Dvorak J. Risk factors for injuries in elite female soccer players. *British Journal of Sports Medicine*. 2006;40(9):785-790.
45. Faude O, Koch T, Meyer T. Straight sprinting is the most frequent action in goal situations in professional football. *Journal of sports sciences*. 2012;30(7):625-631.
46. Fedie R, Carlstedt K, Willson JD, Kernozek TW. Effect of attending to a ball during a side-cut maneuver on lower extremity biomechanics in male and female athletes. *Sports Biomechanics*. 2010;9(3):165-177.
47. Fox A. Change-of-Direction Biomechanics: Is What's Best for Anterior Cruciate Ligament Injury Prevention Also Best for Performance? *Sports Med*. 2018;48(8):1799-1807.
48. Fox A, Spittle M, Otago L, Saunders N. Offensive agility techniques performed during international netball competition. *International Journal of Sports Science & Coaching*. 2014;9(3):543-552.
49. Frank B, Bell DR, Norcross MF, Blackburn JT, Goerger BM, Padua DA. Trunk and hip biomechanics influence anterior cruciate loading mechanisms in physically active participants. *The American journal of sports medicine*. 2013;41(11):2676-2683.
50. Green BS, Blake C, Caulfield BM. A comparison of cutting technique performance in rugby union players. *The Journal of Strength & Conditioning Research*. 2011;25(10):2668-2680.
51. Greska EK, Cortes N, Ringleb SI, Onate JA, Van Lunen BL. Biomechanical differences related to leg dominance were not found during a cutting task. *Scandinavian Journal of Medicine & Science in Sports*. 2016;27(11):1328-1336.

52. Hader K, Mendez-Villanueva A, Palazzi D, Ahmaidi S, Buchheit M. Metabolic Power Requirement of Change of Direction Speed in Young Soccer Players: Not All Is What It Seems. *PloS one*. 2016;11(3):e0149839.
53. Hader K, Palazzi D, Buchheit M. Change of Direction Speed in Soccer: How Much Braking is Enough? *Kineziologija*. 2015;47(1):67-74.
54. Hamill J, Haddad JM, McDermott WJ. Issues in quantifying variability from a dynamical systems perspective. *Journal of applied biomechanics*. 2000;16(4):407-418.
55. Hamill J, Palmer C, Van Emmerik REA. Coordinative variability and overuse injury. *Sports Medicine, Arthroscopy, Rehabilitation, Therapy & Technology*. 2012;4(1):45.
56. Hart NH, Newton RU, Weber J, et al. Functional Basis of Asymmetrical Lower-Body Skeletal Morphology in Professional Australian Rules Footballers. *Journal of strength and conditioning research*. 2018:Published Ahead of Print.
57. Hart NH, Spiteri T, Lockie RG, Nimphius S, Newton RU. Detecting deficits in change of direction performance using the preplanned multidirectional Australian Football League agility test. *The Journal of Strength & Conditioning Research*. 2014;28(12):3552-3556.
58. Havens K, Sigward SM. Whole body mechanics differ among running and cutting maneuvers in skilled athletes. *Gait & posture*. 2014;42(3):240-245.
59. Havens K, Sigward SM. Cutting mechanics: relation to performance and anterior cruciate ligament injury risk. *Medicine and science in sports and exercise*. 2015;47(4):818-824.
60. Havens K, Sigward SM. Joint and segmental mechanics differ between cutting maneuvers in skilled athletes. *Gait & posture*. 2015;41(1):33-38.
61. Hewett T. Preventive biomechanics: A paradigm shift with a translational approach to biomechanics. *Am J Sports Med*. 2017;45(11):2654-2664.
62. Hewett T, Myer GD, Ford KR, et al. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes a prospective study. *The American journal of sports medicine*. 2005;33(4):492-501.
63. Hewett TE, Ford KR, Hoogenboom BJ, Myer GD. Understanding and preventing acl injuries: current biomechanical and epidemiologic considerations-update 2010. *North American journal of sports physical therapy: NAJSPT*. 2010;5(4):234.
64. Jamison ST, Pan X, Chaudhari AMW. Knee moments during run-to-cut maneuvers are associated with lateral trunk positioning. *Journal of biomechanics*. 2012;45(11):1881-1885.
65. Johnston JT, Mandelbaum BR, Schub D, et al. Video analysis of anterior cruciate ligament tears in professional American football athletes. *The American journal of sports medicine*. 2018;46(4):862-868.
66. Jones P, Herrington L, Graham-Smith P. Technique determinants of knee joint loads during cutting in female soccer players. *Human movement science*. 2015;42:203-211.
67. Jones P, Herrington L, Graham-Smith P. Braking characteristics during cutting and pivoting in female soccer players. *Journal of Electromyography and Kinesiology*. 2016;30:46-54.
68. Jones P, Herrington L, Graham-Smith P. Technique determinants of knee abduction moments during pivoting in female soccer players. *Clinical Biomechanics*. 2016;31:107-112.
69. Jones P, Thomas C, Dos'Santos T, McMahon J, Graham-Smith P. The Role of Eccentric Strength in 180° Turns in Female Soccer Players. *Sports*. 2017;5(2):42.
70. Jones PA, Bampouras TM. A comparison of isokinetic and functional methods of assessing bilateral strength imbalance. *The Journal of Strength & Conditioning Research*. 2010;24(6):1553-1558.
71. Karcher C, Buchheit M. On-court demands of elite handball, with special reference to playing positions. *Sports Medicine*. 2014;44(6):797-814.
72. Khalid AJ, Ian Harris S, Michael L, Joseph H, Qu X. Effects of neuromuscular fatigue on perceptual-cognitive skills between genders in the contribution to the knee joint loading during side-stepping tasks. *Journal of sports sciences*. 2015;33(13):1322-1331.

73. Kiapour AM, Demetropoulos CK, Kiapour A, et al. Strain response of the anterior cruciate ligament to uniplanar and multiplanar loads during simulated landings: implications for injury mechanism. *The American journal of sports medicine*. 2016;44(8):2087-2096.
74. Kim JH, Lee K-K, Kong SJ, An KO, Jeong JH, Lee YS. Effect of anticipation on lower extremity biomechanics during side-and cross-cutting maneuvers in young soccer players. *The American journal of sports medicine*. 2014;42(8):1985-1992.
75. Kimura K, Sakurai S. A Sidestep Cut Preparation Strategy Decreases the External Load Applied to the Knee Joint. *International Journal of Sport and Health Science*. 2013;11:109-117.
76. Koga H, Nakamae A, Shima Y, et al. Mechanisms for noncontact anterior cruciate ligament injuries knee joint kinematics in 10 injury situations from female team handball and basketball. *The American journal of sports medicine*. 2010;38(11):2218-2225.
77. Kristianslund E, Faul O, Bahr R, Myklebust G, Krosshaug T. Sidestep cutting technique and knee abduction loading: implications for ACL prevention exercises. *British journal of sports medicine*. 2014;48(9):779-783.
78. Kyritsis P, Bahr R, Landreau P, Miladi R, Witvrouw E. Likelihood of ACL graft rupture: not meeting six clinical discharge criteria before return to sport is associated with a four times greater risk of rupture. *Br J Sports Med*. 2016;50:946-951.
79. Langford JL, Webster KE, Feller JA. A prospective longitudinal study to assess psychological changes following anterior cruciate ligament reconstruction surgery. *British journal of sports medicine*. 2009;43(5):377-378.
80. Lee MJC, Lloyd DG, Lay BS, Bourke PD, Alderson JA. Effects of different visual stimuli on postures and knee moments during sidestepping. *Med Sci Sports Exerc*. 2013;45(9):1740-1748.
81. Lohmander LS, Englund PM, Dahl LL, Roos EM. The long-term consequence of anterior cruciate ligament and meniscus injuries. *The American journal of sports medicine*. 2007;35(10):1756-1769.
82. Maloney SJ. The Relationship Between Asymmetry and Athletic Performance: A Critical Review. *Journal of strength and conditioning research*. 2018:Published ahead of print.
83. Maloney SJ, Richards J, Jelly L, Fletcher IM. Unilateral Stiffness Interventions Augment Vertical Stiffness And Change Of Direction Speed. *The Journal of Strength & Conditioning Research*. 2018:Published ahead of print.
84. Maloney SJ, Richards J, Nixon DG, Harvey LJ, Fletcher IM. Do stiffness and asymmetries predict change of direction performance? *Journal of sports sciences*. 2016;35(6):547-556.
85. Markolf KL, Burchfield DM, Shapiro MM, Shepard MF, Finerman GAM, Slauterbeck JL. Combined knee loading states that generate high anterior cruciate ligament forces. *Journal of Orthopaedic Research*. 1995;13(6):930-935.
86. Marshall B, Franklyn-Miller A, Moran K, et al. Biomechanical symmetry in elite rugby union players during dynamic tasks: an investigation using discrete and continuous data analysis techniques. *BMC sports science, medicine and rehabilitation*. 2015;7(1):13.
87. Marshall BM, Franklyn-Miller AD, King EA, Moran KA, Strike S, Falvey A. Biomechanical factors associated with time to complete a change of direction cutting maneuver. *The Journal of Strength & Conditioning Research*. 2014;28(10):2845-2851.
88. Matava MJ, Freehill AK, Grutzner S, Shannon W. Limb dominance as a potential etiologic factor in noncontact anterior cruciate ligament tears. *The journal of knee surgery*. 2002;15(1):11.
89. Maulder PS. Dominant limb asymmetry associated with prospective injury occurrence. *South African Journal for Research in Sport, Physical Education and Recreation*. 2013;35(1):121-131.
90. McGrath TM, Waddington G, Scarvell JM, et al. The effect of limb dominance on lower limb functional performance-- a systematic review. *Journal of sports sciences*. 2016;34(4):289-302.
91. McLean B, Tumilty D. Left-right asymmetry in two types of soccer kick. *British Journal of Sports Medicine*. 1993;27(4):260-262.

92. McLean SG, Huang X, van den Bogert AJ. Association between lower extremity posture at contact and peak knee valgus moment during sidestepping: implications for ACL injury. *Clinical Biomechanics*. 2005;20(8):863-870.
93. McLean SG, Lipfert SW, Van den Bogert AJ. Effect of gender and defensive opponent on the biomechanics of sidestep cutting. *Medicine and science in sports and exercise*. 2004;36(6):1008-1016.
94. McLean SG, Walker KB, van den Bogert AJ. Effect of gender on lower extremity kinematics during rapid direction changes: an integrated analysis of three sports movements. *Journal of Science and Medicine in Sport*. 2005;8(4):411-422.
95. McMahan JJ, Rej SJE, Comfort P. Sex differences in countermovement jump phase characteristics. *Sports*. 2017;5(1):8.
96. Mendiguchia J, Martinez-Ruiz E, Edouard P, et al. A multifactorial, criteria-based progressive algorithm for hamstring injury treatment. *Medicine & Science in Sports & Exercise*. 2017;49(7):1482-1492.
97. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS medicine*. 2009;6(7):e1000097.
98. Mok K-M, Bahr R, Krosshaug T. Reliability of lower limb biomechanics in two sport-specific sidestep cutting tasks. *Sports Biomechanics*. 2018;17:157-167.
99. Montgomery C, Blackburn J, Withers D, Tierney G, Moran C, Simms C. Mechanisms of ACL injury in professional rugby union: a systematic video analysis of 36 cases. *British Journal of Sports Medicine*. 2018;52(15):944-1001.
100. Mullineaux DR, Bartlett RM, Bennett S. Research design and statistics in biomechanics and motor control. *Journal of sports sciences*. 2001;19(10):739-760.
101. Mullineaux DR, Wheat J. Research methods: sample size and variability effects on statistical power. In: Payton CJ, Burden A, eds. *Biomechanical Evaluation of Movement in Sport and Exercise: The British Association of Sport and Exercise Sciences Guide*; Routledge; 2017.
102. Nedelec M, McCall A, Carling C, Legall F, Berthoin S, Dupont G. The influence of soccer playing actions on the recovery kinetics after a soccer match. *The Journal of Strength & Conditioning Research*. 2014;28(6):1517-1523.
103. Nedergaard NJ, Kersting U, Lake M. Using accelerometry to quantify deceleration during a high-intensity soccer turning manoeuvre. *Journal of sports sciences*. 2014;32(20):1897-1905.
104. Negrete RJ, Schick EA, Cooper JP. Lower-limb dominance as a possible etiologic factor in noncontact anterior cruciate ligament tears. *Journal of strength and conditioning research*. 2007;21(1):270-273.
105. Nimphius S. Increasing Agility. In: Joyce D, Lewindon D, eds. *High-Performance Training for Sports*. Champaign, IL.: Human Kinetics; 2014:185-198.
106. Nimphius S, Callaghan SJ, Sptieri T, Lockie RG. Change of direction deficit: A more isolated measure of change of direction performance than total 505 time. *The Journal of Strength & Conditioning Research*. 2016;30:3024-3032.
107. Oh YK, Lipps DB, Ashton-Miller JA, Wojtys EM. What strains the anterior cruciate ligament during a pivot landing? *The American journal of sports medicine*. 2012;40(3):574-583.
108. Olsen O-E, Myklebust G, Engebretsen L, Bahr R. Injury mechanisms for anterior cruciate ligament injuries in team handball a systematic video analysis. *The American journal of sports medicine*. 2004;32(4):1002-1012.
109. Padua DA, DiStefano LJ, Hewett TE, et al. National Athletic Trainers' Association Position Statement: Prevention of Anterior Cruciate Ligament Injury. *Journal of athletic training*. 2018;53:5-19.
110. Pappas E, Carpes FP. Lower extremity kinematic asymmetry in male and female athletes performing jump-landing tasks. *Journal of Science and Medicine in Sport*. 2012;15(1):87-92.

111. Pappas E, Shiyko MP, Ford KR, Myer GD, Hewett TE. Biomechanical Deficit Profiles Associated with ACL Injury Risk in Female Athletes. *Medicine and science in sports and exercise*. 2016;48(1):107-113.
112. Park E-J, Lee J-H, Ryue J-J, Sohn J-H, Lee K-K. Influence of anticipation on landing patterns during side-cutting maneuver in female collegiate soccer players. *Korean Journal of Sport Biomechanics*. 2011;21(4):391-395.
113. Pataky TC. Generalized n-dimensional biomechanical field analysis using statistical parametric mapping. *Journal of biomechanics*. 2010;43(10):1976-1982.
114. Pataky TC, Robinson MA, Vanrenterghem J. Vector field statistical analysis of kinematic and force trajectories. *Journal of biomechanics*. 2013;46(14):2394-2401.
115. Pataky TC, Robinson MA, Vanrenterghem J, Savage R, Bates KT, Crompton RH. Vector field statistics for objective center-of-pressure trajectory analysis during gait, with evidence of scalar sensitivity to small coordinate system rotations. *Gait & posture*. 2014;40(1):255-258.
116. Paterno MV, Schmitt LC, Ford KR, et al. Biomechanical measures during landing and postural stability predict second anterior cruciate ligament injury after anterior cruciate ligament reconstruction and return to sport. *The American journal of sports medicine*. 2010;38(10):1968-1978.
117. Pollard CD, Norcross MF, Johnson ST, Stone AE, Chang E, Hoffman MA. A biomechanical comparison of dominant and non-dominant limbs during a side-step cutting task. *Sports Biomechanics*. 2018:1-9.
118. Preatoni E, Hamill J, Harrison AJ, et al. Movement variability and skills monitoring in sports. *Sports Biomechanics*. 2013;12(2):69-92.
119. Quatman CE, Hewett TE. The anterior cruciate ligament injury controversy: is “valgus collapse” a sex-specific mechanism? *British journal of sports medicine*. 2009;43(5):328-335.
120. Quatman CE, Quatman-Yates CC, Hewett TE. A ‘Plane’Explanation of Anterior Cruciate Ligament Injury Mechanisms. *Sports Medicine*. 2010;40(9):729-746.
121. Richter C, Marshall B, Moran K. Comparison of discrete-point vs. dimensionality-reduction techniques for describing performance-related aspects of maximal vertical jumping. *Journal of biomechanics*. 2014;47(12):3012-3017.
122. Richter C, O’Connor NE, Marshall B, Moran K. Analysis of characterizing phases on waveforms: an application to vertical jumps. *Journal of applied biomechanics*. 2014;30(2):316-321.
123. Robinson G, O’Donoghue P, Nielson P. Path changes and injury risk in English FA Premier League soccer. *International Journal of Performance Analysis in Sport*. 2011;11(1):40-56.
124. Robinson MA, Vanrenterghem J, Pataky TC. Statistical Parametric Mapping (SPM) for alpha-based statistical analyses of multi-muscle EMG time-series. *Journal of Electromyography and Kinesiology*. 2015;25(1):14-19.
125. Rouissi M, Chtara M, Berriri A, Owen A, Chamari K. Asymmetry of the Modified Illinois Change of Direction Test Impacts Young Elite Soccer Players Performance. *Asian journal of sports medicine*. 2016;7(2): e33598.
126. Ruedl G, Webhofer M, Helle K, et al. Leg dominance is a risk factor for noncontact anterior cruciate ligament injuries in female recreational skiers. *The American journal of sports medicine*. 2012;40(6):1269-1273.
127. Sasaki S, Nagano Y, Kaneko S, Sakurai T, Fukubayashi T. The relationship between performance and trunk movement during change of direction. *Journal of sports science & medicine*. 2011;10(1):112-118.
128. Sayers M, Washington-King J. Characteristics of effective ball carries in Super 12 rugby. *International Journal of Performance Analysis in Sport*. 2005;5(3):92-106.
129. Schot P, Dart J, Schuh M. Biomechanical analysis of two change-of-direction maneuvers while running. *J Orthop Sports Phys Ther*. Dec 1995;22(6):254-258.

130. Schreurs MJ, Benjaminse A, Lemmink KA. Sharper angle, higher risk? The effect of cutting angle on knee mechanics in invasion sport athletes. *Journal of Biomechanics*. 2017;63:144-150.
131. Sell TC, Ferris CM, Abt JP, et al. Predictors of proximal tibia anterior shear force during a vertical stop-jump. *Journal of Orthopaedic Research*. 2007;25(12):1589-1597.
132. Sheppard JM, Dawes JJ, Jeffreys I, Spiteri T, Nimphius S. Broadening the view of agility: A scientific review of the literature. *J Aust Strength Conditioning*. 2014;22(3):6-25.
133. Shin CS, Chaudhari AM, Andriacchi TP. Valgus plus internal rotation moments increase anterior cruciate ligament strain more than either alone. *Medicine and science in sports and exercise*. 2011;43(8):1484-1491.
134. Sigward SM, Cesar GM, Havens KL. Predictors of frontal plane knee moments during side-step cutting to 45 and 110 degrees in men and women: implications for anterior cruciate ligament injury. *Clinical Journal of Sport Medicine*. 2015;25(6):529-534.
135. Sigward SM, Pollard CD, Havens KL, Powers CM. The influence of sex and maturation on knee mechanics during side-step cutting. *Medicine and science in sports and exercise*. 2012;44(8):1497-1503.
136. Sigward SM, Powers CM. Loading characteristics of females exhibiting excessive valgus moments during cutting. *Clinical Biomechanics*. 2007;22(7):827-833.
137. Stergiou N, Decker LM. Human movement variability, nonlinear dynamics, and pathology: is there a connection? *Human Movement Science*. 2011;30(5):869-888.
138. Stergiou N, Harbourne RT, Cavanaugh JT. Optimal movement variability: a new theoretical perspective for neurologic physical therapy. *Journal of Neurologic Physical Therapy*. 2006;30(3):120-129.
139. Stuelcken MC, Mellifont DB, Gorman AD, Sayers MG. Mechanisms of anterior cruciate ligament injuries in elite women's netball: a systematic video analysis. *J Sports Sci*. Aug 2016;34(16):1516-1522.
140. Suchomel TJ, Nimphius S, Bellon CR, Stone MH. The Importance of Muscular Strength: Training Considerations. *Sports Medicine*. 2018;48(4):765-785.
141. Suchomel TJ, Nimphius S, Stone MH. The Importance of Muscular Strength in Athletic Performance. *Sports Medicine*. 2016;46(10):1419-1449.
142. Sweeting AJ, Aughey RJ, Cormack SJ, Morgan S. Discovering frequently recurring movement sequences in team-sport athlete spatiotemporal data. *Journal of Sports Sciences*. 2017;35(24):2439-2445.
143. Taylor JB, Wright AA, Dischiavi SL, Townsend MA, Marmon AR. Activity Demands During Multi-Directional Team Sports: A Systematic Review. *Sports Medicine*. 2017;47(12):2533-2551.
144. Thomas C, Comfort P, Dos'Santos T, Jones PA. Determining bilateral strength imbalances in youth basketball athletes. *International journal of sports medicine*. 2017;38(09):683-690.
145. Thomas C, Dos'Santos T, Kyriakidou I, Cuthbert M, Fields C, Jones PA. An investigation into the effect of limb preference on knee mechanics and braking strategy during pivoting in female soccer players: an exploratory study. *The 8th Annual Strength and Conditioning Student Conference*. Middlesex, London, England; 2017.
146. Vanrenterghem J, Venables E, Pataky T, Robinson MA. The effect of running speed on knee mechanical loading in females during side cutting. *Journal of biomechanics*. 2012;45(14):2444-2449.
147. Walden M, Krosshaug T, Bjorneboe J, Andersen TE, Faul O, Hagglund M. Three distinct mechanisms predominate in non-contact anterior cruciate ligament injuries in male professional football players: a systematic video analysis of 39 cases. *Br J Sports Med*. Nov 2015;49(22):1452-1460.
148. Wheeler KW, Askew CD, Sayers MG. Effective attacking strategies in rugby union. *European Journal of Sport Science*. 2010;10(4):237-242.

149. Young WB, Dawson B, Henry GJ. Agility and change-of-direction speed are independent skills: Implications for training for agility in invasion sports. *International Journal of Sports Science and Coaching*. 2015;10(1):159-169.
150. Young WB, James R, Montgomery I. Is muscle power related to running speed with changes of direction? *Journal of Sports Medicine and Physical Fitness*. 2002;42(3):282-288.
151. Zahidi NNM, Ismail SI. Notational analysis of evasive agility skills executed by attacking ball carriers among elite rugby players of the 2015 Rugby World Cup. *Movement, Health & Exercise*. 2018;7(1):99-113.