

**Please cite the Published Version**

Comfort, P, Dos'Santos, T, Thomas, C, McMahon, JJ and Suchomel, TJ (2018) An investigation into the effects of excluding the catch phase of the power clean on force-time characteristics during isometric and dynamic tasks: An intervention study. *Journal of Strength and Conditioning Research*, 32 (8). pp. 2116-2129. ISSN 1064-8011

**DOI:** <https://doi.org/10.1519/JSC.0000000000002656>

**Publisher:** National Strength and Conditioning Association

**Version:** Accepted Version

**Downloaded from:** <https://e-space.mmu.ac.uk/626032/>

**Usage rights:** © In Copyright

**Additional Information:** This is an Author Accepted Manuscript of a paper accepted for publication in *Journal of Strength and Conditioning Research* published by and copyright National Strength and Conditioning Association.

**Enquiries:**

If you have questions about this document, contact [openresearch@mmu.ac.uk](mailto:openresearch@mmu.ac.uk). Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/325679974>

# An Investigation Into the Effects of Excluding the Catch Phase of the Power Clean on Force–Time Characteristics During Isometric and Dynamic Tasks: An Intervention Study

Article in *The Journal of Strength and Conditioning Research* · June 2018

DOI: 10.1519/JSC.0000000000002656

CITATIONS

12

READS

835

5 authors, including:



**Paul Comfort**

University of Salford

215 PUBLICATIONS 2,264 CITATIONS

SEE PROFILE



**Thomas Dos'Santos**

Manchester Metropolitan University

66 PUBLICATIONS 508 CITATIONS

SEE PROFILE



**Christopher Thomas**

56 PUBLICATIONS 609 CITATIONS

SEE PROFILE



**John J McMahon**

University of Salford

99 PUBLICATIONS 651 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



The effect of training with a force-velocity overload stimulus using weightlifting derivatives [View project](#)



Performance variables in different counter-movement jump strategies between volleyball players with contrasting take-off velocities [View project](#)

**An investigation into the effects of excluding the catch phase of the power clean on force-time characteristics during isometric and dynamic tasks: an intervention study**

Paul Comfort<sup>1#</sup>, Thomas Dos'Santos<sup>1</sup>, Christopher Thomas<sup>1,2</sup>, John. J. McMahon<sup>1</sup> & Timothy. J. Suchomel<sup>3</sup>.

Author affiliations: <sup>1</sup> Directorate of Sport, Exercise and Physiotherapy, University of Salford, Frederick Road, Salford, Greater Manchester. UK. <sup>2</sup> University of South Wales. Pontypridd. Cardiff, Wales. UK. <sup>3</sup> Department of Human Movement Sciences, Carroll University, Waukesha, WI, USA

# Corresponding author: [p.comfort@salford.ac.uk](mailto:p.comfort@salford.ac.uk)

**Running header: Effects of excluding the catch phase of the power clean: an intervention study**

## **An investigation into the effects of excluding the catch phase of the power clean on force-time characteristics during isometric and dynamic tasks: an intervention study**

### **Abstract**

The aims of this study were to compare the effects of the exclusion or inclusion of the catch phase, during power clean (PC) derivatives, on force-time characteristics during isometric and dynamic tasks, after two, four-week mesocycles of resistance training. Two strength matched groups, completed the twice weekly training sessions, either including the catch phase of the PC derivatives (Catch:  $n = 16$ ; age  $19.3 \pm 2.1$  years; height  $1.79 \pm 0.08$  m; body mass  $71.14 \pm 11.79$  kg; PC one repetition maximum [1-RM]  $0.93 \pm 0.15$  kg.kg<sup>-1</sup>) or excluding the catch phase (Pull:  $n = 18$ ; age  $19.8 \pm 2.5$  years; height  $1.73 \pm 0.10$  m; body mass  $66.43 \pm 10.13$  kg; PC 1RM  $0.91 \pm 0.18$  kg.kg<sup>-1</sup>). The Catch and Pull groups both demonstrated significant ( $p \leq 0.007$ , power  $\geq 0.834$ ) and meaningful improvements in countermovement jump (CMJ) height ( $10.8 \pm 12.3\%$ ,  $5.2 \pm 9.2\%$ ), isometric mid-thigh pull (IMTP) performance (force [F]100:  $14.9 \pm 17.2\%$ ,  $15.5 \pm 16.0\%$ , F150:  $16.0 \pm 17.6\%$ ,  $16.2 \pm 18.4\%$ , F200:  $15.8 \pm 17.6\%$ ,  $17.9 \pm 18.3\%$ , F250:  $10.0 \pm 16.1\%$ ,  $10.9 \pm 14.4\%$ , PF:  $13.7 \pm 18.7\%$ ,  $9.7 \pm 16.3\%$ ) and PC 1RM ( $9.5 \pm 6.2\%$ ,  $8.4 \pm 6.1\%$ ), pre- to post-intervention, respectively. In contrast to the hypotheses, there were no meaningful or significant differences in percentage change, for any variables, between groups. This study clearly demonstrates that neither the inclusion nor exclusion of the catch phase of the PC derivatives result in any preferential adaptations over two 4-week, in-season strength and power, mesocycles.

**Key Words:** Countermovement Jump; Weightlifting; Performance; Training

### **INTRODUCTION**

Weightlifting exercises (snatch and clean and jerk) and their derivatives are commonly performed in athletes' training programs, with performance in such exercises reported to be related to athletic tasks, such as sprint, agility and jump performances (29, 40). These positive associations to performances in athletic tasks may be due to the previously reported similarity in kinetics between weightlifting derivatives (hang snatch) and jump performances (4), with similar observations reported between the second pull phase of the snatch and jump performances by Garhammer and Gregor (18).

Observations of weightlifting performances have established that the second pull phase of the clean and snatch elicits the greatest peak power, compared to the other phases of the lifts (18), albeit using barbell velocity and inverse dynamics to assess peak power applied to it. Furthermore, peak force (PF) and rate of force development (RFD) have also been shown to occur during the second pull phase of the clean and clean pull (16, 39). More recently, the

mid-thigh power clean (PC) and mid-thigh pull have been shown to result in significantly greater ( $p < 0.001$ ) PF, peak RFD (5) and peak power applied to the lifter plus bar system (6) when compared to the hang power clean and PC. Moreover, no significant ( $p > 0.05$ ) differences were observed between these lifts irrespective of the inclusion or exclusion of the catch phase (5, 6). In addition, Suchomel et al. (47) reported that the jump shrug, (similar to the mid-thigh pull but initiated with a countermovement and the athlete actually leaves the ground) resulted in significantly ( $p < 0.05$ ) greater PF, peak velocity, and peak power compared to the hang power clean and hang high pull across all loads (30, 45, 65, 80% one repetition maximum [1RM] hang clean), indicating that the removal of a catch phase during a PC derivative is not detrimental to the peak power achieved. Similarly, additional studies by Suchomel et al. (45, 46) also reported greater relative PF, power, impulse, work, and peak RFD in the jump shrug compared to the hang power clean and hang high pull across loads (30, 45, 65, 80% 1RM hang clean). More recently, researchers have examined these differences at the joint-level, with Kipp et al. (32) indicating that the jump shrug produces greater magnitudes of joint work and power compared to the hang power clean across several loads.

Recent reviews of weightlifting derivatives also suggested that variations of the PC, which omit the catch phase, namely the clean pull, mid-thigh pull, jump shrug and hang high pull, may be advantageous when training athletes who are less proficient with full weightlifting movements that include the catch phase (41, 43). This is supported by additional research that has suggested the use of associate exercises that enhance explosive strength during the second pull movement in less skillful athletes (25). Based on the kinetic similarities of the propulsion phases of the clean derivatives performed with and without the catch phase, it would be feasible to suggest that the elimination of the catch phase should not be detrimental during a training program. In fact, the elimination of the catch phase may provide the opportunity for the athlete to ensure full triple extension of the hips, knees and ankles (plantar flexion), without the possibility of terminating the propulsion phase early to initiate the catch. Ultimately, this may lead to superior training adaptations with regard to PF, RFD, and power during the triple extension movement.

Additionally, the catch phase of the weightlifting derivatives has been suggested to be potentially beneficial in terms of training deceleration and eccentric loading; however, the loading during the catch has been reported to only be comparable to landing during a drop jump (36). More recently, the clean pull from the knee was shown to result in greater mean forces during the load absorption phase compared to the clean and PC from the knee (11). Similarly, Suchomel et al. (44) recently reported greater mean forces during the load absorption phase of the jump shrug compared to the hang high pull and hang power clean. The findings of these studies refute the notion that the catch phase of the clean provides effective eccentric loading. To date, however, there are no published intervention studies that compare the effectiveness of including or excluding the catch during weightlifting derivatives on strength and power characteristics.

The aims of this study, therefore, were to compare the effects of the exclusion or inclusion of the catch phase, during PC derivatives, on force-time characteristics during isometric and dynamic tasks, after two, four-week mesocycles of resistance training. It was hypothesized that both groups would improve across all variables, but that the Pull group (elimination of the catch phase) would result in greater improvements in force-time characteristics assessed during isometric and dynamic performance between groups, compared to the Catch group.

## METHODS

### EXPERIMENTAL APPROACH TO THE PROBLEM

To determine the effect of the training interventions, on force-time characteristics during isometric and dynamic tasks, a repeated-measures within subject design was utilized, with subjects assessed twice at baseline (48-72 hours apart) to determine reliability, after the initial four week mesocycle, and again after the second four week mesocycle (Figure 1). Furthermore, a between-subjects experimental approach was used to determine differences in changes between intervention groups (Pull vs. Catch). All testing and training occurred in-season, during the middle of the season for each sport. Data was collected across multiple venues, using the same portable equipment, by the same group of researchers.




Figure 1: Summary of testing schedule

### Subjects

Professional youth soccer players ( $n = 18$ ) and collegiate athletes ( $n = 26$ ), from the United Kingdom, initially volunteered to participate in this investigation. All subjects were experienced (training age:  $3.1 \pm 1.2$  years) and competent in each of the lifts performed in the interventions, as determined by a certified strength and conditioning specialist. After baseline testing subjects were divided into the two groups by matching relative 1RM PC performances, with an equal number of athletes from each sport in both groups. Due to injury from competition and or illness across the duration of the intervention the number of subjects to complete the entire study reduced to 11 professional male soccer players and 23 collegiate athletes who participated in a variety of sports (BMX, rowing, field hockey). Due to drop out, the final mean 1RM PC performance for the groups differed slightly; Catch ( $n = 16$ , 12 male, 4 female [5 soccer, 3 BMX, 6 rowing, 2 field hockey]; age  $19.3 \pm 2.1$  years; height  $1.79 \pm 0.08$  m; body mass  $71.14 \pm 11.79$  kg; 1RM PC  $0.93 \pm 0.15$  kg.kg<sup>-1</sup>) Pull ( $n = 18$ , 14 male, 4 female [6 soccer,

2 BMX, 7 rowing, 2 field hockey]; age  $19.8 \pm 2.5$  years; height  $1.73 \pm 0.10$  m; body mass  $66.43 \pm 10.13$  kg; 1RM PC  $0.91 \pm 0.18$  kg.kg<sup>-1</sup>). A minimum of 11 subjects per groups was required for an *a priori* power  $\geq 0.80$ , at an alpha level of  $p \leq 0.05$ , with post hoc power presented in the results section. This study was approved by the institutional review board, in accordance with the declaration of Helsinki. All subjects provided written informed consent, or parental assent as appropriate.

## PROCEDURES

Prior to testing subjects performed a non-fatiguing standardized warm up consisting of body weight squats, forward and reverse lunges, submaximal squat jumps (SJ) and countermovement jumps (CMJ). Further familiarization and warm up trials were performed prior to the maximal isometric mid-thigh pull (IMTP) and 1RM PC as described below. After the completion of the warm up subjects performed the SJ, CMJ, IMTP and 1RM PC as described below; with testing performed in this sequence to minimize the risk of fatigue or potentiation (Figure 2). All subjects were familiar with all testing procedures as these were included in their 'normal' testing and monitoring procedures. All assessments were conducted by the same experienced researchers.




Figure 2: Testing sequence

## Jump Performances

Both SJ and CMJ performances were assessed with subjects standing on a Kistler force platform, sampling at 1000 Hz, with data collected via Bioware 5.11 software (type 9286AA, Kistler Instruments Inc., Amherst, NY, USA). Subjects were instructed to stand still for the initial one second of data collection (35, 38) to enable the subsequent determination of body weight (vertical force averaged over one second). Subjects performed three maximal efforts SJ and CMJ, with a one-minute rest between trials and a three-minute rest between the SJ and CMJ. Raw unfiltered, force-time data was exported for subsequent analysis.

For the SJ, subjects placed their hands akimbo, squatted down to a self-selected depth of approximately 90° knee joint angle, paused for 3 seconds and then jumped as high as possible after a countdown of, '3, 2, 1, jump'. If there was any obvious countermovement, following visual inspection of the force-time data the jump was excluded, and the subject preformed an additional trial after a one-minute rest.

For the CMJ, subjects were instructed to perform the jumps as fast and as high as possible, whilst keeping their arms akimbo. Any jumps that were inadvertently performed with the inclusion of arm swing or leg tucking during the flight phase were omitted and additional jumps were performed after one minute of rest.

### ***Isometric Mid-thigh Pull Assessment***

For the IMTP, the procedures previously described by Haff et al. (20, 21) were used. The minor differences in knee joint angle, which result from differences in ankle dorsiflexion, have been shown to have minimal effect on kinetic variables during the IMTP (7). It was ensured, however, that each subject adopted the posture that they would use for the start of the second pull phase of the clean resulting in knee and hip angles of  $133.1 \pm 6.6^\circ$  and  $145.6 \pm 4.8^\circ$  respectively, in line with previous research (3, 21). Individual joint angles were recorded and standardized between testing sessions, in line with previous suggestions (3, 15). Briefly, for this test, an immovable cold rolled steel bar was positioned at a height, which replicates the start of the second pull phase of the clean, with the bar fixed above the force platform to accommodate different sized participants. Once the bar height was established, the subjects' stood on the force platform with their hands strapped to the bar in accordance with previously established methods (2). Each participant performed two warm-up pulls, one at 50%, and one at 75% of the participant's perceived maximum effort, separated by one minute of rest.

Once body position was stabilized (verified by watching the participant and force trace), the participants were given a countdown of "3, 2, 1, Pull!". Minimal pre-tension was permitted to ensure there was no slack in the participant's body prior to initiation of the pull, with the instruction to pull against the bar "as fast and hard as possible" (24), and push the feet down into the force plate; this instruction has been previously found to produce optimal testing results (23). Each IMTP trial was performed for approximately five seconds, and all participants were given strong verbal encouragement during each trial. Participants performed three maximal IMTP trials interspersed with two minutes of rest between trials. If PF during all trials did not fall within 250 N of each other, the trial was discounted and repeated after a further two minutes of rest, in line with previous recommendations (19, 21).

Vertical ground reaction force data for the IMTP was collected using a portable force plate sampling at 1000 Hz (Kistler Instruments, Winterthur, Switzerland), interfaced with a laptop computer and specialist software (Bioware 5.11, Kistler Instruments, Winterthur, Switzerland) that allows for direct measurement of force-time characteristics. Raw unfiltered, force-time data was exported for subsequent analysis.

### ***One Repetition Maximum Power Clean***

The 1RM PC performances were determined based on the standardized NSCA protocol (1). Briefly, subjects performed warm-up PC sets using sub maximal loads prior to performing a maximal attempt, with a progressive increase in loading during the maximal attempts (International Weightlifting Federation, accredited bars and plates were used throughout). Any



power clean repetition caught with the top of the subject's thighs below parallel was ruled as an unsuccessful attempt.

## DATA ANALYSIS:

### *Kinetic and Kinematic Variables*

Raw force-time data for both the jumps and the IMTP were analyzed in Microsoft Excel (Excel 2016, Microsoft, Washington, USA). Jump height was calculated from velocity of center of mass at take-off, for both the SJ and CMJ (35). Center of mass velocity was determined by dividing vertical force data (minus body weight) by body mass and then integrating the product using the trapezoid rule. The start of the CMJ was identified in line with current recommendations (38). Take-off was identified when vertical force decreased below five times the standard deviation of the force during the flight phase (residual force) (34).

Reactive strength index modified (RSImod) was calculated using the methods described by previous research (34), where jump height is divided by time to take off ([TTT] combined countermovement, braking and propulsion phase time) during the CMJ.

The maximum forces recorded from the force-time curve during the IMTP trials were reported as the PF and subsequently ratio scaled (PF / body mass). The onset of force production was defined as an increase in force greater than five standard deviations of force during the period of quiet standing (13), and subsequently force at 100-, 150-, 200- and 250 ms (F100, F150, F200, F250) were also determined and ratio scaled. The average value of the three trials was used for statistical analyses.

## INTERVENTION

Participants were divided into either the Pull group or Catch group and performed the prescribed training on two days per week, under the supervision of certified strength and conditioning specialists. The program consisted of two, 4-week mesocycles (Tables 1 & 2). The relative training intensity for each group was matched in an attempt to equate the volume-load completed by each group. The loads prescribed for all pulling and catching derivatives were based on the subjects' 1RM PC. The loads prescribed for the remaining exercises were based on predicted 1RM loads based on the subject's previous 5RM performances as determined at the end of their previous phase of training. The volume load during the second session was reduced, as this was the session closest to the subjects' day of competition. All training sessions were supervised by at least one of the authors, who were qualified strength and conditioning coaches (either as a certified strength and conditioning coach with the National Strength and Conditioning Association, an accredited strength and conditioning coach with the United Kingdom Strength and Conditioning Association, or both), to ensure consistency of performance.

The rowers and professional youth soccer players performed between 10-14 hours of skill and conditioning based training per week, in addition to the intervention; while the other subjects performed between 5-8 hours per week of additional training, dependent on their competition schedule, hence initially dividing the subjects equally across groups.

267 Table 1: Training sessions, weeks 1-4

Mesocycle 1: Day 1				
Exercise	Week 1	Week 2	Week 3	Week 4
Back Squat	3 x 5 @ 75%	3 x 5 @ 80%	3 x 5 @ 82.5%	3 x 5 @ 67.5%
<b>Power Clean / Clean Pull<sup>a</sup></b>	3 x 5 @ 75%	3 x 5 @ 80%	3 x 5 @ 82.5%	3 x 5 @ 67.5%
Push Press	3 x 5 @ 70%	3 x 5 @ 72.5%	3 x 5 @ 75%	3 x 5 @ 60%
Nordic Lowers	2 x 3 BW	3 x 3 BW	3 x 3 BW	3 x 3 BW
Mesocycle 1: Day 2				
<b>Mid-thigh Power Clean / Mid-thigh Pull<sup>b</sup></b>	3 x 5 @ 60%	3 x 5 @ 65%	3 x 5 @ 70%	3 x 5 @ 55%
RDL	3 x 5 @ 70%	3 x 5 @ 75%	3 x 5 @ 77.5%	3 x 5 @ 62.5%
Sets x Repetitions @ 1RM %				
BW = Body Weight				
<sup>a</sup> Power clean for the Catch group / Clean pull for the Pull group				
<sup>b</sup> Mid-thigh power clean for the Catch group / Mid-thigh pull for the Pull group				

268

269 Table 2: Training sessions, weeks 6-9

Mesocycle 2: Day 1				
Exercise	Week 1	Week 2	Week 3	Week 4
<b>Power Clean / Clean Pull<sup>a</sup></b>	3 x 3 @ 80%	3 x 3 @ 85%	3 x 3 @ 90%	3 x 3 @ 75%
Push Press	3 x 3 @ 80%	3 x 3 @ 82.5%	3 x 3 @ 85%	3 x 3 @ 75%
Back Squat	3 x 3 @ 82.5%	3 x 3 @ 87.5%	3 x 3 @ 90%	3 x 3 @ 75%
Nordic Lowers	2 x 3 BW	3 x 3 BW	3 x 3 BW	3 x 3 BW
Mesocycle 2: Day 2				
<b>Mid-thigh Power Clean / Mid-thigh Pull<sup>b</sup></b>	3 x 3 @ 80%	3 x 3 @ 82.5%	3 x 3 @ 85%	3 x 3 @ 70%
RDL	3 x 3 @ 80%	3 x 3 @ 85%	3 x 3 @ 87.5%	3 x 3 @ 72.5%
Sets x Repetitions @ 1RM %				
BW = Body Weight				
<sup>a</sup> Power clean for the Catch group / Clean pull for the Pull group				
<sup>b</sup> Mid-thigh power clean for the Catch group / Mid-thigh pull for the Pull group				

270

271 **Statistical Analyses**

272 Normality of all data was determined via Shapiro-Wilk's test of normality, with all variables  
 273 being normally distributed. Baseline measures were compared to determine within- and  
 274 between-session reliability, as appropriate, using two-way random effects model intraclass  
 275 correlation coefficients (ICC) and 95% confidence intervals. To assess the magnitude of the  
 276 ICC, the values were interpreted as low (<0.30), moderate (0.30-0.49), high (0.50-0.69), very  
 277 high (0.70-0.89), nearly perfect (0.90-0.99), and perfect (1.0) (28). Percentage coefficient of  
 278 variation (%CV) was also calculated to determine the within session variability, with <10%  
 279 classified as acceptable (12). In addition, t-tests were performed and Cohen's *d* effect sizes  
 280 calculated to determine if there were any significant or meaningful differences between the  
 281 baseline testing sessions.

A series of two-way repeated-measures analyses of variance (3 x 2; time x group), with Bonferroni post-hoc analysis, were performed to determine changes in the aforementioned kinetic and kinematic variables at each time point. A series of t-tests were performed to determine differences in the percentage change between phases (pre-mid, mid-post, pre-post) and between groups (Catch vs. Pull), for each variable. An *a priori* alpha level was set at  $p \leq 0.05$ . Further, the magnitude of any changes were determined via the calculation of effect sizes (Cohen's  $d$ ), classified as trivial ( $\leq 0.19$ ), small (0.20 – 0.59), moderate (0.60 – 1.19), large (1.20 – 1.99), and very large (2.0 – 4.0) (27). All statistical analyses were performed using SPSS (Version 23. IBM, New York, NY).

## Results

Between session 1RM PC performances were highly reliable (ICC = 0.997, 0.998) with a very low variability (CV = 0.23%, 0.13%) between sessions one ( $67.58 \pm 23.06$  kg;  $0.94 \pm 0.19$  kg.kg<sup>-1</sup>) and two ( $67.36 \pm 22.59$  kg;  $0.93 \pm 0.19$  kg.kg<sup>-1</sup>), for both absolute and relative performances, respectively.

Reliability of all jump variables demonstrated was very high to nearly perfect both within (ICC = 0.819-0.976) and between (ICC = 0.870-0.981) sessions, with low variability (CV = 0.27-5.96%) between trials. Furthermore, differences between sessions were trivial to small ( $d = 0.03$ -0.22) and not significant (Table 3).

Reliability of all IMTP variables demonstrated was very high to nearly perfect both within (ICC = 0.879-0.983) and nearly perfect (ICC = 0.966-0.981) between sessions, with acceptable variability (CV = 5.36-12.78%) between trials, with the variability reducing progressively with the time-point at which force was assessed. Furthermore, differences between sessions were trivial ( $d = 0.03$ -0.22) and non-significant ( $p > 0.05$ ) (Table 4).

Table 3: Within and between session reliability (ICC (95% confidence intervals)) and variability (% coefficient of variation) of jump performance variables

Variable		Session 1	Session 2
SJ Height (m)	Mean	0.281	0.266
	SD	0.069	0.068
	Within	0.944	0.962
	Session ICC	(0.881-0.977)	(0.920-0.984)
	Between	0.870	
	Session ICC	(0.661-0.951)	
	%CV	5.06	0.27
	<i>d</i>	0.22	
CMJ Height (m)	Mean	0.316	0.318
	SD	0.072	0.071
	Within	0.954	0.981
	Session ICC	(0.903-0.981)	(0.959-0.992)
	Between	0.971	
	Session ICC	(0.925-0.989)	
	%CV	4.15	2.78
	<i>d</i>	0.03	
CMJ TTT (s)	Mean	0.73	0.72
	SD	0.08	0.10
	Within	0.819	0.854
	Session ICC	(0.652-0.921)	(0.710-0.937)
	Between	0.893	
	Session ICC	(0.719-0.960)	
	%CV	3.06	2.86
	<i>d</i>	0.13	
CMJ RSI <sub>mod</sub>	Mean	0.44	0.45
	SD	0.10	0.11
	Within	0.906	0.940
	Session ICC	(0.809-0.960)	(0.875-0.975)
	Between	0.976	
	Session ICC	(0.933-0.991)	
	%CV	5.96	5.04
	<i>d</i>	0.12	
SJ: squat jump, CMJ: countermovement jump, TTT: time to take-off, RSI <sub>mod</sub> : reactive strength index modified, SD: standard deviation, ICC: intraclass correlation coefficient, %CV: percentage coefficient of variation, <i>d</i> : Cohen's <i>d</i> effect size			

Table 4: Within and between session reliability (ICC (95% confidence intervals)) and variability (% coefficient of variation) of IMTP variables

Variable		Session 1	Session 2
F100 ms (N.kg <sup>-1</sup> )	Mean	20.32	20.35
	SD	6.23	5.20
	Within	0.937	0.908
	Session ICC	(0.869-0.974)	(0.798-0.963)
	Between	0.980	
	Session ICC	(0.945-0.992)	
	%CV	5.50	12.78
F150 ms (N.kg <sup>-1</sup> )	<i>d</i>	0.01	
	Mean	25.18	25.01
	SD	7.92	6.15
	Within	0.925	0.903
	Session ICC	(0.845-0.969)	(0.786-0.961)
	Between	0.966	
	Session ICC	(0.909-0.987)	
F200 ms (N.kg <sup>-1</sup> )	%CV	6.28	11.62
	<i>d</i>	0.02	
	Mean	28.73	28.28
	SD	8.72	6.76
	Within	0.935	0.812
	Session ICC	(0.865-0.973)	(0.64-0.918)
	Between	0.967	
F250 ms (N.kg <sup>-1</sup> )	Session ICC	(0.913-0.988)	
	%CV	5.82	8.94
	<i>d</i>	0.05	
	Mean	30.32	30.06
	SD	9.05	7.40
	Within	0.953	0.879
	Session ICC	(0.902-0.981)	(0.761-0.949)
Peak Force (N.kg <sup>-1</sup> )	Between	0.978	
	Session ICC	(0.941-0.992)	
	%CV	5.36	6.19
	<i>d</i>	0.03	
	Mean	38.19	38.91
	SD	12.24	11.70
	Within	0.983	0.968
	Session ICC	(0.964-0.993)	(0.930-0.987)
	Between	0.981	
	Session ICC	(0.950-0.993)	
	%CV	3.44	4.29
	<i>d</i>	0.06	

SD: standard deviation, ICC: intraclass correlation coefficient, %CV: percentage coefficient of variation, *d*: Cohen's *d* effect size

## 328 JUMP PERFORMANCES

329 Sphericity was assumed via Mauchley's test for all jump variables. The Catch group achieved  
 330 significant ( $p < 0.001$ ; power = 0.794) improvements in SJ height across the duration of the  
 331 intervention, with moderate and significant increase ( $12.6 \pm 10.2\%$ ,  $p < 0.001$ ) from pre- to  
 332 post-intervention. In contrast, post-hoc analysis demonstrated that changes were small and  
 333 non-significant ( $p > 0.05$ ) between pre- and mid-intervention and mid- and post-intervention.  
 334 There was only a trivial and non-significant increase ( $2.1 \pm 11.8\%$ ,  $p > 0.05$ ) in SJ performance  
 335 for the Pull group (Table 5). The Catch group exhibited greater improvements in SJ height pre-  
 336 to mid-intervention ( $8.8 \pm 13.1\%$ ), mid- to post-intervention ( $4.1 \pm 7.9\%$ ), or pre- to post-  
 337 intervention ( $12.6 \pm 10.2\%$ ), compared to the Pull group ( $2.1 \pm 11.8\%$ ,  $1.9 \pm 12.8\%$ ,  $4.0 \pm$   
 338  $17.6\%$ , respectively), although these were small and not significantly different ( $d = 0.20$ - $0.59$ ;  
 339  $p > 0.05$ ) (Figure 3a).

340 The Catch group and Pull groups both achieved significant ( $p < 0.001$ ; power = 0.980;  $p = 0.04$ ;  
 341 power = 0.810, respectively) improvements in CMJ height across the duration of the  
 342 intervention. The results of post-hoc analysis demonstrated that changes were small and non-  
 343 significant ( $p > 0.05$ ) between pre- and mid-intervention and mid- to post-intervention for the  
 344 Catch group, with a small yet significant ( $10.8 \pm 12.3\%$ ,  $p = 0.007$ ) increase from pre- to post-  
 345 intervention. The Pull group achieved trivial and non-significant increases between pre- and  
 346 mid-intervention and mid- to post-intervention, with small but significant increases ( $5.2 \pm 9.2\%$ ,  
 347  $p = 0.04$ ) pre- to post-intervention (Table 5). The Catch group exhibited greater improvements  
 348 in CMJ height pre- to mid-intervention ( $5.4 \pm 9.6\%$ ), mid- to post-intervention ( $5.1 \pm 6.5\%$ ), or  
 349 pre-to post-intervention ( $10.8 \pm 12.3\%$ ), compared to the Pull group ( $3.7 \pm 8.0\%$ ,  $1.6 \pm 7.2\%$ ,  
 350  $5.2 \pm 9.2\%$ , respectively), although these were trivial to small and non-significant ( $d = 0.19$ -  
 351  $0.52$ ;  $p > 0.05$ ) (Figure 3b).




Figure 3: Comparison of percentage change in jump variables, across time points, for the Catch and Pull groups (SJ = squat jump; CMJ = countermovement jump; RSI<sub>mod</sub> = reactive strength index modified)

For CMJ TTT there were trivial to small non-significant differences for both the Catch and Pull groups across all time points. There were trivial to small and non-significant differences ( $p > 0.05$ ) in percentage change TTT pre- to mid-intervention ( $1.2 \pm 8.8\%$ ,  $-0.4 \pm 12.2\%$ ,  $d = 0.15$ ), mid- to post-intervention ( $3.5 \pm 11.0\%$ ,  $2.9 \pm 10.6\%$ ,  $d = 0.06$ ), and pre-post ( $4.6 \pm 13.5\%$ ,  $2.0 \pm 12.0\%$ ,  $d = 0.20$ ), between the Catch and Pull groups, respectively (Table 5, Figure 3c). There were only trivial to small changes in RSI<sub>mod</sub> for both groups across all time points (Table 5), with trivial to small and non-significant differences ( $p > 0.05$ ) in percentage change in RSI<sub>mod</sub> across phases (pre-mid:  $4.6 \pm 10.0\%$ ,  $4.9 \pm 10.1\%$ ,  $d = 0.03$ , mid-post:  $2.4 \pm 10.4\%$ ,  $0.0 \pm 13.7\%$ ,  $d = 0.20$ , pre-post:  $7.0 \pm 13.4\%$ ,  $4.4 \pm 14.1\%$ ,  $d = 0.19$ ), between the Catch and Pull groups, respectively (Figure 3d).

Table 5: Changes in jump performance

Variable	Group	Catch			Pull		
		Pre	Mid	Post	Pre	Mid	Post
SJ Height (m)	Mean	0.283	0.305	0.317	0.283	0.287	0.289
	SD	0.052	0.048	0.053	0.061	0.057	0.055
	%CV	4.40	4.95	2.74	5.64	4.06	3.36
	$d$	0.44		0.05		0.05	
	$d$	0.64*		0.10		0.10	
CMJ Height (m)	Mean	0.327	0.341	0.360	0.313	0.324	0.328
	SD	0.064	0.056	0.066	0.062	0.068	0.062
	%CV	4.05	3.12	2.78	3.29	3.92	2.36
	$d$	0.24		0.17		0.05	
	$d$	0.50*		0.23*		0.23*	
CMJ TTT (s)	Mean	0.71	0.72	0.74	0.76	0.75	0.77
	SD	0.09	0.10	0.09	0.09	0.09	0.10
	%CV	2.80	3.28	3.16	3.60	3.69	3.23
	$d$	0.07		0.08		0.19	
	$d$	0.29		0.11		0.11	
RSI <sub>mod</sub>	Mean	0.46	0.48	0.49	0.42	0.43	0.43
	SD	0.09	0.09	0.09	0.09	0.09	0.09
	%CV	6.73	6.24	6.69	6.10	5.89	4.00
	$d$	0.20		0.20		0.05	
	$d$	0.11		0.16		0.16	

\*=significant ( $p < 0.05$ ) increase pre to post intervention

SJ: squat jump, CMJ: countermovement jump, TTT: time to take-off, RSI<sub>mod</sub>: reactive strength index modified, SD: standard deviation, %CV: percentage coefficient of variation,  $d$ : Cohen's  $d$  effect size

#### ISOMETRIC MID-THIGH PULL

Sphericity was assumed via Mauchley's test for all IMTP variables. The Catch and Pull groups both demonstrated significant ( $p < 0.001$ ; power = 0.931) increases in F100. Both groups showed trivial non-significant ( $p > 0.05$ ) changes pre- to mid-intervention, with small significant



(Catch:  $17.3 \pm 22.0\%$ ,  $p = 0.03$  Pull:  $11.5 \pm 21.4\%$ ,  $p = 0.04$ ) increases mid- to post-intervention and pre- to post-intervention (Catch:  $14.9 \pm 17.2\%$ ,  $p = 0.011$  Pull:  $15.5 \pm 16.0\%$ ,  $p = 0.03$ ) (Table 6). Trivial to small and non-significant differences ( $d = 0.08$ - $0.23$ ,  $p > 0.05$ ) in percentage change F100 across phases (pre-mid:  $-0.7 \pm 13.5\%$ ,  $3.7 \pm 15.9\%$ , mid-post:  $17.3 \pm 22.0\%$ ,  $11.5 \pm 21.4\%$ , pre-post:  $14.9 \pm 17.2\%$ ,  $13.5 \pm 16.0\%$ ), were evident between the Catch and Pull groups, respectively (Figure 4a).

Both groups demonstrated significant ( $p = 0.005$ ; power =  $0.855$ ) increases in F150, with both groups showing trivial to small non-significant ( $p > 0.05$ ) changes pre- to mid-intervention, with the Catch group demonstrating small significant ( $16.5 \pm 20.4\%$ ,  $p = 0.022$ ) increases mid- to post-intervention and the Pull group demonstrating small but non-significant ( $12.0 \pm 22.9\%$ ,  $p > 0.05$ ) increases mid- to post-intervention. Both groups demonstrated moderate and significant increases (Catch:  $16.0 \pm 17.6\%$ ,  $p = 0.003$  Pull:  $16.2 \pm 18.4\%$ ,  $p = 0.01$ ) in F150 pre- to post-intervention (Table 6). Trivial to small and non-significant differences ( $d = 0.01$ - $0.31$ ,  $p > 0.05$ ) in percentage change F150 across phases (pre-mid:  $0.9 \pm 14.9\%$ ,  $5.9 \pm 17.5\%$ , mid-post:  $16.5 \pm 17.6\%$ ,  $12.0 \pm 22.9\%$ , pre-post:  $16.0 \pm 17.6\%$ ,  $16.2 \pm 18.4\%$ ), were evident between the Catch and Pull groups, respectively (Figure 4b).




Figure 4: Comparison of percentage change in isometric mid-thigh pull time specific force variables, across time points, for the Catch and Pull groups

Both groups demonstrated significant ( $p = 0.007$ ; power = 0.842) increases in F200. Both groups showed trivial to small non-significant ( $p > 0.05$ ) changes pre- to mid-intervention, with small non-significant (Catch:  $16.6 \pm 17.9\%$ , Pull:  $12.9 \pm 16.8\%$ ,  $p > 0.05$ ) increases mid- to post-intervention and small, significant increases pre- to post-intervention (Catch:  $15.8 \pm 17.6\%$ ,  $p = 0.017$  Pull:  $17.9 \pm 18.3\%$ ,  $p = 0.02$ ) (Table 6). The Pull group demonstrated small yet significantly greater ( $d = 0.38$ ,  $p = 0.002$ ) increases in F200 pre- to mid-intervention ( $5.3 \pm 14.0\%$ ) compared to the Catch group ( $0.1 \pm 13.2\%$ ). There were, however, only trivial to small and non-significant differences ( $d = 0.12$ - $0.21$ ,  $p > 0.05$ ) in percentage change F200 mid- to post-intervention ( $16.6 \pm 17.9\%$ ,  $12.9 \pm 16.8\%$ ) or pre- to post-intervention ( $15.8 \pm 17.6\%$ ,  $17.9 \pm 18.3\%$ ), between the Catch and Pull groups, respectively (Figure 4c).

Both groups demonstrated significant ( $p = 0.007$ ; power = 0.834) increases in F250, with the Catch group showing a trivial non-significant ( $p > 0.05$ ) decrease pre- to mid-intervention, while the Pull group showed a small but non-significant increase ( $p > 0.05$ ). The Catch group demonstrated a small significant ( $12.0 \pm 16.6\%$ ,  $p = 0.045$ ) increase mid- to post-intervention and small significant increase pre- to post-intervention ( $10.0 \pm 16.1\%$ ,  $p = 0.025$ ), while the Pull group demonstrated a small significant ( $6.5 \pm 13.4\%$ ,  $p = 0.045$ ) increase mid- to post-intervention and small significant increase pre- to post-intervention ( $10.9 \pm 14.4\%$ ,  $p = 0.025$ ) (Table 6). Trivial to small and non-significant differences ( $d = 0.06$ - $0.47$ ,  $p > 0.05$ ) in percentage change F250 were evident, across phases (pre-mid:  $-1.0 \pm 12.5\%$ ,  $4.7 \pm 11.7\%$ , mid-post:  $12.0 \pm 16.6\%$ ,  $6.5 \pm 13.4\%$ , pre-post:  $10.0 \pm 16.1\%$ ,  $10.9 \pm 14.4\%$ ), between the Catch and Pull groups, respectively (Figure 4d).

Both groups demonstrated significant ( $p = 0.001$ ; power = 0.869) and progressive increases in relative PF, with the Catch group showing a trivial non-significant ( $p > 0.05$ ) increase pre- to mid-intervention, while the Pull group showed a small but significant increase ( $p = 0.017$ ). In contrast the Catch group demonstrated a small significant ( $8.4 \pm 10.8\%$ ,  $p = 0.028$ ) increase mid- to post-intervention while the Pull group demonstrated a trivial non-significant ( $p > 0.05$ ) increase in relative PF. Both groups demonstrated small significant increases (Catch:  $13.7 \pm 18.7\%$ ,  $p = 0.021$ ; Pull:  $9.7 \pm 16.3\%$ ,  $p = 0.045$ ) in relative PF pre- to post-intervention (Table 6). The Catch group demonstrated a moderately and significantly greater ( $d = 0.84$ ,  $p = 0.014$ ) increase in PF mid- to post-intervention ( $8.4 \pm 10.8\%$ ) compared to the Pull group ( $0.2 \pm 8.5\%$ ). There were, however, only small and non-significant differences ( $d = 0.23$ - $0.45$ ,  $p > 0.05$ ) in percentage change PF pre- to mid-intervention ( $4.6 \pm 9.6\%$ ,  $9.8 \pm 13.1\%$ ) or pre- to post-intervention ( $13.7 \pm 18.7\%$ ,  $9.7 \pm 16.3\%$ ), between the Catch and Pull groups, respectively (Figure 5a).

460 Table 6: Changes in isometric mid-thigh pull performance

	Group	Catch			Pull		
		Pre	Mid	Post	Pre	Mid	Post
F100 ms (N.kg <sup>-1</sup> )	Mean	20.00	19.95	22.92	17.93	18.49	20.14
	SD	5.07	4.52	5.94	3.74	4.06	4.11
	%CV	5.48	8.68	7.76	6.68	9.30	8.20
		0.01			0.14		
	d	0.46*			0.40*		
		0.45*			0.56*		
F150 ms (N.kg <sup>-1</sup> )	Mean	24.76	25.11	28.67	22.07	23.28	25.21
	SD	6.23	5.49	6.61	5.44	6.22	5.37
	%CV	5.66	8.79	5.83	9.26	10.84	8.75
		0.06			0.21		
	d	0.59*			0.33		
		0.61*			0.58*		
F200 ms (N.kg <sup>-1</sup> )	Mean	28.20	28.22	31.36	25.42	26.74	28.54
	SD	6.22	5.41	6.68	5.51	6.47	5.95
	%CV	4.75	7.76	4.04	7.56	9.52	8.95
		0.03			0.23		
	d	0.52*			0.29		
		0.49*			0.54*		
F250 ms (N.kg <sup>-1</sup> )	Mean	29.72	29.27	32.47	26.90	28.16	29.67
	SD	6.30	5.31	6.31	5.45	6.36	6.53
	%CV	4.18	6.99	2.89	5.75	7.32	8.54
		0.00			0.21		
	d	0.47*			0.23		
		0.44*			0.46*		
Peak Force (N.kg <sup>-1</sup> )	Mean	36.83	38.18	41.20	34.69	37.94	37.98
	SD	8.00	7.02	7.51	5.66	6.67	7.95
	%CV	3.72	3.21	3.74	3.58	2.99	3.06
		0.18			0.53*		
	d	0.42*			0.01		
		0.56*			0.48*		
*= significant ( <i>p</i> <0.05) increase							

461

## 462 POWER CLEAN

463 For the relative PC, sphericity was assumed via Mauchley's test, with both groups  
 464 demonstrating significant ( $p < 0.001$ ; power = 1.00) increases in relative PC 1RM. The Catch  
 465 group showed small significant ( $d = 0.44$ ,  $p = 0.01$ ) increases pre- ( $0.93 \pm 0.15 \text{ kg.kg}^{-1}$ ) to mid-  
 466 intervention ( $0.99 \pm 0.12 \text{ kg.kg}^{-1}$ ), with trivial non-significant ( $d = 0.15$ ,  $p = 0.14$ ) increases mid-  
 467 to post-intervention ( $1.01 \pm 0.14 \text{ kg.kg}^{-1}$ ), resulting in a small significant ( $d = 0.55$ ,  $p < 0.001$ )  
 468 increase pre- to post-intervention (Figure 5b). The Pull group showed small significant ( $d =$   
 469  $0.23$ ,  $p = 0.001$ ) increases pre- ( $0.91 \pm 0.18 \text{ kg.kg}^{-1}$ ) to mid-intervention ( $0.95 \pm 0.17 \text{ kg.kg}^{-1}$ ),

with trivial, yet significant ( $d = 0.17$ ,  $p = 0.015$ ) increases mid- to post-intervention ( $0.98 \pm 0.18$  kg.kg<sup>-1</sup>), resulting in a small significant ( $d = 0.39$ ,  $p < 0.001$ ) increase pre- to post-intervention. There were small non-significant differences ( $p > 0.05$ ) in percentage change in relative PC performance pre- to mid-intervention ( $7.4 \pm 5.0\%$ ,  $5.4 \pm 5.4\%$ ,  $d = 0.38$ ) mid- to post-intervention ( $1.9 \pm 0.8\%$ ,  $2.9 \pm 4.1\%$ ,  $d = 0.34$ ) and only trivial differences pre- to post intervention ( $9.5 \pm 6.2\%$ ,  $8.4 \pm 6.1\%$ ,  $d = 0.18$ ) between the Catch and Pull groups, respectively (Figure 5b).




Figure 5: Comparison of percentage change in isometric mid-thigh pull peak force and relative one repetition maximum power clean performances, across time points, for the Catch and Pull groups

There were no significant ( $p > 0.05$ ) changes in body mass for either the Catch (Pre  $71.14 \pm 11.79$  kg; Mid  $71.03 \pm 11.48$  kg; Post  $70.95 \pm 11.07$  kg) or the Pull group (Pre  $66.43 \pm 10.13$  kg; Mid  $66.64 \pm 9.97$  kg; Post  $66.68 \pm 10.11$  kg) across the duration of the intervention.

## Discussion

This is the first study to compare the effects of including or excluding the catch phase of PC derivatives, on training adaptations, in terms of force-time characteristics during dynamic and isometric tasks. Both groups demonstrated improvements in CMJ height, IMTP variables and PC performance pre- to post-intervention, as hypothesized. In contrast to the hypotheses, the Catch group increased SJ height, whereas there was no change in the Pull group. Also in contrast to the hypotheses, there was no difference in percentage change, in any variables, between groups, which may be attributed to the comparable training stimulus during the propulsion phase of each exercise along with the identical volume load.

The Catch group achieved moderate improvements in SJ height (12.6%) across the duration of the intervention, whereas the Pull group only demonstrated trivial increases (2.1%). It is possible that this difference is due to the requirement to rapidly produce force to arrest motion during the Catch, whereas a greater time is available to decelerate the barbell and the system center of mass during the pulling derivatives. The Catch group also exhibited greater improvements in CMJ height (10.8%), compared to the Pull group (5.2%) across the duration of the study, although improvements in both groups were small and significant, the difference in improvements between groups was small yet not significant. To achieve the CMJ heights, there were no meaningful or significant changes in TTT, implying that an increase in jump height must have been a result of an increase in force applied, resulting in an increased impulse and therefore velocity at take-off. The lack of change in TTT, combined with the increase in jump height, resulted in favorable, yet small and non-significant increases in RSImod for both the Catch (7.0%) and Pull (4.4%) groups (Figure 3). The small magnitudes of increases in jump performance are in line with previous findings, reported after a 10-week training intervention comparing the training effects of hang high pulls and hexagonal barbell jump squats, in collegiate swimmers (37). In addition, the transfer of weightlifting style training, has recently been reported to result in only small changes in jump performance over relatively short training periods (26), as observed here. In contrast however, traditional resistance training combined with weightlifting derivatives has been shown to enhance longitudinal maximal strength and jump performance (30).

Both groups demonstrated trivial to small and non-significant increases in time-specific force values during the initial four weeks (pre- to mid- intervention), with small to moderate and significant increases in the final four weeks (mid- to post-intervention). This resulted in small to moderate increases in F100 (14.9%; 15.5%), F150 (16.0%; 16.2%), F200 (15.8%; 17.9%) and F250 (15.8%; 17.9%) for the Catch and Pull groups respectively. The greater increases

in time-specific force production, during the second four weeks of training, may be due the higher intensities used, resulting in the subjects having to ensure a maximal intent and rapid force production to adequately accelerate the barbell. The Pull group consistently demonstrated a greater percentage change in all time-specific forces although these differences were small and non-significant (Figure 4). These observations are similar to those previously reported by Oranchuk et al. (37) who also reported no meaningful differences in relative PF and time-specific force variables, after 10-weeks of hang high pull versus hexagonal barbell jump squat training.

In contrast to the changes in IMTP time-specific forces, PF increased to the greatest extent during the first four weeks (pre-mid), with the Catch group demonstrating greater improvements (13.7%) compared to the Pull group (9.7%), although the differences between groups were trivial. Interestingly, PC performances exhibited similar trends, with the greatest improvements occurring during the first 4 weeks, and the Catch group demonstrating slightly greater improvements (9.5%) compared to the Pull group (8.4%). It is likely that similarity in these adaptations are due to the strong relationships between IMTP PF and PC performance previously reported (33). These greater increases in PC performance, during the first four weeks, may be due to the slightly greater volume of power clean derivatives performed during this phase, compared to the second phase. The magnitude of the changes in PC performance is also greater than the smallest worthwhile change previously reported to indicate meaningful changes for the PC (9, 14) and the IMTP (7, 14).

Both the groups improved their 1RM PC over the course of the training interventions. Interestingly, the Pull group were able to improve their 1RM PC to a similar extent compared to the Catch group despite not training with the catch phase. This is important to note considering not all individuals are able to adequately perform the catch phase due to poor technique, inflexibility or previous or current injury. Thus, training with pulling derivatives may provide an effective training stimulus for improving maximal dynamic strength, which is comparable to the use of weightlifting catching derivatives. As mentioned above, each training group exhibited small, significant training effects over the course of the study, with only a trivial difference, in the percentage increase in performance, between groups. From a specificity standpoint, this finding is unsurprising given that this group performed submaximal training with the PC exercise. These improvements in PC (9, 14, 17) and IMTP (7, 14) performance were also greater than the between session smallest detectable differences previously reported.

A potential limitation to the current study was the use of identical loading procedures between the Catch and Pull groups. In an effort to equalize training volume, each group was prescribed the same relative intensity and volume load, during each training block. While this may make sense from a research standpoint, the pulling derivatives implemented within the current study (e.g. clean grip mid-thigh pull and pull from the floor) are typically implemented using loads in excess of an athlete's 1RM PC (i.e. > 100%) (8, 10, 22, 31), while additional repetitions may be able to be performed at submaximal loads, compared to catch variations. Thus, the loads implemented for these exercises may not have provided an adequate load or volume stimulus to the Pull group, which may have prevented them from displaying greater training benefits compared to the Catch group. Given that weightlifting pulling derivatives may produce greater force and velocity characteristics, dependent on the load used (43), researchers may consider investigating the training effects of weightlifting pulling derivatives that use loads which

emphasize either a force or velocity overload stimulus, as described by Suchomel et al., (43), compared to training with weightlifting catching derivatives.

It is also worth noting, that as this was an in-season training intervention, with relatively low training volumes, to minimize any potentially negative impact on the athletes' competitive performances, a future study conducted in pre-season, is recommended, where higher training volume loads and, or relative intensities (based on 1RM PC performance) can be incorporated.

### **Practical Application**

The results of this study indicate that training with either weightlifting catching or weightlifting pulling derivatives improved the athletes' performance across a spectrum of variables. It is important to note, however that trivial to small differences existed between training groups when examining every variable, indicating that catching and pulling derivatives may provide a similar training stimulus when the same relative intensity (based on 1RM PC) and volume loads are implemented during an in-season training program. Thus, both catching and pulling derivatives may provide an effective training stimulus when training to improve strength-power characteristics. It is suggested, therefore, that strength and conditioning coaches and athletes should appropriately periodize the use of weightlifting derivatives, and that pulling and catching derivatives can be used interchangeable to achieve similar goals, when performed using the same relative intensity and volume loads.

*Acknowledgements: The authors would like to acknowledge Matthew Cuthbert, Rachel Higgins and Daniel Traynor for their assistance during the coaching sessions. We would also like to thank the participants for their time and efforts, and the NSCA for supporting this investigation as part of the International Collaboration Grant.*

*Funding: This study was funded by the NSCA foundation, as part of the International Collaboration Grant scheme (awarded 2016).*

*The authors report no conflict of interest and the finding of the study do not represent the views of the NSCA.*



## 625 References

- 626 1. Baechle TR, Earle RW, and Wathen D. Resistance Training, in: *Essentials of Strength Training*  
627 *and Conditioning*. TR Baechle, Earle, R. W, ed. Champaign, Illinois: Human Kinetics, 2008, pp  
628 381-412.
- 629 2. Beckham G, Mizuguchi S, Carter C, Sato K, Ramsey M, Lamont H, Hornsby G, Haff G, and  
630 Stone M. Relationships of isometric mid-thigh pull variables to weightlifting performance. *J*  
631 *Sports Med Phys Fitness* 53: 573-581, 2013.
- 632 3. Beckham GK, Sato K, Mizuguchi S, Haff GG, and Stone MH. Effect of Body Position on Force  
633 Production During the Isometric Mid-Thigh Pull. *J Strength Cond Res*. Publish Ahead of Print,  
634 2017.
- 635 4. Canavan PK, Garrett GE, and Armstrong LE. Kinematic and Kinetic Relationships Between an  
636 Olympic-Style Lift and the Vertical Jump. *J Strength Cond Res*. 10: 127-130, 1996.
- 637 5. Comfort P, Allen M, and Graham-Smith P. Comparisons of peak ground reaction force and  
638 rate of force development during variations of the power clean. *J Strength Cond Res* 25:  
639 1235-1239, 2011.
- 640 6. Comfort P, Graham-Smith P, and Allen M. Kinetic comparisons during variations of the  
641 Power Clean. *J Strength Cond Res* 25: 3269-3273, 2011.
- 642 7. Comfort P, Jones PA, McMahon JJ, and Newton R. Effect of knee and trunk angle on kinetic  
643 variables during the isometric midthigh pull: test-retest reliability. *Int J Sports Physiol*  
644 *Perform* 10: 58-63, 2015.
- 645 8. Comfort P, Jones PA, and Udall R. The effect of load and sex on kinematic and kinetic  
646 variables during the mid-thigh clean pull. *Sports Biomech* 14: 139-156, 2015.
- 647 9. Comfort P and McMahon JJ. Reliability of Maximal Back Squat and Power Clean  
648 Performances in Inexperienced Athletes. *J Strength Cond Res*. 29: 3089-3096, 2015.
- 649 10. Comfort P, Udall R, and Jones P. The affect of loading on kinematic and kinetic variables  
650 during the mid-thigh clean pull. *J Strength Cond Res* 26: 1208-1214, 2012.
- 651 11. Comfort P, Williams R, Suchomel TJ, and Lake JP. A comparison of catch phase force-time  
652 characteristics during clean derivatives from the knee. *J Strength Cond Res*. 21 (7): 2017.
- 653 12. Cormack SJ, Newton RU, McGuigan MR, and Doyle TL. Reliability of measures obtained  
654 during single and repeated countermovement jumps. *Int J Sports Physiol Perform* 3: 131-144,  
655 2008.
- 656 13. Dos'Santos T, Jones PA, Comfort P, and Thomas C. Effect of Different Onset Thresholds on  
657 Isometric Mid-Thigh Pull Force-Time Variables. *J Strength Cond Res* Publish Ahead of Print,  
658 2017.
- 659 14. Dos'Santos T, Thomas C, Comfort P, McMahon JJ, Jones PA, Oakley NP, and Young AL.  
660 Between-Session Reliability Of Isometric Mid-Thigh Pull Kinetics And Maximal Power Clean  
661 Performance In Male Youth Soccer Players. *J Strength Cond Res* Published ahead of print,  
662 2017.
- 663 15. Dos'Santos T, Thomas C, Jones PA, McMahon JJ, and Comfort P. The Effect Of Hip Joint Angle  
664 On Isometric Mid-Thigh Pull Kinetics. *J Strength Cond Res*. Publish Ahead of Print, 2017.
- 665 16. Enoka RM. The pull in olympic weightlifting. *Med Sci Sports* 11: 131-137, 1979.
- 666 17. Faigenbaum AD, McFarland JE, Herman RE, Naclerio F, Ratamess NA, Kang J, and Myer GD.  
667 Reliability of the one-repetition-maximum power clean test in adolescent athletes. *J*  
668 *Strength Cond Res* 26: 432-437, 2012.
- 669 18. Garhammer J and Gregor R. Propulsion Forces as a Function of Intensity for Weightlifting  
670 and Vertical Jumping. *J Strength Cond Res*. 6: 129-134, 1992.
- 671 19. Haff GG, Carlock JM, Hartman MJ, Kilgore JL, Kawamori N, Jackson JR, Morris RT, Sands WA,  
672 and Stone MH. Force-time curve characteristics of dynamic and isometric muscle actions of  
673 elite women olympic weightlifters. *J Strength Cond Res* 19: 741-748, 2005.

- 674 20. Haff GG, Ruben RP, Lider J, Twine C, and Cormie P. A Comparison of Methods for  
675 Determining the Rate of Force Development During Isometric Mid-Thigh Clean Pulls. *J*  
676 *Strength Cond Res* 29: 386-395, 2015.
- 677 21. Haff GG, Stone M, O'Bryant HS, Harman E, Dinan C, Johnson R, and Han K-H. Force-Time  
678 Dependent Characteristics of Dynamic and Isometric Muscle Actions. *J Strength Cond Res* 11:  
679 269-272, 1997.
- 680 22. Haff GG, Whitley A, McCoy LB, O'Bryant HS, Kilgore JL, Haff EE, Pierce K, and Stone MH.  
681 Effects of Different Set Configurations on Barbell Velocity and Displacement During a Clean  
682 Pull. *J Strength Cond Res* 17: 95-103, 2003.
- 683 23. Halperin I, Williams KJ, Martin DT, and Chapman DW. The Effects of Attentional Focusing  
684 Instructions on Force Production During the Isometric Midthigh Pull. *J Strength Cond Res* 30:  
685 919-923, 2016.
- 686 24. Halperin I, Williams KJ, Martin DT, and Chapman DW. The Effects of Attentional Focusing  
687 Instructions on Force Production During the Isometric Midthigh Pull. *J Strength Cond Res*. 30:  
688 919-923, 2016.
- 689 25. Harbili E and Alptekin A. Comparative kinematic analysis of the snatch lifts in elite male  
690 adolescent weightlifters. *J Sports Sci Med* 13: 417-422, 2014.
- 691 26. Helland C, Hole E, Iversen E, Olsson MC, Seynnes O, Solberg PA, and Paulsen G. Training  
692 Strategies to Improve Muscle Power: Is Olympic-style Weightlifting Relevant? *Med Sci Sports*  
693 *Exerc* 49: 736-745, 2017.
- 694 27. <http://sportsci.org/resource/stats/index.html>. Accessed 08/05/15/2015.
- 695 28. Hopkins WG, Marshall SW, Batterham AM, and Hanin J. Progressive statistics for studies in  
696 sports medicine and exercise science. *Med Sci Sports Exerc* 41: 3-13, 2009.
- 697 29. Hori N, Newton RU, Andrews WA, Kawamori N, McGuigan MR, and Nosaka K. Does  
698 Performance of Hang Power Clean Differentiate Performance of Jumping, Sprinting, and  
699 Changing of Direction? *J Strength Cond Res* 22: 412-418 2008.
- 700 30. Kavanaugh AA, Mizuguchi S, Sands WA, Ramsey MW, and Stone MH. Long-Term Changes In  
701 Jump Performance And Maximum Strength In A Cohort Of Ncaa Division I Women's  
702 Volleyball Athletes. *J Strength Cond Res* Publish Ahead of Print, 2017.
- 703 31. Kawamori N, Rossi SJ, Justice BD, Haff EE, Pistilli EE, O'Bryant HS, Stone MH, and Haff GG.  
704 Peak Force and Rate of Force Development During Isometric and Dynamic Mid-Thigh Clean  
705 Pulls Performed At Various Intensities. *J Strength Cond Res* 20: 483-491, 2006.
- 706 32. Kipp K, Malloy PJ, Smith J, Giordanelli MD, Kiely MT, Geiser CF, and Suchomel TJ. Mechanical  
707 Demands of the Hang Power Clean and Jump Shrug: A Joint-level Perspective. *J Strength*  
708 *Cond Res* Publish Ahead of Print, 2017.
- 709 33. McGuigan M and Winchester JB. The relationship between isometric and dynamic strength  
710 in collegiate football players. *J Sports Sci Med* 7: 101-105, 2008.
- 711 34. McMahon JJ, Jones PA, Suchomel TJ, Lake J, and Comfort P. Influence of Reactive Strength  
712 Index Modified on Force- and Power-Time Curves. *Int J Sports Physiol Perform* E-pub ahead  
713 of print: 1-24, 2017.
- 714 35. Moir GL. Three Different Methods of Calculating Vertical Jump Height from Force Platform  
715 Data in Men and Women. *Measurement in Physical Education and Exercise Science* 12: 207-  
716 218, 2014.
- 717 36. Moolyk AN, Carey JP, and Chiu LZF. Characteristics of Lower Extremity Work During the  
718 Impact Phase of Jumping and Weightlifting. *J Strength Cond Res* 27: 3225-3232, 2013.
- 719 37. Oranchuk DJ, Robinson TL, Switaj ZJ, and Drinkwater EJ. Comparison of the Hang High-Pull  
720 and Loaded Jump Squat for the Development of Vertical Jump and Isometric Force-Time  
721 Characteristics. *J Strength Cond Res*, 2017.
- 722 38. Owen NJ, Watkins J, Kilduff LP, Bevan HR, and Bennett MA. Development of a criterion  
723 method to determine peak mechanical power output in a countermovement jump. *J*  
724 *Strength Cond Res* 28: 1552-1558, 2014.

39. Souza AL, Shimada SD, and Koontz A. Ground reaction forces during the power clean. *J Strength Cond Res* 16: 423-427, 2002.
40. Stone MH, Sanborn KIM, O'Bryant HS, Hartman M, Stone ME, Proulx C, Ward B, and Hruby JOE. Maximum Strength-Power-Performance Relationships in Collegiate Throwers. *J Strength Cond Res* 17: 739-745, 2003.
41. Suchomel T, Comfort P, and Stone M. Weightlifting Pulling Derivatives: Rationale for Implementation and Application. *Sports Medicine* 45: 823-839, 2015.
42. Suchomel TJ, Bailey CA, Sole CJ, Grazer JL, and Beckham GK. Using reactive strength index-modified as an explosive performance measurement tool in Division I athletes. *J Strength Cond Res* 29: 899-904, 2015.
43. Suchomel TJ, Comfort P, and Lake JP. Enhancing the Force-Velocity Profile of Athletes Using Weightlifting Derivatives. *Strength & Conditioning Journal* 39: 10-20, 2017.
44. Suchomel TJ, Lake JP, and Comfort P. Load absorption force-time characteristics following the second pull of weightlifting derivatives. *J Strength Cond Res* 31: 1644-1652, 2017.
45. Suchomel TJ and Sole CJ. Force-Time Curve Comparison Between Weightlifting Derivatives. *International Journal of Sports Physiology and Performance* Published ahead of print, 2017.
46. Suchomel TJ and Sole CJ. Power-Time Curve Comparison between Weightlifting Derivatives. *J Sports Sci Med* 16: 407-413, 2017.
47. Suchomel TJ, Wright GA, Kernozek TW, and Kline DE. Kinetic Comparison of the Power Development Between Power Clean Variations. *J Strength Cond Res* 28: 350-360, 2014.

#### Figure and Table Legends:

Figure 1: Summary of testing schedule

Figure 2: Testing sequence (SJ: squat jump, CMJ: countermovement jump, IMTP: Isometric Mid-Thigh Pull)

776 Figure 3: Comparison of percentage change in jump variables across time points for the Catch  
777 and Pull groups

778 Figure 4: Comparison of percentage change in isometric mid-thigh pull time specific force  
779 variables, across time points, for the Catch and Pull groups

780 Figure 5: Comparison of percentage change in isometric mid-thigh pull peak force and relative  
781 one repetition maximum power clean across time points for the Catch and Pull groups

782

783

784 Table 1: Training sessions, weeks 1-4

785 Table 2: Training sessions, weeks 6-9

786 Table 3: Within and between session reliability (ICC (95% confidence intervals)) and variability  
787 (% coefficient of variation) of jump performance variables

788 Table 4: Within and between session reliability (ICC (95% confidence intervals)) and variability  
789 (% coefficient of variation) of IMTP variables

790 Table 5: Changes in jump performance

791 Table 6: Changes in isometric mid-thigh pull performance