

Please cite the Published Version

Joanisse, Sophie ^(D), Ashcroft, Stephen, Wilkinson, Daniel J, Pollock, Ross D, O'Brien, Katie A, Phillips, Bethan E, Smith, Ken, Lazarus, Norman R, Harridge, Stephen DR, Atherton, Philip J and Philp, Andrew (2020) High Levels of Physical Activity in Later Life Are Associated With Enhanced Markers of Mitochondrial Metabolism. The Journals of Gerontology: Series A, 75 (8). pp. 1481-1487. ISSN 1079-5006

DOI: https://doi.org/10.1093/gerona/glaa005

Publisher: Oxford University Press (OUP)

Version: Accepted Version

Downloaded from: https://e-space.mmu.ac.uk/626373/

Usage rights: O In Copyright

Additional Information: This is an Author Accepted Manuscript of a paper accepted for publication in The Journals of Gerontology: Series A, published by and copyright Oxford University Press (OUP).

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines)

High levels of physical activity in later life are associated with enhanced markers of mitochondrial metabolism

Sophie Joanisse PhD^{1,2}, Stephen Ashcroft MPhil¹, Daniel J Wilkinson PhD³, Ross D Pollock PhD⁴, Katie A O'Brien PhD⁴, Bethan E Phillips PhD³, Ken Smith PhD³, Norman R Lazarus PhD⁴, Stephen DR Harridge PhD⁴, Philip J Atherton PhD³, Andrew Philp PhD^{1,5,6*}

1. School of Sport, Exercise and Rehabilitation Sciences, University of Birmingham, United Kingdom.

2. Department of Kinesiology, McMaster University, Hamilton, Ontario Canada.

3. Division of Medical Sciences and Graduate Entry Medicine, University of Nottingham, United Kingdom.

4. Centre for Human and Applied Physiological Sciences, King's College London, United Kingdom.

5. Garvan Institute of Medical Research, Darlinghurst, NSW, 2010, Australia.

6. St Vincent's Clinical School, UNSW Medicine, UNSW Sydney, NSW, 2010, Australia

* Address for correspondence:

Dr Andrew Philp

Mitochondrial Metabolism and Ageing Laboratory

Division of Diabetes and Metabolism

Garvan Institute of Medical Research

384 Victoria Street, Darlinghurst, NSW, 2010, Australia.

Email: <u>a.philp@garvan.org.au</u>

Phone: +61 9295 8249

Abstract

The age-associated reduction in muscle mass is well characterised, however less is known regarding the mechanisms responsible for the decline in oxidative capacity also observed with advancing age. The purpose of the current study was therefore to compare mitochondrial gene expression and protein content between young and old recreationally active, and older highly active individuals. Muscle biopsies were obtained from the vastus lateralis of young males (YG: 22±3 years) and older (OG: 67±2 years) males not previously engaged in formal exercise and older male master cyclists (OT: 65±5 years) who had undertaken cycling exercise for 32±17 yrs. Comparison of gene expression between YG, OG and OT groups revealed greater expression of mitochondrial-related genes, namely electron transport chain (ETC) complexes II, III, IV (p<.05) in OT compared to YG and OG. Gene expression of mitofusion (MFN)-1/2, mitochondrial fusion genes, were greater in OT compared to OG (p<.05). Similarly, protein content of ETC complexes I, II and IV were significantly greater in OT compared to both YG and OG (p<.001). Protein content of peroxisome proliferatoractivated receptor gamma, coactivator 1 α (PGC-1 α), was greater in OT compared to YG and OG (p<.001). Our results suggest that the aging process per se, is not associated with a decline in gene expression and protein content of ETC complexes. Mitochondrial-related gene expression and protein content is substantially greater in OT, suggesting that exercisemediated increases in mitochondrial content can be maintained into later life.

Keywords: human, athlete, muscle, mitochondria

Running title: Markers of mitochondrial biology are enhanced in master athletes

Introduction

Skeletal muscles of most older individuals are characterized by several adverse morphological changes¹⁻⁴, a phenotype directly influenced by physical activity ⁵⁻⁷. Some groups report an age-associated decline in fibre cross sectional area in sprint-trained older individuals⁸ whereas this association is not observed in highly trained master cyclists⁶. This observation therefore questions the extent to which these adverse changes to skeletal muscle health are a function of the aging process or the interaction with age and physical inactivity⁸.

Considerable attention has been given to determining the effects of aging on mitochondrial function in skeletal muscle⁹. In addition to being associated with a decline in muscle fatigue resistance, aberrant mitochondrial function is proposed as a major contributor to sarcopenia¹⁰. In support of this motion, previous research has reported an age-associated decline in mtDNA¹¹, in addition to a reduction in gene expression of mitochondrial transport chain subunits¹². The reduction in mitochondrial related gene expression translates to the decline in mitochondrial related protein content observed with aging¹¹, although this apparent reduction is not observed in middleaged adults where protein content of electron transport chain (ETC) complex I is greater when compared to young¹³. In addition, aging is associated with an increase in mtDNA mutationdeletions¹⁴, an increase in oxidative damage¹² and altered protein content of factors involved in mitochondrial dynamics¹⁵. These findings however are not universal, as other groups have reported that aging had no effect on mitochondrial associated gene networks¹⁶, mitochondrial respiration and protein content of ETC subunits¹⁹ and regulators of mitochondrial dynamics¹⁷. Contrasting results may be due to a number of factors, one of which may be the physical activity status of old individuals included in these studies. For example, Disteffano and colleagues recently demonstrated that cardiorespiratory fitness has a greater influence on mitochondrial respiration than age¹⁷. Although sedentary aging is associated with a variety of metabolic perturbations, it is suggested that these may be related to the low levels of physical activity rather than to the aging process per se^{6} . Or

alternatively, physical activity remains beneficial even in advancing age. This latter notion is supported by studies demonstrating that older athletes, when compared to sedentary counterparts, have greater mitochondrial content, function²⁰, gene expression of ETC subunits¹⁹, and protein content related to mitochondrial dynamics¹⁷. In older individuals, mitochondrial volume density is correlated with exercise capacity suggesting a role for physical activity to maintain mitochondrial content in this population²¹. Importantly, older individuals retain the ability to positively respond to aerobic exercise training^{21,22}, placing regular physical exercise as an effective measure for improving mitochondrial content and function throughout the lifespan.

Several studies have sought to characterize the differences in gene and protein expression between either young and sedentary older adults or older sedentary and older athletes. However, few have examined the differences between young and old recreationally active and old highly active master athletes, thereby allowing the combined comparison of age and physical activity levels. Therefore, the purpose of the current study was to compare mitochondrial-related gene expression and protein abundance in the quadriceps muscle between recreationally active young (YG), old individuals who were healthy, but did not undertake regular exercise (OG) and older individuals who had been highly active through cycling exercise for many years (OT). We determined whether genes and proteins related to whole muscle metabolism, mitochondrial biogenesis, dynamics and mitophagy were differentially expressed between groups to further characterize the aging muscle phenotype and the impact of physical activity.

Methods

Ethical Approval.

Prior to participation written informed consent was obtained from all subjects. Procedures were approved by the National Health Service Wandsworth Research Ethics Committee (reference number 12/LO/0457) and the University of Nottingham Faculty of Medicine and Health Science Research Ethics Committee (B/10/2010), with all procedures conforming to the Declaration of Helsinki. All human tissue collected, stored and analysed was done so in accordance with the Human Tissue Act.

Participants.

The OT group is a subset from a large cross-sectional study previously described by Pollock et al. (2015) in which the physiological function of a group of amateur non-elite cyclists aged 55-79 years was evaluated²³. The inclusion criteria of males was the ability to cycle 100 km in under 6.5h and subjects were required to have undertaken this task twice in the 3 weeks prior to testing²⁰. Subjects included in the OG group (67±2years; 27±2.8kg/m²; n=8, for subsequent gene expression and protein content analysis) were selected to have similar ages and body mass index (BMI) to those of the OT group (65±5years; 25.9±2.1kg/m²; n=7, for subsequent gene expression analysis, n=8 for subsequent protein content analysis). Subjects included in the YG group (21±3years; 23.9±2.5 kg/m²; n=6, for subsequent gene expression analysis, n=9 for subsequent protein content analysis) were selected to best match BMI from the older groups. Both YG and OG were recreationally active and performed activities of daily living but none were involved in a formal exercise training programme nor did they participate in more than 2 sessions of purposeful exercise each week. No differences between BMI are observed between groups and no differences between ages are observed between the OT and OG groups. Due to lack of tissue the sample size was smaller for gene expression analysis as stated above.

Muscle biopsy sampling.

Following administration of local anaesthetic (1% lidocaine for the YG and OG; 2% lidocaine for the OT) a muscle sample was obtained from the *vastus lateralis* of all participants using the conchotome technique²⁴ for the YG and OG groups and the Bergstrom needle technique with applied suction for the OT group⁶. All participants in the YG and OG groups were instructed not to exercise for 72h prior to their biopsy which was taken after an overnight fast (~12 h) with water *ad libitum*. Participants in the OT group were asked to maintain habitual levels of physical activity the day prior to their biopsy.

RNA isolation and Reverse Transcription

RNA was isolated from 15 – 20 mg of powdered muscle tissue using TRI Reagent/ReliaPrep spin columns. Briefly, all samples were homogenized in 1ml of TRI Reagent (Sigma Aldrich, Gillingham, UK) with the FastPrep 24 5G (MP Biomedicals, Santa Ana, California, USA) at a speed of 6 m/s for 40 seconds. 200ul of chloroform was added to each sample and mixed vigorously for 15 sec then incubated at room temperature for 5 min, then centrifuged at 12000g for 10 min at 4°C. The RNA (aqueous phase) was purified using Reliaprep spin columns (Promega, Madison, Wisconsin, USA) as per manufacturer's instructions. RNA concentrations were determined using the LVis function of the FLUOstar Omega microplate reader. RNA was diluted to 20 μ g/ μ L and reverse transcribed to cDNA using the RT² First Strand kit (Qiagen, Manchester, UK).

Quantitative RT-PCR

Quantitative analysis of 84 genes was completed using custom designed 384-well RT² PCR Profiler Array (Qiagen) and RT² SYBR Green Mastermix (Qiagen) on a CFX384 Real-Time PCR Detection System following manufacturer's instructions (Bio-Rad). Briefly, 2.8 ng of cDNA was added to each well. The absence of genomic DNA, the efficiency of reverse-transcription and the efficiency of the PCR assay were assessed on each plate and conformed to the manufacturer's limits in each case. The C(t) values for housekeeper genes beta actin (Refseq# NM_001101), heat shock protein 90 (Refseq# NM_007355) and beta-2-microglobulin (Refseq# NM_004048) showed no statistical differences between groups. Therefore, the geometric mean C(t) of all three housekeeper genes was used as an internal control²³. Statistical analysis was carried out on the $\Delta\Delta$ C(t) (Δ C(t) gene of interest - Δ C(t) mean of gene of interest of the YG). Data is presented as a fold change from YG as determined using the 2^{- $\Delta\Delta$ C(t)} method**26²⁷**.

Immunoblotting

Approximately 25mg of powdered muscle tissue was homogenized in 300 ul of ice-cold sucrose lysis buffer (50 mM Tris, 1 mM EDTA, 1 mM EGTA, 50 mM NaF, 5 mM Na₄P₂O7-10H₂O, 270 mM sucrose, 1 M Triton-X, 25 mM β -glycerophosphate, 1 μ M Trichostatin A, 10 mM Nicatinamide, 1mM 1,4-Dithiothreitol, 1% Phosphatase Inhibitor Cocktail 2; Sigma, 1% Sigma Phosphatase Inhibitor Cocktail 2; Sigma, 4.8% cOmplete Mini Protease Inhibitor Cocktail; Roche) using the FastPrep 24 5G (MP Biomedicals, Santa Ana, California, USA) at a speed of 6 m/s for 40 seconds and repeated 3 times. Samples were then centrifuged at 4°C at a speed of 8000 g for 10 minutes to remove insoluble material. Protein content was determined from the DC protein assay (Bio-Rad, Hercules, California, USA) using FLUOstar Omega at an absorbance of 750nm). Laemmli samples buffer was added and samples were boiled for 5 minutes, equal amounts of protein (20-50 ug) was separated by SDS-PAGE on 8 – 12.5% gels at a constant current of 23 mA per gel. Proteins were transferred to BioTrace NT nitrocellulose membranes (Pall Life Sciences, Pensacola, Florida, USA) via wet transfer at 100 V for one hour on ice. Membranes were stained with Ponceau S (Sigma-Aldrich, Gillingham, UK) and imaged to assure even loading and for future normalization. Membranes were blocked in 3% drymilk in tris-buffered saline with tween (TBST) for one hour prior to an overnight primary antibody incubation at 4°C. Membranes were washed in TBST three times prior to incubation in appropriate horse radish peroxidase-conjugated secondary antibody at room temperature for one hour. Membranes were then washed in TBST three times prior to antibody detection via enhanced chemiluminescence horseradish peroxidase substrate detection kit (Millipore, Watford, UK). Imaging

and band quantification were undertaken using a G:Box Chemi-XR5 (Syngene, Cambridge, UK). Mean pixel intensity was determined for each band. All bands were normalized to a gel control in addition to the corresponding ponceau image. Statistical analysis was carried out on normalized values for each protein of interest.

Antibodies

All primary antibodies were prepared in TBST. Antibodies used were as follows: total OXPHOS human WB antibody cocktail (ab11041; abcam, 1:1000 in unboiled samples), PGC-1Antibody (AB3242; Merck, 1:1000), pyruvate dehydrogenase antibody (PDH; #2784; Cell signalling technology, 1:1000), Citrate synthase (SAB2701077; MitoSciences, 1:1000), Sirtuin3 (SirT3 D22A3; Cell signalling technology, 1:1000). Very long chain acyl CoA dehydrogenase (VLCAD; 1:5000), long chain acyl CoA dehydrogenase (LCAD; 1:5000), medium chain acyl-CoA dehydrogenase (MCAD; 1:5000), were kind gifts from Prof Jerry Vockley, University of Pittsburgh, USA. Appropriate secondary antibodies either anti-rabbit (7074) or anti-mouse (7076) from Cell Signaling Technology were used at a concentration of 1:10000 in TBST.

Statistical Analysis

Statistical analysis was performed using Prism 7 (GraphPad Software Incorporated). To assess differences between YG, OG and OT groups a one-way analysis of variance (ANOVA) was completed on individual genes and proteins independently, multiple comparisons were assessed using Tukey's test. The normal distribution of data was verified using the Shapiro-Wilk normality test. Values are presented as mean ± standard deviation.

Results

Mitochondrial related genes expression profile is altered in master athletes

The mRNA expression of 84 target genes was assessed in YG (n=6), OG (n=8) and OT (n=7) groups, gene expression is presented as a fold change from the YG group (Supplemental Table 1). Of the 84 genes assessed 23 were differentially expressed (p<.05) (Supplemental Table 1). Differentially expressed genes of mitochondrial related proteins and transcriptional regulators are presented in Figure 1. Differentially expressed genes related to skeletal muscle structural remodelling are presented in Figure 2. Differentially expressed genes related to substrate metabolism are presented in Figure 3.

Mitochondrial related protein content.

Protein content was assessed in YG (n=9), OG (n=8) and OT (n=8). Protein content of each complex of the electron transport chain was assessed via immunoblotting. There was no difference between YG and OG for any protein assessed. Protein content of complex I, II and IV were 7.1-, 1.9- and 1.3fold greater in OT compared to OG and 5.6-, 2.1- and 1.2-fold greater compared to YG (p<.05) (Fig 4a). There are no differences in protein content of complex III and V across groups (p>.05). Protein content of PGC-1 α was 2.9-fold greater in OT compared to OG (p<.05) and 5.4-fold greater compared to YG (p<.001) (Fig 4b). Protein content of CS was 1.9-fold greater in the OT compared to both the OG (p<.001) and 1.3-fold greater in the OT compared to the YG (p<.001) (Fig 4c). Additionally, protein content of SIRT3 was 1.8-fold greater in OT compared to both OG (p<.001) and to YG (p<.001) (Fig 4d).

Protein content related to substrate metabolism.

Protein content of pyruvate dehydrogenase (PDH), LCAD and MCAD was 3-, 1.8- and 1.6-fold greater, respectively in OT compared to OG and 2.8-, 1.9- and 2.1-fold greater, respectively,

compared to YG (p<.05) (Fig 5a,b,c). Protein content of VLCAD was 1.7-fold greater in OT compared to OG (p<.05) (Fig 5d).

Discussion

Many studies have sought to determine the effect of aging on mitochondrial content and function^{11-15,17-19}, however few have accounted for the divergent levels of physical activity in the older population. Here we have directly compared young and old recreationally active adults in addition to old highly aerobically trained (master cyclists) adults. The inclusion of these three distinct groups allows the effects of activity to be established and therefore what extent declines in function can be attributed to chronological aging⁸. The primary finding of this study is that highly active older adults, have greater mitochondrial related protein content and gene expression in comparison to recreationally active untrained young and older adults. Importantly, no observable age-associated reduction in mitochondrial related protein content and gene expression (except for CS mRNA expression) was observed in the recreationally active older group in which the negative effects of inactivity are eliminated. In addition, we report no change in OXPHOS, PGC-1α or mitochondrial fusion and fission-related gene expression with age.

Previous work has described a reduction in mtDNA, mitochondrial protein content and ATP production in older adults who had similar physical activity levels to young counterparts (<30 mins, <2 days/week)¹¹. However more recent work suggests that mitochondrial related protein content may not be affected by aging¹⁷. Our results are in line with the latter in which chronological age does not impact mitochondrial related protein content (Fig 4a). A lower expression in OG compared to YG is only observed with respect to CS (Fig 1c) and UQCRC1 (Fig 1d) gene expression. However, this was not observed at the protein level (Fig 4c). Although chronological aging *per se* did not impact mitochondrial related protein content and gene expression in our hands, the impact of high levels of physical activity in advanced age was substantial. We report greater expression of genes related to the electron transport chain (SDHA, SDHB, UQCRC1, UQCRC2 and COX4l2) in the OT compared to

both the YG and OG groups (Fig 1). These results are supported by greater protein content of complex I, II and IV of the electron transport chain in the OT compared to YG and OG groups (Fig 4a). Larsen and co-workers have reported an association between protein content of complex II and V and total mitochondrial content measured via mitochondrial volume by transmission electron microscopy²⁹. Although not a direct measure of mitochondrial content, our results are in accordance with previous work demonstrating a greater mitochondrial volume density and content of ETC proteins in old active compared to sedentary adults^{20,21}. Given that endurance exercise training is well established as a potent activator of mitochondrial biogenesis³⁰, the greater mitochondrial related gene expression and protein content in the OT group is likely due to their high physical activity levels and the metabolic demands placed on skeletal muscle.

In addition, we report greater PGC-1 α gene expression in the OT compared to the OG group (Fig 1b) and greater PGC-1 α protein content in OT compared to both the YG and OG groups (Fig 4b). These results are in partial contrast with previous work reporting a reduction in PGC-1 α protein content with aging in both 'high' and 'low' functioning sedentary older adults²⁸. As PGC-1 α is an important regulatory signalling node for the initiation of mitochondrial biogenesis, a greater expression in the OT group may reflect a greater mitochondrial mass due to greater habitual physical activity levels in this cohort. Thus, our results suggest that mitochondrial biogenesis is primarily affected by activity status rather than aging. PGC-1 α also regulates the gene expression of SIRT3³¹, a NAD*-dependent deacetylase highly involved in metabolism³². Accordingly, we report greater expression (Fig 1a) and protein content (Fig 4d) of SIRT3, in the OT group compared to both the YG and the OG groups. Again, this observation suggests that aging *per se* is not associated with a reduction in markers of mitochondrial metabolism; rather that exercise training has a positive effect on SIRT3 induction irrespective of age.

Skeletal muscle mitochondrial function is believed, in part, to be regulated by the tumour suppressor p53³⁵. Here we demonstrate that gene expression of p53 targets, SESN2 and PMPAIP1,

are increased in the OT group although not affected by aging (Fig 2). ZMAT3 another target of p53, has a greater expression in both OG and OT (Fig 2g,h). Interestingly, no change in p53 expression was observed between groups (p>.05) (Supplemental Table 1). With the design of the current study we are unable to state whether the increase in p53 target genes are beneficial or detrimental in maintaining muscle health.

Information regarding content of proteins involved in mitochondrial dynamics in aging skeletal muscle are equivocal with some reports indicating either no change or a reduction^{15,17,28}. Recently, Tezze et al.(2017), have reported greater expression of OPA1, MFN1/2 and DRP1 in "older sportsmen" and young individuals compared to old inactive individuals¹⁵. Balan et al. (2019) report greater mitochondrial protein content of MFN2 and OPA1 in active young and older adults in comparison to sedentary age-matched subjects however, they do not observe an age-associated reduction in protein content¹⁹. partially confirmed by our findings where expression of MFN1 and MFN2, were greater in OT compared to OG, however, in our hands aging did not result in a reduced expression (Fig 2e,f). This may relate to possible differences in the activity status of the recreational active older participants in the present study and the "sedentary seniors" in the study of Tezze et al. (2017)¹⁵. Additionally, we did not observe differences in expression of other genes related to mitochondrial dynamics, such as OPA1 and FIS1 (Supplemental Table 1). Our results suggest that aging does not impair the dynamic remodelling of the mitochondrial reticulum but highly active older adults may require enhanced mitochondrial remodelling to support the stresses placed on the skeletal muscle as evidenced by greater gene expression of MFN1/2.

Although we report very few changes associated with chronological aging with respect to protein and gene expression, it is important to note that skeletal muscle of older individuals remains a highly plastic tissue and responds positively to exercise training³⁷⁻⁴¹. Although, older individuals' response to resistance exercise training may be attenuated in comparison to young adults, they are able to increase muscle fibre size⁴¹. In line with this, endurance exercise training in older individuals' results in improved maximal oxygen uptake, mitochondrial enzyme activities, and increase in muscle

fibre size⁴²⁻⁴⁴. Importantly, a recent analysis of cross-sectional data demonstrates that when older adults commence intense training and competition in later life (>50y), no differences are observed with respect to performance and body composition (fat mass and leg lean mass) compared to adults having trained their entire adult life²². Thus, further highlighting the ability of older individuals to positively respond to exercise.

One of the limitations of the present study was the lack of precise quantification of levels of physical activity in the groups that described themselves as being recreationally active. It is likely however, that the OG did not reflect the majority of older people who are known to engage in reduced levels of physical activity⁴⁶. Thus, comparison with the OT group is, if anything, most likely to be an underestimate of the differences in phenotype between lifelong exercisers and most of the older population.

In summary, we observed greater mitochondrial related gene expression and protein content in highly active older individuals compared to young and aged-matched recreationally active individuals. Chronological aging in recreationally active individuals, did not affect mitochondrial related gene and protein expression. Collectively these data suggests that high levels of physical activity, even in advancing age, results in greater mitochondrial-related gene expression and protein content compared to young and age matched untrained individuals. Further, our results highlight that chronological aging does not result in a reduction in mitochondrial content and function and supports the notion that skeletal muscle retains the ability to positively respond to stimuli even in advancing age.

CONFLICTS OF INTEREST

The authors have no conflicts of interest to disclose.

ACKNOWLEDGMENTS

The authors would like to thank the research participants for their time and effort. All authors gave their final approval of the version of the article to be published. P.J.A, S.D.R and A.P designed the study. P.J.A, S.D.R, R.D.P, N.R.L, D.J.W, B.E.P and K.S organized and carried out sample collection. S.J, S.A and K.A.O performed experiments, S.J, S.A and A.P completed data analysis. S.J and A.P performed the statistical analysis of the data. S.J and A.P wrote the manuscript.

References

1. Lexell J. Human aging, muscle mass, and fiber type composition. *J Gerentol A Biol Sci Med Sci.* 1995;50 Spec No:11-6. doi: https://doi.org/10.1093/gerona/50a.special_issue.11

2. McKay BR, Ogborn DI, Bellamy LM, et al.Myostatin is associated with age-related human muscle stem cell dysfunction. *FASEB J.* 2012;26(6):2509-2521. doi: https://doi.org/10.1096/fj.11-198663

3. Nederveen JP, Joanisse S, Snijders T, et al. Skeletal muscle satellite cells are located at a closer proximity to capillaries in healthy young compared with older men. *J Cachexia Sarcopenia Muscle*. 2016;7(5):547-554. doi: https://doi.org/10.1002/jcsm.12105

4. Piasecki M, Ireland A, Piasecki J, et al. Failure to expand the motor unit size to compensate for declining motor unit numbers distinguishes sarcopenic from non-sarcopenic older men. *J Physiol*. 2018;596(9):1627-1637. doi: https://doi.org/10.1113/jp275520

5. Klitgaard H, Mantoni M, Schiaffino S, et al. Function, morphology and protein expression of ageing skeletal muscle: a cross-sectional study of elderly men with different training backgrounds. *Acta Physiol Scand.* 1990;140(1), 41-54. doi: https://doi.org/10.1111/j.1748-1716.1990.tb08974.x

6. Pollock RD, O'Brien KA, Daniels LJ, et al. Properties of the vastus lateralis muscle in relation to age and physiological function in master cyclists aged 55-79 years. *Aging Cell.* 2018;17(2). doi: https://doi.org/10.1111/acel.12735

7. McKendry J, Joanisse S, Baig S, et al. Superior Aerobic Capacity and Indicies of Skeltal Muscle Morphology in Chronically Trained Master Endurance Athletes Compared with Untrained Older Adults. *J Gerontol A Biol Sci Med Sci*. 2019;glz142. doi: https://doi.org/10.1093/gerona/glz142

8. Lazarus NR, Harridge DR. Declining performance of master athletes: silhouette of the trajectory of healthy human ageing? *J Physiol*. 2017;595(9): 2941-2948. doi: https://doi.org/10.1113/jp272443

9. Gouspillou G, Hepple RT. Editorial: Mitochondria in Skeletal Muscle Health, Aging and Diseases. *Front Physiol*. 2016;6(7):446. doi: https://doi.org/10.3389/fphys.2016.00446

10. Peterson CM, Johannsen DL, Ravussin E (2012) Skeletal muscle mitochondria and aging: a review. J Aging Res 2012:194821. doi: https://doi.org/10.1155/2012/194821

11. Short KR, Bigelow ML, Kahl J, et al. Decline in skeletal muscle mitochondrial function with aging in humans. *Proc Natl Acad Sci U S A.* 2005;102(15):5618-5623. doi: https://doi.org/10.1073/pnas.0501559102

12. Zahn JM, Sonu R, Vogel H, et al.Transcriptional profiling of aging in human muscle reveals a common aging signature. *PLoS Genet.* 2006;2(7):e115. doi: https://doi.org/10.1371/journal.pgen.0020115.eor

13. Larsen S, Hey-Mogensen M, Rabol R, et al. (2012a) The influence of age and aerobic fitness: effects on mitochondrial respiration in skeletal muscle. *Acta Physiol (Oxf)*. 2012a; 205(3):423-432. doi: https://doi.org/10.1111/j.1748-1716.2012.02408.x

14. Bua E, Johnson J, Herbst A, et al. Mitochondrial DNA-deletion mutations accumulate intracellularly to detrimental levels in aged human skeletal muscle fibers. *Am J Hum Genet*. 2006;79(3):469-480. doi: https://doi.org/10.1086/507132

15. Tezze C, Romanello V, Desbats MA, et al. Age-Associated Loss of OPA1 in Muscle Impacts Muscle Mass, Metabolic Homeostasis, Systemic Inflammation, and Epithelial Senescence. *Cell Metab.* 2017;25(6):1374-1389. doi: https://doi.org/10.1016/j.cmet.2017.04.021

16. Phillips BE, Williams JP, Gustafsson T, et al. Molecular networks of human muscle adaptation to exerccise and age. *PLoS Genet*. 2013;9(3):e1003389. doi: https://doi.org/10.1371/journal.pgen.1003389

17. Distefano G, Standley RA, Dube JJ, et al. Chronological Age Does not Influence Ex-vivo Mitochondrial Respiration and Quality Control in Skeletal Muscle. *J Gerontol A Biol Sci Med Sci.* 2017;72(4):535-542. doi: https://doi.org/10.1093/gerona/glw102

18. Balan E, Schwalm C, Naslain D, et al. Regular endurance exercise promotes fission, mitophagy, and oxidative phosphorylation in human skeletal msucle independetly of age. *Front Physiol.* 2019;10:1088. doi: https://doi.org/10.3389/fphys.2019.01088

19. Seo DY, Lee SR, Kim N, et al. Age-related changes in skeletal muscle mitochondria: the role of exercise. *Integr Med Res*. 2016;5(3):182-186. doi: https://doi.org/10.1016/j.imr.2016.07.003

20. Broskey NT, Boss A, Fares EJ, et al. Exercise efficiency relates with mitochondrial content and function in older adults. *Physiol Rep.* 2015;3(6). pii. e12418. doi: https://doi.org/10.14814/phy2.12418

21. Broskey NT, Greggio C, Boss A, et al. Skeletal muscle mitochondria in the elderly: effects of physical fitness and exercise training. *J Clin Endocr Metab*. 2014;99(5):1852-1861. doi: https://doi.org/10.1210/jc.2013-3983

22. Piasecki J, Ireland A, Piasecki M, et al. Comparison od muscle function, bone mineral density and body composition of early starting and later staring older masters athletes. *Front Physiol*. 2019;10:1050. doi: doi: 10.3389/fphys.2019.01050

23. Pollock RD, Carter S, Velloso CP, et al. (2015) An investigation into the relationship between age and physiological function in highly active older adults. *J Physiol*. 2015;593(3):657-680. doi: https://doi.org/10.1113/jphysiol.2014.282863

24. Dietrichson P, Coakley J, Smith PE. et al. Conchotome and needle percutaneous biopsy of skeletal muscle. *J Neurol Neurosur PS*. 1987;50:1461-1467. doi: https://doi.org/10.1136/jnnp.50.11.1461

25. Vandesompele J, De Preter K, Pattyn F, et al. Accurate normalization of real-time quantitative R-PCR data by geometric averaging of multiple internal control genes. *Genome Biol.* 2002;3(7): research0034. doi: https://doi.org/10.1186/gb-2002-3-7-research0034

26. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods.* 2001;25(4):402-408. doi: https://doi.org/10.1006/meth.2001.1262

27. Ghosh S, Lertwattanarak R, Lefort N, et al. Reduction in reactive oxygen species production by mitochondria from elderly subjects with normal and impaired glucose tolerance. *Diabetes*. 2011;60(8):2051-2060. doi: https://doi.org/10.2337/db11-0121

28. Joseph AM, Adhihetty PJ, Buford TW, et al. The impact of aging on mitochondrial function and biogenesis pathways in skeletal muscle of sedentary high- and low-functioning elderly individuals. *Aging Cell*. 2012;11(5):801-809. doi: https://doi.org/10.1111/j.1474-9726.2012.00844.x

29. Larsen S, Nielsen J, Hansen CN, et al. Biomarkers of mitochondiral conent in skeletal muscle of healthy young human subjects. *J Physiol.* 2012b;590(14):3349-3360. doi: https://doi.org/10.1113/jphysiol.2012.230185

30. Gibala MJ, Little JP, MacDonald MJ, Hawley JA. Physiological adaptations to low-volume, highintensity interval training in health and disease. *J Physiol.* 2012;590(5):1077-1084. doi: https://doi.org/10.1113/jphysiol.2011.224725

31. Kong X, Wang R, Xue Y, et al. Sirtuin 3, a new target of PGC-1 α , plays an important role in the suppression of ROS and mitochondrial biogenesis. *PLoS One.* 2010;5(7):e11707. doi: https://doi.org/10.1371/journal.pone.0011707

32. Kincaid B, Bossy-Wetzel E. Forever young: SIRT3 a shield against mitochondrial meltdown, aging, and neurodegeneration. *Front Aging Neurosci.* 2013;6(5):48 doi: https://doi.org/10.3389/fnagi.2013.00048

33. Lanza IR, Short DK, Short KR, et al. Endurance exercise as a countermeasure for aging. *Diabetes*. 2008;57(11):2933-2942. doi: https://doi.org/10.2337/db12-er10

34. Jing E, O'Neill BT, Rardin MJ, et al. Sirt3 regulates metabolic flexibility of skeletal muscle through reversible enzymatic deacetylation. *Diabetes*. 2013;2(10):3404-3417. doi: https://doi.org/10.2337/db12-1650

35. Matoba S, Kang JG, Patino WD, et al. p53 regulates mitochondrial respiration. *Science*. 2006;312(5780):1650-1653. doi: https://doi.org/10.1126/science.1126863

36. Marzetti E, Leeuwenburgh C. Skeletal muscle apoptosis, sarcopenia and frailty at old age. *Exp Gerontol*. 2006; 41(12):1234-128. doi: https://doi.org/10.1016/j.exger.2006.08.011

37. Gustafsson T, Bodin K, Sylven C, et al. Increased expression of VEGF following exercise training in patients with heart failure. *Eur J Clin Invest.* 2001;31(4):362-366. doi: https://doi.org/10.1046/j.1365-2362.2001.00816.x

38. Leenders M, Verdijk LB, van der Hoeven L, et al. Elderly men and women benefit equally from prolonged resistance-type exercise training. *J Gerentol A Biol Sci Med Sci.* 2013;68(7):769-779. doi: https://doi.org/10.1093/gerona/gls241

39. Melov S, Tarnopolsky MA, Beckman K, et al. Resistance exercise reverses aging in human skeletal muscle. *PLoSOne*. 2007;2(5):e465. doi: https://doi.org/10.1371/journal.pone.0000465

40. Suetta C, Aagaard P, Rosted A, et al. Training-induced changes in muscle CSA, muscle strength, EMG, and rate of force development in elderly subjects after long-term unilateral disuse. *J Appl Physiol* (1985). 2004;97(5):1954-1961. doi: https://doi.org/10.1152/japplphysiol.01307.2003

41. Verdijk LB, Gleeson BG, Jonkers RA, et al. Skeletal muscle hypertrophy following resistance training is accompanied by a fiber type-specific increase in satellite cell content in elderly men. *J Gerentol A Biol Sci Med Sci.* 2009;64(3):332-339. doi: https://doi.org/10.1093/gerona/gln050

42. Charifi N, Kadi F, Féasson L, Denis C. Effects of endurance training on satellite cell frequency in skeletal muscle of old men. *Muscle Nerve*. 2003;28(1):87-92. doi: https://doi.org/10.1002/mus.10394

43. Coggan AR, Spina RJ, King DS, et al. Skeletal muscle adaptations to endurance training in 60- to 70-yr-old men and women. *J Appl Physiol*.1992;72(5):1780-1786. doi: https://doi.org/10.1152/jappl.1992.72.5.1780

44. Verney J, Kadi F, Charifi N, et al. Effects of combined lower body endurance and upper body resistance training on the satellite cell pool in elderly subjects. *Muscle Nerve*. 2008;38(3):1147-1154. doi: https://doi.org/10.1002/mus.21054

45. Aagaard P, Magnusson PS, Larsson B, et al. Mechanical muscle function, morphology, and fiber type in lifelong trained elderly. *Med Sci Sport Exer.* 2007;39(11):1989-1996. doi: https://doi.org/10.1249/mss.0b013e31814fb402

46. DiPietro L. Physical Activity in Aging: Changes in Patterns and Their Relationship to Health and Function. *J Gerontol A Biol Sci Med Sci.* 2001;2:13-22. doi: https://doi.org/10.1093/gerona/56.suppl 2.13

Figure Captions

Fig. 1 Mitochondrial related gene expression is greater in highly active older individuals. mRNA of genes related to mitochondrial content in recreationally active young (YG n=6; black bars), old individuals who were healthy, but did not undertake regular exercise (OG n=8; grey bars), older individuals who had been highly active through cycling exercise for many years (OT n=7; open bars). Data are presented as a fold change from YG. (a) sirtuin 3 (SIRT3) (b) peroxisome proliferator-activated receptor gamma, coactivator 1 alpha (PGC-1 α) (c) citrate synthase (CS) (d) ubiquinol-cytochrome c reductase core protein I (UQCRC1) (e) ubiquinol-cytochrome c reductase core protein I (UQCRC2) (f) cytochrome c oxidase subunit IV isoform 1 (COX412) (g) cytochrome c oxidase assembly homolog 10 (COX10) (h) succinate dehydrogenase complex, subunit A, flavoprotein (SDHA) (i) succinate dehydrogenase complex, subunit B, iron sulfur (SDHB) (j) translocase of inner mitochondrial membrane 8 homolog A (TIMM8A). ^a significantly different from YG, ^b significantly different from OG.

Fig. 2 Skeletal muscle related gene expression is greater in highly active older individuals. mRNA expression of genes related to skeletal muscle remodelling in recreationally active young (YG n=6; black bars), old individuals who were healthy, but did not undertake regular exercise (OG n=8; grey bars), older individuals who had been highly active through cycling exercise for many years (OT n=7; open bars). Data are presented as a fold change from YG. (a) vascular endothelial growth factor A (VEGFA) (b) vascular endothelial growth factor B (VEGFB) (c) myogenic differentiation 1 (MyoD1) (d) BTG family, member 2 (BTG2) (e) mitofusin 1 (MFN1) (f) mitofusin 2 (MFN2) (g) sestrin 2 (SESN2) (h) phorbol-12-myristate-13-acetate-induced protein 1 (PMAIP1) (i) zinc finger, matrin-type 3 (ZMAT3).

Fig. 3 Substrate metabolism related gene expression is greater in highly active older individuals. mRNA expression of genes related to substrate metabolism in recreationally active young (YG n=6; black bars), old individuals who were healthy, but did not undertake regular exercise (OG n=8; grey bars), older individuals who had been highly active through cycling exercise for many years (OT n=7; open bars). Data are presented as a fold change from YG. (a) isocitrate dehydrogenase 2 (NADP+) (IDH2) (b) malate dehydrogenase 1, NAD (MDH1) (c) pyruvate dehydrogenase kinase, isozyme 4 (PDK4) (d) nuclear receptor subfamily 1, group D, member 2 (NR1D2). ^a significantly different from YG QG, ^b significantly different from OG.

Fig. 4 Mitochondrial related protein content is greater in highly active older individuals. Protein content in recreationally active young (YG n=9; black bars), old individuals who were healthy, but did not undertake regular exercise (OG n=8; grey bars), older individuals who had been highly active through cycling exercise for many years (OT n=8; open bars). (a) Protein content of mitochondrial enzymes (complexes I-V), (b) peroxisome proliferator-activated receptor gamma, coactivator 1 alpha (PGC-1 α), (c) citrate synthase (CS) and (d) sirtuin 3 (SIRT3). (e) Representative images of protein content of mitochondrial enzymes (complexes I-V), PGC-1 α , CS and SIRT3 in YG, OG and OT. All values were normalized to their respective ponceau stain. Data are presented as mean ± standard deviation. ^b significantly different from OG and YG

Fig. 5 Substrate metabolism related protein content is greater in highly active older individuals. Protein content in recreationally active young (YG n=9; black bars), old individuals who were healthy, but did not undertake regular exercise (OG n=8; grey bars), older individuals who had been highly active through cycling exercise for many years (OT n=8; open bars). (a) protein content of pyruvate dehydrogenase (PDH), (b) long chain acyl CoA dehydrogenase (LCAD), (c) medium chain acyl-CoA dehydrogenase (MCAD), and (d) very long chain acyl CoA dehydrogenase (VLCAD). (e) Representative images of protein content of PDH, LCAD, MCAD and VLCAD. All values were normalized to their respective ponceau stain. Data are presented as mean ± standard deviation. ^a significantly different from OG, ^b significantly different from OG and YG

eTable 1. mRNA expression of genes included on the designed RT2 Profiler PCR Array.

Name	Abbreviation	Refseq #	YG	OG	ОТ
sirtuin 1	SIRT1	NM_001142498	1.00	1.11	1.13
sirtuin 3	SIRT3	NM_001017524	1.00	1.04	1.53 (c)
sirtuin 6	SIRT6	NM_001193285	1.00	1.04	1.36
transcription factor A,	TFAM	NM_003201	1.00	0.85	1.05
transcription factor B1, mitochondrial	TFB1M	NM_016020	1.00	0.96	1.22
transcription factor B2, mitochondrial	TFB2M	NM_022366	1.00	1.15	1.19
vascular endothelial growth factor A	VEGFA	NM_001025366	1.00	1.08	2.04 (b)
vascular endothelial growth factor B	VEGFB	NM_001243733	1.00	0.92	1.32 (a)
vascular endothelial growth factor C	VEGFC	NM_005429	1.00	1.00	1.64
BCL2-associated agonist of cell death	BAD	NM_004322, NM 032989	1.00	0.81	0.99
BCL2-antagonist/killer 1	BAK1	NM_001188	1.00	0.97	1.13
BCL2-associated X protein	BAX	NM_004324, NM_138761, NM_138763, NM_138764, NM_138765, NR_027882	1.00	1.03	1.12
BCL2 binding component 3	BBC3	 NM_001127240	1.00	1.36	0.98
BTG family, member 2	BTG2	NM_006763	1.00	2.27 (d)	0.73
caspase 9, apoptosis- related cysteine peptidase	CASP9	NM_001229	1.00	1.14	0.98
Fas cell surface death receptor	FAS	NM_000043	1.00	0.81	0.62
phorbol-12-myristate-13- acetate-induced protein 1	PMAIP1	NM_021127	1.00	1.66	3.30 (b)
zinc finger, matrin-type 3	ZMAT3	NM_152240	1.00	1.28 (c)	1.30 (c)
breast cancer 1, early onset	BRCA1	NM_007294	1.00	1.02	1.25
epidermal growth factor receptor	EGFR	NM_005228	1.00	1.00	0.83
growth arrest and DNA- damage-inducible, alpha	GADD45A	NM_001199741	1.00	0.87	1.01
proliferating cell nuclear antigen	PCNA	NM_002592	1.00	0.99	1.05
pituitary tumor- transforming 1	PTTG1	NM_004219	1.00	1.55	1.41
retinoblastoma 1	RB1	NM_000321	1.00	0.95	0.98

sestrin 2	SESN2	NM_031459	1.00	1.42	2.09 (c)
beclin 1, autophagy	BECN1	NM_003766	1.00	0.93	1.00
related					
Nome	Abbrovistion	Defees #	VC	00	OT
Name	Abbreviation	Refseq #	YG	UG	01
BH3 interacting domain death agonist	BID	NM_001196, NM_001244567	1.00	1.01	1.15
acam agemet		NM 001244569,			
		NM_001244570,			
		NM_001244572,			
		NM_197966,			
BCI 2/adenovirus E1B	BNIP3	NM 004052	1.00	0.80	0.91
19kDa interacting protein	DIVILO	1111_004002	1.00	0.00	0.01
3					
caspase 3, apoptosis-	CASP3	NM_004346,	1.00	0.84	1.01
related cysteine		NM_032991			
peptidase			1.00	0.07	1 15
related cysteine	CASPO	NIVI_001060124	1.00	0.97	1.15
peptidase					
v-myc avian	MYC	NM_002467	1.00	1.03	0.78
myelocytomatosis viral					
oncogene homolog	001/40.0		4.00	0.75	4.05
cytochrome c oxidase	COX10 §	NM_001303	1.00	0.75	1.05
(veast)					
COX18 cytochrome C	COX18	NM 001033760	1.00	0.99	1.41
oxidase assembly factor		_			
fission 1 (mitochondrial	FIS1	NM_016068	1.00	0.86	1.03
outer membrane)					
mitofusin 1	MEN1	NM 017927	1 00	0.88	1 12 (a)
mitofusin 1			1.00	0.00	1.12 (a)
		NM_001127660	1.00	0.97	1.46 (a)
optic atrophy 1 (autosomal dominant)	OPA1	NM_015560	1.00	0.96	1.16
translocase of outer	TOMM20	NM 014765	1.00	0.89	0.92
mitochondrial membrane				0.00	0.01
20 homolog (yeast)					
translocase of outer	TOMM22	NM_020243	1.00	0.92	0.99
mitochondrial membrane					
translocase of inner	TIMM8A	NM 001145951	1 00	0.82	1 55 (a)
mitochondrial membrane			1.00	0.02	1.00 (u)
8 homolog A (yeast)					
translocase of inner	TIMM9	NM_012460	1.00	0.73	0.96
mitochondrial membrane					
9 nomolog (yeast)		NM 012261	1.00	0.72	1 42 (2)
activated receptor	PGC-Iu	INIVI_013201	1.00	0.75	1.42 (a)
gamma, coactivator 1					
alpha					
peroxisome proliferator-	PPARGC1B	NM_001172699,	1.00	0.78	1.30
activated receptor		NM_133263			
beta					
citrate synthase	CS	NM_004077	1.00	0.77 (d)	1.28

hexokinase 2	HK2	NM_000189	1.00	1.53	2.50
pyruvate dehydrogenase kinase, isozyme 4	PDK4	NM_002612	1.00	2.88	0.43 (a)
solute carrier family 2 (facilitated glucose	SLC2A4	NM_001042	1.00	0.94	1.22
Name	Abbreviation	Refseq #	YG	OG	ОТ
myogenic differentiation 1	MYOD1	NM_002478	1.00	0.82	1.73 (b)
histone deacetylase 3	HDAC3	NM_003883	1.00	0.99	0.96
histone deacetylase 5	HDAC5	NM_001015053	1.00	1.22	1.19
peroxisome proliferator- activated receptor alpha	PPARA	NM_001001928	1.00	1.06	1.29
peroxisome proliferator- activated receptor delta	PPARD	NM_001171818	1.00	0.70	0.86
NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 1, 7.5kDa	NDUFA1	NM_004541	1.00	1.10	1.10
NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 1, 7kDa	NDUFB1	NM_004545	1.00	0.88	1.16
NADH dehydrogenase (ubiquinone) 1, subcomplex unknown, 1, 6kDa	NDUFC1	NM_001184986	1.00	1.03	1.12
succinate dehydrogenase complex, subunit A, flavoprotein (Fp)	SDHA	NM_004168	1.00	0.98	1.42 (b)
succinate dehydrogenase complex, subunit B, iron sulfur (Ip)	SDHB	NM_003000	1.00	0.91	1.34 (a)
cytochrome c-1	CYC1	NM_001916	1.00	0.89	1.28
ubiquinol-cytochrome c reductase core protein l	UQCRC1	NM_003365	1.00	0.79 (c)	1.34 (b)
ubiquinol-cytochrome c reductase core protein II	UQCRC2	NM_003366	1.00	0.87	1.28 (a)
cytochrome c oxidase subunit IV isoform 1	COX4I1	NM_001861	1.00	0.85	1.23
cytochrome c oxidase subunit IV isoform 2 (lung)	COX4I2	NM_032609	1.00	1.00	1.54 (b)
ATP synthase, H+ transporting, mitochondrial F1 complex, alpha subunit 1, cardiac muscle	ATP5A1	NM_001001935	1.00	0.86	1.18
ATPase, H+ transporting, lysosomal V0 subunit a2	ATP6V0A2	NM_012463	1.00	1.04	1.08
isocitrate dehydrogenase 1 (NADP+), soluble	IDH1	NM_005896	1.00	0.96	1.07
isocitrate dehydrogenase 2 (NADP+), mitochondrial	IDH2	NM_002168	1.00	0.90	1.34 (a)
malate dehydrogenase 1, NAD (soluble)	MDH1§	NM_001199111	1.00	1.04	1.67
malate dehydrogenase 2, NAD (mitochondrial)	MDH2	NM_005918	1.00	0.79	0.98
poly (ADP-ribose) polymerase 1	PARP1	NM_001618	1.00	0.96	0.97

poly (ADP-ribose)	PARP2	NM_001042618	1.00	0.86	1.05
polymerase 2					
K(lysine)	KAT2A	NM_021078	1.00	1.11	1.12
acetyltransferase 2A					
K(lysine)	KAT2B	NM_003884	1.00	1.03	0.90
acetyltransferase 2B					
Name	Abbreviation	Refseq #	YG	OG	ОТ
nuclear receptor	NCOA1	NM_003743	1.00	1.09	1.04
coactivator 1					
nuclear receptor	NCOA3	NM_001174087	1.00	0.96	0.92
coactivator 3					
nuclear receptor	NR1D2	NM_001145425	1.00	0.99	0.66 (b)
subfamily 1, group D,					
member 2					
estrogen receptor 1	ESR1	NM_000125	1.00	0.71	0.82
estrogen receptor 2 (ER	ESR2	NM_001040275	1.00	2.28	2.54
beta)					
E1A binding protein p300	EP300	NM_001429	1.00	0.96	0.94
CREB binding protein	CREBBP	NM_001079846	1.00	0.96	0.74
cyclin-dependent kinase	CDKN1A	NM_000389	1.00	2.34	1.62
inhibitor 1A (p21, Cip1)					
tumor protein p53	TP53	NM_000546	1.00	1.03	1.12

Note: YG = young; OG = old; OT = old highly active. Data are expressed as fold change from the YG group. Full names and abbreviations of each gene are included along with the manufacturers Refseq #.

(a) significantly different from OG, (b) significantly different from OG and YG, (c) significantly different from YG, (d) significantly different from YG and OT, §main effect.



FIG 1





Figure 2

















