


Please cite the Published Version

Mazuquin, Bruno , Sunemi, Mariana Maia de Oliveira, e Silva, Marcela Ponzio Pinto, Sarian, Luís Otávio Zanatta, Williamson, Esther and Bruce, Julie (2021) Current physical therapy care of patients undergoing breast reconstruction for breast cancer: a survey of practice in the United Kingdom and Brazil. *Brazilian Journal of Physical Therapy*, 25 (2). pp. 175-185. ISSN 1413-3555

DOI: <https://doi.org/10.1016/j.bjpt.2020.05.010>

Publisher: Elsevier BV

Version: Accepted Version

Downloaded from: <https://e-space.mmu.ac.uk/626525/>

Usage rights:  [Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0](https://creativecommons.org/licenses/by-nc-nd/4.0/)

Additional Information: This is an Author Accepted Manuscript of a paper accepted for publication in *Brazilian Journal of Physical Therapy*, published by and copyright Elsevier.

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

1 **Current physical therapy practice of the care of patients undergoing breast**
2 **reconstruction for breast cancer: a survey in the United Kingdom and**
3 **Brazil**

4 **Running Title: Physical therapy for breast reconstruction for breast cancer**

5
6 Bruno Mazuquin¹, Mariana Maia de Oliveira Sunemi², Marcela Ponzio Pinto e Silva², Luís Otávio
7 Zanatta Sarian², Esther Williamson³, Julie Bruce¹

8 ¹ Warwick Clinical Trials Unit, Warwick Medical School, University of Warwick, UK.

9 ² Centre for Integral Attention to Women's Health, Universidade Estadual de Campinas, Campinas,
10 SP, Brazil

11 ³ Nuffield Department of Orthopedics Rheumatology and Musculoskeletal Sciences, University of
12 Oxford, UK.

13

14 **Declarations of interest:** None

15

16 **Contact author:** Bruno Mazuquin, bruno.mazuquin@warwick.ac.uk, Clinical Trials Unit, Medical
17 School, University of Warwick, Gibbet Hill Road, CV4 7AL, Coventry, UK

18

19

20 **ABSTRACT**

21 **Background:** In both the United Kingdom (UK) and Brazil, women undergoing mastectomy should
22 be offered breast reconstruction. Patients may benefit from physical therapy to prevent and treat
23 muscular deficits. However, there are uncertainties regarding which physical therapy program to
24 recommend.

25 **Objective:** The aim was to investigate the clinical practice of physical therapists for patients
26 undergoing breast reconstruction for breast cancer. A secondary aim was to compare physical therapy
27 practice across UK and Brazil.

1 **Methods:** We undertook an online survey with physical therapists in both countries. We asked about
2 physical therapists' clinical practice.

3 **Results:** 181 physical therapists completed the survey, the majority were from Brazil (77%).
4 Respondents reported that only half of women having breast reconstruction were routinely referred to
5 physical therapy postoperatively. Contact with patients varied widely between countries, the mean
6 number of postoperative sessions was 5.7 in the UK and 15.1 in Brazil. The exercise programs were
7 similar for different reconstruction operations. Therapists described a progressive loading structure
8 over time: range of motion (ROM) was restricted to 90° of arm elevation in the first two postoperative
9 weeks; by 2-4 weeks ROM was unrestricted; at 1-3 months muscle strengthening initiated, and after
10 three months the focus was on sports-specific activities.

11 **Conclusion:** Only half of patients having a breast reconstruction are routinely referred to physical
12 therapy. Patients in Brazil have more intensive follow-up, with up to three times more face-to-face
13 contact with a physical therapist than in the UK. Current practice broadly follows programs for
14 mastectomy care rather than being specific to reconstruction surgery.

15

16

17 **Keywords:** Breast cancer; Breast reconstruction; Exercise; Physical therapy; Survey.

18

19 **Highlights**

- 20
- Patients having breast reconstruction are not routinely referred to physical therapy
 - The main reason for referral to physical therapy is complication after surgery
 - Current practice does not consider limitations specific to each reconstruction type
 - Patients in Brazil have three times more treatment sessions compared to those in the UK
- 21
- 22
- 23

24

25

1 INTRODUCTION

2 Breast cancer is the most common type of cancer in women with 55,439 and 85,620 new cases
3 diagnosed in 2018 in the UK and in Brazil, respectively.¹ However, patients are now living longer;
4 five-year survival has improved to 86% in the UK and 75% in Brazil.² Patients treated for breast
5 cancer can experience a long, complex, and distressing healthcare journey, which may include
6 surgery, chemotherapy, radiotherapy, immunotherapy, and endocrine treatment.³ These treatment can
7 cause long-term problems; approximately 35% of patients report pain and limitations of shoulder and
8 arm function more than three years after their breast cancer treatment.⁴ Almost all women will require
9 surgical treatment and up to 40% undergo mastectomy.⁵ Mastectomy can negatively affect body
10 image, self-esteem, and health-related quality of life (QoL).⁶ Several studies have demonstrated that
11 breast reconstruction may improve patients' psychological and emotional wellbeing.^{6,7} Current UK
12 guidance recommends that all women undergoing mastectomy should be offered either immediate or
13 delayed breast reconstruction.⁸ Since 2013, patients with breast cancer in Brazil have the right to
14 request breast reconstruction in the public health service.⁹ The rates of breast reconstruction are
15 increasing yearly with a quarter of women in the UK now electing to undergo immediate
16 reconstruction. In Brazil, there has been a 58% increase, from 2008 to 2014, in the number of breast
17 reconstructions performed in the public health service.¹⁰

18 Reconstruction surgery is complex and different approaches are used; techniques can broadly be
19 divided into procedures using implants or autologous tissue.¹¹ Each type of breast reconstruction can
20 affect postoperative function differently. For instance, women undergoing latissimus dorsi
21 reconstruction are at greater risk of shoulder range of motion (ROM) deficits; with up to 73% of
22 patients reporting postoperative difficulties with daily activities involving arm movement.¹² Implant-
23 based reconstruction may impair pectoralis muscle strength¹³ and abdominal flaps may reduce trunk
24 muscle strength by 23%.¹⁴ There is some evidence to suggest that newer muscle sparing approaches
25 using perforators flaps (e.g. deep or superficial inferior epigastric artery flaps) may result in better
26 functional outcomes.^{11,15} The most common type of reconstruction in the UK involves the use of
27 implants (36%) for immediate reconstruction and autologous tissue (32%) for delayed
28 reconstructions.¹⁶ In Brazil, the most common method of reconstruction involves autologous tissue
29 (66%).¹⁰

30 Physical therapy and structured home-based exercise programs may help to prevent pain and
31 morbidities related to the arm, shoulder, and other joints and improve pain.^{17,18} However, there is still
32 uncertainty regarding the optimal content and timing of exercise prescription after reconstruction.
33 Physical therapy may also support patients in meeting the minimum amount of physical activity
34 recommended per week. A systematic review (n=6 studies, 1607 participants) found that two years

1 post-diagnosis, most women (91%) did not meet the recommended guidelines for weekly physical
2 activity.¹⁹

3 The current guidelines of the UK-based Association of Breast Surgery (ABS) and the British
4 Association of Plastic Reconstructive and Aesthetic Surgeons (BAPRAS) state that all patients should
5 have early access to specialist physical therapy, including pre-operatively, to prevent and treat upper
6 limb morbidities, particularly when extensive reconstruction surgery is required.²⁰ However, these
7 societies also acknowledge that there is very limited evidence for physical therapy after reconstruction
8 surgery and therefore current guidelines and recommendations are largely based on expert opinion
9 and clinical experience.²⁰

10 In Brazil, the national policy for cancer care states that high-complexity cancer care units must have
11 multidisciplinary teams, which may involve physical therapists.²¹ In 2003, the Brazilian Ministry of
12 Health published a consensus document called Breast Cancer Control. The document brings
13 information on prevention, early diagnosis, and treatment. It recommends that physical therapy should
14 be offered before surgery to identify risk factors for postoperative complications. Physical therapy
15 should then continue in the immediate post-operative stage and during adjuvant therapy to identify,
16 prevent, and mitigate problems such as acute and chronic pain, lymphoedema, functional impairment,
17 and respiratory complications.²² It also suggests that home-based exercises and self-massage should
18 be used to control pain. However, there is no specific advice for the care of patients undergoing breast
19 reconstruction, and similar to the UK guidelines, these recommendations were based on expert
20 opinion only.²² The lack of information about current physical therapy care after breast reconstruction
21 may contribute to inconsistency in patterns of care within and across different healthcare systems.

22 Given the lack of evidence, we wanted to identify whether a) patients undergoing breast
23 reconstruction are routinely referred to physical therapy; b) the content, timing of delivery, and
24 structure of physical therapy-led rehabilitation programs; c) the setting and format of current physical
25 therapy care; and d) recommendations for exercise progression over the postoperative period.
26 Therefore, the aim of this cross-sectional study was to investigate the characteristics and content of
27 physical therapy-led rehabilitation programs delivered to patients undergoing breast reconstruction. A
28 secondary objective was to compare the characteristics of physical therapy care in two healthcare
29 settings, the UK and Brazil. Both countries offer universal health coverage, have a free-at-point of
30 care national health service (NHS), and recommend that breast reconstruction should be offered to
31 women undergoing mastectomy.

32

33

34

1 **METHOD**

2 We carried out an open, voluntary online survey in the UK (September – October 2018) and in Brazil
3 (January – March 2019). The survey was developed in three steps: first, the authors reviewed the
4 literature and consulted with at least one specialist cancer physical therapist from each nation to
5 identify the key elements included in standard perioperative care. Based on this initial step, a first
6 draft of the questionnaire was developed. Second, the survey was piloted with a small sample of
7 clinicians involved with the care of this patient population. Adaptations were made based on clinician
8 feedback. The third stage was another pilot phase to test the online platform and layout; four physical
9 therapists were involved across multiple development stages. Using their feedback, we then created
10 the final version of the questionnaire. This study was given **ethical** approval in the UK by the
11 University of Warwick, Coventry (REGO-2018-2217) and by the Universidade de Campinas,
12 Campinas, São Paulo, Brazil (03872318.3.0000.5404). Before starting the survey, participants
13 consented to take part in the study by ticking a box. We followed the Checklist for Reporting Results
14 of Internet E-Surveys (CHERRIES) to report this survey.²³


15

16 Two versions of the questionnaires (English and Portuguese, available at
17 <https://wrap.warwick.ac.uk/123565>) were produced and data were stored via the online platform
18 Qualtrics (Qualtrics, Provo, UT, USA). The final survey comprised 41 questions, across 21
19 screens/pages (one to two questions per page), focusing on physical therapists' clinical practice for the
20 management of patients undergoing breast reconstruction. Questions and items followed the same
21 order for every participant. We did not randomize or alternate items nor did we record the time taken
22 to complete the survey. We included three example clinical cases to standardize the context for
23 responses for three commonly used reconstruction procedures. Physical therapists were invited to
24 describe the routine care of women undergoing either silicone implant, latissimus dorsi (LD) or deep
25 inferior epigastric artery (DIEP) flaps. Each case was presented as follows:

26 *“A woman, 55 years, had a unilateral breast (dominant side) reconstruction using a _____. Her*
27 *overall health is good and she had no postoperative complications.”*

28 Therapists working across both public and private sectors could respond separately where usual care
29 pathways differed. Before submitting their answers, participants could review their responses by using
30 the ‘left arrow/back’ button. Given that no identifiable data could be collected, it was not possible to
31 check for duplicate submissions. However, if the participant decided to start the survey but complete
32 at a later date, the online platform would automatically take the participant back to the relevant
33 section. The online platform had the option of a Survey ID cookie to minimize duplications. We
34 manually checked for completeness once questionnaires were submitted.

1 **Sampling**

2 Given the difficulties in determining the sampling frame of total number of physical therapists
3 involved with the care of patients with breast cancer across both countries, it was not possible to
4 calculate a sample size. Hence, we used a convenience sampling. An invitation to complete the online
5 survey was posted on the UK Chartered Society of Physical therapy Oncology group website and
6 advertised on the news section of the UK Association of Chartered Physical therapists in Oncology
7 and Palliative Care website. In Brazil, it was posted on the Brazilian Society of Physical Therapy in
8 Women’s Health website. These websites are the main source of news related for physical therapists
9 working in oncology and palliative care. In addition to the websites, social media was used to
10 broaden the reach of the survey in both countries. No invitations were sent by standard post, we only
11 used online advertisement. The  used for advertising the survey is provided as Supplemental
12 material.

13 **Eligibility criteria**

14 To be eligible to complete the survey, participants had to be a registered physical therapist, currently
15 practicing either in the UK or in Brazil and be involved with the care for patients undergoing breast
16 reconstruction. The first section of the survey included a mandatory screening section to confirm
17 inclusion criteria and consent to take part in the study. Participants did not receive any incentives to
18 participate or complete the survey. At the end of the survey, participants had the option of
19 downloading a copy of their responses.

20 **Data Analysis**

21 Data were imported from Qualtrics into Excel (Microsoft Corp, Redmond, WA, USA) for descriptive
22 statistical analyses. No statistical methods were used to adjust for representativeness. Partial and
23 complete surveys were included in the final analysis with number of missing data reported. We used
24 radar graphs to illustrate findings for each of the three reconstruction methods (silicone implant, LD,
25 and DIEP). No Log file analysis was performed.

26

1 RESULTS

2 Response rates

3 A total of 265 (Brazil n=200; UK n=65) accesses to the questionnaire link were recorded; however,
4 181 entries were logged. The majority of respondents were from Brazil (139/181; 77%). In the UK, a
5 greater proportion of responding physical therapists worked in the public sector (30/42; 71%), while
6 in Brazil, responding therapists were more equally distributed between public (61/136; 45%) and
7 private sectors (71/136; 52%) (Table 1). We did not record the number of unique site visitors,
8 therefore, we could not compare survey view rates, participation, and completion rates.

9 Clinical practice characteristics

10 Most clinicians treated, on average, less than 20 patients with breast cancer per year (76/173; 44%)
11 although a quarter of respondents (44/173) treated more than 50 patients per annum (Table 2). Half of
12 responding physical therapists from each country stated that despite caring for patients with breast
13 cancer, fewer than half of their patients routinely had breast reconstruction. Half of therapists (80/164;
14 49%) responded that patients were routinely referred to physical therapy postoperatively but only 7%
15 (12/164) were routinely referred for care both pre and postoperatively. Approximately one third
16 (61/164; 37%) reported that patients were not routinely referred for physical therapy at all. The main
17 reasons for referral to physical therapy were for management of complications after surgery (53/116;
18 46%), followed by referrals for patients with a history of shoulder problems before having a breast
19 reconstruction (30/116; 26%).

20 In both countries, the most frequent method of breast reconstruction was implants, followed by LD
21 flaps (Table 3). We observed a marked difference in the mean number of face-to-face treatment
22 sessions between countries; in the UK, the mean number of sessions was 5.7 compared to 15.1 in
23 Brazil. Physical therapists from Brazil reported that the main barriers to providing adequate
24 postoperative care after breast reconstruction were the delayed start of physical therapy (85/271; 31%)
25 and limited patient finances (60/271; 22%). In the UK, physical therapists cited limited number of
26 appointments (13/55; 25%) and other factors (11/55; 21%), such as lack of referral to physical
27 therapy, patient fatigue, and postoperative complications.

28 Characteristics of physical therapy programs

29 Clinicians from both countries were very similar with regards to the types and format of exercises
30 prescribed at each postoperative phase, therefore, we present the combined data. The physical therapy
31 programs for the three reconstruction procedures are displayed in Figures 1-3. In the first two weeks
32 postoperatively, exercises for shoulder mobility were restricted to below 90° of arm elevation; posture
33 correction and manual therapy were the most frequent components covered in these early therapy
34 sessions. Between two to four weeks postoperatively, exercises for shoulder mobility above 90° and
35 shoulder-specific stretching were more common. From one to three months after surgery, therapists

1 would then prescribe general strengthening, shoulder-specific, and core-specific strengthening
2 exercises. After three months, the focus was on sport-specific activities with less attention to other
3 modalities.

4 5 **DISCUSSION**

6 This is the first survey to our knowledge to investigate the characteristics of physical therapy care
7 across the different postoperative stages after breast reconstruction for breast cancer. We explored
8 physical therapy care in two different countries where the public health service has a broadly similar
9 organizational structure, despite variation in underlying patient biosociodemographic characteristics
10 and size of the population covered (the UK-NHS serves 67 million people compared to 211 million in
11 Brazil.)²⁴

12 Overall, we found important gaps regarding physical therapy care, highlighted by the high number of
13 patients who are not routinely referred to physical therapy care after breast reconstruction surgery,
14 despite increasing numbers of women undergoing these procedures. Unlike mastectomy and breast
15 conserving surgery, breast reconstructions are associated with prolonged hospital stays, commonly up
16 to one week.²⁵ We found that patients were only referred for physical therapy if a surgical
17 complication had developed or if a history of shoulder problem was identified.

18 In the UK, the reason for the lack of referral to physical therapy may be due to clinical teams adhering
19 to the current National Institute for Health and Care Excellence (NICE) guidelines. These guidelines
20 recommend that patients with pre-existing shoulder conditions should be identified preoperatively to
21 inform treatment decisions; if these patients present with a persistent reduction in arm and shoulder
22 mobility after treatment, they should then be referred to physical therapy.⁸ However, Woo et al.²⁶
23 observed 420 patients following breast reconstruction over a mean follow-up of 52 months; they
24 identified various other risk factors for shoulder problems besides a preoperative history of shoulder
25 problems. These included reconstruction procedure, older age, and neoadjuvant chemotherapy. The
26 authors also reported that initiating physical therapy within two months of reconstruction surgery
27 reduced the risk of sustained shoulder morbidity in the longer term (odds ratio: 7.2, 95% CI: 1.4,
28 36.7).²⁶ Although UK health care teams may follow NICE recommendations, the advice for
29 postoperative care from both the ABS and BAPRAS contrast with those of NICE.²⁰ Their latest
30 guidelines state that all patients should have early access to physical therapy to prevent and treat
31 upper limb morbidities. Therefore, these conflicting guidelines lead to uncertainties and inequalities in
32 the care of patients with breast cancer. This was recently highlighted by a report from the UK All-
33 party Parliamentary Group Report.²⁷

1 In Brazil, the Comissão Nacional de Incorporação de Tecnologias (CONITEC) guidelines for breast
2 cancer treatment and management do not contain any specific recommendation or information
3 regarding physical therapy care.⁹ The consensus document from the Brazilian Ministry of Health
4 recommends that physical therapy should be routinely incorporated before and after breast cancer
5 surgery. However, there is no specific advice for the care of patients undergoing breast
6 reconstruction.²² There is a lack of information in the Brazilian guidelines, in addition to the limited
7 information regarding the number of physical therapists working with oncology patients in Brazil.
8 This may impact the number of referrals made to physical therapists by oncology teams and may
9 contribute to inequalities in patient care. Better integration of physical therapists within the
10 multidisciplinary oncology team and into the treatment pathway of patients having breast
11 reconstruction could improve awareness, increase referral rates, and potentially reduce complications
12 after surgery.²⁸

13 One of the main differences we found between the healthcare systems was the number of face-to-face
14 appointments offered to patients. Brazilian physical therapists have, on average, three times the
15 number of contacts with patients compared to those in the UK, even within the public sector. The
16 limited access to physical therapy sessions was highlighted by UK physical therapists as the main
17 barrier to care, which may be due to the increasing demand and pressure on the financially
18 constrained NHS^{29,30}. Physical therapy services in the UK are centrally funded by the Department of
19 Health. Thus, the proposed number of sessions is largely based on what is possible for delivery within
20 the NHS. There is currently limited evidence regarding what rehabilitation programs should offer,
21 and similarly, limited or no evidence to support the hypothesis that a higher number of sessions is
22 substantially more clinically and cost-effective long-term.^{31,32}

23 Although the number of sessions offered and available to patients is considerably higher in Brazil, the
24 main barriers to providing care included the delayed presentation to physical therapy and limited
25 ability to pay for private treatment. These barriers may be linked to the lack of physical therapists
26 working in primary care and outpatient settings in the Brazilian public sector.³³ According to Rodes et
27 al.³³, there is a higher concentration of physical therapists working in medium to high complexity
28 services (secondary care) and a shortage of professionals in primary care. Therefore, patients seek out
29 private treatment and thus finances may constrain access to appropriate physical therapy treatment.
30 Physical therapy in oncology is a relatively recent clinical specialty in Brazil; it was accredited in
31 2009 by The Brazilian Federal Council for Physical Therapy and Occupational Therapy (COFFITO).
32 Our data reflect the recent accreditation of this physical therapy specialty; less than half the
33 respondents from Brazil had fewer than five years of experience of treating patients with breast
34 cancer.³⁴

1 The physical therapy programs described by the survey participants were broadly similar for the
2 different reconstruction methods and followed similar principles of exercise progression for patients
3 undergoing mastectomy.^{35,36} For instance, therapists reported that they limited shoulder mobility
4 exercises to 90° or arm elevation in the first two weeks after surgery to avoid wound healing
5 complications while maintaining shoulder mobility.³⁶ Once wound healing had occurred, care from
6 weeks two to four postoperatively allowed unrestricted shoulder movement aiming to improve
7 shoulder ROM³⁵. By one month postoperatively, strengthening exercises were allowed and after three
8 months, sports activities were actively encouraged. This progressive approach is consistent with what
9 is reported in other studies.^{18,37-39} A progressive exercise protocol is advocated to be better than usual
10 care for improving function and pain at six months,³⁸ muscle strength at 12 months,³⁷ and does not
11 increase the risk of complications, such as lymphoedema at 12 months.³⁹ Although there are studies
12 suggesting the benefits of exercise for patients with breast cancer, these studies are generally of low
13 methodological quality and they do not include patients undergoing breast reconstruction.^{36,40}

14 The evidence for physical therapy following breast reconstruction is scarce; to our knowledge, there
15 are no high-quality systematic reviews or randomized controlled trials investigating the clinical and
16 cost-effectiveness of physical therapy for this patient population. A literature review from Teixeira
17 and Sandrin⁴¹ assessed physical therapy care following oncological breast reconstruction; however,
18 the review did not follow the PRISMA⁴² statement and was methodologically limited. The review did
19 not include a clear definition of their patient population, interventions, comparisons, and outcomes.
20 Furthermore, risk of bias of included studies was not undertaken.

21 One randomized controlled trial from Futter et al.⁴³ investigated the effect of pre-operative abdominal
22 strengthening to prevent abdominal complications in 93 women undergoing a DIEP flap. They found
23 that abdominal exercises had a positive impact on well-being before surgery; however, this trial also
24 lacked methodological rigor. Physical therapy following breast reconstruction should not only follow
25 the general principles of rehabilitation following mastectomy, clinicians must be aware of the
26 specificities of each reconstruction method to tailor exercises accordingly. Patients having an LD
27 reconstruction have a higher risk of developing shoulder problems than other methods of
28 reconstruction using implants or autologous tissue. Additional exercises may be needed for the
29 abdominal and back muscles after transverse rectus abdominis (TRAM) reconstruction.⁴⁴ Rindom et
30 al.⁴⁵ randomized 50 women to either a LD or a thoracodorsal artery perforator (TAP) flap
31 reconstruction; patients allocated to the TAP group showed better shoulder function at 12 months.
32 Woo et al.²⁶ found similar results with their cohort of 430 patients. Forty-three percent of patients who
33 had a LD reconstruction developed shoulder morbidity at four years post-surgery, compared to 23%
34 for expander-implant and 14% for DIEP. Further high-quality randomized clinical trials are needed to
35 assess the clinical and cost-effectiveness of physical therapy programs designed for each
36 reconstruction surgery.

1 **Limitations**

2 The limitations of our study include the low overall number of survey respondents and a discrepancy in
3 the proportion of respondents from each country. However, it is difficult to accurately estimate the
4 number of physical therapists who regularly treat patients with breast cancer in both countries. It is,
5 therefore, challenging to estimate sample representativeness with any certainty. Donnelly et al.⁴⁶
6 conducted a survey to investigate physical therapy management of cancer related-fatigue for various
7 types of cancer; the authors identified 102 physical therapists from the UK who stated that they would
8 use exercises for patients with breast cancer. Similarly, O'Hanlon and Kennedy⁴⁷ completed a survey
9 of the Irish members of the Chartered Physical therapists in Oncology and Palliative Care. This survey
10 investigated exercise prescription for cancer patients but the findings were based on only 35 responses.
11 We found no publications or examples of questionnaire surveys of physical therapists working within
12 oncology in Brazil. According to Matsumura et al.⁴⁸, there were 206,170 registered physical therapists
13 in Brazil in 2016, however, oncology rehabilitation is likely to be a smaller specialized subset of all
14 registrations. Another factor that may have affected the number of respondents was the length of the
15 survey, which may have impacted the response rate. Nevertheless, this is the first survey to our
16 knowledge to investigate practice of physical therapists caring for patients undergoing oncological
17 breast reconstruction. There is a need to design and test physical therapy programs for patients
18 undergoing breast reconstruction.

19

20 **CONCLUSION**

21 The majority of physical therapists caring for patients having a breast reconstruction treat a low
22 number of cases per year and overall referral to physical therapy services is low given the increasing
23 volume of breast reconstruction surgeries. Patients in Brazil are more likely to have a higher number
24 of sessions with therapists compared to the UK, with a three-fold difference in face-to-face
25 appointments. The most frequent method of breast reconstruction reported by physical therapists
26 across both countries was silicone implants. Current physical therapy programs follow the same
27 general principles of postoperative care after mastectomy.

Acknowledgements

We would like to thank Shelley Potter, Clare Lait and Catherine Hegarty for their feedback on early versions of the survey. We also would like to thank the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP process number 2018/06161-0).

Table 1. Clinical experience and training of responding physical therapists

	UK N (%)	Brazil N (%)	Total N (%)
Completed training in breast cancer care			
Yes	34 (80.9)	120 (86.4)	154 (85.1)
No	8 (19.1)	18 (12.9)	26 (14.4)
Missing	0 (0)	1 (0.7)	1 (0.5)
Total	42 (100)	139 (100)	181 (100)
Type of training*			
Continuous Professional Development events	28 (48.3)	143 (51.1)	171 (50.6)
Post-graduate modules	7 (12.0)	96 (34.3)	103 (30.6)
Post-graduate courses (MSc or PhD)	4 (6.9)	32 (11.4)	36 (10.6)
Other (short duration courses or in-house training)	19 (32.8)	9 (3.2)	28 (8.2)
Total	58 (100)	280 (100)	338 (100)
Experience of treating patients with breast cancer (years)			
< 5 years	11 (26.2)	60 (43.2)	71 (39.2)
Between 6 and 10 years	16 (38.1)	36 (25.9)	52 (28.7)
Between 11 and 15 years	7 (16.7)	12 (8.6)	19 (10.5)
>15 years	8 (19.0)	31 (22.3)	39 (21.6)
Missing	0 (0)	0 (0)	0 (0)
Total	42 (100)	139 (100)	181 (100)
Work sector			
Public service	30 (71.4)	61 (43.9)	91 (50.3)
Private practice	11 (26.2)	71 (51.1)	82 (45.3)
Both	1 (2.4)	4 (2.9)	5 (2.8)
Missing	0 (0)	3(2.1)	3 (1.6)
Total	42 (100)	139 (100)	181 (100)
Number of patients with breast cancer treated per year			
< 20	16 (38.0)	60 (43.2)	76 (42.0)
Between 20 and 50	12 (28.6)	41 (29.5)	53 (29.3)
>50	13 (31.0)	31 (22.3)	44 (24.3)
Missing	1 (2.4)	7 (5.0)	8 (4.4)
Total	42 (100)	139 (100)	181 (100)

Table 2. Clinical practice characteristics of responding physical therapists

	UK N (%)	Brazil N (%)	Total N (%)
Proportion of patients with breast cancer with reconstruction treated per year			
Less than half	19 (45.2)	64 (46.0)	83 (45.8)
Half	16 (38.1)	35 (25.2)	51 (28.2)
More than half	5 (12.0)	30 (21.6)	35 (19.3)
Missing	2 (4.7)	10 (7.2)	12 (6.7)
Total	42 (100)	139 (100)	181 (100)
Routine referral to physical therapy			
Yes, preoperatively only	2 (4.8)	0 (0)	2 (1.1)
Yes, pre and postoperatively	3 (7.1)	9 (6.5)	12 (6.6)
Yes, postoperatively only	14 (33.3)	66 (47.5)	80 (44.2)
No, not routinely seen	16 (38.0)	45 (32.4)	61 (33.7)
I don't know	5 (12.0)	4 (2.9)	9 (5.0)
Missing	2 (4.8)	15 (10.7)	17 (9.4)
Total	42 (100)	139 (100)	181 (100)
Reasons for referral to physical therapy (if not routinely seen by a physical therapist)*			
Shoulder problems before surgery	6 (24.0)	24 (26.4)	30 (25.9)
Complications after surgery	13 (52.0)	40 (44.0)	53 (45.7)
Other physical problems	3 (12.0)	19 (20.9)	22 (19.0)
Age	0 (0)	4 (4.4)	4 (3.4)
Other	3 (12.0)	4 (4.4)	7 (6.0)
Total	25 (100)	91 (100)	116 (100)
Average number of face-to-face appointments per patient			
Public; mean (SD)	4.3 ± 2.7	13.7 ± 8.1	-
Private; mean (SD)	7.2 ± 3.5	16.6 ± 8.7	-
Overall; mean (SD)	5.7 ± 3.1	15.1 ± 8.4	-

Continue

Table 2 (continued). Clinical practice characteristics of responding physical therapists

	UK N (%)	Brazil N (%)	Combined N (%)
Main barriers when caring for patients with breast reconstruction*			
Patients psychological health	9 (17.3)	22 (8.1)	31 (9.6)
Lack of patient compliance	7 (13.5)	51 (18.8)	58 (18.0)
Lack of time	2 (3.8)	10 (3.7)	12 (3.7)
Lack of training	4 (7.7)	23 (8.5)	27 (8.4)
Lack of research evidence	4 (7.7)	16 (5.9)	20 (6.2)
Limited number of appointments	13 (25.0)	0 (0)	13 (4.0)
Patient unable to pay for physical therapy	1 (1.9)	60 (22.1)	61 (18.9)
Delayed start of physical therapy	1 (1.9)	85 (31.4)	86 (26.6)
Other (e.g. lack of referral to physical therapy, patient fatigue, or postoperative complications)	11 (21.2)	4 (1.5)	15 (4.6)
Total	55 (100)	271 (100)	323 (100)

* Multiple responses could be selected.



Table 3. Frequency of types of breast reconstruction seen in clinic.

Frequency	Country	Reconstruction method n (%)						
		Implant	LD	TRAM	DIEP	SIEA	TUG or TMG	SGAP or IGAP
Do not see	UK	4 (9.5)	0 (0)	8 (19.0)	12 (28.5)	26 (61.9)	28 (66.6)	30 (71.4)
	BR	0 (0)	10 (7.2)	24 (17.2)	81 (58.3)	83 (59.7)	91 (65.5)	98 (70.5)
Rarely	UK	3 (7.1)	5 (11.9)	18 (42.9)	9 (21.4)	5 (11.9)	3 (7.1)	2 (4.7)
	BR	10 (7.2)	45 (32.3)	56 (40.3)	22 (15.9)	21 (15.1)	17 (12.2)	8 (5.8)
Sometimes	UK	18 (42.9)	18 (42.9)	5 (11.9)	6 (14.3)	2 (4.7)	1 (2.4)	1 (2.4)
	BR	39 (28.0)	42 (30.2)	20 (14.4)	6 (4.3)	2 (1.5)	1 (0.7)	1 (0.7)
Often	UK	4 (9.5)	6 (14.2)	2 (4.7)	6 (14.3)	0 (0)	1 (2.4)	0 (0)
	BR	45 (32.4)	17 (12.3)	11 (8.0)	0 (0)	1 (0.7)	0 (0)	1 (0.7)
Very often	UK	4 (9.5)	4 (9.5)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
	BR	25 (18.0)	1 (0.7)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Missing	UK	9 (21.5)	9 (21.5)	9 (21.5)	9 (21.5)	9 (21.5)	9 (21.5)	9 (21.5)
	BR	20 (14.4)	24 (17.3)	28 (20.1)	30 (21.5)	32 (23.0)	30 (21.6)	31 (22.3)
	UK total	42 (100)	42 (100)	42 (100)	42 (100)	42 (100)	42 (100)	42 (100)
	BR total	139 (100)	139 (100)	139 (100)	139 (100)	139 (100)	139 (100)	139 (100)

LD: latissimus dorsi, TRAM: transverse rectus abdominis, DIEP: deep inferior epigastric perforator, SIEA: superficial inferior epigastric artery, TUG: transverse upper gracilis, TMG: transverse musculocutaneous gracilis, SGAP: superior gluteal artery perforator, IGAP: inferior gluteal artery perforator.

Figure 1. Usual physical therapy program for a patient having a breast reconstruction with an implant.




Figure 2. Usual physical therapy program for a patient having a breast reconstruction with a latissimus dorsi flap.





Figure 3. Usual physical therapy program for a patient having a breast reconstruction with a deep inferior epigastric perforator flap.



REFERENCES

1. International agency for research on cancer. World Health Organization. Accessed 28 September, 2018.
2. Allemani C, Matsuda T, Di Carlo V, et al. Global surveillance of trends in cancer survival 2000-14 (CONCORD-3): analysis of individual records for 37 513 025 patients diagnosed with one of 18 cancers from 322 population-based registries in 71 countries. *Lancet*. 2018;391(10125):1023-1075. doi: 10.1016/S0140-6736(1017)33326-33323. Epub 32018 Jan 33331.
3. Freedman RA, Partridge AH. Management of breast cancer in very young women. *Breast*. 2013;22:S176-S179.
4. Feiten S, Dünnebacke J, Heymanns J, et al. Breast cancer morbidity: questionnaire survey of patients on the long term effects of disease and adjuvant therapy. *Dtsch Arztebl Int*. 2014;111(31-32):537-544.
5. Potter S, Conroy EJ, Cutress RI, et al. Short-term safety outcomes of mastectomy and immediate implant-based breast reconstruction with and without mesh (iBRA): a multicentre, prospective cohort study. *Lancet Oncol*. 2019;20(2):254-266.
6. Fang S-Y, Shu B-C, Chang Y-J. The effect of breast reconstruction surgery on body image among women after mastectomy: a meta-analysis. *Breast Cancer Research and Treatment*. 2013;137(1):13-21.
7. Chen W, Lv X, Xu X, Gao X, Wang B. Meta-analysis for psychological impact of breast reconstruction in patients with breast cancer. *Breast Cancer*. 2018;25(4):464-469. doi: 10.1007/s12282-018-0846-8.
8. *National Institute for Health and Care Excellence. Early and locally advanced breast cancer: diagnosis and management*. NG101. In: London: National Institute for Health and Care Excellence; 2018.
9. CONITEC. Diretrizes diagnósticas e terapêuticas - carcinoma de mama. In:2019.
10. Freitas-Júnior R, Gagliato DM, Moura Filho JWC, et al. Trends in breast cancer surgery at Brazil's public health system. *J Surg Oncol*. 2017;115(5):544-549. doi: 10.1002/jso.24572.
11. Mennie JC, Mohanna PN, O'Donoghue JM, Rainsbury R, Cromwell DA. National trends in immediate and delayed post-mastectomy reconstruction procedures in England: A seven-year population-based cohort study. *Eur J Surg Oncol*. 2017;43(1):52-61.
12. Nelson JA, Lee IT, Disa JJ. The Functional Impact of Breast Reconstruction: An Overview and Update. *Plast Reconstr Surg Glob Open*. 2018;6(3):e1640-e1640.
13. Hage JJ, van der Heeden JF, Lankhorst KM, et al. Impact of Combined Skin Sparing Mastectomy and Immediate Subpectoral Prosthetic Reconstruction on the Pectoralis Major Muscle Function: A Preoperative and Postoperative Comparative Study. *Ann Plast Surg*. 2014;72(6):631-637.
14. Leonardis JM, Lyons DA, Giladi AM, Momoh AO, Lipps DB. Functional integrity of the shoulder joint and pectoralis major following subpectoral implant breast reconstruction. *J Orthop Res*. 2019;37(7):1610-1619. doi: 10.1002/jor.24257.
15. Atisha D, Alderman AK. A systematic review of abdominal wall function following abdominal flaps for postmastectomy breast reconstruction. *Ann Plast Surg*. 2009;63(2):222-230. doi: 10.1097/SAP.0b013e31818c4a9e
16. Jeevan R, Cromwell DA, Browne JP, et al. Findings of a national comparative audit of mastectomy and breast reconstruction surgery in England. *J Plast Reconstr Aesthet Surg*. 2014;67(10):1333-1344. doi: 10.1016/j.bjps.2014.04.022.
17. Nijs J, Wijma AJ, Leysen L, et al. Explaining pain following cancer: a practical guide for clinicians. *Braz J Phys Ther*. 2019;23(5):367-377. doi: 10.1016/j.bjpt.2018.12.003.

18. Bruce J, Williamson E, Lait C, et al. Randomised controlled trial of exercise to prevent shoulder problems in women undergoing breast cancer treatment: study protocol for the prevention of shoulder problems trial (UK PROSPER). *BMJ Open*. 2018;8(3):e019078. doi: 10.1136/bmjopen-2017-019078.
19. Lin K-Y, Edbrooke L, Granger CL, Denehy L, Frawley HC. The impact of gynaecological cancer treatment on physical activity levels: a systematic review of observational studies. *Braz J Phys Ther*. 2019;23(2):79-92. doi: 10.1016/j.bjpt.2018.11.007.
20. Cutress RI, Summerhayes C, Rainsbury R. Guidelines for oncoplastic breast reconstruction. *Ann R Coll Surg Engl*. 2013;95(3):161-162. doi: [10.1308/003588413X13511609957696](https://doi.org/10.1308/003588413X13511609957696)
21. Ministério da Saúde. Política nacional de atenção oncológica. In:2005.
22. Brasil. Ministério da Saúde. Secretaria de Atenção à Saúde. Instituto Nacional de C. Controle do Câncer de Mama: documento de consenso. In:2004.
23. Eysenbach G. Improving the Quality of Web Surveys: The Checklist for Reporting Results of Internet E-Surveys (CHERRIES). *J Med Internet Res*. 2004;6(3):e34.
24. Angst F, Schwyzer HK, Aeschlimann A, Simmen BR, Goldhahn J. Measures of adult shoulder function: Disabilities of the Arm, Shoulder, and Hand Questionnaire (DASH) and its short version (QuickDASH), Shoulder Pain and Disability Index (SPADI), American Shoulder and Elbow Surgeons (ASES) Society standardized shoulder assessment form, Constant (Murley) Score (CS), Simple Shoulder Test (SST), Oxford Shoulder Score (OSS), Shoulder Disability Questionnaire (SDQ), and Western Ontario Shoulder Instability Index (WOSI). *Arthritis Care Res (Hoboken)*. 2011;63 Suppl 11:S174-188.
25. Silva AK, Lapin B, Yao KA, Song DH, Sisco M. The Effect of Contralateral Prophylactic Mastectomy on Perioperative Complications in Women Undergoing Immediate Breast Reconstruction: A NSQIP Analysis. *Ann Surg Oncol*. 2015;22(11):3474-3480. doi: 10.1245/s10434-015-4628-7.
26. Woo K-J, Lee K-T, Mun G-H, Pyon J-K, Bang SI. Effect of breast reconstruction modality on the development of postmastectomy shoulder morbidity. *J Plast Reconstr Aesthet Surg*. 2018;71(12):1761-1767. doi: 10.1016/j.bjps.2018.07.033.
27. All-Party Parliamentary Group on Breast Cancer. A mixed picture: An Inquiry into Geographical Inequalities and Breast Cancer. In:2018.
28. Smith-Turchyn J, Richardson J, Tozer R, McNeely M, Thabane L. Physical Activity and Breast Cancer: A Qualitative Study on the Barriers to and Facilitators of Exercise Promotion from the Perspective of Health Care Professionals. *Physiother Can*. 2016;68(4):383-390. doi: [10.3138/ptc.2015-84](https://doi.org/10.3138/ptc.2015-84)
29. Irving G, Neves AL, Dambha-Miller H, et al. International variations in primary care physician consultation time: a systematic review of 67 countries. *BMJ Open*. 2017;7(10):e017902.
30. Owen K, Hopkins T, Shortland T, Dale J. GP retention in the UK: a worsening crisis. Findings from a cross-sectional survey. *BMJ Open*. 2019;9(2):e026048.
31. Fitzgerald GK, Fritz JM, Childs JD, et al. Exercise, manual therapy, and use of booster sessions in physical therapy for knee osteoarthritis: a multi-center, factorial randomized clinical trial. *Osteoarthritis Cartilage*. 2016;24(8):1340-1349. doi: 10.1016/j.joca.2016.03.001.
32. Young JL, Rhon DI, Cleland JA, Snodgrass SJ. The Influence of Exercise Dosing on Outcomes in Patients With Knee Disorders: A Systematic Review. *J Orthop Sports Phys Ther*. 2018;48(3):146-161. doi: 10.2519/jospt.2018.7637.
33. Rodes CH, Kurebayashi R, Kondo VE, Luft VD, Góes ÂBd, Schmitt ACB. O acesso e o fazer da reabilitação na Atenção Primária à Saúde. *Fisioter Pesqui*. 2017;24:74-82.

34. Harrington S, Michener LA, Kendig T, Miale S, George SZ. Patient-reported upper extremity outcome measures used in breast cancer survivors: a systematic review. *Arch Phys Med Rehabil*. 2014;95(1):153-162. doi: 10.1016/j.apmr.2013.07.022.
35. Richmond H, Lait C, Srikesavan C, et al. Development of an exercise intervention for the prevention of musculoskeletal shoulder problems after breast cancer treatment: the prevention of shoulder problems trial (UK PROSPER). *BMC Health Serv Res*. 2018;18(1):463. doi: 10.1186/s12913-018-3280-x.
36. McNeely ML, Binkley JM, Pusic AL, Campbell KL, Gabram S, Soballe PW. A prospective model of care for breast cancer rehabilitation: postoperative and postreconstructive issues. *Cancer*. 2012;118(8 Suppl):2226-2236.
37. Ammitzboll G, Johansen C, Lanng C, et al. Progressive resistance training to prevent arm lymphedema in the first year after breast cancer surgery: Results of a randomized controlled trial. *Cancer*. 2019;125(10):1683-1692.
38. Beurskens CH, van Uden CJ, Strobbe LJ, Oostendorp RA, Wobbes T. The efficacy of physiotherapy upon shoulder function following axillary dissection in breast cancer, a randomized controlled study. *BMC Cancer*. 2007;7:166.
39. Schmitz KH, Courneya KS, Matthews C, et al. American College of Sports Medicine Roundtable on Exercise Guidelines for Cancer Survivors. *Med Sci Sports Exerc*. 2010;42(7):1409.
40. McNeely ML, Campbell K, Ospina M, et al. Exercise interventions for upper-limb dysfunction due to breast cancer treatment. *Cochrane Database Syst Rev*. 2010(6):CD005211. doi: 10.1002/14651858.CD005211.pub2.
41. Teixeira LFN, Sandrin F. The role of the physiotherapy in the plastic surgery patients after oncological breast surgery. *Gland Surg*. 2014;3(1):43-47.
42. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ*. 2009;339.
43. Futter CM, Weiler-Mithoff E, Hagen S, et al. Do pre-operative abdominal exercises prevent post-operative donor site complications for women undergoing DIEP flap breast reconstruction? A two-centre, prospective randomised controlled trial. *Br J Plast Surg*. 2003;56(7):674-683.
44. Rietjens M, De Lorenzi F, Andrea M, et al. Technique for Minimizing Donor-site Morbidity after Pedicled TRAM-Flap Breast Reconstruction: Outcomes by a Single Surgeon's Experience. *Plast Reconstr Surg Global Open*. 2015;3(8):e476-e476.
45. Rindom MB, Gunnarsson GL, Lautrup MD, et al. Shoulder-related donor site morbidity after delayed breast reconstruction with pedicled flaps from the back: An open label randomized controlled clinical trial. *J Plast Reconstr Aesthet Surg*. 2019;72(12):1942-1949. doi: 10.1016/j.bjps.2019.07.027
46. Donnelly CM, Lowe-Strong A, Rankin JP, Campbell A, Allen JM, Gracey JH. Physiotherapy management of cancer-related fatigue: a survey of UK current practice. *Support Care Cancer*. 2010;18(7):817-825. doi: 10.1007/s00520-009-0715-2.
47. O'Hanlon E, Kennedy N. Exercise in cancer care in Ireland: a survey of oncology nurses and physiotherapists. *Eur J Cancer Care (Engl)*. 2014;23(5):630-639.
48. Matsumura ESS, Sousa Júnior AS, Guedes JA, Teixeira RC, Kietzer KS, Castro LSF. Distribuição territorial dos profissionais fisioterapeutas no Brasil. *Fisiot Pesqui*. 2018;25:309-314.