

**Please cite the Published Version**

Mahamued, BA, Donald, PF, Collar, NJ, Marsden, SJ, Ndang'ang'a, PK, Wondafrash, M, Abebe, YD, Bennett, JAMES, Wotton, SR, Gornall, D and Lloyd, HUW (2022) Rangeland loss and population decline of the critically endangered Liben Lark *Heteromirafraga archeri* in southern Ethiopia. Bird Conservation International, 32 (1). pp. 64-77. ISSN 0959-2709

**DOI:** <https://doi.org/10.1017/S0959270920000696>

**Publisher:** Cambridge University Press

**Version:** Accepted Version

**Downloaded from:** <https://e-space.mmu.ac.uk/627173/>

**Usage rights:**



[Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0](#)

**Additional Information:** This is an Author Accepted Manuscript of a paper accepted for publication in Bird Conservation International, published by and copyright Cambridge University Press.

**Enquiries:**

If you have questions about this document, contact [openresearch@mmu.ac.uk](mailto:openresearch@mmu.ac.uk). Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

# Rangeland loss and population decline of the critically endangered Liben Lark *Heteromiraфра archeri* in southern Ethiopia

BRUKTAWIT A. MAHAMUED<sup>1,2</sup>, PAUL F. DONALD<sup>3,4\*</sup>, NIGEL J. COLLAR<sup>4</sup>,  
STUART J. MARSDEN<sup>1</sup>, PAUL KARIUKI NDANG'ANG'A<sup>5</sup>,  
MENGISTU WONDAFRASH<sup>6</sup>, YILMA DELLELEGN ABEBE<sup>6</sup>, JAMES BENNETT<sup>7</sup>,  
SIMON R. WOTTON<sup>3</sup>, DANIEL GORNALL<sup>1</sup> and HUW LLOYD<sup>1</sup>

<sup>1</sup> Department of Natural Sciences, Manchester Metropolitan University, Chester Street, Manchester M1 5GD, UK.

<sup>2</sup> Current address: Department of Biology, Kotebe Metropolitan University, P.O. Box 31248, Addis Ababa, Ethiopia.

<sup>3</sup> RSPB Centre for Conservation Science, The Lodge, Sandy, Bedfordshire SG19 2DL, UK.

<sup>4</sup> BirdLife International, David Attenborough Building, Pembroke Street, Cambridge CB2 3QZ, UK.

<sup>5</sup> BirdLife International Africa Partnership Secretariat, P.O. Box 3502 – 00100, Nairobi, Kenya.

<sup>6</sup> Ethiopian Wildlife and Natural History Society, Bole Sub City, P. O. Box 13303, Addis Ababa, Ethiopia.

<sup>7</sup> Centre for Agroecology, Water and Resilience, Coventry University, Ryton Gardens, Wolston Lane, Coventry CV8 3LG, UK.

\* Author for correspondence; e-mail: [paul.donald@birdlife.org](mailto:paul.donald@birdlife.org)

(Received 21 September 2020; revision accepted 06 December 2020)

## Summary

Liben Lark *Heteromiraфра archeri* is a 'Critically Endangered' species threatened by the loss and degradation of grassland at the Liben Plain, southern Ethiopia, one of only two known sites for the species. We use field data from nine visits between 2007 and 2019 and satellite imagery to quantify changes over time in the species' abundance and in the extent and quality of its habitat. We estimate that the population fell from around 279 singing males (95% CL: 182–436) in 2007 to around 51 (14–144) in 2013, after which too few birds were recorded to estimate population

size. Arable cultivation first appeared on the plain in the early 1990s and by 2019 more than a third of the plain had been converted to crops. Cultivation was initially confined to the fertile

black soils but from 2008 began to spread into the less fertile red soils that cover most of the plain.

Liben Larks strongly avoided areas with extensive bare ground or trees and bushes, but the extent of these did not change significantly over the survey period. A plausible explanation for the species' decline is that grassland degradation, caused before 2007 by continuous high-pressure grazing by livestock, reduced its rates of reproduction or survival to a level that could not support its previous population. Since 2015, communal *kalos* (grazing exclosures) have been established to generate forage and other resources in the hope of also providing breeding habitat for Liben Larks. Grass height and density within four grassland *kalos* in 2018 greatly exceeded that in the surrounding grassland, indicating that the plain retains the potential to recover rapidly if appropriately managed. Improvement of grassland structure through the restitution of

traditional and sustainable rangeland management regimes and the reversion of cereal agriculture to grassland are urgently needed to avert the species' extinction.

**Keywords:** Ethiopia, extinction, birds, Alaudidae

## Introduction

The Borana rangelands of southern Ethiopia were once among the most productive and culturally important in Africa (Hogg 1997), but in recent decades they have become increasingly degraded due to the encroachment of woody vegetation onto grassland, a decline in grass quality through overgrazing, and the loss of grassland to arable agriculture (e.g. Abate and Angassa 2016, Tilahun *et al.* 2017, Yassin 2019, Tolera and Senbeta 2020). These changes have been driven by a number of complex interacting processes, including the erosion of the *gada* system (a sophisticated traditional rangeland management regime based on indigenous ecological knowledge and strong social and cultural structures; Dalle *et al.* 2005), increasing frequency of drought, reduced food security, expanding human populations, land-grabbing, the prohibition of burning as a management tool, government policies aimed at settling nomadic pastoralists and the encroachment of neighbouring ethnic groups (Coppock 1994, Tache and Oba 2010, Adugna 2011, Galaty 2013, Reda 2016, Gilo and Kelkay 2017, Liao *et al.* 2018, Tekla *et al.* 2018).

Perhaps because of its cultural significance to the Borana people (Bassi 2005), the Liben Plain (5°15'N, 39°41'E; Figure 1) is one of the largest surviving remnants of open rangeland in southern Ethiopia. The plain was traditionally used by the Borana for grazing during the two annual wet seasons (e.g. Coppock 1994), but over the past 40 years there has been livestock grazing through the year due to the provision of permanent water sources, which remove the need to move cattle out to deep wells or rivers in the dry season, and increasing human settlement. Whistling Thorn *Vachellia drepanolobium* is encroaching the edges of the plain due to overgrazing and the cessation of burning, particularly along the eastern edge (Gemedo-Dalle *et al.* 2006, Tefera *et al.* 2007), and some parts of the plain are now suffering severe soil erosion. In the 1970s, a shift in government policy encouraged permanent settlement in place of transhumance, thereby increasing grazing pressure and eventually triggering cereal cultivation (Tilahun *et al.* 2017). Anecdotal evidence, particularly the testimony of elders we have interviewed, indicated that by 2010 the grassland on the plain was already much degraded in terms of structure and grass species diversity compared to a generation earlier (Donald *et al.* 2010).

The Liben Plain is one of only two known sites for Liben Lark *Heteromirafrarcheri*, also known as Archer's or Sidamo Lark (Collar 2009, Spottiswoode *et al.* 2009, 2013, Donald *et al.* 2010). A limited survey in October 2019 of the second site, near Jijiga in north-eastern Ethiopia, suggested that very few birds remain there (Negussie Toye and Abiy Dagne pers. comm.), and the Liben Plain may therefore support the more viable of the two populations. The species formerly occurred in Somaliland, but is probably extinct there (Mills *et al.* 2015), and distribution models have failed to identify suitable areas elsewhere (Spottiswoode *et al.* 2013). It is listed as 'Critically Endangered' on the IUCN Red List due to its small and declining population (BirdLife International 2020). A 40% decline in the number of Liben Larks recorded along the same transect routes between 2007 and 2009 coincided with changes in vegetation structure and loss of grassland to cereal crop production (Donald *et al.* 2010).

Between 2015 and 2018, a project funded by the UK government's Darwin Initiative supported local people to establish four *kalos* (grassland reserves) covering 259 ha (Figure 1) and opened 950 ha of grassland by clearing encroaching scrubland (<https://www.darwininitiative.org.uk/documents/22015/24529/22-015%20FR%20edited.pdf>). The purpose of *kalos* is to create fenced areas of tall, dense heterogeneous grass as both dry-season cattle forage and roofing materials for 1,300 households and to provide suitable breeding habitat for the Liben Lark. Although a series of censuses was undertaken between 2009 and 2019, these data have not hitherto been analysed to




Figure 1. Satellite image of the Liben Plain, southern Ethiopia (January 2014, 30-m NDVI from Landsat). The large dashed rectangle indicates the 286-km<sup>2</sup> area within which long-term crop expansion was measured from satellite imagery. White dots show the 252 transect waypoints. White asterisks indicate the locations of all Liben Larks recorded over the nine transect surveys. The white polygon indicates the 87.1-km<sup>2</sup> area across which density estimates from distance sampling were extrapolated to generate population estimates (see text). The four smaller hatched polygons indicate the *kalos* mentioned in the text. In the satellite base map, cultivation appears darker than grassland.

assess long-term changes in habitat area, habitat selection, grassland condition or population size. Furthermore, the effects on grassland structure of reducing or removing grazing pressure through the creation of fenced grassland reserves (*kalos*) have not yet been assessed.

We therefore (1) quantify long-term (1994–2019) changes in the area of grassland and crop farming using satellite imagery, (2) assess trends in a number of measures of grassland condition between 2007 and 2019, (3) estimate changes in the population size of Liben Lark between 2007 and 2019, (4) assess habitat selection by Liben Larks with respect to available habitats (recognising that none may now be optimal), and (5) compare measurements of grassland structure inside and outside four *kalos* created as part of an effort to seek conservation solutions for the Liben Lark that also benefit local people.

## Methods

### Study site

The Liben Plain is a relatively flat area of grassland at 1,500–1,550 m elevation, starting 10 km south-east of the town of Negele (Figure 1). It has a mean annual temperature of 20°C and a mean

annual rainfall of 700 mm confined to two rainy seasons, a longer (main) one in March–June (traditionally known as *ganna*) and a shorter one in September–November (*hagaya*) (Ethiopian Wildlife and Natural History Society 1996). The main soil types comprise relatively infertile iron-rich red sandy loams ('red soils') and more fertile black clay and volcanic light-coloured silty clay, silt and vertisols ('black soils') (Coppock 1994).

### *Transect surveys*

All field data were collected along 20 transects established during the first fieldwork on the Liben Plain in 2007 (Spottiswoode *et al.* 2009); details of how count and habitat data were collected and analysed are given below. These transects were broken up into 250-m lengths by a series of 252 waypoints (Figure 1), at which vegetation data were collected. In 2007 all 252 waypoints were located in grassland, but over the years an increasing number were converted to agriculture (see Results), after which visits were not made to the lost waypoints and no measures of vegetation were taken to avoid damaging the crop. Despite this, we spent much time observing cereal crops as we walked around them to regain the transects, and were convinced that Liben Larks were not present in cropland. Several singing (therefore male) Liben Larks were seen over a newly sprouting patch of cereal near a grassland edge in 2011 (Collar and Donald 2018), but this was the only time the species was observed using any habitat other than open grassland, and it may be the birds were singing over territories that had been established on grassland before it was ploughed and sown.

Field data were collected by the authors and others in various combinations in June 2007 (survey 1), May 2009 (survey 2), June 2010 (survey 3), May 2011 (survey 4), November 2012 (survey 5), May 2013 (survey 6), October–November 2013 (survey 7), June 2018 (survey 8) and May–June 2019 (survey 9). Surveys were timed to coincide with one of the two annual rainy seasons, as this was likely to coincide with peak Liben Lark breeding activity and song (Collar and Donald 2018) and hence greatest detectability.

### *Measuring loss of grassland area*

To assess long-term land use change on the Liben Plain and its environs using satellite imagery, we selected a rectangular area of 286 km<sup>2</sup>, centred on the remnant Liben Lark population but also extending out into areas where grassland was present in 2007 but larks were not recorded (Figure 1). Black soil within this area covers an area of 103 km<sup>2</sup> (36.0%), estimated from Landsat imagery (from which the different soil types can readily be distinguished by eye; Figure S1 in the online supplementary material); the rest of the area is covered by red soil. We downloaded 43 freely available Landsat 1–8 satellite images of the study area from the Global Visualization Viewer (GLOVIS) website from the years 1972–1995 and 2000–2019 (obstruction-free images from 1996–1999 were unavailable) at 30-m resolution. The images were stacked and clipped to the area of interest in ArcGIS Desktop (version 10.2). No pre-processing of the images was required as the maps were used simply to guide manual digitisation. The arable lands on red and black soils were digitised separately for each year and the cultivated area each year was then calculated in ArcMap using a Mollweide equal area projection.

### *Vegetation surveys and vegetation change*

During each field survey, grassland structure variables were recorded at each waypoint and at each location at which Liben Larks were first detected during transect surveys using the methods described in Spottiswoode *et al.* (2009) and Donald *et al.* (2010). The methods were designed to be highly repeatable across observers and years. Within a 25-m radius of the waypoint or bird location we recorded the number of trees, Giant Fennel *Ferula communis* plants (the only common herbaceous plant that stood much taller than the grass), paths (as an index of human disturbance) and harvester ant *Messor cephalotes* nests (which are large, circular bare patches in the grassland

and form the only naturally occurring areas of bare ground on the plain). Within two 5-m radius circles centred 10 m either side of the waypoint we counted the number of bushes and cowpats (as an index of grazing pressure), and estimated by eye the percentage of bare ground and the percentage cover of four grass height categories (<5 cm, 5–15 cm, 15–40 cm and >40 cm) (Spottiswoode *et al.* 2009). The four estimates of grass cover by height class were combined into a single grass height index for analysis by multiplying the percentage cover of each height class by the mid-points 2.5, 10, 28 and 40, respectively, and summing them (range of possible values: 0–4,000). At the start of each survey, all observers undertook a number of habitat assessments together to ensure comparable recording.

For analysis of changes in grassland condition we only included data from the 168 (66.7%) transect waypoints which remained as grassland throughout the entire period of coverage and for which data were therefore available in each of the nine field seasons. The remaining 84 transect points were converted to cropland during the period of coverage and so lacked measurements of grass condition for some or most years (nine returned to grassland after the abandonment of cultivation). Owing to the uneven distribution and relatively small number of visits across the survey period, we did not attempt to fit time series models to examine temporal patterns of change, but instead treated visit as a fixed factor and assessed significant differences between visits. Counts of fennel plants, trees, bushes, ant nests and paths were heavily skewed by a large number of zeros and non-zero counts were generally low (typically 1 or 2 for most variables), so we analysed changes over time in the presence or absence of these variables using generalised linear mixed models in which the presence or absence of each variable was modelled as an effect of survey, with transect waypoint nested within transect fitted as a random effect to control non-independence of repeat measures from the same points (Bolker *et al.* 2009). The count of cowpats present was modelled using Poisson errors with a log link, whereas the percentage of bare ground and the log of grass height index were approximately normally distributed and modelled using general linear mixed models and an identity link; again, waypoint and transect were fitted as nested random effects. Pairwise differences between visits were assessed from the models with *post hoc* Tukey HSD tests using the *glht* function in the R package *multcomp* (Hothorn *et al.* 2008). All models were checked for overdispersion using a function that compares residual degrees of freedom with residual variance. Analyses were undertaken using R v. 4.0.1 (R Core Team 2020).

### *Liben Lark population estimates*

We surveyed Liben Larks using the transect methods described by Spottiswoode *et al.* (2009) and Donald *et al.* (2010). Similar methods have subsequently been shown to be efficient for surveying other rare larks in large, open areas (Pérez-Granados and López-Iborra 2016). Transect surveys were undertaken from 06h30 to 11h00 only in suitable weather conditions. Along each transect observers recorded the number and activity of all Liben Larks seen or heard, and GPS coordinates were taken where each individual was first sighted. Singing birds were assumed to be males, as singing in females is very rare in larks (Donald and Alström 2021 in press). For each observation, the perpendicular distance to the bird from the transect line was estimated in GIS for distance sampling (Buckland *et al.* 2001). In June 2018 we had sufficient time and capacity to walk the transects twice, but on neither occasion did we record sufficient birds for distance sampling (see below).

Transect count data were analysed using the Conventional Distance Sampling (CDS) engine of Distance v. 6.2 software (Thomas *et al.* 2010). Density was expressed as number of individuals per km<sup>2</sup>. The shape criteria of distance histograms were examined for heaping or cluster bias and where necessary any outliers were right-hand truncated (Thomas *et al.* 2010). The actual values for truncation and subsequent grouping of records into distance bands were determined through visual inspection of histograms (Thomas *et al.* 2010). Data were truncated to 300 m to improve modelling of the detection function, thus minimising any bias caused by the few data points recorded beyond this distance. Three key functions (uniform, half-normal and hazard rate, all with cosine series



adjustment) were considered for each analysis (Buckland *et al.* 2001). Key function selection was evaluated using Akaike's Information Criterion (AIC), and a  $\chi^2$  statistic was used to assess the goodness of fit of each function (Buckland *et al.* 2001). We estimated the population density of Liben Larks first by using a global detection function based on all data pooled from all surveys, and then estimated densities per survey year by post-stratifying the same global detection function by survey. This assumes—reasonably (but necessarily, due to small sample sizes)—that detectability did not differ significantly across years. The reliability of density estimates was provided using the percentage coefficient of variation (% CV) and 95 % confidence intervals (95 % CI).

Population estimates for each visit were derived by multiplying the estimated densities from distance sampling by the area of a minimum convex polygon encompassing all transect survey points, from which two areas in the centre of the plain that were already under crops at the time of the first survey were excised, and which was then buffered by 300 m to account for the transect truncation distance (total area = 87.1 km<sup>2</sup>; Figure 1). This was then multiplied by the proportion of the 252 transect waypoints that had not been converted from grassland to cropland in the respective year, to account for changing grassland area. Extensive searches outside this polygon, particularly in apparently suitable grassland at the eastern end of the plain (Figure 1), failed to record any birds there and we are confident that the clipped minimum convex polygon contained most or all individuals in this population since 2007. In 2018 and 2019, too few birds were recorded during the transect counts to calculate density estimates effectively.

### *Habitat selection*

We assessed habitat selection of Liben Larks by comparing field measurements taken at locations where birds were first encountered during transects with those taken at transect waypoints, which were assumed to constitute a representative random sample of conditions across the plain. The presence or absence of larks was fitted as a binary variable in a generalised linear mixed model with binomial errors and a logit link function. Transect nested within survey was fitted as random effects to control the non-independence of observations along transects and within years (lark locations were assigned to the transect from which they were observed). The presence/absence variable was modelled as a function of the log of the grass height index, the percentage of bare ground and the number of bushes, ant nests, paths and fennel plants. All explanatory variables were scaled to zero mean and unit variance. We used variance inflation factors to assess the degree of collinearity in the predictors and, as these were always less than 1.5, we then estimated the model-averaged coefficients for each predictor across all possible subsets of the full model, including the full model itself, to identify predictors with model-averaged coefficients whose upper and lower 95 % CI did not span zero.

## **Results**

### *Changes in grass and crop cover across the Liben Plain*

Within the 286-km<sup>2</sup> rectangle containing the Liben Plain and its immediate environs (Figure 1) arable cropland first appeared in 1994, after which it expanded rapidly to cover a total area of 110.2 km<sup>2</sup>, or 38.5% of the area, by 2019 (Figure 2, Figure S2). By 2006 most of the more fertile black soils were under permanent cultivation, after which there was a sharp increase in cropland on the less fertile red soils, such that by 2019 the areas of red and black soils under cropland were almost equal. There was no evidence that the rate of conversion of grassland to crops has slowed over time, although field observations from 2019 indicated that some abandoned cropland was reverting to grassland. A similar rate of conversion of grassland was apparent along transects, where 75 (29.8%) of the 252 transect waypoints established in grassland in 2007 had been converted to cropland by 2019 (by which year nine transect points previously converted to crops had reverted to grassland).




Figure 2. Changes in the area of cropland within a 286-km<sup>2</sup> rectangle (see Figure 1) enclosing the Liben Plain, 1990–2019, taken from visual digitisation in GIS of Landsat imagery. No obstruction-free images were available for 1996–1999. Analysis of images back to 1972 (data not shown here) confirmed that crops first appeared in the area in 1994. The areas of cropland on red and black soils are shown separately and summed.

### *Trends in grassland condition*

Field measurements of grassland structure and condition were uncorrelated except for a strong negative correlation between grass height index and percentage of bare ground. There were significant pairwise differences between surveys in the proportion of points at which ant nests, fennel, paths, trees, and bushes were recorded, but no clear trend over time in any of them (Figure 3). Similarly, there were significant pairwise differences between surveys in number of cowpats recorded per point, percentage of bare ground, and grass height index, but again with no clear trend over time in any of them (Figure 4).

### *Habitat selection*

Model-averaged parameters from models of the presence or absence of larks indicated their strong avoidance of bushes and trees and of areas with high cover of bare ground (Figure 5). There was also some evidence for a selection for areas around ant nests and a weak avoidance of paths and tracks. Interpretation of these models is discussed further below.

### *Liben Lark population estimates*

The overwhelming majority of birds recorded along transects were singing males, so population estimates are more reflective of the number of territorial males than the number of individuals overall. The number of birds counted along the fixed transect routes varied between 10 (in 2018) and 68 (2007) (Figure 6). Densities of Liben Larks within the buffered MCP described by the transect points, estimated from distance sampling, varied between 3.2 singing birds/km<sup>2</sup> (95% CL: 2.1–5.0) in 2007 to 0.7 singing birds/km<sup>2</sup> (95% CL: 0.2–2.0) in 2013. These translated into overall population estimates of 279 singing birds (95% CL: 182–436) in 2007 to 51 (14–144) in 2013, accounting for changes in both density and grassland area (Figure 6).






Figure 3. Changes over time (nine surveys from the first in 2007 to the most recent in 2019) in the modelled probability of the presence or absence ( $\pm 95\%$  CL) at fixed transect waypoints across the Liben Plain of (a) fennel *Ferula* plants, (b) nests of harvester ants, (c) paths or tracks, (d) trees and bushes. Surveys sharing one or more letters did not differ from each other at  $P < 0.05$  in the respective attribute.

### Grassland structure in kalos

In 2018, the grassland height index was significantly higher in all four *kalos* and the percentage of bare ground significantly lower than in grassland outside the *kalos* (Figure 7). In 2019, however, long after the fences had been broken and cattle allowed entry to the *kalos*, the grassland structure in the *kalos* was much more similar to that of the grassland across the Liben Plain: the grass height index was still higher and the area of bare ground lower than grassland across the plain in two *kalos*, but not significantly different in the other two (Figure 7).




Figure 4. Changes over time (nine surveys from the first in 2007 to the most recent in 2019) in (a) number of cowpats, (b) percentage of bare ground and (c) grass height index (logged) recorded at fixed transect waypoints. Data are shown as boxplots. The associated tests of significant pairwise differences are derived from generalised linear mixed models (see text). Surveys sharing one or more letters did not differ from each other at  $P < 0.05$  in the respective attribute.

## Discussion

Our results suggest that the population of Liben Larks on the Liben Plain, as estimated from raw counts of birds along transects, declined by over 80% between 2007 and 2019 and may now comprise fewer than 50 territorial males, although the precision of our population estimates is very low. As there is evidence that the adult sex ratio is heavily skewed towards males (Collar and Donald 2018), as is usually the case in declining and threatened bird populations (Donald 2007), the effective population may be considerably lower than 50 pairs. A rapid survey in 2019 of the only other known population, near Jijiga in north-eastern Ethiopia, found only small numbers in




Figure 5. Model-averaged parameters ( $\pm 95\%$  CI), estimated over all subsets of the full model, of the presence or absence of Liben Larks (see text for full details). Negative parameter estimates indicate negative association of the variable with the presence of Liben Larks. Model-averaged parameters shown with dotted lines differed significantly from zero at  $P < 0.0001$ ; all others were not significantly different from zero ( $P > 0.05$ ).




Figure 6. Population estimates (squares,  $\pm 95\%$  CL, left axis) of territorial Liben Larks, 2007–2013, and counts of larks (dots, right axis) recorded along transects, 2007–2019.




Figure 7. Boxplots of log-transformed grass height index (a) and percentage of bare ground (b) measured at grassland transects across the Liben Plain (open bars) and along transects within each of four *kalos* (K1–K4, shaded bars; Fig. 1) in 2018 and 2019.

remaining patches of grassland, suggesting it is similarly threatened there and indicating the urgent need for a comprehensive survey. On this evidence the Liben Lark is poised to become mainland Africa's first recorded bird extinction. The only congeneric species, Rudd's Lark *H. ruddi* of South Africa, is also highly threatened and declining (Gush *et al.* 2019), so that the survival of an entire genus of birds is at risk.

Our results show that the Liben Plain has been affected by the same changes that have afflicted grasslands across Ethiopia in the past 50 years: despite frequent harvest failures (Tolera and Abebe 2007), the impetus for arable cultivation is such that even infertile red soils are being converted to cereal production, with communal land often being appropriated illegally by agricultural investors from elsewhere (Boru *et al.* 2015). The decline in the Liben Lark's population on the Liben Plain since the first survey in 2007 is correlated with an increase in cereal coverage that satellite imagery shows started in 1994, around the time that Borana pastoralists and Arsi agropastoralists first became permanently settled on the plain.

However, the approximately 80% decline in the lark's population between 2007 and 2019 greatly exceeds the less than 40% loss of grassland over the same period, suggesting that conversion to cropland is not the only (and perhaps not even the main) driver of the decline. We detected no long-term trends in a number of other metrics of grassland use or condition over the same period, including the prevalence of bushes or bare ground, which were negatively associated with the presence of Liben Larks. We consider that changes in grassland management and structure that

occurred before our first survey in 2007, and have not subsequently been reversed, may have reduced the species' productivity or survival such that the population has entered long-term decline. Examples from elsewhere in Africa have shown that poorly managed rangelands can transition from productive to less productive, yet nevertheless stable, states (e.g. Briske *et al.* 2005) and our data suggest that this is what has happened on the Liben Plain. Although too few nests have been found to estimate breeding success, it appears to be very low, probably because the short swards of the heavily overgrazed grassland offer few safe nest sites (Collar and Donald 2018). That Liben Lark is a species adapted to heavily grassed habitats is evidenced by its long hallux (hind-claw) (Green *et al.* 2009), and by its construction of a nest covered by a grass dome (Collar and Donald 2018). In contrast, Somali Short-toed Lark *Alaudala somalica*, which is abundant on the Liben Plain, has (as its common name implies) the short hallux of a species adapted to bare ground, on which it builds an open nest.

If the extinction of the Liben Lark is to be averted, means must be sought to improve the condition and extent of grassland on the Liben Plain, to the mutual benefit of both birds and local people. Crop field abandonment following harvest failures could present an opportunity to restore smaller patches of the Liben Plain to suitable areas for dry-season grazing and, perhaps, suitable vegetation for the Liben Lark, but this alone is unlikely to be sufficient to avert the lark's extinction.

The tall, dense grass cover that developed rapidly in the *kalos* certainly indicated that the Liben Plain retains the potential to recover quickly if grazing pressure can be controlled. However, it appeared to be too tall and dense to allow occupation and breeding by Liben Larks, which we never saw in fully tended *kalos*, although a *kalo* half-grazed by intruding cattle was found to host a breeding pair of Liben Larks (seen carrying food to hidden young). This suggested that a controlled grazing regime may be required within *kalos* to develop an optimal vegetation structure for the species, as might be expected to have existed when wild ungulates roamed the Liben Plain. Socioeconomic pressures led to the partial removal of *kalo* fencing early in 2019, and the entry of cattle caused the previously dense grassland within the *kalos* to revert to the heavily grazed structure of the rest of the plain. Nevertheless, a survey of 200 households found that 87% reported benefiting from the brief presence of the *kalos* (SRW and colleagues, unpubl. data). Thus, while initial results were discouraging, there is reason to think that more extensive *kalos* within which grazing pressure is managed at levels that promote a suitably heterogeneous sward for breeding and foraging Liben Larks, represent the best option for the conservation of the species in the short term. These are likely to be more effective if embedded within a participatory land-use plan for the plain that considers the views and needs of all relevant stakeholders. However, the long-term survival of the species, and of the traditional Borana way of life, is likely to depend on the restoration and expansion of rangeland across the entire plain through the reversion of cropland to grassland, a reduction in grazing pressure (including the reintroduction of lengthy resting periods), the clearance of encroaching scrub around the edges of the plain and a revival of the *gada* system. Examples of such restoration at large scales are unfortunately very rare, and the future of the Liben Lark remains very uncertain despite ongoing conservation efforts.

## Supplementary Material

To view supplementary material for this article, please visit <http://dx.doi.org/10.1017/S0959270920000696>.

## Acknowledgements

Our work on the Liben Plain was funded over the years by various sources acknowledged in earlier papers, but notably by the UK Darwin Initiative, the A. G. Leventis Foundation and Julian Francis. Field data were collected by ourselves and the late M. Bekele, K. Blomerley, A. Dagne, M. Davies, L. D. C. Fishpool, M. N. Gabremichael, M. Kariuki, S. Jervis, D. Lindo, A. Morris, K. Mwangi, A. D. Ruffo, W. Sara, G. Sherkete, M. Smith, R. Spencer, C. N. Spottiswoode, and S. Zekele. We have

been greatly assisted for years by Abiy Dagne, and we dedicate this paper to the late Negussie Toye, a great supporter of Liben Lark conservation. We thank the Associate Editor and two anonymous reviewers for helpful comments.

## References

- Abate, T. and Angassa, A. (2016) Conversion of savanna rangelands to bush dominated landscape in Borana, Southern Ethiopia. *Ecol. Processes* 5: 6.
- Adugna, F. (2011) Overlapping nationalist projects and contested spaces: The Oromo–Somali borderlands in Southern Ethiopia. *J. Eastern Afr. Stud.* 5: 773–787.
- Bassi, M. (2005) *Decisions in the shade: Political and juridical processes among the Oromo–Borana*. Trenton, New Jersey: Red Sea Press.
- BirdLife International (2020) Species fact-sheet: *Heteromirafra archeri*. <http://www.birdlife.org>.
- Bolker, B. M., Brooks, M. E., Clark, C. J., Geange, S. W., Poulsen, J. R., Stevens, M. H. H. and White, J.-S. S. (2009) Generalized linear mixed models: a practical guide for ecology and evolution. *Trends Ecol. Evol.* 24: 127–135.
- Boru, D., Schwartz, M., Kam, M. and Degen, A.A. (2015) Effects of family size and wealth on size of land cultivated by Borana pastoralists in southern Ethiopia. *Human Ecol.* 4: 15–28.
- Briske, D. D., Fuhlendorf, S. D. & Smeins, F. E. (2005) State-and-transition models, thresholds, and rangeland health: A synthesis of ecological concepts and perspectives. *Rangeland Ecol. Manage.* 58: 1–10.
- Buckland, S. T., Anderson, D. R., Burnham, K. P., Laake, J. L., Borchers, D. L. and Thomas, L. (2001) *Introduction to distance sampling: Estimating abundance of biological populations*. New York: Oxford University Press.
- Collar, N. J. (2009) Conservation-driven changes in English bird names, and the case of the Liben Lark. *Bull. Afr. Bird Club* 16: 245.
- Collar, N. J. and Donald, P. F. (2018) Notes on breeding activity, nesting habitat and sex ratio in the critically endangered Liben Lark *Heteromirafra archeri*. *Bull. Afr. Bird Club* 25: 45–53.
- Coppock, D. L. (ed.) (1994) *The Borana Plateau of southern Ethiopia: synthesis of pastoral research, development and change, 1980–91*. Addis Ababa, Ethiopia: International Livestock Centre for Africa (ILCA).
- Dalle, G., Isselstein, J. and Maass, B. L. (2005) Indigenous ecological knowledge of Borana pastoralists in southern Ethiopia and current challenges. *Internatn. J. Sustain. Devel. World Ecol.* 12: 1–18.
- Donald, P. F. (2007) Adult sex ratios in wild bird populations. *Ibis* 149: 671–692.
- Donald, P. F., Buchanan, G. M., Collar, N. J., Abebe, Y., Gabremichael, M., Mwangi, M., Ndang’ang’a, P. K., Spottiswoode, C. N. and Wondafrash, M. (2010) Rapid declines in habitat quality and population size of the Liben (Sidamo) Lark *Heteromirafra sida-moensis* necessitate immediate conservation action. *Bird Conserv. Internatn.* 20: 1–12.
- Donald, P. F. and Alström, P. (2021, in press) *Larks of the world*. London: Helm.
- Ethiopian Wildlife and Natural History Society (1996) *Important Bird Areas of Ethiopia*. Addis Ababa: EWNHS.
- Galaty, J. G. (2013) Land grabbing in the eastern African rangelands. Pp. 143–153 in A. Cately, J. Lind and I. Scoones, eds. *Pastoralism and development in Africa: dynamic change at the margins*. Oxford, UK: Routledge.
- Gemedo-Dalle, Maass, B. L. and Isselstein, J. (2006) Rangeland condition and trend in the semi-arid Borana lowlands, southern Oromia, Ethiopia. *Afr. J. Range Forage Sci.* 23: 49–58.
- Gilo, B. N. and Kelkay, T. Z. (2017) Changes in vegetation structure and aboveground biomass in response to traditional rangeland management practices in Borana, southern Ethiopia. *Afr. J. Range Forage Sci.* 34: 21–31.
- Green, R. E., Barnes, K. N. and Brooke, M. d. L. (2009) How the longspur won its spurs: a study of claw and toe length in ground-dwelling passerine birds. *J. Zool.* 277: 126–133.
- Gush, W., Maphisa, D. H., Reynolds, C., Donald, P. F. and Spottiswoode, C. N. (2019) Declines of the globally threatened



- Rudd's Lark *Heteromirafr ruddi* in one of its last remaining strongholds. *Bird Conserv. Internatn.* 29: 644–656.
- Hogg, R. (1997) Introduction. Pp. 1–22 in R. Hogg, ed. *Pastoralists, ethnicity and the state in Ethiopia*. London: Haan Publishing.
- Hothorn, T., Bretz, F. and Westfall, P. (2008) Simultaneous Inference in General Parametric Models. *Biometric. J.* 50: 346–363.
- Liao, C., Clark, P. E. and DeGloria, S.D. (2018) Bush encroachment dynamics and rangeland management implications in southern Ethiopia. *Ecol. Evol.* 8: 11694–11703.
- Mills, M. S. L., Cohen, C., Francis, J. and Spottiswoode, C. N. (2015) A survey for the Critically Endangered Liben Lark *Heteromirafr archeri* in Somaliland, north-western Somalia. *Ostrich* 86: 291–294.
- Pérez-Granados, C. and López-Iborra, G. M. (2016) Assessment of counting methods used for estimating the number of territorial males in the endangered Dupont's Lark. *Ardeola* 64: 75–84.
- R Core Team (2020) *R: A language and environment for statistical computing*. Vienna, Austria: Foundation for Statistical Computing.
- Reda, K. T. (2016) Dynamics in pastoral resource management and conflict in the Borana rangelands of southern Ethiopia. *Afr. Security Rev.* 25: 31–43.
- Spottiswoode, C., Wondafrash, M., Gabremichael, M., Abebe, Y., Mwangi, M., Collar, N. and Dolman, P. (2009) Rangeland degradation is poised to cause Africa's first recorded avian extinction. *Anim. Conserv.* 12: 249–257.
- Spottiswoode, C. N., Olsson, U., Mills, M. S. L., Cohen, C., Francis, J. E., Tøye, N., Hoddinott, D., Dagne, A., Wood, C., Donald, P. F., Collar, N. J. and Alström, P. (2013) Rediscovery of a long-lost lark reveals the conspecificity of endangered *Heteromirafr* populations in the Horn of Africa. *J. Ornithol.* 154: 813–825.
- Tache, B. and Oba, G. (2010) Is poverty driving Borana herders in Southern Ethiopia to crop cultivation? *Human Ecol.* 38: 639–649.
- Tefera, S., Snyman, H. A. and Smit, G. N. (2007) Rangeland dynamics in southern Ethiopia:(3). Assessment of rangeland condition in relation to land-use and distance from water in semi-arid Borana rangelands. *J. Environ. Manage.* 85: 453–460.
- Teka, H., Madakadze, C. I., Botai, J. O., Has-sen, A., Angassa, A. and Mesfin, Y. (2018) Evaluation of land use land cover changes using remote sensing Landsat images and pastoralists' perceptions on range cover changes in Borana rangelands, Southern Ethiopia. *Internatn. J. Biodivers. Conserv.* 10: 1–11.
- Thomas, L., Buckland, S. T., Rexstad, E. A., Laake, J. L., Strindberg, S., Hedley, S. L., Bishop, J. R. B., Marques, T. A. and Burnham, K. P. (2010) Distance software: design and analysis of distance sampling surveys for estimating population size. *J. Appl. Ecol.* 47: 5–14.
- Tilahun, A., Teklu, B. and Hoag, D. (2017) Challenges and contributions of crop production in agro-pastoral systems of Borana Plateau, Ethiopia. *Pastoralism* 7. DOI: [10.1186/s13570-016-0074-9](https://doi.org/10.1186/s13570-016-0074-9).
- Tolera, A. and Abebe, A. (2007) Livestock production in pastoral and agro-pastoral production systems of southern Ethiopia. *Livestock Res. Rural Devel.* 19: 177.
- Tolera, T. and Senbeta, F. (2020) Pastoral system in the face of climate variability: household adaptation strategies in Borana Rangelands, Southern Ethiopia. *Environ., Devel. Sustainabil.* 22: 3133–3157.
- Yassin, I. M. (2019) Bush encroachment in Borana rangeland in the case of Southern Ethiopia: Causes, impacts and management implications. *Internatn. J. Agric. Innovat. Res.* 7: 420–428.