


**Please cite the Published Version**

Craig, Hayley, Antwis, Rachael E, Cordero, Irene, Robinson, Claire H, Osborne, Tod Z, Bardgett, Richard D, Rowntree, Jennifer  and Simpson, Lorea T (2020) Nitrogen addition alters composition, diversity and functioning of microbial communities in mangrove soils: an incubation experiment. *Soil Biology and Biochemistry*, 153. p. 108076. ISSN 0038-0717

**DOI:** <https://doi.org/10.1016/j.soilbio.2020.108076>

**Publisher:** Elsevier

**Version:** Published Version

**Downloaded from:** <https://e-space.mmu.ac.uk/627211/>

**Usage rights:**  [Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0](https://creativecommons.org/licenses/by-nc-nd/4.0/)

**Additional Information:** This is an Open Access article published in *Soil Biology and Biochemistry*, published by Elsevier, copyright The Author(s).

**Enquiries:**

If you have questions about this document, contact [openresearch@mmu.ac.uk](mailto:openresearch@mmu.ac.uk). Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)



## Nitrogen addition alters composition, diversity, and functioning of microbial communities in mangrove soils: An incubation experiment

Hayley Craig<sup>a,\*</sup>, Rachael E. Antwis<sup>b</sup>, Irene Cordero<sup>a</sup>, Deborah Ashworth<sup>a</sup>, Clare H. Robinson<sup>a</sup>, Todd Z. Osborne<sup>c</sup>, Richard D. Bardgett<sup>a</sup>, Jennifer K. Rowntree<sup>d</sup>, Loraé T. Simpson<sup>c</sup>

<sup>a</sup> Department of Earth and Environmental Sciences, The University of Manchester, Oxford Road, Manchester, M13 9PT, United Kingdom

<sup>b</sup> School of Science, Engineering and Environment, University of Salford, The Crescent, Salford, M5 4WT, United Kingdom

<sup>c</sup> Whitney Laboratory for Marine Bioscience, University of Florida, Ocean Shore Boulevard, St. Augustine, FL, 32080, USA

<sup>d</sup> Ecology and Environment Research Centre, Department of Natural Sciences, Manchester Metropolitan University, Chester Street, Manchester, M1 5GD, United Kingdom

### ARTICLE INFO

#### Keywords:

Mangroves  
Coastal wetlands  
Nutrient pollution  
Bacterial diversity

### ABSTRACT

Mangrove ecosystems are important for carbon storage due to their high productivity and low decomposition rates. Waterways have experienced increased nutrient loads as a result of anthropogenic activities and it is unclear how this may affect carbon and nutrient cycles in downstream mangroves that receive these nutrient-rich waters. Using a laboratory-based incubation experiment, this study aimed to assess the effects of nutrient addition on the diversity and structure of mangrove soil bacterial communities, as well as biomass and activity of the soil microbial community, under different oxygen conditions. Bacterial community diversity and composition was characterised using 16S rRNA gene sequencing and microbial activity was examined through the measurement of microbial respiration and the activities of enzymes associated with organic matter decomposition. Nitrogen addition caused clear shifts in bacterial community composition, with decreases in bacterial diversity and the abundance of sulfate-reducing bacteria. Microbial biomass also decreased with nitrogen addition under reduced oxygen incubations. Changes in bacterial community structure were accompanied by changes in the activity of some enzymes involved in carbon, nitrogen, and phosphorus cycling. Under reduced oxygen conditions, nitrogen addition resulted in a significant increase in the microbial metabolic quotient but no accompanying change in microbial respiration, which was explained by a decrease in microbial biomass. The findings of this study indicate that nitrogen loading has potential implications for microbial communities and carbon and nutrient cycling in mangrove environments that warrant further investigation under field conditions.

### 1. Introduction

Nitrogen (N), phosphorus (P), and potassium (K) are essential nutrients for plants and microbes, and limitations in these nutrients can constrain growth and functioning. Anthropogenic activities have altered fluxes of these growth-limiting nutrients in freshwater and coastal marine ecosystems (Smith 2003), which can have detrimental impacts on water quality and the organisms that live in these environments (Smith et al., 1999). In many tropical and sub-tropical coastal regions, mangrove forests are often found in estuaries at the receiving end of nutrient enriched riverine waters. As mangrove ecosystems are important coastal carbon (C) stores (Donato et al., 2011; Alongi 2012), there is a need to better understand how elevated nutrient levels may impact ecosystem functioning, such as C and nutrient cycling, in these

environments.

Several field experiments have assessed mangrove plant responses to N and P addition. Generally, they have found that when plants were given a limiting nutrient, growth increased significantly, both above-ground (Feller 1995; Feller et al., 2003; Lovelock et al., 2004; Dangremond et al., 2019) and belowground (McKee et al., 2007), which in turn should increase leaf and root litter inputs to the soil. Chemical and structural changes have also been observed in the leaves of these plants, with reduced sclerophylly (leaf toughness), and altered concentrations of nutrients (Keuskamp et al., 2015a; Simpson et al., 2020) and phenolic compounds with fertilisation (Feller 1995; Feller et al., 1999; Keuskamp et al. 2015a, 2015b). Increases in litter quantity and changes in litter quality could have implications for soil processes, particularly those related to the C cycle (e.g. rates of decomposition, C storage, and surface

\* Corresponding author. NatureMetrics, CABI Site, Bakeham Lane, Egham, Surrey, TW20 9TY, United Kingdom.

E-mail address: [hcraig@hotmail.com.au](mailto:hcraig@hotmail.com.au) (H. Craig).

<https://doi.org/10.1016/j.soilbio.2020.108076>

Received 25 May 2020; Received in revised form 4 October 2020; Accepted 13 November 2020

Available online 18 November 2020

0038-0717/© 2020 The Authors.

Published by Elsevier Ltd.

This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

accretion). Further, nutrient enrichment will likely impact mangrove soil bacterial communities, which play an important role in C flux processes in mangrove soils (Holguin et al., 2001), and account for most of the total soil microbial biomass (Chen et al., 2016) in these systems. Despite this, few studies have investigated the impacts of nutrient enrichment on mangrove soil bacterial communities.

Several recent meta-analyses and large-scale multi-site studies have assessed overall trends of soil microbial responses to nutrient enrichment across a range of terrestrial ecosystems. Nitrogen addition has been associated with decreases in soil microbial diversity (Wang et al., 2018) and microbial biomass (Treseder 2008; Ramirez et al., 2012), while P addition can increase soil microbial biomass (Yue et al., 2018). Many of these terrestrial studies have also observed changes in soil bacterial community composition with nutrient enrichment (Leff et al., 2015; Ramirez et al., 2012; Wang et al., 2018), which may contribute to observed changes in soil functioning. Soil respiration responses to N and P addition vary across ecosystem types and have been associated with changes in microbial biomass, root biomass, and soil C (Zhou et al., 2014; Feng and Zhu 2019). Phosphorus addition has also been shown to suppress activity of P acquiring enzymes (Marklein and Houlton 2012; Xiao et al., 2018) and N addition can stimulate C and P acquiring enzymes, but suppress lignin-modifying oxidase activity (Chen et al., 2018a,b; Xiao et al., 2018). This decrease in oxidase activity, as well as increased litter input, can result in increased soil C storage in N-amended soils (Chen et al., 2018a,b).

Despite differences in environmental conditions compared to terrestrial soils, similar microbial responses to nutrient addition have been observed in mangrove and marine environments. Nitrogen addition to mangrove and marine sediments has been found to decrease microbial biomass (Keuskamp et al., 2015b), bacterial abundance (Luo et al., 2017), and bacterial diversity (Aoyagi et al., 2015). The redox conditions and low N levels in mangrove soils mean that the majority of the organic matter decomposition in mangroves occurs via sulfate reduction (Kristensen et al., 1991). However, studies that have experimentally increased levels of the more thermodynamically favourable nitrate have observed increases in the abundance of *Sulfurimonas* (Aoyagi et al., 2015; Wang et al., 2016), nitrate-reducing genus of bacteria. These shifts in soil bacterial communities could lead to changes in soil functioning, as well as increased emissions of greenhouse gases including carbon dioxide (CO<sub>2</sub>), nitrous oxide (N<sub>2</sub>O), and methane (CH<sub>4</sub>), as has been observed in some studies (Muñoz-Hincapié et al., 2002; Kreuzwieser et al., 2003; Chen et al., 2011). Changes in potential enzyme activity have also been observed in response to nutrient addition (Tam 1998; Luo et al., 2017), although responses can vary between vegetation zones within the mangrove ecosystem (Keuskamp et al., 2015b). When combined with changes in plant productivity, these nutrient-induced changes in soil functioning could affect C cycling and decomposition processes leading to changes in both the rate and direction of surface elevation change (McKee et al., 2007), an important factor when considering resistance and resilience of these ecosystems to rises in sea level.

The aim of this experiment was to assess the effects of N addition alone, and in combination with P and K, on the diversity, structure, and function of mangrove soil microbial communities, with a focus on bacterial communities, being the dominant component of the microbial biomass of mangrove soils (Chen et al., 2016). A laboratory-based, controlled soil incubation experiment enabled us to focus on soil microbial responses to nutrient enrichment by removing plant-mediated responses and potential impacts of other fluctuating environmental variables on microbial communities. We sought to identify which bacterial groups change with nutrient addition, and to link this with observed changes in microbial functioning. As mangrove soils experience both aerobic and anaerobic conditions, each with their own distinct microbial communities, we incubated soil in two different oxygen environments to compare responses. This information will improve our understanding of microbial responses to elevated nutrients and the

potential implications for C storage and nutrient cycling in mangrove environments.

This study was conducted with soils collected near a long-term field fertilisation experiment in northeast Florida where plant growth responses indicate that the site is limited by N, but not P (Dangremond et al., 2019). However, concentrations of leaf N:P in the control plots (Simpson et al., 2020) suggest that plant growth could be limited by either N, P, or both (Koerselman and Meuleman 1996). A decomposition study at the site found initial decay rates of leaf litter were higher in the nutrient addition treatments but stabilised across all treatments within six months (Simpson et al., 2020). The soil microbial communities of this site had not yet been investigated, but we hypothesised that N addition would: (1) decrease bacterial diversity, and (2) decrease abundance of sulfate-reducing bacteria. In terms of soil function, we hypothesised that N addition would: (3) increase microbial respiration and the activity of a P acquiring enzyme, and (4) decrease activity of lignin-modifying enzymes and an N acquiring enzyme. If the microbial community is also P and/or K limited, the addition of PK in combination with N could have additive effects on microbial responses. However, if the microbes are also limited by available C sources, the response to nutrient addition may be limited.

## 2. Methods

### 2.1. Study site and soil sampling

The soil for this experiment was obtained from Guana Tolomato Matanzas National Estuarine Research Reserve, St John's County, Florida (29.7291°N, 81.2415°W). The site is located within the saltmarsh-mangrove ecotone near the northernmost range limit of mangroves in Florida (Giri and Long 2016) and consists of black mangrove (*Avicennia germinans*) shrubs amongst *Batis maritima* and *Sarcocornia perennis* saltmarsh plants. Soil was collected in April 2018 at low tide when the soil surface was exposed. Approximately 20 L of soil was collected into a clean bucket using a trowel to a depth of 10 cm underneath unfertilised *A. germinans* shrubs, across an area of approximately 100 m<sup>2</sup>. Oxygen concentrations rapidly decline with depth from the surface in mangroves (Booth et al., 2019) and we targeted the top 10 cm to ensure we collected a mix of aerobic and anaerobic communities. Due to the peaty nature of the soil, instead of sieving it was sorted by hand in the lab to remove leaf litter, woody roots, sticks, shells, and other large debris, and then thoroughly mixed to homogenise. Soil was homogenised to minimise variation amongst mesocosms prior to nutrient application so differences at the end of the experiment could be attributed to treatments.

### 2.2. Incubation experiment

Soil mesocosms were set up in 473 mL (16 oz.) Ball® smooth sided jars (Rubbermaid Inc., Atlanta, GA, USA) with approximately 72 g (fresh weight) of the homogenised soil pressed down into the bottom of the jar. The lids of the jars were modified with stop cock valves and gas tubing to enable gas measurements. Soils were processed and weighed into jars on the day of collection. Mesocosms were randomly assigned to one of four nutrient treatments (control, +N, +PK, +NPK), and one of two oxygen treatments (ambient or reduced oxygen incubation) in a fully factorial design, with five replicates for each treatment combination (total n = 40). To obtain a reduced oxygen environment, jars were flushed with N<sub>2</sub> gas and the valves in the lid were then closed. Ambient oxygen jars were lidded, but with the valves left open. The mesocosms were left to incubate for four days to allow the soil to settle prior to administering the nutrient treatments.

Soil samples collected from the site in December 2017 indicated background soil concentrations of approximately 10.8 g/kg total N and 0.387 g/kg total P on a dry weight basis. This experiment aimed to double those concentrations by adding 2 mL of 4.95 M ammonium

nitrate ( $\text{NH}_4\text{NO}_3$ ) solution as the N source, and 1 mL of 0.16 M potassium phosphate dibasic ( $\text{KH}_2\text{PO}_4$ ) as the source of P and K. Deionised water was added to mesocosms as required to ensure that 3 mL of liquid was held constant in each treatment. The liquid was mixed into the soil to ensure even distribution, then the lids were refitted and the reduced oxygen mesocosms flushed again with  $\text{N}_2$  gas. The average soil moisture content (wet weight basis) of three samples taken from the homogenised soil before the start of the experiment was 82.2% and addition of the nutrients in liquid form would have increased this to ~82.9%. The mesocosms were kept in a dark room at around 22 °C when not being measured which is within the temperature range that would be experienced in the field at that time of year. The mesocosms were kept in the dark to prevent growth of algae or seeds ensuring that responses could be linked predominantly to microbial communities. To maintain soil moisture, after 11 days deionised water was added to each mesocosm until the starting weight was reached. The experiment ran for 15 days in April 2018, and all mesocosms were then destructively harvested for soil analyses.

### 2.3. Soil analyses

Soil pH was measured with an accumet pH meter (Fisher Scientific, Pittsburgh, PA, USA) in a 1:2 fresh soil to deionised water slurry. Total soil C and N were measured by dry combustion of dried, ground soil using a Vario EL cube CN analyser (Elementar Analysensysteme GmbH, Hanau, Germany). Soil moisture content was calculated gravimetrically on a wet weight basis from soil that had been oven-dried at 105 °C for 48 h. Organic matter content and total soil P were determined using a loss on ignition (LOI) method (Kuo 1996) by combusting soils at 550 °C for 4 h then using 0.5 M  $\text{H}_2\text{SO}_4$  as an extractant. Phosphate in the acid extracts was measured colourimetrically on a CLARIOstar microplate reader (BMG LABTECH, Ortenberg, Germany) using an ascorbic acid method adapted from Kuo (1996), with volumes scaled down for use in a microplate. All results are presented as means  $\pm$  1 standard error.

### 2.4. Microbial biomass and activity

Microbial biomass C was determined for each sample using the chloroform fumigation-extraction method (Vance et al., 1987) with 0.5 M  $\text{K}_2\text{SO}_4$  as an extractant. The standard 24 h fumigation time was used as this has previously been identified as suitable for water-saturated soils from rice fields (Witt et al., 2000). Total C was measured in the extracts with a TOC-L analyser (Shimadzu, Kyoto, Japan) and an extraction efficiency coefficient of 0.45 (Joergensen et al., 2011) was applied to calculate microbial biomass C.

$\text{CO}_2$  flux was measured daily with an LI-8100A automated soil  $\text{CO}_2$  flux system (Li-Cor Inc., Lincoln, Nebraska, USA) to assess microbial respiration. The first measurement from each mesocosm was taken within 30 min of the addition of the nutrients (or water) and each morning thereafter. Four consecutive flux readings of 60 s were measured. The initial reading was not included in the analysis to allow sufficient time for flushing of the gas lines and an average was calculated from the last three readings. All reduced oxygen mesocosms were flushed with  $\text{N}_2$  gas immediately prior to measurement to flush out any respired gases that had accumulated while sitting. The ambient oxygen mesocosms had their lids removed for a few minutes prior to analysis and then refitted immediately prior to measurement to ensure that ample fresh air was getting into the mesocosms. When water was added mid-experiment to the mesocosms to maintain soil moisture content, the reduced oxygen mesocosms required more water than the ambient mesocosms, which was likely due to the daily flushing of  $\text{N}_2$  gas. This led to a significant change in  $\text{CO}_2$  flux readings in the reduced oxygen mesocosms that was not observed in the ambient mesocosms. For this reason, we excluded the flux readings made after that point from the respiration analyses for both oxygen treatments. One of the ambient oxygen + PK samples was removed from the respiration analysis as it

had unusual readings for the first few days, most likely due to a suspected air leak in the lid during measurements. The metabolic quotient ( $q\text{CO}_2$ ) was calculated as the ratio of microbial respiration to microbial biomass C on the final day of the experiment (Anderson and Domsch 1985).

Potential extracellular enzyme activity was measured in soil that had been stored at 4 °C using assays for two oxidative enzymes (lignin peroxidase (PER) and phenol oxidase (POX)), and six hydrolytic enzymes (urease (URE), cellobiohydrolase (CBH),  $\beta$ -N-acetylglucosaminidase (NAG), acid phosphatase (AP), xylosidase (XYL), and  $\beta$ -glucosidase (GLC)). All enzyme activity assays were performed within two weeks of sampling following the methods from Jackson et al. (2013) for pNP-linked substrates (CBH, NAG, AP, XYL and GLC) and Cordero et al. (2019) for URE. L-DOPA assays were used to evaluate POX and PER activity following the methods in De Long et al. (2019) with double the amount of fresh soil added to the soil slurry due to high soil moisture content. Saturating substrate concentrations for pNP-linked substrates and URE were determined by first testing a range of concentrations on three samples until absorbance values did not increase. Incubation times were increased if absorbance values were still low after the initial incubation time from the referenced protocols. Details of the final assays are presented in Table 1. 50 mM sodium acetate buffer pH 5.0 was used for all assays. All enzyme assays were performed in microplates and measured colourimetrically using an EZ Read 400 microplate reader (Biochrom Ltd., Cambridge, United Kingdom). Absolute enzyme activities were calculated as potential activity per gram of dry weight soil and specific enzyme activities were calculated by dividing absolute enzyme activities by microbial biomass C (Moorhead et al., 2013).

### 2.5. Bacterial community diversity and composition

A metabarcoding approach was used to examine soil bacterial community diversity and composition. DNA was extracted the day after the end of the experiment from 0.5 g fresh weight soil from three of the five mesocosms within each treatment group using the PowerSoil PowerLyser DNA Isolation kit (MoBio Laboratories Inc., Carlsbad, CA, USA). An extraction blank containing no soil was also included to identify any contamination during the extraction process. DNA yield and quality were checked using a NanoDrop 1000 spectrophotometer (Thermo Fisher Scientific, Wilmington, DE, USA). A paired-end dual-index sequencing approach was used. The V4 region of the 16S rRNA gene was amplified with primers 515F (5'-GTGCCAGCMGCCGCGGTAA-3') and 806R (5'-GGACTACHVGGGTWTCTAAT-3') from Caporaso et al. (2011) containing Illumina adapter sequences and 8 bp index sequences following a one-step PCR protocol (Kozich et al., 2013; Griffiths et al., 2018). PCRs were performed in 15  $\mu\text{L}$  reactions containing 20 ng DNA, 1 x KAPA HiFi HotStart ReadyMix, and 0.2  $\mu\text{M}$  of each primer with the following cycle conditions: 95 °C for 5 min; followed by 28 cycles of 95 °C for 20 s, 55 °C for 15 s, 72 °C for 30 s; followed by 5 min extension at 72 °C. All PCRs were conducted in duplicate and each plate contained a negative control as well as a mock community control (Microbial Mock Community B Staggered, Low Concentration, v5.2L for 16S Gene sequencing, HM-783D, BEI Resources, Manassas, VA, USA). PCR duplicates were pooled, then cleaned using Ampure XP beads (Beckman Coulter Inc., Indianapolis, IN, USA). Fragment size was checked on a 4200 TapeStation (Agilent Technologies, Santa Clara, CA, USA). Samples were normalised using a titration run, in which equal volumes of each sample were combined and sequenced using v2 nano (2 x 150bp) chemistry on a MiSeq (Illumina, Inc., San Diego, CA, USA) at the University of Salford. Based on the number of reads assigned to each sample, an equimolar pool of all samples was created and sequenced on a MiSeq using 250 bp paired-end v2 chemistry.

Sequencing data were processed in R v3.5.2 (R Core Team 2018) with DADA2 v1.10.1 (Callahan et al., 2016) using the default pipeline. After paired-end reads had been merged, the modal sequence length was 253 bp. Ten sequence variants (SVs) of length >260 bp (0.002% of total

**Table 1**  
Details of the soil enzyme activity assays conducted in this study.

Enzyme	Code	Related nutrient cycle	Substrate	Slurry concentration	Incubation time (h)
Phenol oxidase	POX	C	L-3,4-dihydroxyphenylalanine (20 mM)	0.5 g soil + 25 mL buffer	20
Lignin peroxidase	PER	C	L-3,4-dihydroxy phenylalanine and H <sub>2</sub> O <sub>2</sub> (20 mM)	0.5 g soil + 25 mL buffer	1.5
Urease	URE	N	Urea (1 M)	2 g soil + 5 mL buffer	2
$\beta$ -glucosidase	GLC	C	<i>p</i> -nitrophenyl- $\beta$ -D- glucopyranoside (30 mM)	3.75 g + 5 mL buffer	2.5
$\beta$ -xylosidase	XYL	C	<i>p</i> -nitrophenyl- $\beta$ -D-xylopyranoside (25 mM)	3.75 g + 5 mL buffer	7
<i>N</i> -acetyl- $\beta$ -glucosaminidase	NAG	C and N	<i>p</i> -nitrophenyl- <i>N</i> -acetyl- $\beta$ -D-glucosaminide (5 mM)	3.75 g + 5 mL buffer	6
Cellobiohydrolase	CBH	C	<i>p</i> -nitrophenyl- $\beta$ -D-cellobioside (4 mM)	3.75 g + 5 mL buffer	8
Acid phosphatase	AP	P	<i>p</i> -nitrophenyl-phosphate disodium salt hexahydrate (30 mM)	3.75 g + 5 mL buffer	0.5

sequences) and 992 bimeras were removed. Sequence variant taxonomies were mapped to genus level using the Silva v132 training set (Quast et al., 2013). 46% of SVs were unknown at the genus level so their assignments were edited from “NA” to their family name so they would not all be lost when later agglomerating to genus. Data were then exported to the phyloseq package (McMurdie and Holmes 2013) for further analysis.

Once eukaryota, archaea, chloroplast, and mitochondria reads had been removed, the number of assigned reads for each sample ranged from 15,606 to 29,600. Rarefaction curves were plotted using the rarecurve function in the vegan package (Oksanen et al., 2019), and these indicated that sampling depth was sufficient for all samples, plateauing at around 5000 reads. The negative control did not contain any reads so no adjustments were made to the libraries to account for reagent contamination. To evaluate the mock community results, precision, recall, and F-scores were calculated using the number of genera correctly identified, the number of false positives, and the number of false negatives (Bokulich et al., 2018).

## 2.6. Statistical analyses

All analyses were conducted in RStudio v1.1.456 (RStudio Team 2016) for R v3.5.2 (R Core Team 2018) unless otherwise stated. Soil variables, microbial biomass C, qCO<sub>2</sub>, enzyme activities, and bacterial diversity were analysed using three-way ANOVAs with the aov package to test for effects of oxygen, N addition, PK addition, and their interactions. Box-cox transformations were applied to total P and some enzymes to meet ANOVA assumptions (Tables S1, S3 and S4) and URE activity was transformed with log (x+1). Microbial biomass C and qCO<sub>2</sub> were not improved by transformations so aligned rank ANOVAs were also performed. The outcome did not vary between the two ANOVA methods and we have presented the three-way ANOVA results from the unranked data (Table S1). Post-hoc pairwise contrasts were performed using the LSMmeans Contrast function in JMP v14.3.0 (SAS Institute Inc., Cary, NC, USA) to test for differences between nutrient treatments within the same oxygen treatment. As pH can strongly influence activity of many enzymes (Sinsabaugh et al., 2008), correlations between absolute enzyme activity and soil pH were tested with the cor.test function in R's base stats package using Spearman's rank correlations. Microbial respiration data were analysed with a linear mixed-effects model, using the lme function in R package nlme (Pinheiro et al., 2018), with oxygen, N addition, and PK addition as fixed effects and sample nested within day of measurement as a random effect representing repeated measures within each mesocosm. An ANOVA table was calculated with Wald chi-square tests for the model using the Anova function in the car package (Fox and Weisberg 2011). Soil variables are presented as means

± 1 standard error through the manuscript.

For bacterial diversity analyses, sequencing libraries were rarefied to the lowest depth (11,993). Shannon and Simpson diversity indices were calculated and plotted using the estimate richness and plot richness functions in phyloseq (McMurdie and Holmes 2013). Diversity analyses were repeated on rarefied datasets in which taxa with fewer than 10 or 100 total reads had been removed, but this did not affect the results (data not presented). To assess composition differences between treatments, read counts were transformed to centred log-ratio (Aitchison 1986) using the transform function in the microbiome package (Lahti et al., 2017). Euclidean distances were calculated using the distance package in phyloseq (McMurdie and Holmes 2013) and the adonis function in vegan (Oksanen et al., 2019) was used to perform permutational multivariate analysis of variance (PERMANOVA) with 1000 permutations to test effects of oxygen, N addition, and PK addition on bacterial community composition. Differences between samples were visualised using principal component analysis (PCA) performed with the rda function in vegan (Oksanen et al., 2019). The envfit function in vegan was also used to find significant soil variables to fit to the PCA plot as vectors. Homogeneity of treatment group dispersion (variance) was measured using the betadisper function and differences between groups were tested using the permutest.betadisper function, both in the vegan package (Oksanen et al., 2019).

Indicator analysis was performed using the multipatt function in the indicpecies package (De Cáceres and Legendre 2009) to identify bacterial taxa associated with nutrient addition. This method uses the relative abundance of a taxa and its relative frequency of occurrence within treatment groups to calculate an indicator value index. The significance of the association between the taxa and the treatment group is then tested using permutation tests (De Cáceres and Legendre 2009). To focus on indicators of nutrient effects, the two oxygen treatments were analysed separately. We tested for indicators within each individual nutrient treatment group as well as within specified group combinations (Control with +PK, +N with +NPK, +PK with +NPK). For this analysis, libraries were normalised using cumulative sum scaling (CSS; Paulson et al., 2013) due to the issues associated with using rarefied data for differential abundance analyses (McMurdie and Holmes 2014). This was performed using the phyloseq\_transform\_css function in the metagMisc package (Mikryukov 2019). To exclude rare taxa from the analysis, SVs with fewer than 100 reads or that were prevalent in fewer than two samples were removed. SVs were agglomerated at the genus level, except for SVs which had not been assigned to genus level which were agglomerated at the family level. CSS-normalised abundance heatmaps were produced for identified indicator taxa using the plot\_heatmap function in phyloseq (McMurdie and Holmes 2013).

For enzymes that changed significantly with nutrient addition, the

associate function in the microbiome package (Lahti et al., 2017) was used to cross-correlate CSS-normalised taxa abundances with enzyme activity using Spearman's rank correlations. As soil redox potential has been found to affect activity of some enzymes associated with C, N, and P cycling (Pulford and Tabatabai 1988) it is possible that responses of taxa could vary between oxygen environments. Therefore, the data were separated by oxygen treatment for these correlations. Significance values were corrected for multiple testing using the false discovery rate method (Benjamini and Hochberg 1995) and heatmaps of correlation coefficients significant at the p-adjusted threshold of 0.05 were produced using the heat function in the microbiome package (Lahti et al., 2017).

## 2.7. Accession numbers

Forward and reverse sequence reads have been deposited in the NCBI Sequence Read Archive under BioProject ID PRJNA632865. All other data has been archived in the Mendeley Data repository (DOI:10.17632/86rpthxb7.1).

## 3. Results

### 3.1. Soil analyses

Experimental treatments resulted in significant differences in some of the measured soil variables (Table 2, Table S1). Addition of PK increased total soil P by 7.4% ( $F_{1,32} = 35.90$ ,  $p < 0.001$ ), with no significant difference between the oxygen treatments ( $F_{1,32} = 0.83$ ,  $p = 0.368$ ). There was a significant interaction effect of oxygen treatment and N addition ( $F_{1,32} = 21.23$ ,  $p < 0.001$ ) on total soil N, with N addition resulting in larger N increases in ambient oxygen (+24.7%) compared to reduced oxygen (+10.1%). The oxygen environment ( $F_{1,32} = 136.17$ ,  $p < 0.001$ ) and N addition treatment ( $F_{1,32} = 19.12$ ,  $p < 0.001$ ) had significant effects on soil pH, with the average pH in reduced oxygen mesocosms 12.6% higher than in the ambient, and average pH in N addition mesocosms 4.5% higher than in those without N addition. Total soil C ( $22.52\% \pm 0.16$ ), moisture content ( $82.7\% \pm 0.05$ ), and organic matter content ( $46.50\% \pm 0.12$ ) did not vary significantly with oxygen or nutrient treatments (Table S1).

### 3.2. Microbial biomass and activity

Oxygen levels within the mesocosms had a significant effect on microbial respiration (Table 3) with higher values recorded in the reduced oxygen treatment (Fig. 1). Nutrient additions did not significantly affect microbial respiration (Table S2), but the +NPK treatment tended to have higher respiration rates than the other treatments in the reduced oxygen mesocosms (Fig. 2). This trend was not observed in the ambient oxygen incubations. There was a significant interaction effect of oxygen treatment and N addition ( $F_{1,32} = 10.97$ ,  $p = 0.002$ , Table S1) on microbial biomass, with lower microbial biomass in the N addition treatments compared to the control in the reduced oxygen treatment only (Fig. 2a). When microbial respiration was considered per unit microbial biomass

**Table 2**


Mean ( $\pm 1$  standard error) measures for soil variables at the end of experiment in ambient and reduced oxygen environments with N and PK addition treatments.

Oxygen environment	Nutrient treatment	Total P %	Total N %	Total C %	pH	Organic matter %	Moisture % wet weight
<b>Reduced</b>	Control	0.068 $\pm$ 0.001 a	1.33 $\pm$ 0.02 a	22.52 $\pm$ 0.49	7.69 $\pm$ 0.2 a	46.35 $\pm$ 0.51	82.86 $\pm$ 0.05
	+ PK	0.074 $\pm$ 0.001 b	1.34 $\pm$ 0.02 ab	22.58 $\pm$ 0.29	7.78 $\pm$ 0.15 a	46.17 $\pm$ 0.41	82.82 $\pm$ 0.08
	+ N	0.068 $\pm$ 0.002 a	1.52 $\pm$ 0.04 c	22.83 $\pm$ 0.24	7.95 $\pm$ 0.11 ab	46.67 $\pm$ 0.32	82.76 $\pm$ 0.12
	+ NPK	0.072 $\pm$ 0.001 b	1.42 $\pm$ 0.04 b	21.48 $\pm$ 0.51	8.12 $\pm$ 0.02 b	46.31 $\pm$ 0.2	82.84 $\pm$ 0.08
<b>Ambient</b>	Control	0.068 $\pm$ 0.0003 a	1.35 $\pm$ 0.04 a	22.3 $\pm$ 0.72	6.83 $\pm$ 0.05 a	46.04 $\pm$ 0.25	82.51 $\pm$ 0.15
	+ PK	0.072 $\pm$ 0.002 b	1.37 $\pm$ 0.01 a	22.75 $\pm$ 0.31	6.80 $\pm$ 0.07 a	46.41 $\pm$ 0.29	82.75 $\pm$ 0.32
	+ N	0.067 $\pm$ 0.001 a	1.68 $\pm$ 0.02 b	22.74 $\pm$ 0.33	7.18 $\pm$ 0.08 b	46.39 $\pm$ 0.38	82.61 $\pm$ 0.14
	+ NPK	0.072 $\pm$ 0.0005 b	1.71 $\pm$ 0.04 b	23 $\pm$ 0.41	7.19 $\pm$ 0.05 b	47.18 $\pm$ 0.18	82.71 $\pm$ 0.17


**Table 3**

Three-way ANOVA results testing effects of oxygen environment, N addition, and PK addition on bacterial community diversity in soil, using Shannon and Simpson diversity indices calculated from rarefied 16S rRNA gene sequencing data.

	Shannon		Simpson	
	$F_{1,16}$	p-value	$F_{1,16}$	p-value
Oxygen	59.03	<0.001	16.80	<0.001
N	90.55	<0.001	45.26	<0.001
PK	2.03	0.174	0.01	0.921
Oxygen:N	0.24	0.632	0.34	0.566
Oxygen:PK	2.27	0.152	2.51	0.133
N:PK	0.20	0.661	1.19	0.292
Oxygen:N:PK	0.96	0.342	0.66	0.430



**Fig. 1.** Microbial respiration response to nutrient addition in ambient and reduced oxygen mesocosms. Data displayed are means  $\pm 1$  standard error and are offset across the X axis to minimise overlap.




**Fig. 2.** Effect of oxygen environment and nutrient addition on a) microbial biomass carbon (MBC) and b) microbial metabolic quotient ( $qCO_2$ ) across oxygen environments and nutrient addition treatments. The boxes show the median between the first and third quartiles and the whiskers extend up to 1.5 x the interquartile range.

(i.e. qCO<sub>2</sub>), there was a significant interaction effect of oxygen treatment and N addition ( $F_{1,31} = 20.47, p < 0.001$ , Table S1) with qCO<sub>2</sub> increasing with N addition in the reduced oxygen environment only (Fig. 2b).

When significant effects of nutrient addition were found, posthoc tests were performed to test for differences between nutrient addition treatments within the same oxygen treatment (results indicated by different lowercase letters).

Absolute activities of all enzymes, except for POX and CBH, were significantly lower in the reduced oxygen mesocosms (Fig. 3; Table S3). Nitrogen addition resulted in significant decreases in absolute activity of PER ( $F_{1,32} = 5.22, p = 0.029$ ), URE ( $F_{1,32} = 26.32, p < 0.001$ ), and AP ( $F_{1,32} = 20.76, p < 0.001$ ). The only enzyme whose absolute activity was significantly affected by PK addition was PER ( $F_{1,32} = 1.19, p = 0.042$ ), but post-hoc tests did not find any differences between PK addition treatments and the control in either oxygen environment. Soil pH had significant negative correlations with the absolute activity of PER (Spearman's Rho = -0.393,  $p = 0.012$ ), URE (Spearman's Rho = -0.571,  $p < 0.001$ ), GLC (Spearman's Rho = -0.477,  $p = 0.002$ ), NAG (Spearman's Rho = -0.691,  $p < 0.001$ ), and AP (Spearman's Rho = -0.883,  $p < 0.001$ ). When enzyme activities were considered per unit microbial biomass, there was a significant interaction of oxygen and N addition on specific activities of PER ( $F_{1,31} = 4.76, p = 0.037$ ), POX ( $F_{1,31} = 10.11, p = 0.003$ ), CBH ( $F_{1,31} = 5.64, p = 0.024$ ), GLC ( $F_{1,31} = 11.25, p = 0.002$ ), XYL ( $F_{1,31} = 15.39, p < 0.001$ ), NAG ( $F_{1,31} = 19.81, p < 0.001$ ), and AP ( $F_{1,31} = 12.97, p = 0.001$ ), all of which increased with N addition in the reduced oxygen environment only (Figure S1; Table S4). Specific activity of URE decreased with N addition ( $F_{1,31} = 15.72, p < 0.001$ ) regardless of oxygen environment.



**Fig. 3.** Absolute soil enzyme activity for each treatment group, measured at the end of the experiment. Bars show mean  $\pm$  1 standard error and different lowercase letters indicate significant difference between nutrient treatments within the same oxygen treatment at  $p \leq 0.05$  using post-hoc pairwise contrasts. All values are untransformed data expressed on a dry weight basis.


### 3.3. Soil bacterial community diversity and composition

Sequence variants (SVs) were assigned to 52 unique phyla with three phyla accounting for 75.9% of the reads that were able to be assigned at the Phylum level (Proteobacteria, Bacteroidetes, Epsilonbacteraota; Fig. 4). At the phylum level, N addition increased relative abundance of Epsilonbacteraota in the ambient oxygen mesocosms, and increased relative abundance of Proteobacteria in the reduced oxygen mesocosms. Across all samples, 37.4% of SVs were shared between the ambient and reduced oxygen mesocosms. Within the ambient oxygen mesocosms, 38.7% of SVs were shared between samples with and without N addition. Within the reduced oxygen mesocosms, there were greater differences between the samples with and without N addition, with only 33.7% of shared SVs. The three most abundant genera in the reduced oxygen mesocosms were *Arcobacter* (8.2% of reads), *Sulfurimonas* (5.9%), and *Marinobacter* (5.0%). The three most abundant genera in the ambient oxygen mesocosms were *Arcobacter* (7.7%), *Ignavibacterium* (3.5%), and *Marinobacterium* (3.4%). 68% of SVs were unable to be assigned at genus level. Of the 17 genera expected in the mock community sample, 15 were detected. The two genera that were not detected were *Propionibacterium* and *Actinomyces* and these were only expected in low concentrations. There were no false positives and the F-score was 0.94.


Bacterial diversity was significantly reduced with N addition in both the ambient and reduced oxygen mesocosms (Fig. 5, Table 3), and was lower in the reduced oxygen mesocosms compared to the ambient ones. There were significant effects of oxygen environment and N addition on bacterial community composition, and a significant interaction was detected between the two (Table 4). Oxygen environment and N addition each accounted for ~14% of the variation in bacterial community composition. These effects are illustrated in Fig. 6, where control and +PK samples group together, and +N and +NPK samples group together in both oxygen treatments, which are themselves distinct groups. Total N content ( $R^2 = 0.77, p < 0.001$ ) and pH ( $R^2 = 0.80, p < 0.001$ ) of the soil were both found to have significant associations with the bacterial community composition. There was no difference in multivariate dispersions between the treatment groups ( $F_{7, 16} = 0.93, p = 0.522$ ).

### 3.4. Bacterial indicators

The bacterial taxa that were identified as indicators for treatments with and without N addition are shown in Figs. 7 and 8 (for further details see Tables S5 and S6). Only one taxa was identified as an indicator of a single treatment group (*Shewanella* in +NPK ambient oxygen), the rest were indicators of a combined N addition group (+N and +NPK)



**Fig. 4.** Relative abundances of bacterial phyla across oxygen and nutrient addition treatments. Only phyla with relative abundances above 1% are plotted.




**Fig. 5.** Effects of nutrient addition on the diversity of soil bacterial communities after 15 days incubation in ambient or reduced oxygen environments. Boxplots of Shannon and Simpson Indices determined from rarefied 16S rRNA gene sequencing data. The boxes show the median between the first and third quartiles and the whiskers extend up to 1.5 x the interquartile range.

**Table 4**


Results of Permutational Multivariate Analysis of Variance (PERMANOVA) to test for effects of oxygen environment, N addition, and PK addition on soil bacterial community composition.

	F <sub>1,16</sub>	R <sup>2</sup>	p-value
Oxygen	4.63	0.146	0.001
N	4.53	0.143	0.001
PK	1.20	0.038	0.187
Oxygen:N	1.98	0.038	0.008
Oxygen:PK	1.04	0.033	0.344
N:PK	1.10	0.035	0.279
Oxygen:N:PK	1.14	0.036	0.221




**Fig. 6.** Principal components analysis (PCA) plot of soil bacterial communities from nutrient addition mesocosms incubated in ambient or reduced oxygen environments. PCA calculated from centred log-ratio transformed 16S rRNA sequencing data. Vectors of significant soil variables are plotted with grey arrows.

or a combined no N addition group (Control and +PK). In the ambient oxygen mesocosms, 26 taxa were identified as indicators of samples without N addition (Control and +PK treatments), and 30 taxa as indicators for samples with N addition (+N and +NPK treatments). In the reduced oxygen mesocosms, 35 taxa were identified as indicators of samples without N addition (Control and +PK treatments), and 18 taxa as indicators for samples with N addition (+N and +NPK treatments). No taxa were identified as indicators of a combined PK addition group (the +PK and +NPK treatments) in either oxygen environment. In both oxygen treatments, N addition drives a decrease in the overall abundance (Fig. 9) of the sulfate-reducing order Desulfobacterales (Kuever




**Fig. 7.** Heatmap showing cumulative sum scaling (CSS) normalised abundances of soil bacteria indicator taxa identified for treatments with and without N addition in ambient oxygen mesocosms. Only taxa significant at  $p \leq 0.01$  are shown. For taxa where genus could not be assigned, the family is shown instead.



**Fig. 8.** Heatmap showing cumulative sum scaling (CSS) normalised abundances of soil bacteria indicator taxa identified for treatments with and without N addition in reduced oxygen incubations. Only taxa significant at  $p \leq 0.01$  are shown. For taxa where genus could not be assigned, the family is shown instead.

et al., 2005), with eight taxa in this order identified as indicators of the group without N addition in the reduced oxygen and six taxa in the ambient oxygen. The N addition group saw an increase in abundance of several taxa of unknown function from the order Oceanospirillales (three taxa in reduced oxygen and five taxa in ambient oxygen). The only N-cycling related taxa that we identified amongst the indicators, was *Nitrocooccus*, a nitrifying bacterium (Watson and Waterbury 1971)






**Fig. 9.** Total abundances for bacteria from the sulfate-reducing order Desulfobacterales within each nutrient addition treatment group in ambient and reduced oxygen mesocosms. Each section within a bar is the number of sequencing reads of the 16S rRNA gene for one sample using rarefied data.

that increased abundance with N addition in the ambient oxygen treatment.

### 3.5. Correlations between bacterial abundance and enzyme activities

In the ambient oxygen mesocosms, no significant correlations were found between abundances of bacterial genera and absolute soil enzyme activities. In the reduced oxygen mesocosms, 33 taxa significantly correlated with URE, nine taxa correlated with AP, and only one taxa correlated with PER activity (Fig. 10). Six taxa significantly correlated with both URE and AP, and all of these were negative correlations. There was some overlap between the indicator taxa and those that significantly correlated with enzyme activities. Eleven of the indicator taxa for the reduced oxygen treatments with N, had significant negative correlations with URE activity, and nine of the indicator taxa for the reduced oxygen treatments without N had significant positive correlations with URE.



**Fig. 10.** Heatmap showing Spearman correlation coefficients for taxa that significantly correlated (FDR  $p$  adjusted value  $\leq 0.05$ ) with absolute enzyme activity of lignin peroxidase (PER), urease (URE), or acid phosphatase (AP) in reduced oxygen mesocosms. For taxa where genera could not be assigned, the lowest assigned taxonomic rank is shown. \* also identified as indicator taxa.

## 4. Discussion

Nitrogen addition drove clear shifts in bacterial community composition, and as hypothesised, resulted in decreased bacterial diversity and decreased abundance of sulfate-reducing bacteria. These changes in bacterial community structure were accompanied by changes in the activity of some enzymes involved in C, N, and P cycling. Although nutrient addition had no effect on overall microbial respiration, N addition decreased microbial biomass and increased microbial metabolic quotient in the reduced oxygen environment. The addition of PK alone did not have any significant effects on bacterial diversity or composition, and soil functions were not altered when P and K were combined with N. Overall, these results indicate that N addition causes shifts in the diversity and composition of mangrove soil bacterial communities with potential consequences for nutrient cycling and decomposition processes.

In contrast to other mangrove soil nutrient addition experiments, where microbial respiration increased with nutrient addition (Chen et al., 2011; Keuskamp et al., 2013; Davies et al., 2017), we did not see significant responses of microbial respiration to either N or PK. However, when respiration was considered per unit of microbial biomass (ie.  $qCO_2$ ), an increase in  $qCO_2$  with N addition was detected in the reduced oxygen environment, suggesting that it was the observed decrease in microbial biomass which contributed to the lack of an overall effect. This decrease in soil microbial biomass with N addition is consistent with the findings of many other studies across a range of biomes, which can have corresponding decreases in soil  $CO_2$  emissions (Treseder 2008). Our findings suggest a change in the physiological status of the microbial community, with increased  $qCO_2$  indicating reduced C use efficiency with N addition in the reduced oxygen soils. In field conditions, we must also consider the indirect effects of nutrient addition on microbial respiration and biomass through increased plant productivity leading to an increased supply of C to the soil microbial community via exudates (Uselman et al., 2000). While plant growth showed strong positive responses to N addition at the field site related to this study (Dangremond et al., 2019), microbial responses to nutrient addition in this incubation study were not so straightforward. Contrasting nutrient limitations are often observed between plants and their soil microbes, with organic carbon typically being the dominant limitation for secondary productivity by heterotrophic microbial communities (Soong et al., 2000). While mangrove soils are often rich in organic carbon, previous studies indicate that their microbial communities can be both C and N limited, with Chen et al. (2011) and Keuskamp et al. (2013) only detecting increases in microbial respiration with nutrient addition when a labile C source was also added.

Nitrogen addition generally suppresses activity of lignin-modifying enzymes such as lignin peroxidase (PER) and phenol oxidase (POX) leading to increases in soil C storage (Chen et al., 2018a,b). Here, N addition suppressed absolute activity of PER, but did not affect absolute activity of POX. Our lack of response in absolute POX activity corresponds with the findings of another N addition incubation experiment of mangrove soils by Luo et al. (2017). We did, however, find an increase in specific activity of POX with N addition under reduced oxygen, and this is likely due to lower microbial biomass in the N addition treatments. Nitrogen addition has also been associated with increased activity of cellulases such as  $\beta$ -glucosidase (GLC),  $\beta$ -xylosidase (XYL), and cellobiohydrolase (CBH) (Chen et al., 2018a,b) but no changes in absolute activity were observed in these three enzymes in this study. Previous N addition incubation experiments with mangrove soils have observed either an increase (Keuskamp et al., 2015) or a decrease (Luo et al., 2017) in absolute GLC activity. Despite the lack of discernible response in the absolute activity of GLC in our study, significant increases were observed in the specific activity of GLC under reduced oxygen. Similar to what was observed with microbial respiration, these findings suggest N addition may have resulted in a physiological response at the microbial level but the accompanying decrease in microbial biomass contributed

to the lack of an overall effect of N addition on GLC activity within the reduced oxygen mesocosms. The variable responses observed here in enzymes involved in C cycling make it difficult to predict how carbon storage rates might be affected.

Under the resource allocation model (Sinsabaugh and Moorhead 1994), N addition should result in decreased activity of N acquiring enzymes. A marked decrease in absolute activity of urease (URE) was observed with N addition in the reduced oxygen treatment, and while the same pattern was observed in the ambient oxygen treatment, the response was more variable. In a previous experiment with mangrove soils (Chen et al., 2015), N addition was found to have a stimulatory effect on soil URE activity (Chen et al., 2015). However, it should be noted that in Chen et al.'s (2015) study the positive effect of N was only observed in soils that had plants growing in them, there was no effect of N on URE activity in the non-planted soils. We observed no change in absolute activity of *N*-acetyl- $\beta$ -glucosaminidase (NAG), but specific activity of NAG was higher with N under reduced oxygen. Nitrogen addition has been found to increase both absolute and specific activity of NAG in waterlogged rice paddies (Zhang et al., 2015) but in other incubation experiments with mangrove soils, absolute activity of NAG has either decreased (Luo et al., 2017) or been unaffected (Keuskamp et al., 2015). Keuskamp et al. (2015) also assessed activities of two other N acquiring enzymes: leucine aminopeptidase (LAP) and glycine aminopeptidase (GAP). They saw increased activity of LAP with N addition in incubation experiments but not in the field, and GAP did not respond to N addition in the field or the incubations (Keuskamp et al., 2015). These conflicting results amongst N-acquiring enzymes support the findings from Sinsabaugh and Follstad Shah's (2012) review paper covering a range of ecosystems that there is no simple relationship between N availability and N acquiring extracellular enzyme activity.

Contrary to our hypothesis and findings of other studies in terrestrial (Marklein 2012) and mangrove soils (Luo et al., 2017), absolute activity of the P acquiring enzyme acid phosphatase (AP) decreased with N addition rather than increased. We found a strong negative correlation between pH and AP activity, which we think accounts for the results that we observed, as pH significantly increased with N addition in our experiment. PER and URE also had significant negative correlations with pH suggesting that some of the enzyme responses that we are seeing could be driven by pH changes in the soil. This is similar to the findings of Liu and Zhang (2019), who found changes in enzyme activity with N addition were related more to changes in soil pH than changes in plant or microbial community composition, although in their study pH decreased rather than increased with N addition. Globally, pH tends to decrease with N addition, rather than increase as observed here, although responses can vary across ecosystem types (Tian and Niu 2015). Similar to our study, Luo et al. (2017) also observed an increase in pH with N addition in their incubation experiment with mangrove soils. We suggest that this increase, as well as the differences in pH between the oxygen treatments, could be related to the unique redox conditions found in wetland soils and the amount of oxidisable constituents present (D'Angelo and Reddy 2003). We suspect that pH changes of the magnitude observed in this incubation experiment would be unlikely in surface soils in the field due to regular tidal flushing. Significant effects of oxygen environment on absolute activity were observed in all enzymes except POX and CBH. This is not unexpected as many studies have found that waterlogging and redox potential can have significant effects on soil enzyme activities (eg. Pulford and Tabatabai 1988; Wang and Lu 2006; Vo and Kang 2013). Our results highlight the complexity of enzyme responses to nutrient amendment, and further research is required to assess enzyme responses and nutrient cycling processes under field conditions.

As hypothesised, soil bacterial diversity, measured by Shannon and Simpson diversity indices, decreased with N addition. Our findings are

supported by other studies within mangroves (Wang et al., 2016), marine sediments (Aoyagi et al., 2015), and other terrestrial environments (Wang et al., 2018). Similar diversity decreases with N addition have also been observed in fungal (Wang et al., 2018) and plant communities (Midolo et al., 2019). There is currently no consensus on what drives this decrease in diversity, but it could be related to changes in competitive interactions, with fast-growing species that are able to take advantage of increased N availability outcompeting slower growing oligotrophic species adapted to low nutrient conditions (Leff et al., 2015). Other studies have suggested that decreases in bacterial diversity with N are indirectly affected by changes in soil pH or plant communities (Zeng et al., 2016; Zhang et al., 2017). The loss of soil microbial diversity can lead to a reduction in soil functioning (Maron et al., 2018; Trivedi et al., 2019) so when considered alongside observed decreases in microbial biomass and the activity of some enzymes (e.g. PER, URE, AP), this suggests that nutrient cycling rates could be altered by N addition in this system.

Bacterial community composition was significantly affected by N addition. *Arcobacter* was the most abundant genus and an indicator taxa of N addition in both oxygen treatments. Species within this genus have previously been identified as indicators of fecal pollution in environmental water samples (Collado et al., 2008). This suggests that the mangrove site from which these soils were sourced may already be experiencing anthropogenic derived nutrient pollution but that nutrient concentrations were not saturated as N addition led to increased *Arcobacter* abundance. In both the ambient and reduced oxygen treatments, N addition resulted in decreased abundance of several genera within the sulfate-reducing order Desulfobacterales, and increased abundance of several genera from Oceanospirillales. This same trend was also reported in a nitrate-addition incubation study of saltmarsh sediment (Bulsecu et al., 2019), which found that although sulfate-reduction remained the dominant process, a shift towards nitrate-reduction allowed for decomposition of organic matter that had previously been inaccessible in sulfate-reducing conditions. Bulsecu et al. (2019) suggested that N induced changes in bacterial community composition could potentially increase decomposition rates as new pools of organic matter become available. While litter decomposition rates over six months did not vary with nutrient addition at the field experiment site associated with our study (Simpson et al., 2020), decomposition rates of older, more recalcitrant carbon sources have yet to be assessed. In contrast to previous studies (Aoyagi et al., 2015; Wang et al., 2016; Bulsecu et al., 2019) we did not observe increased abundance of nitrate-reducing bacteria such as *Sulfurimonas*. This genus actually decreased in abundance with N addition in the reduced oxygen treatment. Of the bacterial taxa whose abundance increased in our study, their functions were generally unknown, so we were unable to confirm a shift towards nitrate-reduction in our study. Further work is needed in the field to assess changes in soil bacterial community structure and function and their potential impacts on nutrient cycling and decomposition in the longer term.

Changes in bacterial community composition drove changes in the activity of some enzymes associated with N addition. PER activity only correlated with abundance changes in one taxa, but 37 genera were identified whose relative abundance correlated with changes in URE and/or AP activity. As correlation does not imply causation, further work is needed to confirm direct linkages between the function of each of these taxa and their response to nutrient addition. Due to the lack of available functional information for many mangrove bacterial taxa, we support the suggestion of Trivedi et al. (2016) that abundances of functional genes may prove more useful than taxonomic abundance for predicting changes in enzyme activity and soil processes.

While this study has improved our understanding of microbial responses to nutrient addition in mangrove soils, there were some

limitations associated with this study that may have influenced the findings. Microbial respiration was significantly lower in the ambient oxygen treatment compared to the reduced oxygen mesocosms. As microbial respiration is generally lower in anaerobic conditions (Reddy and DeLaune, 2008), we suggest that the high soil moisture content in the mesocosms provided a poor environment for aerobic microbes, but that the oxygen levels in the ambient mesocosms were sufficient to inhibit obligate anaerobic microbes. As such, responses of microbes in our ambient mesocosms to N addition may have been muted in comparison with the responses observed in the reduced oxygen mesocosms. Future mangrove incubation experiments could potentially improve aerobic conditions by increasing the soil surface area in the mesocosms relative to volume. Addition of PK had no significant effect on any of the measured microbial response variables. This could be because the microbial community in this soil was not P limited, but it could also be that the amount of P added was insufficient to induce a microbial response. This study targeted bacteria for community characterisation but the microbial activities measured would include responses from other components of the microbiome, such as archaea and fungi. Further research is needed to determine how these different kingdoms contribute to the overall response. We acknowledge that soil disturbance during the experimental setup may have altered bacterial community composition in the mesocosms compared to the field. Future nutrient incubation experiments may wish to consider the use of intact cores or characterising the microbial community at different time points, over a longer duration, to track community changes and resilience.

## 5. Conclusions

This study highlights the potential vulnerabilities of mangrove microbial communities to N enrichment. We found that N addition caused significant shifts in the diversity and composition of soil bacterial communities and impacts on microbial biomass and functioning. This has potential implications for C and nutrient cycling which warrant further investigation in the field. While many mangrove habitats have already been subjected to increasing nutrient levels over the last few decades, additional nutrients may not elicit much response. However, some relatively pristine habitats still remain whose water catchments will see increased human impacts in the coming years. The findings from this study contribute to a better understanding of the effects of elevated nutrients on mangrove soil microbial communities which could help focus management of these environments more effectively.

## Author contributions

HC, LTS, and TZO conceived the study, and HC and LTS designed and conducted the experiment. HC, IC, and DA performed nutrient analyses and enzyme assays, HC and REA performed microbial sequencing work, and HC analysed the data. JKR, RDB, CHR, and TZO supervised the project and provided guidance on the framing of questions and interpretation of findings. HC prepared the manuscript with all authors contributing to the drafts.

## Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Acknowledgements

The study contained in this publication was supported by Natural Environment Research Council (NERC) EAO Doctoral Training Partnership (NE/L002469/1) funding to H. Craig, and I. Cordero was supported by a Ramon Areces Foundation Research Fellowship and BBSRC Discovery Fellowship (BB/S010661/1). Soil samples were transferred to

the UK for nutrient and enzyme analyses under DEFRA licence number 116619/225604/2. We wish to thank Prof. Richard Preziosi for his invaluable assistance with statistical analyses, Anna Thornton for her assistance with CO<sub>2</sub> flux measurements, and Guana Tolomato Matanzas National Estuarine Research Reserve for their permission to conduct research at the site.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.soilbio.2020.108076>.

## References

- Aitchison, J., 1986. *The Statistical Analysis of Compositional Data*, Monographs on Statistics and Applied Probability. Chapman & Hall Ltd., London.
- Alongi, D.M., 2012. Carbon sequestration in mangrove forests. *Carbon Management* 3 (3), 313–322.
- Anderson, T.-H., Domsch, K.H., 1985. Determination of ecophysiological maintenance carbon requirements of soil microorganisms in a dormant state. *Biology and Fertility of Soils* 1, 81–89.
- Aoyagi, T., Kimura, M., Yamada, N., Navarro, R.R., Itoh, H., Ogata, A., Sakoda, A., Katayama, Y., Takasaki, M., Hori, T., 2015. Dynamic transition of chemolithotrophic sulfur-oxidizing bacteria in response to amendment with nitrate in deposited marine sediments. *Frontiers in Microbiology* 6, 426.
- Benjamini, Y., Hochberg, Y., 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B* 57 (1), 289–300.
- Bokulich, N.A., Kaehler, B.D., Rideout, J.R., Dillon, M., Bolyen, E., Knight, R., Huttley, G. A., Caporaso, J.G., 2018. Optimizing taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier plugin. *Microbiome* 6, 90.
- Booth, J.M., Fusi, M., Marasco, R., Mboho, T., Daffonchio, D., 2019. Fiddler crab bioturbation determines consistent changes in bacterial communities across contrasting environmental conditions. *Scientific Reports* 9, 3749.
- Bulseco, A.N., Giblin, A.E., Tucker, J., Murphy, A.E., Sanderman, J., Hiller-Bittrolf, K., Bowen, J.L., 2019. Nitrate addition stimulates microbial decomposition of organic matter in salt marsh sediments. *Global Change Biology* 25 (10), 3224–3241.
- Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes, S.P., 2016. DADA2: high-resolution sample inference from Illumina amplicon data. *Nature Methods* 13 (7), 581–583.
- Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Lozupone, C.A., Turnbaugh, P.J., Fierer, N., Knight, R., 2011. Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proceedings of the National Academy of Sciences* 108 (1), 4516–4522.
- Chen, G.C., Tam, N.F.Y., Wong, Y.S., Ye, Y., 2011. Effect of wastewater discharge on greenhouse gas fluxes from mangrove soils. *Atmospheric Environment* 45 (5), 1110–1115.
- Chen, J., Zhou, H.C., Wang, C., Zhu, C.Q., Tam, N.F.-Y., 2015. Short-term enhancement effect of nitrogen addition on microbial degradation and plant uptake of polybrominated diphenyl ethers (PBDEs) in contaminated mangrove soil. *Journal of Hazardous Materials* 300, 84–92.
- Chen, Q., Zhao, Q., Li, J., Jian, S., Ren, H., 2016. Mangrove succession enriches the sediment microbial community in South China. *Scientific Reports* 6, 27468.
- Chen, H., Li, D., Zhao, J., Xiao, K., Wang, K., 2018. Effects of nitrogen addition on activities of soil nitrogen acquisition enzymes: a meta-analysis. *Agriculture, Ecosystems & Environment* 252, 126–131.
- Chen, J., Luo, Y., Groenigen, K.J. Van, Hungate, B.A., Cao, J., Zhou, X., Wang, R., 2018. A keystone microbial enzyme for nitrogen control of soil carbon storage. *Science Advances* 4 (8), eaq1689.
- Collado, L., Inza, I., Guarro, J., Figueras, M.J., 2008. Presence of *Arcobacter* spp. in environmental waters correlates with high levels of fecal pollution. *Environmental Microbiology* 10 (6), 1635–1640.
- Cordero, I., Snell, H., Bardgett, R.D., 2019. High throughput method for measuring urease activity in soil. *Soil Biology and Biochemistry* 134, 72–77.
- D'Angelo, E., Reddy, K.R., 2003. Effect of aerobic and anaerobic conditions on chlorophenol sorption in wetland soils. *Soil Science Society of America Journal* 67 (3), 787.
- Dangremond, E.M., Simpson, L.T., Osborne, T.Z., Feller, I.C., 2019. Nitrogen enrichment accelerates mangrove range expansion in the temperate-tropical ecotone. *Ecosystems*. <https://doi.org/10.1007/s10021-019-00441-2>.
- Davies, T.K.R., Lovelock, C.E., Pettit, N.E., Grierson, P.F., 2017. Short-term microbial respiration in an arid zone mangrove soil is limited by availability of gallic acid, phosphorus and ammonium. *Soil Biology and Biochemistry* 115, 73–81.
- De Cáceres, D.M., Legendre, P., 2009. Associations between species and groups of sites: indices and statistical inference. *Ecology* 90 (12), 3566–3574.
- De Long, J.R., Semchenko, M., Pritchard, W.J., Cordero, I., Fry, E.L., Jackson, B.G., Kurnosova, K., Ostle, N.J., Johnson, D., Baggs, E.M., Bardgett, R.D., 2019. Drought soil legacy overrides maternal effects on plant growth. *Functional Ecology* 33 (8), 1400–1410.
- Donato, D.C., Kauffman, J.B., Murdiyarso, D., Kurnianto, S., Stidham, M., Kanninen, M., 2011. Mangroves among the most carbon-rich forests in the tropics. *Nature Geoscience* 4, 293.

- Feller, I.C., 1995. Effects of nutrient enrichment on growth and herbivory of dwarf Red Mangrove (*Rhizophora mangle*). *Ecological Monographs* 65 (4), 477–505.
- Feller, I.C., Whigham, D.F., O'Neill, J.P., McKee, K.L., 1999. Effects of nutrient enrichment on within-stand cycling in a mangrove forest. *Ecology* 80 (7), 2193–2205.
- Feller, I.C., Whigham, D.F., McKee, K.L., Lovelock, C.E., 2003. Nitrogen limitation of growth and nutrient dynamics in a disturbed mangrove forest, Indian River Lagoon, Florida. *Oecologia* 134, 405–414.
- Feng, J., Zhu, B., 2019. A global meta-analysis of soil respiration and its components in response to phosphorus addition. *Soil Biology and Biochemistry* 135, 38–47.
- Fox, J., Weisberg, S., 2011. *An R Companion to Applied Regression*, second ed. Thousand Oaks, Sage, CA.
- Giri, C., Long, J., 2016. Is the geographic range of mangrove forests in the conterminous United States really expanding? *Sensors* 16 (12), 2010.
- Griffiths, S.M., Harrison, X.A., Weldon, C., Wood, M.D., Pretorius, A., Hopkins, K., Fox, G., Preziosi, R.F., Antwis, R.E., 2018. Genetic variability and ontogeny predict microbiome structure in a disease-challenged montane amphibian. *The ISME Journal* 12, 2506–2517.
- Holguin, G., Vazquez, P., Bashan, Y., 2001. The role of sediment microorganisms in the productivity, conservation, and rehabilitation of mangrove ecosystems: an overview. *Biology and Fertility of Soils* 33 (4), 265–278.
- Jackson, C.R., Tyler, H.L., Millar, J.J., 2013. Determination of microbial extracellular enzyme activity in waters, soils, and sediments using high throughput microplate assays. *Journal of Visualized Experiments* 80, 50399.
- Joergensen, R.G., Wu, J., Brookes, P.C., 2011. Measuring soil microbial biomass using an automated procedure. *Soil Biology and Biochemistry* 43 (5), 873–876.
- Keuskamp, J.A., Schmitt, H., Laanbroek, H.J., Verhoeven, J.T.A., Hefting, M.M., 2013. Nutrient amendment does not increase mineralisation of sequestered carbon during incubation of a nitrogen limited mangrove soil. *Soil Biology and Biochemistry* 57, 822–829.
- Keuskamp, J.A., Hefting, M.M., Dingemans, B.J.J., Verhoeven, J.T.A., Feller, I.C., 2015a. Effects of nutrient enrichment on mangrove leaf litter decomposition. *The Science of the Total Environment* 508, 402–410.
- Keuskamp, J.A., Feller, I.C., Laanbroek, H.J., Verhoeven, J.T.A., Hefting, M.M., 2015b. Short- and long-term effects of nutrient enrichment on microbial exoenzyme activity in mangrove peat. *Soil Biology and Biochemistry* 81, 38–47.
- Koerselman, W., Meuleman, A.F.M., 1996. The vegetation N:P ratio: a new tool to detect the nature of nutrient limitation. *Journal of Applied Ecology* 33 (6), 1441–1450.
- Kozich, J.J., Westcott, S.L., Baxter, N.T., Highlander, S.K., Schloss, P.D., 2013. Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the miseq illumina sequencing platform. *Applied and Environmental Microbiology* 79 (17), 5112–5120.
- Kreuzwieser, J., Buchholz, J., Rennenberg, H., 2003. Emission of methane and nitrous oxide by Australian mangrove ecosystems. *Plant biology* 5 (4), 423–431.
- Kristensen, E., Holmer, M., Bussarawit, N., 1991. Benthic metabolism and sulfate reduction in a southeast Asian mangrove swamp. *Marine Ecology Progress Series* 73 (1), 93–103.
- Kuever, J., Rainey, F.A., Widdel, F., 2005. Order III. Desulfobacterales ord. nov. The Proteobacteria, part C (The Alpha-, Beta-, Delta-, and Epsilonproteobacteria. In: Brenner, D.J., Krieg, N.R., Staley, J.T., Garrity, G.M. (Eds.), *Bergey's Manual of Systematic Bacteriology*, second ed., vol. 2. Springer, New York, p. 959.
- Kuo, S., 1996. Phosphorus. In: Sparks, D.L. (Ed.), *Methods of Soil Analysis: Part 3 – Chemical Methods*. Soil Science Society of America Book Series No. 5. Soil Science Society of America, Madison, pp. 869–919.
- Lahti, L., Shetty, S., Blake, T., Salojärvi, J., 2017. *Microbiome R package*. <http://microbiome.github.io>.
- Leff, J.W., Jones, S.E., Prober, S.M., Barberán, A., Borer, E.T., Firm, J.L., Harpole, W.S., Hobbie, S.E., Hofmöckel, K.S., Knops, J.M.H., McCulley, R.L., La Pierre, K., Risch, A. C., Seabloom, E.W., Schütz, M., Steenbock, C., Stevens, C.J., Fierer, N., 2015. Consistent responses of soil microbial communities to elevated nutrient inputs in grasslands across the globe. *Proceedings of the National Academy of Sciences* 112 (35), 10967–10972.
- Liu, X., Zhang, S., 2019. Nitrogen addition shapes soil enzyme activity patterns by changing pH rather than the composition of the plant and microbial communities in an alpine meadow soil. *Plant and Soil* 440 (1–2), 11–24.
- Lovelock, C.E., Feller, I.C., McKee, K.L., Engelbrecht, B.M.J., Ball, M.C., 2004. The effect of nutrient enrichment on growth, photosynthesis and hydraulic conductance of dwarf mangroves in Panama. *Functional Ecology* 18 (1), 25–33.
- Luo, L., Meng, H., Wu, R., Gu, J., 2017. Impact of nitrogen pollution/deposition on extracellular enzyme activity, microbial abundance and carbon storage in coastal mangrove sediment. *Chemosphere* 177, 275–283.
- Marklein, A.R., Houlton, B.Z., 2012. Nitrogen inputs accelerate phosphorus cycling rates across a wide variety of terrestrial ecosystems. *New Phytologist* 193 (3), 696–704.
- Maron, P., Sarr, A., Kaisermann, A., Lévêque, J., Mathieu, O., Guigues, J., Karimi, B., Bernard, L., Dequiedt, S., Terrat, S., Chabbi, A., Ranjard, L., 2018. High microbial diversity promotes soil ecosystem functioning. *Applied and Environmental Microbiology* 84 (9), e02738-17.
- McKee, K.L., Cahoon, D.R., Feller, I.C., 2007. Caribbean mangroves adjust to rising sea level through biotic controls on change in soil elevation. *Global Ecology and Biogeography* 16 (5), 545–556.
- McMurdie, P.J., Holmes, S., 2013. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS One* 8 (4), e61217.
- McMurdie, P.J., Holmes, S., 2014. Waste not, want not: why rarefying microbiome data is inadmissible. *PLoS Computational Biology* 10 (4), e1003531.
- Midolo, G., Alkemade, R., Schipper, A.M., Benítez-López, A., Perring, M.P., De Vries, W., 2019. Impacts of nitrogen addition on plant species richness and abundance: a global meta-analysis. *Global Ecology and Biogeography* 28, 398–413.
- Mikryukov, V., 2019. *metagMisc: miscellaneous functions for metagenomic analysis*. <https://rdr.io/github/vmik/metagMisc/>.
- Moorhead, D.L., Rinkes, Z.L., Sinsabaugh, R.L., Weintraub, M.N., 2013. Dynamic relationships between microbial biomass, respiration, inorganic nutrients and enzyme activities: informing enzyme-based decomposition models. *Frontiers in Microbiology* 4, 223.
- Muñoz-Hincapié, M., Morell, J.M., Corredor, J.E., 2002. Increase of nitrous oxide flux to the atmosphere upon nitrogen addition to red mangroves sediments. *Marine Pollution Bulletin* 44 (10), 992–996.
- Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGinn, D., Minchin, P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szocs, E., Wagner, H., 2019. *Vegan: community ecology package*. <https://cran.r-project.org/package=vegan>.
- Paulson, J.N., Stine, O.C., Bravo, H.C., Pop, M., 2013. Differential abundance analysis for microbial marker-gene surveys. *Nature Methods* 10, 1200–1202.
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., R Core Team, 2018. *Nlme: Linear and Nonlinear Mixed Effects Models*. R Package Version 3.1-137. <https://CRAN.R-project.org/package=nlme>.
- Pulford, I.D., Tabatabai, M.A., 1988. Effect of waterlogging on enzyme activities in soils. *Soil Biology and Biochemistry* 20 (2), 215–219.
- Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glöckner, F.O., 2013. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Research* 41, D590–D596.
- R Core Team, 2018. *R: a language and environment for statistical computing*. <https://www.r-project.org/>.
- RStudio Team, 2016. *RStudio. Integrated Development for R*. RStudio, Inc., Boston, MA. <http://www.rstudio.com/>.
- Ramirez, K.S., Craine, J.M., Fierer, N., 2012. Consistent effects of nitrogen amendments on soil microbial communities and processes across biomes. *Global Change Biology* 18 (6), 1918–1927.
- Reddy, K.R., DeLaune, R.D., 2008. *Biogeochemistry of Wetlands: Science and Applications*. CRC Press, Boca Raton, pp. 160–167.
- Simpson, L.T., Lovelock, C.E., Cherry, J.A., Feller, I.C., 2020. Short-lived effects of nutrient enrichment on *Avicennia germinans* decomposition in a saltmarsh-mangrove ecotone. *Estuarine, Coastal and Shelf Science* 235, 106598.
- Sinsabaugh, R.L., Moorhead, D.L., 1994. Resource allocation to extracellular enzyme production: a model for nitrogen and phosphorus control of litter decomposition. *Soil Biology and Biochemistry* 26 (10), 1305–1311.
- Sinsabaugh, R.L., Lauber, C.L., Weintraub, M.N., Ahmed, B., Allison, S.D., Crenshaw, C., Contosta, A.R., Cusack, D., Frey, S., Gallo, M.E., Gartner, T.B., Hobbie, S.E., Holland, K., Keeler, B.L., Powers, J.S., Stursova, M., Takacs-Vesbach, C., Waldrop, M. P., Wallenstein, M.D., Zak, D.R., Zeglin, L.H., 2008. Stoichiometry of soil enzyme activity at global scale. *Ecology Letters* 11 (11), 1252–1264.
- Sinsabaugh, R.L., Follstad Shah, J.J., 2012. Eoenzymatic stoichiometry and ecological theory. *Annual Review of Ecology Evolution and Systematics* 43, 313–343.
- Smith, V.H., Tilman, G.D., Nekola, J.J., 1999. Eutrophication: impacts of excess nutrient inputs on freshwater, marine, and terrestrial ecosystems. *Environmental Pollution* 100 (1–3), 179–196.
- Smith, V.H., 2003. Eutrophication of freshwater and coastal marine ecosystems: a global problem. *Environmental Science and Pollution Research* 10, 126–139.
- Soong, J.L., Fuchslueger, L., Marañon-Jimenez, S., Torn, M.S., Janssens, I.A., Penuelas, J., Richter, A., 2020. Microbial carbon limitation: the need for integrating microorganisms into our understanding of ecosystem carbon cycling. *Global Change Biology* 26, 1953–1961.
- Tam, N.F., 1998. Effects of wastewater discharge on microbial populations and enzyme activities in mangrove soils. *Environmental Pollution* 102 (2–3), 233–242.
- Tian, D., Niu, S., 2015. A global analysis of soil acidification caused by nitrogen addition. *Environmental Research Letters* 10, 024019.
- Treseder, K.K., 2008. Nitrogen additions and microbial biomass: a meta-analysis of ecosystem studies. *Ecology Letters* 11 (10), 1111–1120.
- Trivedi, C., Delgado-Baquerizo, M., Hamonts, K., Lai, K., Reich, P.B., Singh, B.K., 2019. Losses in microbial functional diversity reduce the rate of key soil processes. *Soil Biology and Biochemistry* 135, 267–274.
- Uelman, S.M., Qualls, R.G., Thomas, R.B., 2000. Effects of increased atmospheric CO<sub>2</sub> temperature, and soil N availability on root exudation of dissolved organic carbon by a N-fixing tree (*Robinia pseudoacacia* L.). *Plant and Soil* 222, 191–202.
- Vance, E.D., Brookes, P.C., Jenkinson, D.S., 1987. An extraction method for measuring soil microbial biomass C. *Soil Biology and Biochemistry* 19 (6), 703–707.
- Vo, N.X.Q., Kang, H., 2013. Regulation of soil enzyme activities in constructed wetlands under a short-term drying period. *Chemistry and Ecology* 29 (2), 146–165.
- Wang, C., Liu, D., Bai, E., 2018. Decreasing soil microbial diversity is associated with decreasing microbial biomass under nitrogen addition. *Soil Biology and Biochemistry* 120 (72), 126–133.
- Wang, L., Huang, X., Zheng, T.L., 2016. Responses of bacterial and archaeal communities to nitrate stimulation after oil pollution in mangrove sediment revealed by Illumina sequencing. *Marine Pollution Bulletin* 109 (1), 281–289.
- Wang, X.-C., Lu, Q., 2006. Effect of waterlogged and aerobic incubation on enzyme activities in paddy soil. *Pedosphere* 16 (4), 532–539.
- Watson, S.W., Waterbury, J.B., 1971. Characteristics of two marine nitrite oxidizing bacteria, *Nitrospina gracilis* nov. gen. nov. sp. and *Nitrococcus mobilis* nov. gen. nov. sp. *Archiv für Mikrobiologie* 77, 203–230.

- Witt, C., Gaunt, J.L., Calicia, C.C., Ottow, J.C.G., Neue, H.-U., 2000. A rapid chloroform-fumigation extraction method for measuring soil microbial biomass carbon and nitrogen in flooded rice soils. *Biology and Fertility of Soils* 30, 510–519.
- Xiao, W., Chen, X., Jing, X., Zhu, B., 2018. A meta-analysis of soil extracellular enzyme activities in response to global change. *Soil Biology and Biochemistry* 123, 21–32.
- Yue, K., Yang, W., Peng, Y., Peng, C., Tan, B., Xu, Z., Zhang, L., Ni, X., Zhou, W., Wu, F., 2018. Individual and combined effects of multiple global change drivers on terrestrial phosphorus pools: a meta-analysis. *The Science of the Total Environment* 630, 189–202.
- Zeng, J., Liu, X., Song, L., Lin, X., Zhang, H., Shen, C., Chu, H., 2016. Nitrogen fertilization directly affects soil bacterial diversity and indirectly affects bacterial community composition. *Soil Biology and Biochemistry* 92, 41–49.
- Zhang, Y., Shen, H., He, X., Thomas, B.W., Lupwayi, N.Z., Hao, X., Thomas, M.C., Shi, X., 2017. Fertilization shapes bacterial community structure by alteration of soil pH. *Frontiers in Microbiology* 8, 1325.
- Zhou, L., Zhou, X., Zhang, B., Lu, M., Luo, Y., Liu, L., Li, B., 2014. Different responses of soil respiration and its components to nitrogen addition among biomes: a meta-analysis. *Global Change Biology* 20 (7), 2332–2343.