


Please cite the Published Version

Altakroni, B, Nevin, C, Carroll, M , Murgatroyd, C, Horne, G, Brison, DR and Povey, AC (2021) The marker of alkyl DNA base damage, N7-methylguanine, is associated with semen quality in men. Scientific Reports, 11. ISSN 2045-2322

DOI: <https://doi.org/10.1038/s41598-021-81674-x>

Publisher: Springer Nature

Version: Published Version

Downloaded from: <https://e-space.mmu.ac.uk/627224/>

Usage rights:  [Creative Commons: Attribution 4.0](https://creativecommons.org/licenses/by/4.0/)

Additional Information: Open Access article, copyright The Authors.

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)



OPEN

The marker of alkyl DNA base damage, N7-methylguanine, is associated with semen quality in men

B. Altakroni^{1,5}, C. Nevin², M. Carroll^{1b}, C. Murgatroyd², G. Horne³, D. R. Brison^{3,4} & A. C. Povey^{1b✉}

Sperm DNA contains a range of DNA base damage that can arise, in part, from exposure to methylating agents. However, the effects are not fully characterized and so the aim of this study was to investigate associations between semen quality and the levels of N7-methyldeoxyguanosine (N7-MedG), a marker of exposure to methylating agents, and other markers of DNA damage and DNA methylation. Sperm samples were collected from 105 men attending an assisted reproduction clinic as part of a couple undergoing treatment for infertility and semen quality assessed manually according to WHO guidelines. Semen levels of N7-MedG, quantified by immunoslotblot, were significantly higher in men with sperm concentration $<15 \times 10^6/\text{ml}$ ($p \leq 0.01$), semen volume $<1.5 \text{ ml}$ ($p \leq 0.05$) and also in men with any aspect of semen quality below WHO reference levels ($p \leq 0.001$). Measures of neutral Comet DNA damage were correlated with semen quality in a univariate analysis but not after adjustment for N7-MedG levels. Sperm concentration was negatively associated with % methylation at the gene for *DAZL* but no other marker of global or gene-specific DNA methylation. Results support the hypothesis that the known toxic and DNA damaging properties of alkylating agent exposure may have direct deleterious consequences on semen quality.

Sperm DNA contains damage that may impact on semen quality and ART outcome^{1,2}. This damage is not fully characterised, with the most widely used assays measuring some loss in DNA integrity such as single and/or double strand DNA breaks, DNA fragmentation or DNA denaturation³. In addition, specific DNA base damage has also been detected in sperm DNA, particularly that arising from reactive oxygen species such as 8-oxoguanine⁴. However, other types of DNA base damage including that arising from methylating agents has also been reported⁵. The potential adverse consequences of exposure to DNA methylating and other alkylating agents on somatic cells i.e. toxicity, mutagenicity and carcinogenicity is well recognised⁶. Such changes are largely ascribed to DNA damage formation⁷, though exposure to alkylating agents can induce other cellular changes including changes in gene expression⁸ or histone phosphorylation⁹. Treatment of male mice with the ethylating agent, *N*-ethyl-*N*-nitrosourea, causes mutations in sperm that has led to the development of many models of human disease^{10,11}. In contrast very little is known about DNA alkylation in human spermatozoa. We reported several years ago that N7-MedG levels, a biomarker of exposure to methylating agents¹², were associated with male factor infertility in a population of men attending as a part of a couple for ART⁵. N7-MedG is quantitatively the most significant methyl DNA adduct, constituting around 70–85% of all adducts produced by methylating agents¹³. Though N7-MedG does not alter DNA duplex structure or coding properties and is neither mutagenic nor toxic¹⁴, N7-methylguanine in DNA is relatively unstable and can depurinate leading to abasic sites that are toxic and mutagenic¹⁵ and can further lead to DNA strand breaks¹⁶. More recently it has been reported that N7-MedG

¹Centre for Occupational and Environmental Health, Epidemiology and Public Health Group, Division of Population Health, Health Services Research and Primary Care, School of Health Sciences, Faculty of Biology, Medicine and Health, Manchester Academic Health Sciences Centre, University of Manchester, Manchester M13 9PL, UK. ²Department of Life Sciences, Faculty of Science and Engineering, Manchester Metropolitan University, John Dalton Building, Chester Street, Manchester M1 5GD, UK. ³Department of Reproductive Medicine, Old Saint Mary's Hospital, Manchester Academic Health Science Centre, Manchester University NHS Foundation Trust, Oxford Road, Manchester M13 9PT, UK. ⁴Division of Developmental Biology & Medicine, School of Medicine, Maternal and Fetal Health Research Centre, Faculty of Biology, Medicine and Health, University of Manchester, Manchester M13 9PL, UK. ⁵B. Altakroni is deceased. ✉email: apovey@manchester.ac.uk

Variable	Value	Mean \pm SD (n)	Range
Age	Years	35.3 \pm 5.8 (105)	26–57
Semen quality	Sperm concentration ($\times 10^6$ /ml)	59.1 \pm 66.2 (105)	0.5–362.0
	Progressive motile sperm (%)	49.0 \pm 17.1 (105)	1.0–77.0
	Immobile sperm (%)	40.9 \pm 14.5 (105)	0.5–99.0
	Semen volume (ml)	3.2 \pm 1.4 (105)	0.5–8.1
Comet assay	Median tail DNA (%)	1.02 \pm 0.76 (105)	0.07–5.28
	Sperm with low DNA damage (LDD) (%)	15.4 \pm 8.3 (105)	2.3–39.2
	Sperm with high DNA damage (HDD) (%)	14.7 \pm 8.6 (105)	4.0–47.2
N7-MedG	fmoles/ μ g DNA	0.55 \pm 0.35 (101)	0.12–2.30
Global methylation	nM	396.1 \pm 71.3 (44)	296–687
Gene Methylation	DAZL (% methylated)	4.6 \pm 5.0 (31)	0.4–20.0
	H19 (% methylated)	40.4 \pm 11.9 (44)	12.8–73.0
	MEG3 (% methylated)	77.4 \pm 27.8 (45)	0–91.3
	SNRPN (% methylated)	3.2 \pm 2.5 (43)	0.7–12.3

Table 1. Descriptive analysis of semen quality and markers of DNA damage and DNA methylation.

adducts can form reversible, but toxic, DNA–protein cross links with histone proteins¹⁷. Furthermore, overexpression of the DNA repair protein, N3-methylpurine-DNA glycosylase, that repairs this adduct, sensitises cells to the genotoxic effects of methylating agents^{14,18}. In addition, the presence of N7-MedG in DNA, depending upon the type of alkylating agent exposure, may also indicate significant levels of the pro-mutagenic, pro-carcinogenic and teratogenic adduct, *O*⁶-methylguanine¹⁹.

Associations between DNA alkylation and other types of DNA damage or DNA modifications, such as epigenetic methylation, in human sperm are unknown. To provide further information on these associations with semen quality, we have investigated these markers in sperm of men attending an infertility clinic as part of a couple.

Results

The study included 105 semen samples with data available for semen quality, DNA damage measured by Comet assay, and N7-MedG (101). Of these, 44 also had DNA methylation data.

Descriptive analysis of semen quality and markers of DNA damage and DNA methylation. The semen quality of the studied population is shown in Table 1 together with a descriptive analysis of markers of DNA damage and DNA methylation. There were 28 men whose sperm concentration was less than 15×10^6 /ml, 18 men with a % progressively motility of <32 , and 11 men whose semen volume was less than 1.5 ml. Overall there were 44 men with a below reference level of either sperm concentration, % progressive motility or semen volume.

% of sperm with low DNA damage (% LDD sperm) was inversely correlated ($r_s = -0.31$, $p < 0.001$ $n = 105$) with % of sperm with high DNA damage (% HDD sperm) and with median % tail DNA ($r_s = -0.79$, $p < 0.001$ $n = 105$) whereas % HDD sperm was positively correlated with median % tail DNA ($r_s = 0.67$, $p < 0.001$ $n = 105$). N7-MedG in sperm DNA was negatively associated with % LDD sperm ($r_s = -0.22$, $p < 0.05$ $n = 101$) and positively associated with median % tail DNA ($r_s = 0.20$, $p < 0.05$ $n = 101$): it was not correlated with % HDD sperm ($r_s = 0.12$, $n = 101$). % DAZL methylation was positively associated with % HDD sperm ($r_s = 0.37$; $p < 0.05$) and with median % tail DNA ($r_s = 0.39$; $p < 0.05$). There was no other association between global methylation and gene-specific methylation or markers of DNA damage.

Relationship between semen quality and markers of DNA damage and DNA methylation. Sperm concentration was negatively associated with N7-MedG levels ($r_s = -0.32$, $p < 0.001$), % HDD sperm ($r_s = -0.41$, $p < 0.001$), median % tail DNA ($r_s = -0.31$, $p < 0.001$) and with % methylation of DAZL ($r_s = -0.56$, $p < 0.001$) but not with global methylation or % methylation of H19CTCF6, SNRPN, or MEG3 (Table 2). The multiple regression model including N7-MedG, %HDD sperm, median % tail DNA and % DAZL was not significant ($R^2 = 0.20$, $p = 0.2$) with DAZL methylation showing a non-significant trend ($p = 0.08$; Table 3).

% progressively motile sperm was positively associated with % LDD sperm ($r_s = 0.34$, $p < 0.001$), but negatively associated with % HDD sperm ($r_s = -0.50$, $p < 0.001$) and median % tail DNA ($r_s = -0.48$, $p < 0.001$). There was no other correlation between semen quality and global methylation or gene-specific DNA methylation (Table 2). The multiple regression model with % LDD sperm, % HDD sperm, median % tail DNA and N7-MedG produced $R^2 = 0.28$, ($p < 0.001$) with %HDD sperm ($p < 0.01$) and N7-MedG ($p < 0.05$) being significant (Table 3).

% immotile sperm was negatively associated with % LDD sperm ($r_s = -0.31$, $p < 0.01$), but positively associated with % HDD sperm ($r_s = 0.41$, $p < 0.001$) and median % tail DNA ($r_s = 0.41$, $p < 0.001$). There was no other correlation between semen quality and global methylation or % methylation of H19CTCF6, SNRPN, or MEG3 (Table 2). The multiple regression model with %LDD sperm, % HDD sperm and median % tail DNA produced

DNA marker	Specific variable	r_s (n) ^a			
		Sperm concentration ($\times 10^6$ /ml)	Progressively motile sperm (%)	Immotile sperm (%)	Semen volume (ml)
Alkylation	N7-MedG (fmoles/ μ g DNA)	-0.32*** (101)	-0.19 [‡] (101)	0.11 (101)	-0.12 (101)
Integrity	% sperm with low DNA damage (LDD)	0.11 (105)	0.34*** (105)	-0.31** (105)	0.09 (105)
Integrity	% sperm with high DNA damage (HDD)	-0.41*** (105)	-0.50*** (105)	0.41*** (105)	0.16 (105)
Integrity	% tail DNA (median)	-0.31*** (105)	-0.48*** (105)	0.41*** (105)	0.04 (105)
Methylation	Global methylation (nM)	0.03 (44)	0.07 (44)	-0.02 (44)	0.06 (44)
	DAZL (% methylated)	-0.56*** (31)	-0.04 (31)	0.06 (31)	0.27 (31)
	H19 (% methylated)	-0.02 (44)	-0.03 (44)	0.04 (44)	-0.06 (44)
	MEG3 (% methylated)	0.01 (45)	0.10 (45)	-0.07 (45)	0.08 (45)
	SNRPN (% methylated)	-0.02 (43)	0.05 (43)	-0.06 (43)	-0.06 (43)

Table 2. Correlations between markers of DNA damage and methylation and semen quality. ^a r_s Spearman rho correlation coefficient; * $p < 0.05$; ** $p < 0.01$; *** $p \leq 0.001$; [‡] $p = 0.053$.

Semen parameter	Model variable	B	SE B	β
Sperm concentration	Constant	132.5	42.4	
	DAZL	-4.27	2.32	-0.33 [‡]
	N7MeG	-45.86	74.91	-0.12
	HDD	-0.33	2.34	-0.04
	Median tail DNA	-12.44	26.91	-0.14
Progressive motility	Constant	62.80	6.84	
	N7MeG	-7.83	3.86	-0.18*
	HDD	-0.79	0.30	-0.38**
	Median tail DNA	-1.56	4.96	-0.06
	LDD	0.26	0.26	-0.14
Immotile sperm	Constant	45.6	5.21	
	HDD	0.93	0.26	0.55***
	Median tail DNA	-8.00	3.74	-0.42*
	LDD	-0.66	0.23	-0.38**

Table 3. Multiple regression analysis of semen quality with DNA damage and DNA methylation markers. [‡] $p < 0.1$; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

$R^2 = 0.19$, ($p < 0.001$) with %HDD sperm ($p < 0.01$), %LDD sperm ($p < 0.01$) and median % tail DNA ($p < 0.05$) all being significant (Table 3). Semen volume was not associated with any of the measured biomarkers.

Associations between semen quality and DNA damage and DNA methylation in men with semen quality above and below WHO referent values. To assess whether DNA damage and DNA methylation levels were associated with clinically more relevant factors, we examined whether levels were different in men whose semen quality was either below or above WHO reference values (Table 4). Specifically, N7-MedG levels, % HDD sperm and median % Tail DNA were higher in those men whose sperm concentration was $< 15 \times 10^6$ ml. In contrast, % LDD sperm was lower and % HDD sperm and median %tail DNA higher, in men with $< 32\%$ progressively motile sperm. Furthermore, N7-MedG levels were higher in those men with a semen volume was less than 1.5 ml. Overall, in any men whose sperm concentration or % progressively motile sperm or semen volume was below the reference category, N7-MedG levels, % HDD sperm and median %tail DNA were significantly higher and % LDD sperm significantly lower compared to men whose semen quality was above the reference value. In this analysis there was no association between any DNA methylation marker and semen quality above or below the WHO reference value.

In a logistic regression analysis (Table 5), N7-MedG levels were significantly associated with low sperm concentration (OR 6.46, 95%CI 1.51–27.7) even after adjustment (OR 6.42, 95%CI 1.47–28.0) for other markers of DNA damage, % HDD sperm and median % tail DNA. Whilst % HDD sperm and % median tail DNA were significantly associated with having low progressive motility in a univariate analysis (Table 5), after adjustment

Semen variable	Category	Marker of DNA damage or methylation. Mean \pm SD (n)			
		N7-MedG	% LDD sperm ^a	% HDD sperm ^b	% Tail DNA
Concentration	> 15 \times 10 ⁶ /ml	0.49 \pm 0.29 (77)	16.2 \pm 8.7 (77)	13.4 \pm 8.2 (77)	0.88 \pm 0.62 (77)
	< 15 \times 10 ⁶ /ml	0.75 \pm 0.47 (24)**	13.3 \pm 7.1 (28)	18.1 \pm 8.9 (28)**	1.39 \pm 0.97 (28)***
Progressive motility	> 32%	0.53 \pm 0.33 (86)	16.3 \pm 8.5 (87)	13.0 \pm 6.9 (87)	0.87 \pm 0.54 (87)
	< 32%	0.70 \pm 0.46 (15)‡	11.0 \pm 6.1 (18)*	22.7 \pm 11.3 (18)***	1.74 \pm 1.16 (18)***
Semen volume	> 1.5 ml	0.51 \pm 0.28 (91)	15.6 \pm 8.5 (94)	14.6 \pm 8.7 (94)	1.02 \pm 0.79 (94)
	< 1.5 ml	0.91 \pm 0.69 (10)*	13.6 \pm 6.8 (11)	15.8 \pm 7.9 (11)	1.03 \pm 0.44 (11)
Any category	> Reference	0.45 \pm 0.20 (61)	17.3 \pm 8.9 (61)	12.2 \pm 7.2 (61)	0.77 \pm 0.53 (61)
	< Reference	0.71 \pm 0.47 (40)***	12.9 \pm 6.8 (44)*	18.1 \pm 9.2 (44)***	1.36 \pm 0.89 (44)***

Table 4. Associations between markers of DNA damage and semen quality categorised by WHO reference values. ^a % sperm with low DNA damage; ^b % sperm with high DNA damage; Statistically significant by Mann Whitney U test: ‡ $p \leq 0.1$; * $p \leq 0.05$; ** $p \leq 0.01$; *** $p \leq 0.001$.

DNA damage marker	OR (95%CI)						
	Semen volume	Concentration		Progressive motility		Any category	
	Unadjusted	Unadjusted	Adjusted	Unadjusted	Adjusted	Unadjusted	Adjusted
N7MedG	7.2 (1.7–30.0)	6.46 (1.51–27.7)	6.42 (1.47–28.0)	2.80 (0.79–9.91)	3.42 (0.89–13.2)	20.7 (2.75–155.6)	19.0 (2.41–150)
% LDD sperm ^b	-	-	-	0.91 (0.84–0.98)	0.98 (0.87–1.10)	0.93 (0.88–0.98)	1.03 (0.94–1.13)
% HDD sperm ^c	-	1.06 (1.01–1.12)	0.99 (0.90–1.08)	1.12 (1.05–1.20)	1.09 (0.97–1.22)	1.10 (1.04–1.17)	1.01 (0.92–1.12)
% Tail DNA	-	2.48 (1.26–4.86)	2.33 (0.75–7.31)	4.54 (1.89–10.9)	1.54 (0.23–10.2)	4.90 (2.03–11.8)	5.17 (0.8–33.2)

Table 5. Unadjusted and adjusted^a associations between DNA damage marker and poor semen quality categorised by WHO reference values. ^aAdjusted for other markers of DNA damage that were significantly associated with semen quality in a univariate analysis; ^b % sperm with low DNA damage; ^c % sperm with high DNA damage.

for % LDD sperm and N7-MeG there were no significant associations. N7-MedG levels were associated with an increased risk of having low progressive motility but this increase was not significant (OR_{adj} 3.42, 95%CI 0.89–13.2). N7-MedG levels were, however, significantly associated with having a semen sample with any parameter below the WHO reference value (OR 20.7, 95%CI 2.75–155.6) and after adjustment with other markers of DNA damage (OR 19.0, 95%CI 2.41–150). No markers of DNA damage were associated with poor semen quality after adjustment.

Discussion

In this study of sperm DNA damage and DNA methylation of men attending for ART as part of a couple, the main finding was that N7-MedG levels were significantly associated with adverse semen quality even after adjustment for other markers of DNA damage as measured by the Comet assay. Measures of DNA damage assessed by the Comet assay were correlated with % progressive motility and % immotile sperm and differed in men whose semen quality was below WHO reference levels in a univariate analysis but not after adjustment for N7-MedG levels. Sperm concentration was negatively associated with % methylation of *DAZZL* but no other marker of DNA methylation was associated with adverse semen quality in this study.

N7-MedG is a marker of exposure to exogenous and endogenous DNA methylating agents and has been detected in a wide range of human tissues¹³. The precise contribution of exogenous (e.g. tobacco specific nitrosamines) and endogenous (e.g. s-adenosylmethionine and other agents formed in situ through *N*-nitrosation of amines) methylating agents to N7-MedG levels is unknown. Smoking is an exclusion criterion for treatment in our NHS centre and none of the men in the study were self-reported current smokers making it very unlikely that smoking was associated with N7-MedG levels.

There is also evidence that obesity is associated with DNA damage levels in that higher levels of DNA fragmentation (ie cells that did not have a detectable nucleus but only a Comet tail as detected by the alkaline Comet assay) were reported in obese men as compared to controls^{20,21}. However, we found no correlation between BMI and DNA damage levels (data not shown). However, the finding in this study that N7-MedG levels were significantly associated with adverse semen quality is consistent with our previous report that N7-MedG levels were higher in men diagnosed with male factor infertility⁵ and implicates DNA base damage, and not just DNA strand breaks, as a contributing cause of male infertility. The mechanistic basis of this association is unclear but it might be related to the known chemical instability of the N7-MedG adduct leading to DNA strand breaks¹⁶ and it is worthy of note that N7-MedG levels were negatively associated with %LDD sperm and positively associated with

median % tail DNA. In this context N7-MedG instability might represent a previously uncharacterised exposure route to increased DNA strand breaks in human sperm. Alternatively, the association between N7-MedG and adverse semen quality may be due to toxic DNA protein cross-links¹⁷ or due to the presence of other more toxic adducts such as O⁶-methylguanine or N3-methyladenine that will be formed at the same time as N7-MedG²². The latter 2 adducts have as far as we can ascertain not yet been reported in human sperm DNA but their presence in sperm DNA may have more deleterious consequences as O⁶-methylguanine is a known pro-mutagenic adduct¹⁹ and N3-methyladenine can induce chromosome aberrations²³. The post-fertilisation effects of these adducts would depend not only on its levels in individual sperm but also the repair capacity of the oocyte.

The Comet assay measures a variety of different types of DNA damage (e.g. single and double strand breaks and apurinic sites) whether induced for example, by oxidative stress, chemical modifications of DNA or DNA repair processes^{24,25}. The precise contribution of these different mechanisms to DNA damage in any one sample is unknown. The neutral Comet assay has been reported to measure principally double-strand breaks as Alu1 induced DNA fragmentation was higher with the neutral Comet assay than the alkaline comet assay²⁶. This, though, is controversial with other authors reporting that the neutral comet assay measures single strand nicks when the DNA remains attached to the nuclear matrix²⁷. As the extent of strand breaks increases, very large stretches of DNA can become detached from the matrix and move out of the nucleus and into the gel²⁸. We refer to these events as DNA damage, recognising that the neutral Comet assay measures a mix of single and double strand breaks. The relationship between DNA damage detected by the neutral Comet assay and semen quality is currently unclear. The DNA fragmentation index in sperm, as assessed by the neutral Comet assay, has been negatively associated with sperm concentration and morphology in one study¹, and with sperm viability, motility and morphology but not sperm concentration²⁹. Similarly, a weak correlation was observed between tail % intensity and % motility, % normal morphology but not sperm concentration³⁰. Conversely, no associations between semen quality and % tail DNA were detected in 80 healthy male volunteers³¹. It is unclear whether these differences result from differences in analysis protocols, descriptors of DNA damage or populations studied, though in one study associations between indices of semen quality varied with how the DNA damage was characterised³². One additional interpretation is that the differences in associations may result from differences in other types of DNA damage e.g. N7-MedG present within human sperm.

Previous studies have reported associations between environmental exposures and/or DNA damage and sperm DNA methylation in specific genes, suggesting, at least, that DNA damage and DNA methylation share a common exposure or cause and potentially that the effects of such exposure are mediated, in part, by their epigenetic effects. Associations between reactive oxygen species and *H19-Igf2* methylation have been described³³ and also methylation levels of *IGF2* and *KCNQ1* but not *H19*, *MEG3*, *PEG3*, *SNRPN* were associated with DNA fragmentation index³⁴. We also examined but found no associations between markers of DNA damage and global DNA methylation and methylation of genes that have been previously associated with male infertility³⁵. This suggests that the effects of alkylating agent exposure may not be mediated through changes in DNA methylation, but rather via the direct effects of DNA damage. The exact source/origin of DNA damage detected by the Comet assay still remains unknown but there is little evidence in the present study of these sources also impacting on the methylation of sperm. If such associations do exist, they will likely depend upon the initial exposures causing the DNA damage. *DAZL* methylation was negatively associated with sperm concentration, consistent with previous studies³⁶. Higher *DAZL* methylation was found in asthenozoospermia and oligozoospermia men³⁷. In contrast, no associations were detected with global methylation or methylation of other genes (*H19*, *MEG3*, *NR3C1*, *SNRPN*) that have previously been associated with semen quality³⁵; this may be due to population or sample size differences.

Like previous studies examining associations between DNA damage and semen quality, the main weakness of this study is that it is cross-sectional in nature and so unable to determine causative associations between the measured biomarkers and semen quality. In addition, there is the possibility that the levels of N7-MedG reported here may reflect levels occurring in contaminating leucocytes and round cells (for example). However, as these samples were used for clinical purposes, numbers of other cells were routinely assessed and samples identified as containing high levels of leucocytes and round cells were omitted from the study. Furthermore, seminal round cells have been reported to be mostly immature post-meiotic germ cells³⁸ and leucocytes account for maximum 5% of total number of round cells in the seminal fluid³⁹. The alternative possibility that low levels of leucocytes and round cells influenced the results is considered unlikely as their N7-MedG levels would have had to have been much higher than observed here, at levels only previously observed in patients receiving methylating agent chemotherapy¹³. We are thus confident that the results herein are representative of sperm N7-MedG levels.

However, our results are consistent with current knowledge regarding the toxicological consequences of alkylating agent exposure in other tissues. This study and other studies indicate that sperm DNA contains a range of different types of DNA damage that might be associated with semen quality and have more deleterious consequences post-fertilisation.

Materials and methods

Study population and ethical approval. 105 men of couples attending the Department of Reproductive Medicine, St Mary's Hospital, Manchester University NHS Foundation Trust, Manchester, for assisted reproduction treatment (ART), participated in this study. All couples had provided written informed consent for research and this study had Local Ethics Committee Approval (Central Manchester REC ERP/91/078, HFEA research licence R0026) and was performed in accordance with relevant guidelines/regulations.

Semen analysis. Participants provided by masturbation a semen sample on the day of ART treatment. After liquefaction, semen volume, sperm concentration and motility were determined manually according to

WHO guidelines⁴⁰ in the hospital by biomedical and clinical scientific staff, and according to the UK NEQAS Reproductive Sciences scheme⁴¹. Samples were to be used for clinical treatment and those with any significant round cell/white blood cell count or samples with leukospermia were omitted from the study. After collecting the remaining sample from the hospital, the neat semen was centrifuged to separate the sperm from the seminal plasma and an aliquot of the neat sperm pellet was directly used to determine the level of DNA damage by using the neutral Comet assay.

Extraction of DNA from sperm. Briefly, genomic DNA was extracted from up to 4×10^7 sperm cells using the Qiagen Blood and cell culture DNA Midi Kit except that the proteinase K and ribonuclease A digestion step was carried out at 4 °C overnight. The following day, 0.5 ml of 5 mM DTT solution was added to each tube, mixed and incubated again in the cold room on a rotor mixer for 1 h, followed by incubation for another 1 h at 37 °C in water bath. Following the manufacturer's instructions, the samples were added into gravity flow columns. Eluted from the columns, DNA was isopropanol and ethanol precipitated and rapidly air dried to remove residual alcohol. The concentration of extracted sperm DNA in 1X TE buffer was measured first by Nano-drop and then by a picogreen assay⁵.

N7-MedG slotblot analysis of sperm DNA. Immunoblot analysis of N7-MedG in sperm DNA has been described previously^{5,12}. Methylated DNA standards were prepared by reacting CT-DNA with *N*-nitroso-*N*-methylurea. Briefly, DNA samples were sonicated (Sonicator, Bandeline electronic sonopuls, UW 2070, Probe: SH70G) for 10 s at 50% power to obtain DNA fragments of size between 400 and 1500 bp and these were incubated in alkaline solution to induce imidazole ring opening. After neutralisation, DNA samples were heat denatured and then 1 µg of DNA immobilised on a nitrocellulose membrane (Whatman: Portran, BA79, pore size 0.1 µm). Membranes were incubated first with a rabbit polyclonal antibody raised against 2,6-diamino-4-hydroxy-5-*N*-methylformamidopyrimidine (me-Fapy) conjugated to keyhole limpet hemocyanin which specifically detected N7-MedG in mice treated with *N*-Nitroso-*N*-methylurea⁴² and then a polyclonal goat anti-rabbit-horseradish peroxidase conjugate. Membranes were incubated with ECL reagent (Amersham Hyperfilm) and exposed to an X-ray film (Fuji) which was subsequently developed and scanned using a HP Scanjet 485. The intensity of the bands for the samples and the standards was quantified by using ImageJ program and a standard curve was generated and subsequently the levels of N7-MedG in the samples were calculated. To confirm DNA binding the membrane was stained with a propidium iodide solution and scanned with a Typhoon scanner (GE Healthcare) at 610 emission and 560 absorption wavelengths and the intensity of DNA bands in the obtained image quantified using ImageJ.

Single-cell gel electrophoresis (Comet) assay. DNA damage was assessed by a neutral comet assay according to previous studies but with minor modifications^{1,43}. Sperm ($\sim 2 \times 10^4$) were suspended in LMP agarose and applied to a microscope slide pre-coated with 1% NMP agarose. After gel setting, the slides were immersed firstly in freshly made precooled lysis buffer (2.5 M NaCl, 100 mM EDTA, 10 mM Tris-base, 10% DMSO and 1% Triton X-100, pH 10) at 4 °C, then in fresh lysis buffer containing 10 mM DTT and finally fresh lysis buffer containing proteinase K (100 µg/ml) to lyse the cells. The slides were then equilibrated for 20 min in cold electrophoresis buffer (300 mM sodium acetate 100 mM Tris, pH9) then electrophoresis was carried out at 13 V, 120–125 mA for 1 h. After neutralisation in cold neutralization buffer (0.4 M Tris-base pH 7.5), slides were stained with SYBR gold (Invitrogen, UK) used according to the manufacturer's instructions and were then examined using an Olympus fluorescence microscope (OLYMPUS, U-RFL-T; Filter: FITC; Camera: Q IMAGING, RETIGA-SRV, FAST 1394; Software: QCapture Pro 6.0) and around 35 images containing in total around 400 sperm (Comet) were obtained. Then, % tail DNA was determined by using TriTeK CometScore™ Freeware version 1.5 (TriTeK Corp, US) in 400 sperm.

Prior to the analysis of the participant samples, we validated our neutral comet assay by incubating sperm with up to 30 µM H₂O₂ and this clearly showed that the assay was able to detect increases in DNA damage induced by a DNA damaging agent. We found a dose-dependent decrease in the % of sperm with few strand breaks ($r^2 = 0.077$; < 2% at the highest H₂O₂ concentration) and a dose-dependent increase in median levels of % tail DNA ($r^2 = 0.91$) and % sperm with high levels of DNA damage ($r^2 = 0.90$; 20% at the highest H₂O₂ concentration). A pilot reproducibility study also showed little variance from day to day.

Global DNA methylation. Genomic DNA was heat-denatured and digested to nucleosides by nuclease P1 (Sigma, UK) and then alkaline phosphatase (Sigma, UK). 5-methylcytosine levels were then quantified using the Global DNA Methylation ELISA Kit (Cambridge Biosciences, UK), following the manufacturer's instructions.

Gene specific methylation. DNA samples underwent bisulphite conversion using the EpiTect Fast DNA Bisulphite Kit (Qiagen, UK), according to the manufacturer's instructions. Pyrosequencing target regions were amplified using the PyroMark PCR Kit (Qiagen, UK) as follows: PCR reactions were performed in a total volume of 25 µl consisting of 12.5 µl 2×PCR master mix, 2.5 µl 10×Coraload, 0.75 µl (3 µM) of each forward and reverse primer, 2 µl (~10 ng) bisulfite-converted DNA and 6.5 µl H₂O. Primer sequences covering the target CpG regions were either purchased as pre-designed CpG assays from Qiagen UK (Supplementary data Table 1), or custom oligonucleotides were designed using the PyroMark Assay Design software (Qiagen, UK) and purchased through Life Technologies (UK) (Supplementary Table 2). Gradient PCR was initially carried out to determine the optimal annealing temperature for amplification ranging from 52 to 60 °C and cycling conditions were carried out as follows: 95 °C for 15 min; followed by 45 cycles of denaturation, 94 °C for 30 s; annealing, 56 °C for 30 s; extension, 72 °C for 30 s. Finally, an extension of 72 °C for 10 min was included. PCR

products were frozen at -20 °C until pyrosequencing. PCR products were immobilised to Streptavidin Sepharose High Performance beads (GE Healthcare Life Sciences, Uppsala, Sweden), denatured and the pyrosequencing primer (Supplementary Table) was annealed to the template. Pyrosequencing was performed on a PyroMark 24 Pyrosequencing system (Qiagen). Results were analysed on the PyroMark Q24 software to determine the % methylation for each CpG site.

Statistical analyses. Semen quality was assessed both as a continuous variable and after dividing the population into two based upon WHO reference values for semen quality⁴⁴ (i.e. sperm concentration > 15 × 10⁶/ml, % progressive motility > 32% and a semen volume > 1.5 ml). We established the Comet thresholds blind to the semen quality status. The distribution of % tail DNA in the sperm samples was not normal with the log % tail DNA showed an almost trimodal distribution in 75% of the samples, ~ 14% of the samples showed bimodal distributions while the remaining 11% samples showed multimodal distribution. It was possible to distinguish two subgroups of sperm; sperm with low levels of DNA damage (% tail DNA < 0.005%, LDD sperm) and sperm with high levels of DNA damage (% tail DNA ≥ 7.5%, HDD sperm). Correlation (Spearman rank analysis) and multiple regression analyses (including those markers whose associations with semen quality had a $p \leq 0.1$) were conducted to examine the relationship between semen quality and markers of DNA damage and DNA methylation (mean methylation of all CpG sites analysed). This regression analysis is of limited clinical value and so biomarker levels were then compared in men whose semen quality were above or below WHO referent levels in both a univariate analysis and by logistic regression after adjustment for other DNA damage markers with $P < 0.1$.

Data availability

The datasets generated during and analysed during the current study are available from the corresponding author on reasonable request.

Received: 20 May 2020; Accepted: 18 November 2020

Published online: 04 February 2021

References

- Morris, I. D., Iltott, S., Dixon, L. & Brison, D. R. The spectrum of DNA damage in human sperm assessed by single gel electrophoresis (Comet assay) and its relationship to fertilization and embryo development. *Hum. Reprod.* **17**, 990–998 (2002).
- Simon, L., Emery, B. R. & Carrell, D. T. Review: Diagnosis and impact of sperm DNA alterations in assisted reproduction. *Best Pract. Res. Clin. Obstet. Gynaecol.* **44**, 38–56 (2017).
- Panner Selvam, M. K. & Agarwal, A. A systematic review on sperm DNA fragmentation in male factor infertility: laboratory assessment. *Arab. J. Urol.* **16**, 65–76 (2018).
- Bisht, S., Faiq, M., Tolahunase, M. & Dada, R. Oxidative stress and male infertility. *Nat. Rev. Urol.* **14**, 470–485 (2017).
- Stocks, S. J. *et al.* Alkylation of sperm DNA is associated with male factor infertility and a reduction in the proportion of oocytes fertilised during assisted reproduction. *Mutat. Res.* **698**, 18–23 (2010).
- Kaina, B., Christmann, M., Naumann, S. & Roos, W. P. MGMT: key node in the battle against genotoxicity, carcinogenicity and apoptosis induced by alkylating agents. *DNA Repair (Amst.)* **6**, 1079–1099 (2007).
- Shrivastav, N., Li, D. & Essigmann, J. M. Chemical biology of mutagenesis and DNA repair: cellular responses to DNA alkylation. *Carcinogenesis* **31**, 59–70 (2010).
- Hebels, D. G. *et al.* Time-series analysis of gene expression profiles induced by nitrosamides and nitrosamines elucidates modes of action underlying their genotoxicity in human colon cells. *Toxicol. Lett.* **207**, 232–241 (2011).
- Chen, K. *et al.* Biphasic reduction of histone H3 phosphorylation in response to N-nitroso compounds induced DNA damage. *Biochim. Biophys. Acta.* **1860**, 1836–1844 (2016).
- Acevedo-Arozena, A. *et al.* ENU mutagenesis, a way forward to understand gene function. *Annu. Rev. Genom. Hum. Genet.* **9**, 49–69 (2008).
- Jamsai, D. & O'Bryan, M. K. Genome-wide ENU mutagenesis for the discovery of novel male fertility regulators. *Syst. Biol. Reprod. Med.* **56**, 246–259 (2010).
- Harrison, K. L. *et al.* Development and application of a sensitive and rapid immunoassay for the quantitation of N7-methyldeoxyguanosine in DNA samples. *Chem. Res. Toxicol.* **14**, 295–301 (2001).
- Bianchini, F. & Wild, C. P. 7-Methyldeoxyguanosine as a marker of exposure to environmental methylating agents. *Toxicol. Lett.* **72**, 175–184 (1994).
- Rinne, M. L., He, Y., Pachkowski, B. F., Nakamura, J. & Kelley, M. R. N-methylpurine DNA glycosylase overexpression increases alkylation sensitivity by rapidly removing non-toxic 7-methylguanine adducts. *Nucleic Acids Res.* **33**, 2859–2867 (2005).
- Loeb, L. A. & Preston, B. D. Mutagenesis by apurinic/aprimidinic sites. *Annu. Rev. Genet.* **20**, 201–230 (1986).
- Gates, K. S., Nooner, T. & Dutta, S. Biologically relevant chemical reactions of N7-alkylguanine residues in DNA. *Chem. Res. Toxicol.* **17**, 839–856 (2004).
- Yang, K., Park, D., Tretyakova, N. Y. & Greenberg, M. M. Histone tails decrease N7-methyl-2'-deoxyguanosine depurination and yield DNA-protein cross-links in nucleosome core particles and cells. *Proc. Natl. Acad. Sci. USA* **115**, E11212–E11220 (2018).
- Coquerelle, T., Dosch, J. & Kaina, B. Overexpression of N-methylpurine-DNA glycosylase in Chinese hamster ovary cells renders them more sensitive to the production of chromosomal aberrations by methylating agents—a case of imbalanced DNA repair. *Mutat. Res.* **336**, 9–17 (1995).
- Margison, G. P., Santibanez-Koref, M. F. & Povey, A. C. Mechanisms of carcinogenicity/ chemotherapy by O⁶-methylguanine. *Mutagenesis* **17**, 483–487 (2002).
- Fariello, R. M. *et al.* Association between obesity and alteration of sperm DNA integrity and mitochondrial activity. *BJU Int.* **110**, 863–867 (2012).
- Ferigolo, P. C. *et al.* Sperm functional aspects and enriched proteomic pathways of seminal plasma of adult men with obesity. *Andrology* **7**, 341–349 (2019).
- Beranek, D. T., Weis, C. C. & Swenson, D. H. A comprehensive quantitative-analysis of methylated and ethylated DNA using high-pressure liquid-chromatography. *Carcinogenesis* **1**, 595–606 (1980).
- Engelward, B. P. *et al.* A chemical and genetic approach together define the biological consequences of 3-methyladenine lesions in the mammalian genome. *J. Biol. Chem.* **273**, 5412–5418 (1998).

24. Azqueta, A. & Collins, A. R. The essential comet assay: a comprehensive guide to measuring DNA damage and repair. *Arch. Toxicol.* **87**, 949–968 (2013).
25. Pu, X., Wang, Z. & Klaunig, J. E. Alkaline Comet assay for assessing DNA damage in individual cells. *Curr. Protoc. Toxicol.* **65**, 3.12.1–3.12.11 (2015).
26. Ribas-Maynou, J. *et al.* Alkaline and neutral Comet assay profiles of sperm DNA damage in clinical groups. *Hum. Reprod.* **27**, 652–658 (2012).
27. Collins, A. R. *et al.* The comet assay: topical issues. *Mutagenesis* **23**, 143–151 (2008).
28. Afanasieva, K. & Sivolob, A. Physical principles and new applications of comet assay. *Biophys. Chem.* **238**, 1–7 (2018).
29. Chi, H. J. *et al.* Integrity of human sperm DNA assessed by the neutral comet assay and its relationship to semen parameters and clinical outcomes for the IVF-ET program. *Clin. Exp. Reprod. Med.* **38**, 10–17 (2011).
30. Duydu, Y. *et al.* Assessment of DNA integrity (COMET assay) in sperm cells of boron-exposed workers. *Arch. Toxicol.* **86**, 27–35 (2012).
31. Schmid, T. E. *et al.* The effects of male age on sperm DNA damage in healthy non-smokers. *Hum. Reprod.* **22**, 180–187 (2007).
32. Trisini, A. T., Singh, N. P., Duty, S. M. & Hauser, R. Relationship between human semen parameters and deoxyribonucleic acid damage assessed by the neutral comet assay. *Fertil. Steril.* **82**, 1623–1632 (2004).
33. Darbandi, M. *et al.* Reactive oxygen species-induced alterations in H19-Igf2 methylation patterns, seminal plasma metabolites, and semen quality. *J. Assist. Reprod. Genet.* **36**, 241–253 (2019).
34. Ni, W. *et al.* Methylation levels of IGF2 and KCNQ1 in spermatozoa from infertile men are associated with sperm DNA damage. *Andrologia* **51**, e13239 (2019).
35. Giacone, F. *et al.* Epigenetics of male fertility: effects on assisted reproductive techniques. *World J. Mens Health* **37**, 148–156 (2019).
36. Navarro-Costa, P. *et al.* Incorrect DNA methylation of the DAZL promoter CpG island associates with defective human sperm. *Hum. Reprod.* **25**, 2647–2654 (2010).
37. Li, B. *et al.* Altered DNA methylation patterns of the H19 differentially methylated region and the DAZL gene promoter are associated with defective human sperm. *PLoS ONE* **8**, e71215 (2013).
38. Palermo, G. D. *et al.* Shedding light on the nature of seminal round cells. *PLoS ONE* **11**, e0151640 (2016).
39. Eggert-Kruse, W., Bellmann, A., Rohr, G., Tilgen, W. & Runnebaum, B. Differentiation of round cells in semen by means of monoclonal antibodies and relationship with male fertility. *Fertil. Steril.* **58**, 1046–1055 (1992).
40. World Health Organization. *WHO Laboratory Manual for the Examination of Human Semen and Sperm-Cervical Mucus Interaction* 4th edn. (Cambridge University Press, Cambridge, 1999).
41. UK NEQAS Reproductive Science. <https://mft.nhs.uk/saint-marys/services/gynaecology/reproductive-medicine/ukneqasreproductive-science/> (2020).
42. Elder, R. H. *et al.* Alkylpurine-DNA-N-glycosylase knockout mice show increased susceptibility to induction of mutations by methyl methanesulfonate. *Mol. Cell. Biol.* **18**, 5828–5837 (1998).
43. Dumax-Vorzet, A. F., Tate, M., Walmsley, R., Elder, R. H. & Povey, A. C. Cytotoxicity and genotoxicity of urban particulate matter in mammalian cells. *Mutagenesis* **30**, 621–633 (2015).
44. Cooper, T. G. *et al.* World Health Organization reference values for human semen characteristics. *Hum. Reprod. Update* **16**, 231–245 (2010).

Acknowledgements

The authors wish to thank research nurses Lucy Dwyer and Claudette Wright and the clinical embryologist Tope Adeniyi for their help in setting up this study.

Author contributions

A.P., M.C., C.M., D.B. drafted the manuscript. A.P., M.C., C.M. and D.B. discussed and performed the statistical analysis. B.A. co-ordinated the study and data management was carried out by B.A., C.N. and A.P. B.A. undertook the lab assays for DNA damage and C.N. for DNA methylation assays. All members of the group contributed to the collection of data for the study and discussions on the design, conduct and interpretation of the results.

Funding

This work was supported by funding from Damascus University and the British Council to BA to support his PhD study and CN was supported by the Faculty of Science and Engineering, Manchester Metropolitan University Ph.D. studentship.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-021-81674-x>.

Correspondence and requests for materials should be addressed to A.C.P.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021