

Please cite the Published Version

Dos'Santos, Thomas ^(D), Thomas, Christopher, Comfort, Paul and Jones, Paul A (2022) Biomechanical Effects of a 6-Week Change of Direction Speed and Technique Modification Intervention: Implications for Change of Direction Side step Performance. Journal of Strength and Conditioning Research, 36 (10). pp. 2780-2791. ISSN 1064-8011

DOI: https://doi.org/10.1519/jsc.000000000003950

Publisher: Lippincott, Williams & Wilkins

Version: Accepted Version

Usage rights: (cc) BY-NC

Downloaded from: https://e-space.mmu.ac.uk/627372/

Creative Commons: Attribution-Noncommercial 4.0

Additional Information: Copyright 2021 National Strength and Conditioning Association

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines)

1	Biomechanical Effects of a Six-Week Change of Direction Speed and Technique
2	Modification Intervention: Implications for Change of Direction Side-Step Performance
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	
16	
17	
18	
19	

20 ABSTRACT

The aim of this study was to evaluate biomechanical effects of change of direction (COD) 21 speed and technique modification training on COD performance (completion time, ground 22 contact time [GCT], and exit velocity) during 45° (CUT45) and 90° (CUT90) side-step cutting. 23 A non-randomized, controlled 6-week intervention study was administrated. 15 male 24 multidirectional sport athletes (age:23.5±5.2 years;height:1.80±0.05 m;mass:81.6±11.4 kg) 25 formed the intervention group (IG) who participated in two 30-minute COD speed and 26 technique modification sessions per week, while 12 male multidirectional sport athletes 27 (age:22.2±5.0 years;height:1.76±0.08 m;mass:72.7±12.4 kg) formed the control group (CG) 28 29 and continued their normal training. All subjects performed six trials of the CUT45 and CUT90 task whereby pre-to-post intervention changes in lower-limb and trunk kinetics and kinematics 30 were evaluated using three-dimensional motion and ground reaction force analysis. Two-way 31 32 mixed analysis of variances revealed significant main effects for time (pre-to-post changes) for CUT45 completion time, exit velocity and CUT90 completion time ($p \le 0.045$, $\eta^2 = 0.152 - 0.539$), 33 and significant interaction effects of time and group were observed for CUT45 completion 34 time, GCT, exit velocity and CUT90 completion time ($p \le 0.010$, $\eta^2 = 0.239 - 0.483$), with the IG 35 displaying superior performance post-intervention compared to the CG ($p \leq 0.109$, g=0.83-36 37 1.35). Improvements in cutting performance were moderately to very largely associated ($p \le 0.078$, r or $\rho = 0.469 - 0.846$) with increased velocity profiles, increased propulsive forces 38 over shorter GCTs, and decreased knee flexion. COD speed and technique modification is a 39 simple, effective training method, requiring minimal equipment, that can enhance COD 40 performance which practitioners should consider incorporating into their pitch- or court-based 41 training programs. 42

43 Keywords: side-step; side-stepping; cutting; kinetics, external cues

44 INTRODUCTION

Change of direction (COD) ability is a fundamental movement associated with successful 45 performance in multidirectional sports (7, 29). Athletes can perform a range of angled 46 directional changes in sport (11, 12), and there are various different COD actions (side-step, 47 crossover cut, split step, pivot, spin maneuverer) performed which have their own distinct 48 49 biomechanical differences, advantages, and disadvantages (11). Specifically, side-step cutting is characterised by a lateral foot plant to push-off towards the opposite direction (11), and this 50 is an important and frequently performed action linked to decisive agility moments, such as 51 52 evading tackles to gain territorial advantage and pressing opponents (defensive agility) in multidirectional sports (11, 16, 40). Additionally, athletes also sprint and side-step to 53 predetermined locations on a pitch or court (29), such as wide-receiver running a pre-planned 54 "slant" route in American Football or a baseball player running around the bases to score a run. 55 Consequently, training interventions that can enhance an athlete's mechanical ability to side-56 step cut are of great interest to practitioners working in cutting dominant sports. 57

Specifically, COD speed is defined as "the ability to decelerate, reverse or change 58 movement direction and accelerate again" (22) and the determinants of COD speed are 59 multifaceted, underpinned by interactions between velocity (linear speed), mechanics (kinetics, 60 61 technique and postures), deceleration (braking strategy), and physical capacity (strength capacity, maximal and rapid force production, neuromuscular control, and muscle activation) 62 (11-13, 24, 28). Results of a recent meta-analysis confirmed that plyometric training, strength 63 training, COD speed training, and combined training methods positively enhance COD speed 64 and manoeuvrability performance (14). However, it worth noting that COD speed and 65 manoeuvrability performance during the successful training interventions is generally assessed 66 67 via completion time using electronic timing gates (31). Thus, because the majority of the time during these assessments is spent linear sprinting, these tasks are biased towards faster athletes 68

69 and confounded by linear speed (31). Importantly, the underlying biomechanical mechanisms which are responsible for enhanced COD speed performance are currently unclear. By only 70 71 assessing COD performance using completion time, it is uncertain whether an athlete's mechanical ability to COD has indeed improved, or whether reductions in completion times 72 have improved due to improvement linear speed capabilities. Consequently, for a better 73 74 understanding of whether training interventions improve an athlete's mechanical ability to change direction, more detailed analysis into COD biomechanics such as kinetics, ground 75 contact time (GCT), and velocity of centre of mass (COM) are required, which are suggested 76 77 to be better measures of COD ability (31, 33).

78 Kinetic, technical, and mechanical determinants of faster side-step cutting (30-90°) 79 performance include greater medio-lateral propulsive forces (MLPF) over short GCTs, greater ankle power and plantar-flexor moments, torso lean and pelvis rotation towards the intended 80 81 direction of travel, greater hip power and extensor moments, rapid knee and hip extension, wider lateral foot plants, reduced hip and knee flexion, low COM, faster velocity profiles, and 82 greater penultimate foot contact (PFC) horizontal braking forces (HBF) (for 90° cuts only) (11, 83 84 13, 19, 24, 27, 28, 37). Therefore, teaching and modifying athletes' side-step cutting technique by promoting the aforementioned favourable postures and mechanics associated with faster 85 86 cutting (e.g., reduced knee flexion, medial trunk lean and pelvis rotation, greater MLPFs, faster velocity profiles, greater PFC braking forces) could be an effective and simple training strategy 87 for enhancing side-step cutting performance (5, 9, 23). For example, in male soccer players, 88 six-weeks' COD speed and technique modification training focusing on earlier and greater PFC 89 braking, trunk lean towards the intended direction of travel, and rapid and forceful push-off 90 improved 70° side-cutting performance completion times and COD deficits (9). Acutely, 91 92 modifying knee flexion range of motion (ROM) has shown to influence GCT and exit velocity 93 during 45°side-step cutting (5), and 6-weeks' 180° COD technique modification training (PFC

94 braking, backwards trunk inclination, and neutral foot position) improved completion times in 95 female netballers (23). Although these results are promising, further research into the 96 biomechanical effects of COD speed and technique modification training is required to assist 97 in the development of more effective pitch- or court-based COD training programs for 98 practitioners working in multidirectional sports.

The aims of this study were two-fold: 1) to evaluate the effectiveness of a 6-week COD speed and technique modification intervention on COD performance (completion time, GCT, and exit velocity) during 45° (CUT45) and 90° (CUT90) side-step cutting; and 2) to identify which kinetic and kinematic factors explain changes in performance. It was hypothesized that a COD speed and technique modification program would improve COD performance in multidirectional athletes, and changes in GCT, velocity at key instances of the COD, propulsive forces, knee flexion ROM, and pelvic rotation will explain improvements in performance.

106 METHODS

107 Experimental approach to the problem

A non-randomized, controlled 6-week intervention study with a repeated measures pre-to-post 108 design was used (Figure 1). Male multidirectional sport athletes were recruited for the 109 intervention group (IG) which consisted of a 6-week COD speed and technique modification 110 training program (Supplemental digital content 1), consisting of two 30-minute sessions per 111 week. Conversely, male multidirectional sport athletes acted as the control group (CG). Pre-to-112 post assessments of 45° and 90° cutting performance and COD biomechanics were assessed 113 114 using three-dimensional (3D) motion and ground reaction force (GRF) analysis to monitor the effectiveness of the training intervention. Pre-and post-intervention testing was performed at 115 the same time of day for each subject to control for circadian rhythm. 116

Insert Figure 1 here

117

118 Subjects

Thirty men from multidirectional sports (amateur/semi-professional level) were recruited and participated in this study. Based on previous work (23), a minimum sample size of 14 per group was determined from an *a priori* power analysis using G*Power (Version 3.1, University of Dusseldorf, Germany) (15). This was based upon an effect size of 0.74 (pre-to-post change completion time), a power of 0.80, and alpha level of 0.05.

Sixteen males (soccer n = 12, rugby n = 4; age: 23.5 ± 5.2 years; height: 1.80 ± 0.05 m; 124 125 mass: 81.6 ± 11.4 kg) were recruited for the IG. Conversely, fourteen men (soccer n = 9, rugby n = 4, field hockey n = 1; age: 22.2 ± 5.0 years; height: 1.76 ± 0.08 m; mass: 72.7 ± 12.4 kg) 126 acted as the CG and continued their normal sport and resistance training sessions. The 127 investigation was approved by the Institutional Ethics Review Board (HSR1617-131), and all 128 subjects were informed of the benefits and risks of the investigation prior to signing an 129 institutionally approved consent form to participate in the study. All subjects from both groups 130 possessed at least five years' training experience in their respective sport and had never 131 sustained a severe knee injury prior to testing. All subjects had minimum one years' resistance 132 training experience, all performed two resistance training sessions a week and were all in a 133 strength mesocycle. At the time of the training intervention, all subjects completed two skills 134 sessions and played one competitive match a week. All of the procedures were carried out 135 during the competitive season to ensure that no large physical changes were made as a result 136 of the conditioning state . To remain as an active subject in the study and used for further 137 analysis, subjects were not allowed to miss more than two of the 12 sessions in total (i.e., \geq 138 83% compliance rate). Subsequently, due to match-related injuries or illness, one and two 139 subjects withdrew from the IG and CG, respectively, resulting in sample sizes of 15 and 12 140

- 141 (Figure 1). Subjects in the IG completed on average 11.9 ± 0.4 sessions (98.3 ± 3.5%), with 12
- subjects completing 12 (100%) sessions and three completing 11 sessions (91.7%).

143 <u>Procedures</u>

The warm up, cut, marker placement, and 3D motion analysis and procedures were based on 144 previously published methodologies (8, 24, 25, 28). In summary, each subject performed six 145 trials of the 45° and 90° (5 m entry and 3 m exit) side-step cut (right limb push- off) as fast as 146 possible and were provided with standardized footwear to control for shoe-surface interface 147 (Balance W490, New Balance, Boston, MA, USA). Completion time was assessed using two 148 sets of Brower timing lights placed at hip height at the start and finish (Draper, UT, USA), and 149 subjects adopted a two-point staggered stance, 0.5 m behind the start line. Marker and force 150 151 data were collected over the PFC and final foot contact (FFC) using ten Qualisys Oqus 7 152 (Gothenburg, Sweden) infrared cameras (240 Hz) operating through Qualisys Track Manager software (Qualisys, version 2.16 (Build 3520), Gothenburg, Sweden) and GRF's were 153 collected from two 600 mm × 900 mm AMTI (Advanced Mechanical Technology, Inc, 154 Watertown, MA, USA) force platforms (Model number: 600900) embedded into the running 155 track sampling at 1200 Hz, respectively. Using the pipeline function in visual 3D, joint 156 coordinate (marker) and force data were smoothed using a Butterworth low-pass digital filter 157 with cut-off frequencies of 15 and 25 Hz, respectively, based on *a priori* residual analysis (39) 158 and visual inspection of motion data. The kinematic model process was based on previous 159 reported methodologies (8, 24, 25, 28). Lower limb joint moments were calculated using an 160 inverse dynamics approach (39) through Visual 3D software (C-motion, version 6.01.12, 161 Germantown, USA) and were defined as external moments and normalised to body mass. Joint 162 kinematics and GRFs were also calculated using Visual 3D, while GRF braking and propulsive 163 characteristics were normalised relative to body weight, with vertical, anterior-posterior, and 164

165

166

medio-lateral corresponding to Fz, Fx, and Fy, respectively. Horizontal COM velocity profiles over key instances of the COD were calculated as described previously (24, 25).

167 Supplemental digital content 2 provides a full description the variables examined, definitions, and calculations. Briefly, the following kinetic and kinematics were examined 168 during the FFC for both tasks: GCT and velocity at FFC (touchdown) and exit velocity (toe-169 off); mean and peak MLPF; hip, knee, ankle dorsi-flexion moments and ROM (sagittal plane 170 joint moment and angles); lateral trunk flexion angle, forward trunk inclination angle, pelvis 171 rotation angle, initial foot progression angle (IFPA), and lateral foot plant distance (trunk, 172 pelvis, and foot variables). Additionally, mean horizontal propulsive force was examined 173 during the FFC for CUT90, mean HBF was examined during the FFC and PFC only during 174 175 CUT90, while PFC GCT and approach velocity (velocity at PFC) was also examined. 176 Completion time, GCT, and exit velocity were the considered primary outcome variables while approach velocity, velocity at FFC, and cut angle were also secondary variables to assess 177 178 cutting performance. Five trials were used in the analysis of each subject based and the average of individual trial peaks for each variable were calculated (8). A subset of the sample (n=10)179 performed the cuts on two separate occasions separated by 7 days to establish between-session 180 reliability. The reliability measures for COD biomechanics data was considered high (intraclass 181 correlation coefficient = 0.704-0.958, coefficient of variation = 2.4-11.9%). 182

183 <u>6-week COD speed and technique modification training intervention</u>

A six-week COD speed and technique modification intervention described in Supplemental digital content 1, was performed by the IG twice a week (30 minutes per session), with minimum 48 hours between sessions. The intervention was slightly adapted from a previously successful six-week COD speed and technique modification training intervention (9), which focused on pre-planned low intensity decelerations, cuts, and turns (weeks 1-2), before 189 progressing intensity via velocity and angle (weeks 3-4) and introducing a stimulus with increased intensity (weeks 3-6) to increase complexity, target technique integrity (reinforce 190 optimal technical), and to provide variation within the COD development framework (7, 11, 191 30). The COD program was in accordance with COD speed training recommendations (7, 13), 192 and the duration, distances, and number of CODs were similar to previous research (9, 23). 193 194 The sessions were led by the principle researcher who is a certified strength and conditioning specialist with extensive experience in coaching COD speed and agility drills. All sessions took 195 place in the Human Performance Laboratory, on the same surface as used for 3D motion 196 197 analysis. Athlete to coach ratios ranged from 5-8:1.

The COD technique modification intervention focused on three aspects outlined in Table 1 based on the success of a previous COD speed and technique modification and training recommendations (7, 9, 11, 23, 28). Subjects were given individual feedback regarding their technique, and external verbal coaching cues were used to promote faster performance and facilitate better motor skill retention (11, 32, 38).

203

Insert Table 1 here

204 <u>Statistical Analyses</u>

All statistical analyses were performed using SPSS v 25 (SPSS Inc., Chicago, IL, USA) and 205 Microsoft Excel (version 2016, Microsoft Corp., Redmond, WA, USA). Normality was 206 inspected for all variables using a Shapiro-Wilks test. A two-way mixed analysis of variance 207 (ANOVA) (group; time) with group as a between-subjects factor measured at 2 levels (IG and 208 CG), and time (pre- and post-training measures) the within-subject factor. This was used to 209 identify any significant main (time) or interaction (group \times time) effects for outcome variables 210 between IG and CG, pre-to-post testing. A Bonferroni-corrected pairwise comparison design 211 was used to further analyze the effect of the group when a significant interaction effect was 212

observed for time and group. Partial eta squared effect sizes were calculated for all ANOVAs with the values of 0.010-0.059, 0.060-0.149, and \geq 0.150 considered as small, medium, and large, respectively (4).

Pre-to-post changes in variables for each group were assessed using paired sample t-216 tests (parametric) and Wilcoxon-sign ranked tests (non-parametric). Magnitudes of differences 217 were assessed using Hedges' g effect sizes with 95% CIs, and interpreted as trivial (≤ 0.19), 218 small (0.20-0.59), moderate (0.60-1.19), large (1.20-1.99), very large (2.00-3.99), and 219 220 extremely large (≥ 4.00) (21). Mean changes were also calculated and interpreted as ratios relative to the smallest worthwhile change (SWC). The SWC was calculated as 0.2 × between-221 subject SD. Comparisons in post-intervention primary outcome variables and changes in 222 outcome variables between the IG and CG were also assessed using independent sample t-tests 223 224 or Mann-Whitney U tests, with effect sizes as outlined above. Furthermore, to link changes in COD performance with cutting kinetic and kinematic changes, Pearson's correlations 225 226 (parametric) or Spearman's correlations (non-parametric) were calculated with 95% CIs, with p values Bonferroni corrected to control for type 1 error. Correlations were interpreted as trivial 227 (0.00-0.09), small (0.10-0.29), moderate (0.30-0.49), large (0.50-0.69), very large (0.70-228 0.89), nearly perfect (0.90–0.99), and perfect (1.00) (20). A correlation cut-off value of ≥ 0.40 229 was considered relevant (36). Thus, correlations greater than this value are only reported. 230 Statistical significance was defined as $p \le 0.05$ for all tests. 231

232 **RESULTS**

Results of the two-way mixed ANOVAs are presented in Tables 2 and 4, and pre-to-post changes in cutting biomechanics for the IG and CG are presented in Tables 3 and 5, containing descriptive statistics, *p* values, effect sizes, and mean differences.

236 <u>CUT45: Pre-to-post changes in cutting biomechanical variables</u>

Insert Table 2 here

238	Large and significant interaction effects of time and group for all CUT45 performance
239	variables were observed (Table 2), with the IG displaying moderately to largely better
240	performance variables post-intervention compared to the CG ($p \le 0.109$, $g = 0.62-1.21$).
241	Moderate and significant improvements in completion time, exit velocity, approach velocity,
242	and velocity at FFC were observed for the IG only, while a small and significantly shorter GCT
243	was also observed (Table 3). Medium to large significant interaction effects of time and group
244	were observed for CUT45 peak and mean MLPF, and knee flexion angle (KFA) ROM (Table
245	2). Intervention group subjects produced significantly small to moderately greater peak and
246	mean MLPFs, greater IFPAs, and smaller knee and ADFA ROMs post-intervention (Table 3).
247	Control group subjects demonstrated significantly greater IFPAs post-intervention only (Table
248	3).
249	***Insert Figure 2 here***
249 250	***Insert Figure 2 here*** ***Insert Table 3 here***
249 250 251	***Insert Figure 2 here*** ***Insert Table 3 here*** <u>CUT90: Pre-to-post changes in performance biomechanical variables</u>
249 250 251 252	<pre>***Insert Figure 2 here***</pre>
249 250 251 252 253	<pre>***Insert Figure 2 here***</pre>
249 250 251 252 253 254	<pre>***Insert Figure 2 here***</pre>
249 250 251 252 253 254 255	<pre>***Insert Figure 2 here*** ***Insert Table 3 here*** CUT90: Pre-to-post changes in performance biomechanical variables ***Insert Table 4 here*** A large and significant interaction effect of time and group for CUT90 completion time was observed (Table 4), with the IG showing significantly faster post-intervention completion times (p=0.031, g=-1.35) compared to the CG. Additionally, a medium and non-significant</pre>
249 250 251 252 253 254 255 256	<pre>***Insert Figure 2 here***</pre>

5 and Figure 3), respectively, for the IG only. Medium to large significant interaction effects

moderate significant improvement was observed for completion time and exit velocity (Table

258

of time and group were observed for CUT90 mean MLPF, peak and mean HPF, FFC mean 260 HBF, FFC peak ADFM, IFPA, and knee and ADFA ROM (Table 4). Intervention group 261 subjects produced significantly small to moderately greater peak ADFMs, greater peak and 262 mean MLPFs, greater peak and mean HPFs, and greater FFC and PFC mean HBFs (Table 5). 263 Additionally, IG subjects demonstrated significantly small to moderately greater forward trunk 264 inclination angles, greater pelvic rotation angles, greater IFPAs, smaller KFA ROM, and 265 greater ADFA ROM post-intervention (Table 5). Control group subjects produced significantly 266 greater peak MLPFs and greater forward trunk inclination angles post-intervention only (Table 267 268 5).

- 269 ***Insert Figure 3 here***
- 270 ***Insert Table 5 here***

Associations between changes in performance primary outcome variables with cutting mechanical and technical variables

Improvements in CUT45 completion time were very largely associated with increased exit 273 velocity; largely associated with decreased FFC GCT; and moderately associated with 274 decreased KFA ROM, increased mean MLPF, and increased FFC velocity (Table 6). 275 Additionally, CUT45 FFC GCT improvements were very largely associated with decreased 276 277 KFA ROM; and largely associated with increased approach and exit velocity (Table 6). Finally, CUT45 exit velocity improvements were very largely associated with increased FFC velocity; 278 and largely associated with decreased KFA ROM and FFC GCT, and increased approach 279 280 velocity (Table 6).

Improvements in CUT90 completion time were largely associated with increased peak
 KFM, increased FFC peak HBF, and decreased PFC GCT; and moderately associated with

increased ADFA ROM (Table 6). Additionally, CUT90 FFC GCT improvements were very
largely associated with increased FFC velocity, and decreased hip and knee flexion ROM; and
largely associated with increased exit velocity and lateral foot plant distance (Table 6). Finally,
CUT90 exit velocity improvements were very largely associated with increased FFC velocity;
largely associated with decreased FFC GCT, decreased HFA ROM, increased PFC mean HBF;
and moderately associated with increased FFC peak KFM, and lateral foot plant distance (Table 6).

290

Insert Table 6 here

291 **DISCUSSION**

The aims of this study were two-fold: 1) to examine the biomechanical effects of a 6-week 292 COD speed and technique modification intervention on side-step cutting performance; and 2) 293 to understand which technical and mechanical changes during cutting explained improvements 294 295 in performance. The primary findings were that COD speed and technique modification resulted in meaningful improvements in cutting performance, which were meaningfully greater 296 than the CG (Tables 2-5 Figures 2-3), supporting the study hypotheses. The positive 297 298 improvements in cutting performance were primarily attributed to increases in velocity at key instances of the PFC and FFC, increases in propulsive force over shorter GCTs, and reduced 299 knee flexion, supporting the study hypotheses. Consequently, COD speed and technique 300 modification appears to be a training strategy effective for improving COD performance, which 301 practitioners can simply and easily administer on the pitch or court. 302

303 COD speed training has been reported to enhance COD completion time (9, 14); 304 however, the technical and biomechanical mechanisms underpinning the improvements in 305 COD performance are relatively unknown, and it is not fully understood whether the 306 improvements in completion time were attributed to improvements in COD ability or linear 307 speed (31). This study is one of the first to provide a novel insight into the mechanical and technical changes underpinning improvements in 45° and 90° cutting performance. Although 308 clear improvements in cutting completion times were observed in the present study (Table 3 & 309 5, Figure 2-3), importantly shorter GCTs, greater propulsive forces, and enhancements in 310 velocity at key instances of the COD were observed. Thus, these results confirm that the 311 312 athletes' mechanical ability to COD has indeed improved via COD speed and technique modification training with external verbal cues. External cueing has been recommended for 313 speed development (38) and has been shown to enhance COD performance (32), while 314 315 promoting changes in side-stepping technique (2). Additionally, external cues are reported to be effective for motor skill development and retention (2, 11, 38). In the present study, we 316 found the specific external cues, in addition to COD speed and technique modification training, 317 "slam on the bakes", "cushion and push the ground away", and "face towards the direction of 318 travel", to be effective in improving side-step performance and promoting favourable 319 techniques and mechanics linked to faster performance (Table 3 & 5). 320

Importantly, the present study was able to explore some of the underlying technical and 321 mechanical mechanisms responsible for enhanced cutting performance. The key and primary 322 changes in cutting mechanics were as follows: increased velocity at key instances of the PFC 323 324 and FFC, increased mean and peak propulsive forces in shorter FFC GCTs, reduced knee flexion ROM, and greater PFC mean HBFs (90° cut only) which were moderately to very 325 largely associated with improvements in cutting performance (Table 6). The fact that the 326 abovementioned variables were associated with improvements in COD performance are 327 unsurprising because they have been identified as key determinants of faster COD performance 328 (11, 13, 19, 27, 28, 37). For example, Hader et al. (18) demonstrated greater minimum 329 330 velocities and velocity maintenance were strong determinants of faster 45° and 90° cut performance, thus increasing an athlete's ability to attain greater horizontal displacements in 331

shorter times. Greater propulsive forces increase impulse which, based on the impulsemomentum relationship, leads to greater changes in momentum, thus velocity (11), and this
was confirmed by the moderate association between MLPF changes and exit velocity (Table
6).

Additionally, shorter GCTs are associated with faster performance (10, 27, 34, 36, 37), 336 which is advantageous because the athletes will spend less time braking and propelling, thus 337 less time performing the COD action, and potentially greater utilisation of the stretch-338 shortening cycle (3, 36). IG subjects demonstrated smaller ADFA ROMs for CUT45 339 performance post-intervention (Table 3). Reduced ankle dorsi-flexion has been reported to play 340 a critical role in early acceleration sprint performance, mostly likely contributing to a stiffer 341 ankle joint; thus, potentially contributing to the faster performance observed in the present 342 study. Finally, reduced KFA ROM is important for reducing GCT and increasing exit velocity, 343 as confirmed by the results from this study and previous research (5). This potentially results 344 in a rapid transition from braking (eccentric) to propulsion (concentric) and therefore 345 facilitating a shorter GCT and greater utilisation of the stretch shortening cycle (increased 346 elastic energy) to permit more effective force generation (3). This positive adaptation could be 347 attributed to the COD speed and technique modification training with the external cue to "push 348 349 or punch the ground away."

High propulsive forces (10, 19, 28, 33, 34, 36) over short GCTs (10, 27, 28, 34, 36, 37) are strong determinants of faster COD performance. The finding of increased propulsive force production in shorter GCTs following COD technique modification training is a positive outcome (Tables 3 & 5), and is similar to de Hoyo et al. (6) who found 10-weeks eccentric overload training resulted in likely to almost certain increases in relative peak braking and propulsive forces, shorter braking and propulsive times thus shorter GCTs, and increases in relative total and braking impulse during side-step cutting. Unfortunately, de Hoyo et al. (6) did not examine completion time or exit velocity; however, improvements in completion time training have been observed following flywheel eccentric training (17, 35); thus, it is speculated that the improved side-step kinetics from flywheel training should improve COD speed performance. Similarly, Bencke et al. (1) reported improvements in vertical propulsive force, GCT, and concentric phase duration following 12-weeks neuromuscular warm-up, corroborating the mechanism of improved completion time in the present study.

To author's knowledge, King et al. (26) is the only other research group to document 363 changes in cutting biomechanics and some performance indices following a training 364 intervention. No significant changes in approach velocity were observed following a 365 multicomponent training programme (resistance training, running drills, and COD drills with 366 focus on intersegmental control); however, a small reduction in GCT was observed which is 367 an important determinant of COD performance. Unfortunately, King et al. (26) did not examine 368 completion time or exit velocity, but did report reductions in lateral trunk flexion, increased 369 pelvic rotation towards the intended direction, increased COM to centre of pressure distance, 370 increased ankle plantar-flexor moments, and decreased peak KFA, which have all been 371 associated with faster cutting completion times (5, 19, 27, 37). Although no significant changes 372 373 in lateral trunk flexion or foot plant distance were observed in the present study (Tables 3 & 5), similar to King et al. (26), subjects post-intervention in the IG demonstrated increased pelvic 374 rotation, decreased KFA ROM, and increased ADFMs, which have previously been associated 375 with faster cutting performance (5, 19, 27, 37). Nevertheless, the results from this study confirm 376 that COD speed and technique modification training with external verbal coaching cues 377 regarding technique result in positive improvements in side-stepping performance. Therefore, 378 379 practitioners should consider implementing this form of training in their pitch- or court-based training programs when working with multidirectional athletes. 380

381 It is worth noting that the present study has several limitations. Firstly, it should be acknowledged that only CUT45 and CUT90 side-stepping was investigated with a right limb 382 push-off. As the biomechanical demands are angle- and task-dependent (12, 31), caution is 383 advised extrapolating the findings from the present study to CODs of different angles, actions 384 and populations. Therefore, further research is necessary that investigates the effect of COD 385 technique modification on sharper CODs and different COD actions (i.e. crossover cut, split 386 step). Similarly, it is worth acknowledging that the cutting actions were examined in a pre-387 planned, controlled laboratory environment. Future research should consider investigating the 388 389 effect of COD speed and technique modification training on agility performance. Lastly, it is unknown whether the technique can be maintained for extensive periods and unknown what 390 happens to cutting biomechanics when this form of training is discontinued. Thus, further 391 investigation into short- and long-term retention of modified cutting technique is required to 392 improve our understanding of training prescription and dosages. Notwithstanding these 393 limitations, COD provides the mechanical and physical basis for agility (29), thus 394 improvements in the mechanical ability to change direction (i.e., fast mover) should partially 395 396 transfer to improved agility (7, 29, 30). Additionally, athletes sprint and change direction to pre-determined locations on a court or pitch (29), thus the findings of this study have important 397 implications for the development of effective COD speed training programs. 398

399 PRACTICAL APPLICATIONS

This is the first study to comprehensively examine the biomechanical effects of COD speed and technique modification on side-step cutting performance, finding meaningful improvements in side-step cutting performance. The positive improvements in performance were primarily attributed to increases in velocity at key instances of the COD, increases in propulsive force production over shorter GCTs, and decreased KFA ROM. Consequently, 6weeks' COD speed and technique modification training with externally directed verbal coaching cueing ("slam on the bakes, cushion and push the ground away, and face towards the direction of travel") is an effective training strategy to enhance COD performance. Practitioners should therefore consider incorporating this form of training (2×30 -minute sessions a week) simply and easily into their pitch- or court-based training programs to enhance side-step cutting performance.

411 ACKNOWLEDGEMENTS

The authors would like to thank Callum Stratford, Alistair McBurnie, Laura Smith, and Steve Horton for their assistance with data collection. This study was funded by the National Strength and Conditioning Association (NSCA) Foundation, as part of the Doctoral Grant Scheme (awarded 2018). Funding was used for subject recruitment only. Thomas Dos'Santos, Christopher Thomas, Paul Comfort, and Paul Jones declare that they have no conflicts or competing interests relevant to the content of this article, and the findings of the study do not represent the views of the NSCA.

419 **REFERENCES**

- Bencke J, Næsborg H, Simonsen EB, and Klausen K. Motor pattern of the knee joint muscles during side-step cutting in European team handball. *Scand J Med Sci Spor* 10: 68-77, 2000.
- Benjaminse A, Welling W, Otten B, and Gokeler A. Transfer of improved movement
 technique after receiving verbal external focus and video instruction. *Knee Surgery, Sports Traumatology, Arthroscopy* 26: 955-962, 2018.
- Bosco C, Komi PV, and Ito A. Prestretch potentiation of human skeletal muscle during
 ballistic movement. *Acta Physiologica Scandinavica* 111: 135-140, 1981.
- 427 4. Cohen J. Statistical analysis for the behavioral sciences. *Hillsdale: Lawrance Erlbaum*, 1988.
- 428 5. Dai B, William EG, Michael TG, Darin AP, Robin MQ, and Bing Y. The Effects of 2
 429 Landing Techniques on Knee Kinematics, Kinetics, and Performance During Stop-Jump and
 430 Side-Cutting Tasks. *Am J Sport Med* 43: 466-474, 2014.
- de Hoyo M, Sañudo B, Carrasco L, Mateo-Cortes J, Domínguez-Cobo S, Fernandes O, Del
 Ojo JJ, and Gonzalo-Skok O. Effects of 10-week eccentric overload training on kinetic
- parameters during change of direction in football players. *J Sports Sci* 34: 1380-1387, 2016.
 DeWeese BH and Nimphius S. Program Design Technique for Speed and Agility Training,
- 435 in: *Essentials of Strength Training and Conditioning*. GG Haff, NT Triplett, eds. Champaign:
 436 Human Kinetics, 2016, pp 521-558.

437	8.	Dos'Santos T, Comfort P, and Jones PA. Average of trial peaks versus peak of average
438		profile: impact on change of direction biomechanics. Sport Biomech 19: 483-492, 2020.
439	9.	Dos'Santos T, McBurnie A, Comfort P, and Jones PA. The Effects of Six-Weeks Change of
440		Direction Speed and Technique Modification Training on Cutting Performance and
441		Movement Quality in Male Youth Soccer Players. Sports 7: 205, 2019.
442	10.	Dos'Santos T, Thomas C, Jones PA, and Comfort P. Mechanical determinants of faster
443		change of direction speed performance in male athletes. J Strength Cond Res 31: 696-705,
444		2017.
445	11.	Dos'Santos T, McBurnie A, Thomas C, Comfort P, and Jones PA. Biomechanical
446		Comparison of Cutting Techniques: A Review and Practical Applications. Strength Cond J
447		41: 40-54, 2019.
448	12.	Dos'Santos T, Thomas C, Comfort P, and Jones PA. The effect of angle and velocity on
449		change of direction biomechanics: an angle-velocity trade-off. Sports Med 48: 2235-2253,
450		2018.
451	13.	Dos'Santos T, Thomas C, Comfort P, and Jones PA. The Role of the Penultimate Foot
452		Contact During Change of Direction: Implications on Performance and Risk of Injury.
453		Strength Cond J 41: 87-104, 2019.
454	14.	Falch HN, Rædergård HG, and van den Tillaar R. Effect of Different Physical Training Forms
455		on Change of Direction Ability: A Systematic Review and Meta-analysis. Sports Medicine-
456		<i>Open</i> 5: 53, 2019.
457	15.	Faul F, Erdfelder E, Buchner A, and Lang A-G. Statistical power analyses using G* Power
458		3.1: Tests for correlation and regression analyses. <i>Behav Res Methods</i> 41: 1149-1160, 2009.
459	16.	Fox AS, Spittle M, Otago L, and Saunders N. Offensive agility techniques performed during
460		international netball competition. Int J Sports Sci Coach 9: 543-552, 2014.
461	17.	Gonzalo-Skok O, Tous-Fajardo J, Valero-Campo C, Berzosa C, Bataller AV, Arjol-Serrano
462		JL, Moras G, and Mendez-Villanueva A. Eccentric-overload training in team-sport functional
463		performance: Constant bilateral vertical versus variable unilateral multidirectional
464		movements. Int J Sports Physiol and Perform 12: 951-958, 2017.
465	18.	Hader K, Palazzi D, and Buchheit M. Change of Direction Speed in Soccer: How Much
466		Braking is Enough? <i>Kineziologija</i> 47: 67-74, 2015.
467	19.	Havens KL and Sigward SM. Cutting mechanics: relation to performance and anterior
468		cruciate ligament injury risk. Med Sci Sports Exerc 47: 818-824, 2015.
469	20.	Hopkins WG. Measures of reliability in sports medicine and science. Sports Med 30: 1-15,
470		2000.
471	21.	Hopkins WG. A scale of magnitudes for effect statistics. A new view of statistics: Retrieved
472		from http://sportsci.org/resource/stats/effectmag.html, 2002.
473	22.	Jones P, Bampouras TM, and Marrin K. An investigation into the physical determinants of
474		change of direction speed. J Sports Med Phys Fitness 49: 97-104, 2009.
475	23.	Jones PA, Barber OR, and Smith LC. Changing pivoting technique reduces knee valgus
476		moments. Free Communication. Journal of Sports Sciences. J Sports Sci (BASES Annual
477		Conference) 33: S62, 2015.
478	24.	Jones PA, Dos' Santos T, McMahon JJ, and Graham-Smith P. Contribution of Eccentric
479		Strength to Cutting Performance in Female Soccer Players. J Strength Cond Res: Published
480		ahead of print 2019.
481	25.	Jones PA, Thomas C, Dos'Santos T, McMahon J, and Graham-Smith P. The Role of
482		Eccentric Strength in 180° Turns in Female Soccer Players. Sports 5: 42, 2017.
483	26.	King E, Franklyn-Miller A, Richter C, O'Reilly E, Doolan M, Moran K, Strike S, and Falvey
484		E. Clinical and biomechanical outcomes of rehabilitation targeting intersegmental control in
485		athletic groin pain: prospective cohort of 205 patients. Br J Sports Med 52: 1054-1062, 2018.
486	27.	Marshall BM, Franklyn-Miller AD, King EA, Moran KA, Strike S, and Falvey A.
487		Biomechanical factors associated with time to complete a change of direction cutting
488		maneuver. J Strength Cond Res 28: 2845-2851, 2014.

- 489 28. McBurnie A, Dos' Santos T, and Jones PA. Biomechanical Associates of Performance and Knee Joint Loads During an 70-90° Cutting Maneuver in Sub-Elite Soccer Players. J Strength 490 Cond Res: Published Ahead of print., 2019. 491 29. Nimphius S. Increasing Agility, in: High-Performance Training for Sports. D Joyce, D 492 493 Lewindon, eds. Champaign, IL.: Human Kinetics, 2014, pp 185-198. 494 30. Nimphius S. Training change of direction and agility, in: Advanced Strength and 495 Conditioning. A Turner, P Comfort, eds. Abdingdon, Oxon, United Kingdom: Routledge,
- 2017, pp 291-308.
 31. Nimphius S, Callaghan SJ, Bezodis NE, and Lockie RG. Change of Direction and Agility
- 498Tests: Challenging Our Current Measures of Performance. Strength Cond J 40: 26-38, 2017.
- 499 32. Porter J, Nolan R, Ostrowski E, and Wulf G. Directing attention externally enhances agility
 500 performance: A qualitative and quantitative analysis of the efficacy of using verbal
 501 instructions to focus attention. *Front Psychol* 1: 216, 2010.
- Spiteri T, Cochrane JL, Hart NH, Haff GG, and Nimphius S. Effect of strength on plant foot
 kinetics and kinematics during a change of direction task. *Eur J Sports Sci* 13: 646-652, 2013.
- Spiteri T, Newton RU, Binetti M, Hart NH, Sheppard JM, and Nimphius S. Mechanical
 determinants of faster change of direction and agility performance in female basketball
 athletes. J Strength Cond Res 28: 2205–2214, 2015.
- Tous-Fajardo J, Gonzalo-Skok O, Arjol-Serrano JL, and Tesch P. Enhancing Change-of Direction Speed in Soccer Players by Functional Inertial Eccentric Overload and Vibration
 Training. Int J Sports Physiol and Perform 11: 66-73, 2015.
- 36. Welch N, Richter C, Franklyn-Miller A, and Moran K. Principal Component Analysis of The
 Associations Between Kinetic Variables In Cutting And Jumping, And Cutting Performance
 Outcome. *J Strength Cond Res*: Published ahead of print, 2019.
- 37. Welch N, Richter C, Franklyn-Miller A, and Moran K. Principal Component Analysis of the
 Biomechanical Factors Associated With Performance During Cutting. *J Strength Cond Res*:
 Published Ahead of Print, 2019.
- 516 38. Winkelman NC. Attentional Focus and Cueing for Speed Development. *Strength Cond J* 40: 13-25, 2018.
- 518 39. Winter DA. *Biomechanics and motor control of human movement*. John Wiley & Sons, 2009.
- 40. Zahidi NNM and Ismail SI. Notational analysis of evasive agility skills executed by attacking
 ball carriers among elite rugby players of the 2015 Rugby World Cup. *Movement Health Ex*7: 99-113, 2018.

522