

Please cite the Published Version

Sanderson, Andrew , Cescon, Corrado, Heneghan, Nicola, Rushton, Alison, Barbero, Marco and Falla, Deborah (2019) What is the Influence of Low Back Pain on Muscle Activity and Movement during a Cyclical Dynamic Task? In: Pain Science in Motion III (PSIM), 31 May 2019 - 02 June 2019, Savona, Italy.

DOI: https://doi.org/10.1097/PR9.000000000000753

Publisher: Lippincott, Williams & Wilkins

Version: Published Version

Downloaded from: https://e-space.mmu.ac.uk/627665/

Usage rights:

(cc) BY

Creative Commons: Attribution 4.0

Additional Information: Abstract

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines)

What is the influence of low back pain on muscle activity and movement during a cyclical dynamic task?

Andy Sanderson^a, Corrado Cescon^b, Nicola Heneghan^a, Alison Rushton^a, Marco Barbero^b, Deborah Falla^a

^aCPR Spine, University of Birmingham, ^bUniversity of Applied Sciences and Arts of Southern Switzerland, SUPSI

Introduction: Low back pain (LBP) is a leading cause of disability. High-density electromyography (HDEMG) has revealed differences in the spatial distribution of back muscle activity during movements. However, these studies have only considered a small portion of the erector spinae (ES) during tasks which were either static or mono-planar.

Methods: This study combines HDEMG and kinematic analysis to investigate the effect of LBP on the spatial distribution of ES activity during a dynamic lifting task. Sixteen people with chronic LBP (8 male, age: 26.9 ± 10.8 years) and 16 age and gender-matched controls (7 male, age: 31.7 ± 14.0 years) completed the study. HDEMG signals from the ES were detected bilaterally by four 64-channel semi-disposable 13×5 electrode grids. Kinematic surface markers were placed over the back in triangular arrangements, creating lumbar and thoracic segments to track movement. HDEMG and kinematic data were recorded continuously during a dynamic task involving the cyclical lifting of a 5 kg box between 6 shelves for 10 cycles (~7 minutes). The shelves were arranged around the participant, at knee and sternal height.

Results: Data analysis is underway; with full results presented at the conference.

Conclusions: This study will impact on our understanding of neuromuscular adaptations to LBP during functional activity.