

Please cite the Published Version

McBurnie, Alistair J and Dos'Santos, Thomas (2022) Multidirectional Speed in Youth Soccer Players: Theoretical Underpinnings. Strength & Conditioning Journal, 44 (1). pp. 15-33. ISSN 1524-1602

DOI: https://doi.org/10.1519/ssc.00000000000658

Publisher: Lippincott, Williams & Wilkins

Version: Accepted Version

Downloaded from: https://e-space.mmu.ac.uk/628047/

Usage rights: Creative Commons

Creative Commons: Attribution-Noncommercial 4.0

Additional Information: This is a non-final version of an article published in final form in McBurnie, Alistair J.; Dos'Santos, Thomas, Multidirectional Speed in Youth Soccer Players: Theoretical Underpinnings, Strength and Conditioning Journal: February 2022 - Volume 44 - Issue 1 - p 15-33 doi: 10.1519/SSC.00000000000658. Copyright 2021 by the National Strength & Conditioning Association.

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines)

Multi-Directional Speed in Youth Soccer Players: Theoretical Underpinnings

1 Abstract

2 This review will provide a definition for 'multi-directional speed' (MDS) and evaluate its 3 technical and mechanical underpinnings. This review will explore each component of MDS while considering unique aspects of youth physiology and epidemiology. With a theoretical 4 5 understanding of MDS, practitioners will be more informed on the planning and periodization of MDS training methods in soccer. MDS comprises of linear speed, change of direction 6 7 speed, curvilinear speed, contextual speed, and agility, which each have distinct physiological, 8 biomechanical, and neuro-cognitive distinctions that can either be differentiated or harmonized 9 to optimise training.

10 Key words: sprinting, change of direction, agility, soccer, youth athletic development.

11

12 Introduction

Soccer is a team sport which is intermittent in nature, where frequent changes in locomotor 13 14 patterns (e.g., walking, running, sprinting), directional changes (e.g., cutting, pivoting), and sports-specific tasks (e.g., tackles, headers, dribbles) occur (19,184). Professional soccer 15 players have become faster over time (84), and although data are highly position-specific 16 17 (22,29,53,54), both the total sprinting distance and number of sprints (> 25.2 km h⁻¹) performed in match-play have increased 35% and 85%, respectively, in the English Premier 18 19 League from the 2006-06 to the 2012-13 season (10). Maximal sprint velocities are typically 20 attained within 30 to 40 m in soccer (198), of which can be initiated from a moving start (198). Subsequently, players can often attain between 85 to 94% of their peak sprint velocities in 21 match-play (76), which suggests that soccer players can often have the opportunity to achieve 22 maximum sprint speeds in games when initiating sprints while already moving. Of note, 23 24 approximately 85% of all maximum velocity actions that are executed in elite soccer present 25 some degree of curvature (31). Another consideration is that of actions considered to be 'lowspeed' activity, that still require high energy outputs (33). The large majority of sprinting actions 26

are typically performed within distances of 20 m (53,184,199) and data have shown that accelerations and decelerations (>2 m·s²) can contribute to as much as 7 to 10% and 5 to 7% of total player game load, respectively (44). Moreover, on a qualitative level, players can perform a mean of 30.7% of game actions and maneuvers in backward, lateral, diagonal and arced directions, and perform an average of 726 ± 203 turns in a game (19).

32 Importantly, it is each of these above-mentioned high-intensity and sprint-type actions that are considered to be vital determinants of successful performance in soccer (34). For example, 33 34 sprinting has been identified as the most frequent action associated with goal scoring scenarios (63). Furthermore, acceleration and maximum sprinting speed capabilities, along 35 with the ability to perform sudden and unpredictable high-intensity change of direction (COD) 36 manoeuvers while in motion, have been shown to differentiate between players from different 37 38 playing standards (84,165). Arguably, then, speed can and should be seen as one of the most sought after physical qualities in sport (136,143,161). Yet these high physiological and 39 40 mechanical demands also present an inherent risk of injury, due to the repeated nature of high intensity movements such as jumping, cutting, and contact with opposition players, particularly 41 42 when fatigued (46). Consequently, it should be a priority for practitioners to develop an athlete's capacity and robustness to perform multi-directional, high-speed actions, while 43 ensuring these manoeuvers are underpinned by movement quality. These conditioning 44 elements should be key in the development of successful performance and injury risk 45 46 mitigation strategies in soccer.

In a male youth soccer context (youth herein referring to both children, boys up to 13 years of age, and adolescents, boys aged 14-18 years) the aims of strength and conditioning methods are to enhance long-term athletic development (LTAD) and subsequent progression into the professional game (35,73,110,165). Furthermore, it has been highlighted in cross-sectional analyses that speed development increases in youth and may in fact plateau early on in a professional player's senior career (e.g., before 20-22 years old) (84). Resultantly, if movement skill is not well developed at late junior or early senior age, it may become more 54 challenging to improve these capabilities to the standard required when performing at the 55 senior level. Practitioners should, however, be aware that training the youth athlete requires 56 careful consideration, given the significant individual variability in factors that affect the pathway of growth, development and maturation within their developmental journey 57 58 (120,121,144). Youth soccer players should not been seen as 'mini adults', as they can 59 possess markedly different physical and physiological processes that attribute to their 60 performance in comparison to their senior peers (155,168). This also extends to injury risk, 61 where youth athletes present maturity-specific mechanisms that underlie certain injury risk 62 factors within the youth soccer setting (163).

In order to programme multi-directional speed training for youth athletes, the age- and 63 maturity-specific movement demands of youth soccer need to be accurately classified to guide 64 65 appropriate training prescription (155). Not only may this allow for specific movement skills to be trained more effectively during different phases of development, but also allow for the dose-66 response relationship of training to be uncovered with a long-term vision in mind. With that 67 said, although research in this area is emerging (9,28,29,80,117,173), there is currently a lack 68 69 of consensus regarding the match activity profiles in youth soccer (155), and furthermore, investigations into the more nuanced components of multi-directional speed (e.g., 70 deceleration, curvilinear speed, 'gamespeed', and agility) in soccer are still in their infancy in 71 72 the senior game (31,65,81,83), let alone with respect to youth soccer.

The aims of this review are to propose a definition for the term, 'multi-directional speed' (MDS), 73 74 underlining how the key components of MDS relate to performance and injury risk, through an 75 exploration of its theoretical underpinnings. Although practitioners are interested in training 76 strategies to enhance MDS (129), it is imperative to understand the mechanical and technical underpinning of MDS to effectively design physical preparation and MDS sessions for the 77 athletes. This review will firstly begin with a discussion on the performance and injury risk 78 79 considerations for the growing athlete. This should enable the subsequent sections in this review to be viewed through the lens of LTAD, in which practitioners are afforded the time to 80

implement foundational, age-appropriate and progressive training methods, as a means of
optimising athletic competency, reduce injury risk, and ensure the attainment of future
potential.

84

85 Movement Characteristics of Youth Soccer

As mentioned previously, an understanding of the movement characteristics in youth soccer 86 match-play will enable more accurate and bespoke training prescription to be delivered with 87 respect to volume and intensity (155). Age-group comparisons of youth match running 88 89 performance (29) indicate that significantly greater absolute high-intensity activity (e.g., > 16.1 km h⁻¹; U13 = 509 ± 156 vs. U18 = 1239 ± 337 m), 'sprinting' distances (e.g., > 19.1 m h⁻¹; 90 U13 = 186 ± 92 m vs. U18 = 666 ± 256 m), and peak game speeds (e.g., U13 = 22.3 ± 1.4 91 km·h⁻¹ vs. U18 = 28.3 ± 2.2 km·h⁻¹) are attained as they progress through the age-groups. 92 93 The aforementioned trends have been reported elsewhere (9,80,117,173), yet to the authors' best knowledge, there currently exists no insights into the mechanical demands (e.g., 94 acceleration and deceleration) of youth soccer match-play which spans and directly compares 95 multiple age groups (e.g., the progression from U12 to U18). Thus, a key area regarding high-96 97 intensity movement in relation to youth soccer currently remains unknown.

It is important to note that playing position plays a pivotal role in player outputs (29), with which 98 99 the progression of players through the age groups may see their playing roles specialise. 100 Furthermore, recent studies have suggested that biological maturation itself can positively impact match running performance for individuals performing within the same age-group 101 102 (28,71,76); however, utilising this information to guide training prescription remains difficult. As with the complexity and diversity of youth soccer environments around the world, factors 103 such as game rules (i.e., age-grouping criteria, substitution policy), match format and 104 105 configuration (i.e., number of players involved, pitch dimensions, match durations and number 106 of intervals), maturity status, playing position, and assignment of different pre-determined 107 movement thresholds means that studies of age-matched players are almost completely 108 unstandardized (155). Resultantly, more detailed investigations into the match-play activity 109 profiles of youth soccer players that are both specific to the game format, playing standard, 110 and maturity status of individuals spanning across an academy are warranted. It is 111 recommended, where possible, that data specific to the individuals in question be collected to 112 inform practitioners in this regard. This will enable more individualised training prescriptions based on population-specific criteria that can be progressed as individuals mature through 113 114 age categories.

115

116 Injury Characteristics of Youth Soccer

Unique to the youth athlete population are the constructs of growth and maturation and their 117 118 potential as risk factors for injury. Briefly, growth rate describes changes in an individual's 119 physical dimension over a given time, whereas maturation refers to the process towards a mature state (120). Due to the large individual variation in the timing and tempo in the process 120 of maturation, the physical and neuro-developmental differences between individuals, such as 121 immature structures and underdeveloped neuromuscular control, and disparities in maturation 122 123 status between age-matched individuals have been identified as mechanisms that may affect the risk of injury (132,160). Insights from research have also suggested that episodes of rapid 124 growth and the period around peak-height velocity (PHV) are associated with increased injury 125 risk in elite sports (79,96,103,127,194). 126

¹²⁷ 'Overuse' injuries (i.e., Osgood-Schlatter's disease (2)) are prevalent in youth soccer ¹²⁸ populations (159) and may be explained by a reduced structural tolerance of growth plates ¹²⁹ and developing bone during rapid growth (200,201). Furthermore, the high exposures to ¹³⁰ repetitive patterning of soccer activities (i.e., kicking) typically seen in highly-specialised ¹³¹ academy soccer players can lead to other morphological maladaptations (i.e., Cam-type ¹³² deformities and femoral acetabular impingement) (3,164,188). The sensitivity to these issues 133 can be exacerbated during the stage of skeletal maturation (i.e., typically in boys between 12 134 and 14 years) when the proximal femoral growth plates are open (3,4,188). Although largely 135 speculative, the concomitant increases in circulating growth hormone and insulin-like growth 136 factor-1 during rapid periods of growth may heighten the bone's osteogenic responsiveness 137 to joint loading (64), which, theoretically, may predispose the young athlete to an increased 138 risk of developing the above-mentioned maladaptations if movement variability is limited. 139 Thus, it is of the author's belief that promoting movement variability throughout training will 140 facilitate a more evenly distributed stress to greater range of anatomical structures by altering 141 the point of force attenuation or production (164). Certainly, this potential link warrants further investigation and may have important implications for the programming of youth activity. 142

Furthermore, the lags in musculoskeletal growth apparent in the growing athlete can lead to 143 144 compromised neuromuscular control during dynamic activities (e.g. running, cutting and 145 landing) and is suggested to be a key mechanism for lower-limb ligament injuries (85,86). As 146 such, a high proportion (20%) of injuries in male youth soccer players are acute traumatic 147 ligament sprains at the ankle and knee (70,159,169), of which the medial collateral ligament 148 and anterior talofibular ligament appear to be the most commonly reported injuries (43,159). 149 The presence of neuromuscular deficits do not necessarily indicate an explicit causative factor for these ligamentous injuries, yet it should be noted that these injuries may occur when active 150 muscular protective mechanisms are unable to adequately attenuate joint torgues during 151 152 dynamic manoeuvers involving deceleration and high forces (158). A possible explanation 153 may be due to delay in growth between muscle length and cross-sectional area may affect the sensorimotor function processes responsible for maintaining optimal alignment during rapid 154 adjustments of the system's centre of mass (COM), which typically occur when performing 155 156 cutting and landing maneuvers (85,215). This is important to consider, as such maneuvers 157 have been identified as key actions that amplify multi-planar knee joint loading (i.e., knee flexion, rotational, and abduction loading) while the foot is planted (51,52,101,128). Multi-158 159 planar knee joint loads are associated with increased strain on the anterior cruciate ligament (ACL) (177,189), while chronic exposure to abhorrent movement patterns may lead to the
 development of patellofemoral pain, a common injury observed in youth soccer (140,151).

Moreover, due to the muscle loading patterns inherent during repetitive training and 162 163 competition in soccer, quadriceps dominance relative to the hamstrings can present as an 164 injury risk factor (90). This can subsequently alter the reciprocal balance of activation between the hamstrings and the quadriceps, reducing the strength and dynamic stabilisation around 165 the knee joint (90). This functional hamstrings:guadriceps ratio (H:Q) has been shown to 166 167 improve with maturity, as youth athletes experience positive shifts in muscle architectural parameters (166), as well as develop co-activation of the hamstrings, which function to reduce 168 anterior tibial translation and high shear forces generated by the quadriceps as a product of 169 high-speed actions (48,90). Much like their senior peers, the requirement for the youth soccer 170 player to be regularly exposed to high-velocity movements, as well as knee flexor 171 strengthening programmes, to reduce the injury risk associated with the hamstrings muscle 172 group is important (123,163). 173

174

175 The Components of Multi-directional Speed

Currently, there is no universally agreed definition for MDS; however, it can be seen as a term 176 which encompasses all the key speed-related qualities that have been shown to be vital for 177 sporting performance in team sports. That is, linear and COD speed; incorporating 178 acceleration, lateral movements, deceleration, and back-pedalling (i.e., backwards running) 179 (116). The ability to decelerate is vital, particularly for sharper COD (57), while having the 180 181 competency and capacity to perform a variety of different maneuvers, across a range of angles and velocities, from both limbs, would undoubtedly be advantageous for soccer performance. 182 183 The authors propose that MDS can be defined as, the competency and capacity to accelerate, 184 decelerate, change direction, and ultimately, maintain speed in multiple directions and movements, within the context of sports-specific scenarios. Fundamentally, given the open 185

nature of team sports, it should be the goal of every coach and practitioner to create fast,
efficient, and robust, '360-degree' athletes. The components of MDS should be firstly broken
down and explained with respect to its components and their subsequent implications for
performance and injury risk.

190 Specific training methods can be utilised to develop characteristic qualities along the continuum of MDS (Figure 1), which can assist in determining which methods could have a 191 greater emphasis with respect to the broad context of the youth soccer player. Research has 192 193 indicated that the key determinants of sprinting and COD performance are the physical capabilities (i.e., impulse [force × time] and power qualities) of the athlete, and the technical 194 ability (i.e., joint kinematics, postures, and placement) to apply the subsequent ground reaction 195 forces (GRFs) effectively intended direction 196 in the of travel (59,61,98– 100,128,138,150,161,181,212). As such, in order to develop the various gualities that 197 underpin MDS, athletes need to be able to produce high forces, rapidly, and transmit them to 198 the ground effectively in the intended direction of travel (59-61,107,128,136,148). 199

200 Linear Speed

During the acceleration phase of a sprint, greater increases in horizontal propulsion are 201 202 required to achieve high acceleration (16,89,102,107,108,136,138,205). This phase is characterized by a greater forward lean of the torso (49) and shank to facilitate horizontal 203 propulsive forces and a longer time for the application of force during the support phase of 204 ground contact (206,214). Conversely, the transition out of acceleration towards the maximal 205 206 velocity phase of a sprint is characterized by a shift towards a more vertically oriented GRF 207 vector (41,205), as the torso and shank (at touch-down) becomes progressively more upright 208 (142,143), maximal stride frequency (SF) is attained, and a marked increase in stride length 209 (SL) is observed as running velocity continues to rise (143). This greater emphasis on 210 vertically oriented GRF during maximum velocity limits the time spent decelerating due to reducing ground contact time (GCT) (41,205). The shorter GCT associated with this higher 211

212 speed (206,214) resultantly limits the mechanical ability to apply such force, given the shorter period of time. Thus, although horizontal GRF is the primary factor governing the rate of 213 214 acceleration (89,108,136,205), it is fundamentally the impulse-momentum relationship which 215 determines the time in which to force is applied to overcome inertia and gravity. As such, the 216 ability to maintain horizontal impulse over increasingly shorter GCT is critical to sprinting performance (89,108,138). Hence, notable differences are observed between the impulse 217 profiles of acceleration (e.g., peak resultant GRF ~ 2.3 N·kg ⁻¹; GCT ~ 200-135 ms) and 218 maximal velocity (e.g., peak resultant GRF ~ 3.7 N·kg ⁻¹; GCT ~ 101-108 ms) phases of 219 220 sprinting (108,141,206,214). These observations point towards the emphasis of kinetic, 221 kinematic and spatiotemporal attributes that operate on a continuum, whereby the generation and transmission of GRF variables are dependent on the certain phase of the sprint (Figure 222 223 1).

224 Pertinently, given the intermittent nature of soccer match-play, the majority of sprint efforts occur within distances of 20 m (53,184,199), which limits the attainment of maximum speed. 225 Consequently, to adhere to the principle of specificity, the development of acceleration in the 226 227 context of soccer is a priority (13,14,185), and has led to the proposal of sub-phases within acceleration, where distinctions in biomechanical characteristics have been identified 228 (11,14,119,141,142). In professional soccer players, Bellon et al. (14) established statistically 229 different transitions in sprint characteristics as the progression within the specific distances 230 that categorise the early (0 - 2.5 m), mid (2.5 - 6 m) and late (6 - 12 m) sub-phases of 231 232 acceleration. Specifically, the early-acceleration phase may be characterized by minimal flight times (FTs) coupled with GCT at their peak, and SL at their lowest with a rapid increase in SF 233 (11,14,143,161). Further, in order to promote a 'pushing motion', limited knee flexion over 234 235 ground support, rapid hip extension velocity, high horizontal GRF behind the COM, and also a rapid rise of the hips with the COM, is reported to occur (142). During the mid-acceleration 236 phase, this antagonistic relationship shifts as GCT declines while FT increases, SL continues 237 to increase and SF typically reaches its maximum, even showing a slight decline prior to 238

239 stabilizing (11,14,143,161). A reduced rate of COM elevation, increased knee extension 240 velocity, as well as increased knee flexion during ground support are also observed, which is 241 likely a result of striking the ground further in front of the body (142). Finally, the approach 242 towards maximum velocity during the late-acceleration phase is characterized by further 243 reductions in GCT, SF plateauing, and maximal values of FT and SL being attained 244 (11,14,143,161). Additionally, the reduced range of motion observed at the hip will see 245 increasing contributions of the lower leg during this phase (142), which may be explained by 246 the greater need for the foot to aggressively strike the ground in a 'backwards' motion to apply 247 high GRFs within close proximity of the COM (89).

248 These aforementioned acceleration sub-phases have been shown to be athletic populationspecific (11,14,119,141,142); however, the evaluation of acceleration phases with a higher 249 250 degree of precision (e.g., tracking GCT, FT, SL, SF, COM trajectory and joint angles) allows 251 strength and speed training interventions to be developed that target distinct kinetic and kinematic variables more specific to soccer across the various sub-phases, which may be 252 sequenced at different times during the season (14). Importantly, during acceleration, it is the 253 254 athlete's technical ability to adopt these phase-specific kinematic postures that facilitate the application of forces in the opposite direction to their COM displacement (17,107,136); an 255 effective horizontal transmission of these forces over increasingly abbreviated GCT as velocity 256 continues to rise has been termed mechanical effectiveness (172). Although it is beyond the 257 258 scope of this review to discuss off-field training methods in relation to MDS, resistance training 259 should be seen as the cornerstone for which to build positive adaptations to the mechanical gualities (i.e., force, velocity and power) that underpin the kinetics specific to each sub-phase 260 of acceleration (i.e., early, mid and late sub-phases). These methods should maintain a high 261 262 priority in any supplementary programme that aims to improve an athlete's mechanical effectiveness, and subsequently acceleration performance (87,172). 263

It is well established that sprinting performance is underpinned by an athlete's mechanical
 force characteristics (89,102,108,136,138,205), but from a physiological perspective, the

266 action of sprinting requires high levels of motor unit activation (137,190,213). The key muscles involved in sprinting are the hamstrings and the gluteus maximus (i.e., hip extension), 267 268 quadriceps (i.e., knee extension) and the gastrocnemius and soleus (i.e., plantar flexion) muscle groups, all of which elicit high levels of activation either just prior to or on the instant 269 270 of ground contact (137). The horizontal orientation of the body combined with the low 271 displacement velocities during acceleration indicates that a large proportion of the stance 272 phase (approx. 87 – 95% of total GCT) is spent applying propulsive GRF (89). Subsequently, 273 this phase is mechanically characterized by an increased activation of the hip, knee and ankle 274 extensors, which work to concentrically contract to produce high levels of force to accelerate 275 the body horizontally (107,137). Conversely, the notable spatiotemporal (i.e., reduced GCT and increased SL), kinetic (i.e., increased vertical GRF) and kinematic (i.e., greater knee and 276 277 ankle angles) shifts in parameters as a consequence of increased sprinting speeds (141,142) 278 place a much greater requirement on the rate of force development (23,38,41,178). This is achieved through an eccentric-concentric coupling action (i.e., stretch-shortening cycle; SSC), 279 whereby the storage and release of elastic energy through the muscles and tendons around 280 the knee and the ankle (i.e., gastrocnemius-soleus-achilles complex; GSA) joints allow high 281 282 propulsive GRF to be transmitted in a limited amount of time (174). Taking these findings together, it may be argued that the characteristics of sprinting may follow a 'proximal-to-distal' 283 nature (107), with which the force-producing capabilities of these key muscle groups play 284 important roles in certain phases of the sprint. 285

Overall, it has been shown that the hip extensor and flexor muscles (i.e., gluteals and hamstrings) play a pivotal role in performance across all sprint phases (12,175) and display increasingly greater activation as running speeds increase to maximal sprinting (174). This is further supported by findings that have shown athletes with a better ability to orientate horizontal GRFs have are able to highly activate their hamstrings just before ground contact and present high hip extensor torque capabilities (137). It appears that the concentric torque and activation of the gluteal muscles appear to be more strongly related to horizontal GRF in the initial acceleration phase (137), whereas eccentric hamstring force and activation are more strongly related to top speed mechanics (137,175,187). Thus, particularly at high running speeds (187), it may appear that the high torque requirements of the hamstrings to both eccentrically reduce the kinetic energy of the lower limb during the late-swing phase (32,56), as well as producing high rates of force during the early-stance phase of ground contact (41,205), can work in combination to reduce deceleration time during ground contact and increase horizontal propulsion (138,187).

300 Notably, it is suggested that the high levels of activation and subsequent torque generated by the hamstring muscles (i.e., as high as eight times bodyweight) can predispose an athlete to 301 an increased risk of hamstring muscle strain injury (HSI) when performing sprint actions 302 (104,187,207,213). Debate exists in relation to whether the mechanistic basis for HSI are 303 304 explained for in the end-swing phase or early-stance phase (104); however, it has been shown 305 that hip extensor concentric (186) and knee flexor eccentric strength (153) are both considered 306 risk factors for HSI. Moreover, the activation levels (45) and force generating capacity (135) 307 of these muscle groups are compromised in individuals who have previously sustained a HSI. 308 Additionally, architectural characteristics within the hamstrings muscle group, namely, shorter biceps femoris fascicles (BFF), have been identified as a risk factor that may predispose an 309 individual to sustaining a HSI (191). Off-field HSI risk mitigation strategies which focus on 310 conditioning the gluteal and hamstring muscle groups (135) as a means of eliciting specific 311 312 architectural adaptations (e.g., fascicle length, pennation angle and cross-sectional area) 313 (6,157), as well as positive morphological (e.g., muscle-tendon interactions) and neurological (i.e., motor unit activation, muscular co-ordination and H:Q) adaptations (18,68), are becoming 314 315 more commonly practised in soccer (130,131). However, none of these methods can replicate 316 the sprint-specific muscular activation patterns experienced during sprinting (190). A comprehensive sprint training programme in complement to regular soccer practise has been 317 shown to induce greater increases in BFF length, alongside improvements in sprint 318 performance and mechanical outcomes, in comparison to isolated knee flexor eccentric 319

320 strengthening (i.e., Nordic hamstring exercise) and sports-specific practise alone (134). Athletes who regularly sprint at near maximal intensities (i.e., > 95% maximum velocity) during 321 322 training have displayed a reduced risk of lower-limb injuries in comparison to their teammates who produced lower running intensities (i.e., < 85% maximum velocity) (123). Further to this 323 324 point, athletes who accumulate larger chronic training exposures to sprinting may also tolerate higher volumes of sprinting (122,123). These findings point towards an approach that features 325 326 frequent exposure to sprinting, and therefore MDS training, as a central component to both 327 improving performance and reducing injury risk in soccer players (122,123,131,134).

- 328
- 329

Insert Figure 1 around here

330

331 Change of Direction Speed

Change of direction is defined as a "reorientation and change in the path of travel of the whole-332 body COM towards a new intended direction" (47,209). COD ability provides the technical, 333 mechanical, and physical basis for effective agility (60,148,212). In soccer, CODs are linked 334 to both attacking and defensive scenarios, such as creating space to receive a pass, evading 335 and dribbling past an opponent, pressing opponents and making key interceptions (1). Soccer 336 players perform a large volume of COD actions (~600 cuts of 0-90°; ~100 turns of 90-180°) 337 during match-play (19), to both left and right directions (19,167), with CODs which are then 338 followed by sprints linked to decisive actions such as goal scoring and assists (63). Recent 339 insights into the movement demands of a professional Russian soccer team revealed that ~ 340 600 CODs were performed during match-play, of which ~60 and ~15 were performed at high 341 (> 16 km.h⁻¹) and maximum intensity (> 21 km.h⁻¹), respectively (74). Interestingly, the authors 342 343 revealed that COD frequencies were typically greater during away matches and when during 344 matches performed on artificial field-turf. As such, COD ability is considered a highly important quality to develop in soccer players (193), yet to optimise COD ability, it is important tounderstand the key biomechanical and physical determinants.

347 The determinants of COD ability are multi-faceted and underpinned by the interaction between speed, deceleration, mechanics, and physical capacity (Figure 2) (60,61,77,97–99,128,181). 348 The COD foot plant instigates a break in cyclical running, and COD can subsequently be 349 divided into four key phases: 1) initial acceleration; 2) preliminary deceleration; 3) COD foot 350 plant; and 4) re-acceleration. High levels of eccentric, isometric, and concentric strength are 351 352 required for the braking, plant, and propulsive phases, respectively (124,182,183,204). Undoubtedly, COD is a multi-planar and multi-step action (8.60.62), with the steps preceding 353 and following the main push-off involved in re-direction. For example, the penultimate foot 354 contact (PFC) plays a key role in deceleration and creating posterior braking impulse to reduce 355 356 momentum prior to changing direction (57,61,62,98,128). The PFC is particularly important for sharper CODs, and is considered a key 'braking step' for facilitating faster COD performance 357 and alleviating potentially high-risk knee joint loads (57,59,61,62,98). Conversely, particularly 358 for shallower CODs, the PFC is considered a key 'positional' or 'preparatory step' for 359 facilitating effective postures during the main COD foot plant to optimise braking and push-off 360 (57,59,61,62,98). Specifically, the COD foot plant requires a manipulation of the base of 361 support relative to COM (e.g., lateral foot plant) to create an external braking impulse to reduce 362 momentum, and a propulsive force and impulse to accelerate into the intended direction of 363 364 travel, within a short GCT (42,60,128). These factors are all significant associates of faster 365 COD performance (59–61,128,181,182,204). The COD foot plant is described as an eccentricconcentric coupling action (i.e., SSC) and consists of two-phases: weight acceptance (i.e., 366 braking) and push-off (i.e., propulsion) (8,60). During these phases, the body typically goes 367 368 from lower-limb triple flexion into a rapid, forceful triple extension (i.e., simultaneous joint movements) (8,60), with successful COD performance influenced by key postures (i.e. trunk 369 positioning, pelvis rotation, hip, knee, ankle range of motion, foot plant distance, etc.) to 370 facilitate effective re-direction and to orientate the GRF towards the intended direction of travel 371

372 for effective net acceleration (60,61,126,128,204). In addition to high magnitudes of braking and propulsive force (impulse) and key technical postures, high levels of muscular activation 373 around the hip, knee, ankle and trunk (i.e., pre- and co-activation) are required to support the 374 375 large multi-planar external moments created when changing direction and to facilitate effective 376 braking and propulsion, particularly around the knee joint (15,109,124,125,146). 377 Fundamentally, an athlete's physical capacity (i.e., neuromuscular control, rapid force 378 production, strength, muscle activation) is considered highly important during the COD foot 379 plant for the following reasons: 1) applying high and rapid levels of force in short GCT; 2) to 380 support and tolerate the large multi-planar joint loads (particularly from faster approaches velocities (57), sharper CODs angles (57), externally directed attention and agility movements 381 (5,27), limited preparation time (69,139), and fatigue (202)); and 3) permit the athlete to adopt 382 383 key postures associated with faster performance (57,58,62,98,99,124,148,183).

384

385

Insert Figure 2 around here

386

Because of the unpredictable nature of soccer, players require the capacity to change direction 387 rapidly and effectively across a spectrum of angles, from both their left and right limbs; 388 possessing the ability to perform CODs from low, moderate, and high approach velocities 389 (19,81,167). In addition, soccer players require 'movement solutions', underpinned by physical 390 literacy, in order to perform a variety of different COD actions in the context of sport-specific 391 scenarios (60). For example, 'crossover cuts' are critical when a player aims to maintain 392 velocity during a COD (i.e., < 45°), such as when a striker performs a curved or arced sprint 393 to evade the offside trap and defensive line. Conversely, a defender may perform a 'side-step' 394 395 cutting action when pressing and reacting to an opposition player. 'V-cuts' are also a common 396 feature in attacking scenarios, whereby midfield players typically accelerate towards the ball, 397 and then pivot or spin away in order to deceive and create separation from an opposing

defender in response to a pass. Finally, 'split-steps' are also performed by attacking players
during dead-ball situations (i.e., corners, free-kicks, throw-in), whereby man-to-man marking
takes place, and the athlete may perform a deceiving maneuver to evade and create
separation from an opponent.

402 It is important to note that biomechanical demands of COD are 'angle-dependent' (57). As such, the acceleration, deceleration, COD foot plant mechanics, and re-acceleration 403 requirements of COD are governed by the approach velocity, intended COD angle, sporting 404 405 scenario (i.e., pre-planned, offensive or defensive agility), and the athletes' physical capacity (57,62). An 'angle-velocity trade-off' concept has been discussed with respect to changing 406 direction, whereby, as the intended COD angle increases, in addition to increased approach 407 velocities, a concurrent reduction in velocity, greater deceleration and braking is required to 408 change inertia and accelerate into the new intended direction of travel (20,57). During shallow 409 CODs (i.e., < 45°), the deceleration and braking requirements are limited, with GCTs (~150-410 200 ms) being relatively shorter for COD (57). The ability to approach fast, attain high minimum 411 412 speeds, and maintain velocity is critical for faster COD performance during shallow cuts 413 (61,78,98,99). This has led to strength and conditioning physical preparation 414 recommendations to focus on the 'velocity' end of the force-velocity curve (FVC), and the utility of ballistic and fast SSC exercises (20,57). Conversely, with sharper CODs (i.e., \geq 60°), 415 although approach velocity is still vital (61,98,99,128), the ability to decelerate by braking hard, 416 417 late, and rapidly is fundamental to faster COD performance (61,62,98,99), with sharper CODs 418 from greater approach velocities typically requiring greater braking distances and longer GCTs (~300-500 ms) during the COD foot plant to facilitate the directional change (57,62,77,78). As 419 such, this has led to the physical preparation recommendations which target the 'force' aspect 420 421 of the FVC through resistance training and utilising slow SSC plyometric exercises (20,57). Resultantly, angle and velocity are two crucial factors which regulate intensity during COD, 422 and should be progressed accordingly when coaching and designing MDS training 423 programmes (Figure 1). 424

425 An athlete's ability to decelerate should be considered as an essential component to COD performance (Figure 2). A recent meta-analysis (81) provided further insights when 426 highlighting that high-intensity (>2.5 m \cdot s⁻²) and very high-intensity (>3.5 m \cdot s⁻²) decelerations 427 were performed more frequently than equivalently intense accelerations, across every team 428 429 sport during competitive match-play, with the exception of American Football. Importantly, soccer was considered the team sport where this difference was the largest (standardized 430 mean difference = -1.74). For example, high-intensity decelerations (>2 m·s⁻²) have been 431 432 shown to occur up to 2.9 times more frequently than accelerations within the same speed 433 intensities in soccer (88). Interestingly, a 'self-regulatory' concept has been discussed with 434 respect to COD (57,98,99), whereby athletes sprint at a velocity based on the deceleration load that they can tolerate. This deceleration ability is fundamentally underpinned by eccentric 435 strength; for example, eccentrically (knee flexor and extensor) stronger female soccer players 436 437 have been shown to approach faster and display greater reductions in velocity and braking forces during 180° CODs (98), while eccentrically stronger (knee flexor and extensor) soccer 438 players were able to maintain velocity, attain higher minimum speeds, and tolerate greater 439 loads during a 90° cutting task (99). Taken together, these findings highlight the interaction 440 441 between an athletes physical capacity and their movement repertoire, which this has led to the strength and conditioning recommendations of developing eccentric strength capacity 442 programming for deceleration performance 443 alongside field-based COD and (36,62,81,83,97,100,128). 444

Although accelerations and decelerations both expose individuals to high levels of physiological and biomechanical stress (92,196), these stressors can be seen as fundamentally different (30,196). Accelerations may have a higher metabolic cost (77) in comparison to decelerations which can elicit higher mechanical demands (44); this heightened mechanical load is likely explained by the high force impact peaks and loading rates which occur as a consequence of often suddenly imposed deceleration actions (197). Specifically, the high magnitude eccentric braking force requirements of deceleration can impart damage 452 on soft-tissue structures (i.e., tendons) through eccentric muscle fibre contractions that can disrupt the integrity of the muscle fibres (75), particularly if there is a reduced capacity to 453 454 effectively attenuate these high force demands (83). In contrast, acceleration has high concentric strength requirements (23,115,147) of which are vital for the production of 455 456 horizontal GRFs (16,89,102,107,138,205). This may bring with it an increased vulnerability of the muscle-tendon tissue properties in handling eccentric braking demands (72,156) that will 457 458 come as a direct consequence of increased movement speeds (57,145). Therefore, 459 interventions should be implemented that "mechanically protect" individuals from the 460 damaging nature of high-intensity decelerations (62,83). Indeed, improving an athlete's acceleration and COD mechanics are important for successful MDS; however, it is 461 emphasised that coaches, 'do not speed up, what an athlete cannot slow down' (81-83). 462 463 Practitioners should ensure that they develop their athletes' braking mechanics, in conjunction 464 with their eccentric strength capacity, for effective deceleration, and subsequently COD, from both performance and injury risk mitigation perspectives (61,62,82,98,99,128). 465

466 As mentioned previously, the compromised neuromuscular control in youth athletes during 467 high-risk maneuvers that involve rapid decelerations and high eccentric forces place this population at a heightened risk of sustaining acute traumatic lower-limb ligament injuries about 468 the ankle and the knee (70,85,86,159,169). As such, correcting hazardous postures (e.g., 469 470 knee valgus) are often encouraged in injury mitigation programmes (154). It has been shown 471 that high gluteal activation is required to oppose knee valgus and rotator moments (125), and 472 from a performance perspective, facilitate effective propulsion (124). Furthermore, in addition to playing a key role during linear sprinting (137), the hamstrings also have an important role 473 in preventing anterior translation of the tibia, reducing anterior tibial shear, and attenuating 474 475 impact GRF during deceleration and COD activities (124,125). Resultantly, training recommendations that focus on improving external hip rotator (gluteal) strengthening and 476 activation (105,106), as well as increasing hamstring strength (203), are recommended. 477 Improving gluteal strength and activation, theoretically, should limit internal hip rotation and 478

479 adduction, and as such, reduce the frontal plane moment arm and knee abduction moments 480 (KAM) during COD actions (86,133). Furthermore, improving hamstring activation and 481 strength will enable greater knee flexion to reduce impact GRF (55), and permit load 482 absorption via the muscular structure, in contrast to passive structures, which can in turn 483 facilitate a reduction in ACL loading (203).

Ultimately, as with many aspects of human movement there will be movement variability. Athletes may display slight differences in movement strategy while still achieving similar outcomes. That said, there will still be key movement principles and subsequent postures and positions which optimise force generation and transmission. The extent of the techniques will differ cross individuals, and will influenced by a myriad of factors, such as anthropometry, physical capacity, training age, contextual and situational variables.

490

491 Curvilinear Speed

492 Although linear speed is indeed important for soccer players (63), in the context of soccer performance, a large proportion of sprints do not occur in a straight line, but in fact are curved 493 or arced with a deviation and curvature of the path of travel (1,24,31,67). Curved sprinting 494 (synonymous with curvilinear speed, curved speed, and arced running), has been recently 495 defined as "the upright running portion of the sprint completed with the presence of some 496 497 degree of curvature" (31), described as a hybrid between linear speed and COD (66). However, a COD requires a clear break from cyclical running and with a clear lateral foot plant 498 499 involved in re-direction, in contrast to curved sprinting which maintains a cyclical running 500 pattern (60). Thus, curved sprinting, theoretically, will most likely share greater biomechanical similarities to linear sprinting opposed to COD. Nevertheless, curved sprinting actions are 501 502 typically performed by soccer players in offensive situations, such as when performing 503 overlapping or channel runs along a defensive line to maintain an onside position, or defensive situations, such as performing recovery and covering runs when marking and channelling 504

505 opposition players or making interceptions (1,19,25,26,31). Consequently, the objective of 506 curved sprinting is to deviate from the path of travel in a curvilinear motion while attaining or 507 maintaining high velocities.

508 Traditionally, emphasis in soccer research has been placed on quantifying and classifying 509 linear speed performance in soccer players (1,21), yet studies examining curved sprinting are scarce. Seminal time-motion analysis from Bloomfield et al. (19) found soccer players perform 510 511 ~10-20 swerves during match-play, but provided limited information as to how these actions 512 were classified. Pilot analysis from Brice et al. (24) reported curved motions of travel in English 513 FA Premier League soccer players ranged from 3.5 -11 metres. More recently, Caldbeck (31) reported ~85% of sprint actions in elite soccer were in fact curvilinear, while in elite youth 514 soccer players matches, Fitzpatrick et al. (67) observed an average sprinting (> 24 km h⁻¹) 515 516 angle and frequency of ~5 degrees and 20, respectively, that were typically performed over distances of 10-20 m. However, it is worth noting that, although not as frequent as the acute 517 sprint angles, all players performed sprint angles of up to 30° (67). Ade et al. (1) reported 518 soccer players perform ~10-30 arced runs pre-, mid-, and post- high intensity effort during 519 520 match-play. From a positional perspective, forwards have been shown to perform a greater number of swerves (19), sharper angled sprints, and arced runs in and out of possession (1), 521 compared to other playing positions (67). This potentially highlights the importance of extreme 522 curved sprinting ability in attacking soccer players and implications for position-specific 523 524 training. Nevertheless, because of the open skilled nature of soccer, irrespective of playing 525 position, it would be advantageous for soccer players to be able to attain and maintain high velocities during curvilinear motion, across a spectrum of varying degrees and radii. 526

527 Despite the importance and frequency of curved sprinting in soccer (19,24–26,31,67), it is 528 somewhat surprising that curved sprinting assessments do not commonly feature in the testing 529 batteries (192). To the best of our knowledge, Fílter et al. (66) are the only researchers that 530 have investigated the isolated ability to curvilinear sprint in soccer players in the field, using a 531 17-m curved sprint with the simple use of the penalty arc (9.15 m radius). The authors 532 observed highly reliable measures of curved sprint speed (ICC \ge 0.89, CV% \le 1.15) in semiprofessional soccer players. However, while the authors observed very strong associations 533 between left and right curved sprinting performance ($r^2 = 0.77$), lower associations were 534 observed between curved and linear sprint speed ($r^2 = 0.34-0.37$). These findings suggest that 535 536 athletes who display superior linear speed performance may not necessarily display fast performance during curvilinear tasks, and vice versa. It appears, therefore, that linear and 537 538 curvilinear sprinting are independent athletic qualities which should be assessed and trained 539 as such. It is currently unknown which training methods are most effective at developing 540 curved sprinting, and it appears that no study has examined the effects of linear speed training on curve sprinting ability in soccer players, and thus, it is a recommended future direction of 541 research. 542

The finding that linear and curved sprinting appear to be independent athletic qualities could 543 be attributed to key kinetic, kinematic, muscle activation, and spatiotemporal differences 544 between the two tasks (7,25,26,37,39,40,65,91,179,192). In contrast to the more upright and 545 sagittal plane approach adopted during linear sprinting, curvilinear sprinting requires the 546 generation of centripetal and medial-lateral (ML) GRF through a medial whole-body lean to 547 548 help counteract a rotating moment and permit the continuation of a curved path of travel (25,26,91). For example, different hip and knee kinematics, foot placement and shank angle, 549 and greater lower-limb ROM (26), have all been observed during curved sprinting compared 550 551 to linear sprinting (7,25,26,37,91). Specifically, the inside leg has been described as a frontal 552 plane stabiliser (7,37), with the lower-limb displaying greater adductor and semitendinosus muscle activation (65), alongside greater lower-limb ROM within longer GCT, compared to the 553 outside leg and normal linear sprinting (26,65). The inside leg also displays greater hip 554 555 adduction, hip flexion, and hip external rotation, alongside greater ankle eversion (i.e., ankleeversion strategy), in comparison to the outside leg and linear sprinting (7,37,39). This posture 556 commutatively lowers the COM, facilitating an inclination of the inside leg and medial body 557 558 lean towards the curvature to assist in centripetal acceleration (7,25,26,37,39,91,179).

559 Conversely, the outside leg has been suggested to play a more important role in terms of propulsion and rotation (7,37,40,179), with greater hip and knee internal rotation, combined 560 561 with greater gluteus medius and biceps femoris muscle activity in comparison to the inside limb (26,37,40,65,91,180). Typically, this will result in an asymmetrical GRF profile between 562 563 limbs, as greater ML GRFs over shorter GCT are observed in the outside limb (26,37,40,65,91,180). Interestingly, it is has been suggested that the GSA complex is pivotal 564 to managing the centripetal forces during curvilinear sprinting and may be the performance-565 566 limiting factor through the generation of plantarflexion moments (118). With curvilinear sprinting drawing more similarities between linear sprinting, opposed to COD, perhaps 567 strengthening the mechanical characteristics of the ankle complex may translate to 568 performance improvements in both qualities of MDS; however, future investigations are 569 570 needed to further examine this speculation.

571

572

Insert Table 1 near here

573

From a kinematic and spatial-temporal perspective, lower velocities have been observed with 574 curvilinear sprinting (e.g., 1 – 15 m radii) compared to linear sprinting (25,26,37,65,91,180). 575 Shorter stride cycle times, reduced SL, and slightly greater SF are required to maintain 576 balance and posture during travel, with these differences amplified with increased sprinting 577 curvature (25,26,37,65,91,180). Additionally, subtle differences in step length and frequency 578 have been observed between inside and outside limbs during curved sprinting (e.g., 1 - 6 m 579 580 radii), with the inside limb typically displaying greater SF and shorter SL compared to the outside limb (37). Reduced FTs have been reported during curved sprints (5 m radius), which 581 582 contributes to the shorter stride cycle times (179). Thus, in combination with the increased 583 step and stride frequencies associated with curved sprinting, this increases the time in contact 584 with the ground for force application and centripetal acceleration (37,179). Additionally, the

585 reduced FTs ensure that the athlete does not further deviate from the curvilinear path of travel 586 because greater airtime results in greater travel of the COM along a path tangential to the 587 curve, which is sub-optimal for curved sprinting (179). It should be noted that the kinetic, 588 kinematic, and spatiotemporal differences between limbs, and compared to linear sprinting, 589 are largely influenced and dependent on the curvature (radius) of the sprint (25,26,37,91,180). 590 Nonetheless, because of the importance of curved sprinting actions in soccer (19,24-591 26,31,67), and the distinct biomechanical differences between curved and linear sprinting 592 (7,25,26,37,39,40,65,91,179,192), it is essential that curved sprinting is assessed and trained 593 as an independent athletic quality, and practitioners are encouraged to design MDS 594 programmes in accordance with the movement characteristics that are specific to the sport.

595

596 Agility & Contextual Speed

597 Agility can be broadly defined as a "rapid whole-body movement with change of velocity or direction in response to a stimulus" (176). An agility action is, therefore, predicated on a 598 perceptual and decision-making process in response to a stimulus, of which the subsequent 599 outcome is a positive or negative change in acceleration, or a COD manoeuvre (176). These 600 601 decision-making processes require perceptual-cognitive factors (i.e., visual scanning, knowledge of situations, pattern recognition, anticipation) to be coupled with a motor response 602 which will produce a whole-body movement (176). As such, although the aforementioned pre-603 planned qualities (i.e., acceleration, deceleration, COD and curvilinear speed) discussed in 604 605 this text are considered essential elements to develop within a soccer strength and conditioning programme, it is the interaction of perceptive, cognitive and motor control 606 607 capabilities with the ability to apply effective movement that wholly contributes to MDS 608 performance (93-95,210).

609

Insert Figure 3 around here

610

612 Specifically, from a soccer perspective, agility may explain the ability to move effectively in the context of a soccer match. The performance of soccer-specific skills require the athlete to 613 614 move towards these specific actions through a perception-action coupling in relation to the 615 constantly changing scenarios on the pitch (95). Therefore, athletes need to be able to recognise and exploit game scenarios in order to utilise efficient movement skills to apply their 616 velocity capacities. This has brought about the development of the 'gamespeed' concept, 617 618 which refers to the ability to exploit the qualities of speed and agility within the context of a soccer match (95). Such an approach requires a blend of skill-related technical and tactical 619 elements of the game to improve sport-specific movement (94). This concept, therefore, 620 should not only be specific to the team sport in question, but should also be specific to the 621 622 soccer team itself, as a means of developing key movements in relation to underpinning 623 technical and tactical principles that guide the club's coaching philosophy. Thus, establishing a club's game model (50) is recommended in order to effectively scaffold a club's coaching 624 philosophy upon the development framework of a MDS programme (Figure 3). For 625 626 practitioners embedded within clubs, this may be achieved through communication with the 627 technical coaching staff to harmonise the themes of soccer skill coaching with the aims of physical training blocks. For a more detailed outline of how such an approach is developed 628 and applied within a soccer academy context, readers are referred to an excellent introductory 629 630 article by Jeffreys et al. (95).

With that said, however, the development of agility is a controversial area, especially in the context of youth athletes. Predominantly, research has investigated the maturational and training-induced changes in strength, SSC function, sprinting and jumping ability (111,113,152,162,170,171). Research regarding growth, maturation and COD performance in youth athletes is lacking (58), and even less is known with regards to agility, which is a task of much greater complexity (114). The underlying issue being it is difficult to evaluate 'agility' performance, and subsequently monitor the effectiveness of MDS training programmes, because it is very difficult to administer a standardised, reliable, sport-specific agility test (149,211). Therefore, working in cohesion with the technical coaching staff may be an effective means of developing sports-specific agility (i.e., gamespeed or contextual speed) that may have a positive effect and translate to performance on the pitch from physical, technical and tactical perspectives.

With that said, the authors advise caution regarding an approach towards early sports 643 specialisation in youth athletes, in particular within the context of agility development, which 644 645 should, arguably, be emphasised during the transition to senior training (i.e., late-adolescence and early adulthood), where there can a be greater emphasis on sports-specificity and tactical 646 training. Certainly, within the early years of development (i.e., < 13 years of age), an emphasis 647 should be placed on developing general agility skills through a diversified, multi-lateral 648 649 approach (112,195,208). Diversifying the youth athlete's exposures to games and scenarios which emphasise different movements, tactical dimensions, and patterns of play in attacking 650 651 and defensive situations, can be achieved through agility games (e.g., tag, evasion, partner follow, agility races) and the sampling of a variety of different sports (e.g., rugby, basketball, 652 653 hockey, and American football) (129). Given the high-intensity nature that is inherent with MDS 654 training, this approach will increase the young athlete's movement variability and subsequently provide a more balanced distribution of stress to greater range of anatomical structures by 655 altering the point of force attenuation or production, thus potentially reducing the risk of 656 657 overuse injury (163,164). This training methodology may be used as a means of promoting 658 athletic skills development, fostering creativity and intuition and reducing the risk of injuries, which will provide the youth soccer player with an expansive foundation of movement skills for 659 when the time comes to transition to the senior game. 660

661 Conclusion

662 In this review article, the authors have defined MDS and explored the scientific underpinnings 663 of each of its components in relation to performance and injury risk in soccer. In order to 664 optimise the development of MDS, it is critical to understand its theoretical underpinnings to appreciate how each MDS quality has its own unique implications for performance and injury 665 risk. Ultimately, effective MDS performance in the context of soccer is underpinned by a 666 perceptual-action (perceptual-cognitive component) response to dynamic, ever-changing 667 668 scenarios within a game. It cannot be dismissed, however, that enhancing an athlete's physical and mechanical ability to accelerate, decelerate, COD, and attain top-end speeds 669 both linearly and curvilinearly, will positively enhance 'agility' and 'game-speed' soccer 670 performance. Fundamentally, the underpinning philosophy regarding the preparation of soccer 671 players should be to generate robust and effective 360° athletes, who have the competency 672 to accelerate, decelerate, and change direction rapidly and effectively from both limbs. 673 Harmonising sporting skill with speed development is, therefore, pivotal to sporting 674 performance. Thus, although each quality can be isolated and trained as such, an optimal 675 676 programme should always seek to combine and develop the two in concert.

677

References

- 1. Ade, J, Fitzpatrick, J, and Bradley, PS. High-intensity efforts in elite soccer matches and associated movement patterns, technical skills and tactical actions: Information for position-specific training drills. *J Sports Sci* 34: 2205–2214, 2016.
- 2. Adirim, TA and Cheng, TL. Overview of injuries in the young athlete. *Sport Med* 33: 75–81, 2003.
- 3. Agricola, R, Bessems, JHJM, Ginai, AZ, Heijboer, MP, Van Der Heijden, RA, Verhaar, JAN, et al. The development of cam-type deformity in adolescent and young male soccer players. *Am J Sports Med* 40: 1099–1106, 2012.
- 4. Agricola, R, Waarsing, JH, Thomas, GE, Carr, AJ, Reijman, M, Bierma-Zeinstra, SMA, et al. Cam impingement: Defining the presence of a cam deformity by the alpha angle. Data from the CHECK cohort and Chingford cohort. *Osteoarthr Cartil* 22: 218–225, 2014.
- 5. Almonroeder, TG, Garcia, E, and Kurt, M. The effects of anticipation on the mechanics of the knee during single-leg cutting tasks: A systematic review. *Int J Sports Phys Ther* 10: 918, 2015.
- 6. Alonso-Fernandez, D, Docampo-Blanco, P, and Martinez-Fernandez, J. Changes in muscle architecture of biceps femoris induced by eccentric strength training with nordic hamstring exercise. *Scand J Med Sci Sport*, 2018.
- 7. Alt, T, Heinrich, K, Funken, J, and Potthast, W. Lower extremity kinematics of athletics curve sprinting. *J Sports Sci* 33: 552–560, 2015.
- 8. Andrews, JR, McLeod, WD, Ward, T, and Howard, K. The cutting mechanism. *Am J Sports Med* 5: 111–121, 1977.
- 9. Atan, SA, Foskett, A, and Ali, A. Motion analysis of match play in New Zealand U13 to U15 age-group soccer players. *J Strength Cond Res* 30: 2416–2423, 2016.
- 10. Barnes, C, Archer, DT, Hogg, B, Bush, M, and Bradley, PS. The evolution of physical and technical performance parameters in the english premier league. *Int J Sports Med* 35: 1–6, 2014.
- 11. Barr, MJ, Sheppard, JM, and Newton, RU. Sprinting kinematics of elite rugby players. *J Aust Strength Cond* 21: 14–20, 2013.
- 12. Bartlett, JL, Sumner, B, Ellis, RG, and Kram, R. Activity and functions of the human gluteal muscles in walking, running, sprinting, and climbing. *Am J Phys Anthropol* 153: 124–131, 2014.
- 13. Bellon, CR. The relationship between strength, power, and sprint acceleration in division 1 men's soccer players. East Tennesse State University, 2016.
- 14. Bellon, CR, Deweese, BH, Sato, K, Clark, KP, and Stone, MH. Defining the early, mid, and late subsections of sprint acceleration in division 1 men's soccer players. *J Strength Cond Res* 33: 1001–1006, 2019.
- 15. Besier, TF, Lloyd, DG, and Ackland, TR. Muscle activation strategies at the knee during running and cutting maneuvers. *Med Sci Sport Exerc* 35: 63–70, 2003.

- 16. Bezodis, NE, North, JS, and Razavet, JL. Alterations to the orientation of the ground reaction force vector affect sprint acceleration performance in team sports athletes. *J Sports Sci* 35: 1817–1824, 2017.
- 17. Bezodis, NE, Trewartha, G, and Salo, AIT. Understanding the effect of touchdown distance and ankle joint kinematics on sprint acceleration performance through computer simulation. *Sport Biomech* 14: 232–245, 2015.
- 18. Blazevich, AJ, Gill, ND, Bronks, R, and Newton, RU. Training-specific muscle architecture adaptation after 5-wk training in athletes. *Med Sci Sports Exerc* 35: 2013–2022, 2003.
- 19. Bloomfield, J, Polman, R, and O'Donoghue, P. Physical demands of different positions in FA Premier League soccer. *J Sport Sci Med* 6: 63, 2007.
- 20. Bourgeois, FA, McGuigan, MR, Gill, ND, and Gamble, P. Physical characteristics and performance in change of direction tasks: A brief review and training considerations. *J Aust Strength Cond* 25: 104–117, 2017.
- 21. Bradley, PS and Ade, JD. Are current physical match performance metrics in elite soccer fit for purpose or is the adoption of an integrated approach needed? *Int J Sports Physiol Perform* 13: 656–664, 2018.
- 22. Bradley, PS, Sheldon, W, Wooster, B, Olsen, P, Boanas, P, and Krustrup, P. Highintensity running in English FA Premier League soccer matches. *J Sports Sci* 27: 159–168, 2009.
- 23. Bret, C, Rahmani, A, Dufour, AB, Messonnier, L, and Lacour, JR. Leg strength and stiffness as ability factors in 100 m sprint running. *J Sports Med Phys Fitness* 42: 274–81, 2002.
- 24. Brice, B, Smith, N, and Dyson, R. Frequency of curvilinear motion during competitive soccer play. *J Sports Sci* 22: 485–593, 2004.
- 25. Brice, P, Smith, N, and Dyson, R. 3 Body segment orientations for curved running in soccer players. *Sci Footb VI* 18, 2008.
- 26. Brice, P, Smith, N, and Dyson, R. Curved running in soccer: kinematic differences between the inside and outside limbs. In: ISBS-Conference Proceedings Archive.2008.
- 27. Brown, SR, Brughelli, M, and Hume, PA. Knee mechanics during planned and unplanned sidestepping: A systematic review and meta-analysis. *Sport Med* 44: 1573–1588, 2014.
- Buchheit, M and Mendez-Villanueva, A. Effects of age, maturity and body dimensions on match running performance in highly trained under-15 soccer players. *J Sports Sci* 32: 1271–1278, 2014.
- 29. Buchheit, M, Mendez-Villanueva, A, Simpson, BM, and Bourdon, PC. Match running performance and fitness in youth soccer. *Int J Sports Med* 31: 818–825, 2010.
- 30. Buchheit, M and Simpson, BM. Player-tracking technology: Half-full or half-empty glass? *Int J Sports Physiol Perform* 12: S2-35, 2017.
- 31. Caldbeck, P. Contextual Sprinting in Football. Liverpool John Moores University, 2019.

- 32. Caldwell, GE and Chapman, AE. Applied muscle modelling: Implementation of muscle-specific models. *Comput Biol Med* 19: 417–434, 1989.
- 33. Carling, C, Bloomfield, J, Nelsen, L, and Reilly, T. The Role of Motion Analysis in Elite Soccer. *Sport Med* 38: 839–862, 2008.
- 34. Carling, C, Le Gall, F, and Dupont, G. Analysis of repeated high-intensity running performance in professional soccer. *J Sports Sci* 30: 325–336, 2012.
- 35. Carling, C, Le Gall, F, Reilly, T, and Williams, AM. Do anthropometric and fitness characteristics vary according to birth date distribution in elite youth academy soccer players? *Scand J Med Sci Sports* 19: 3–9, 2008.
- 36. Chaabene, H, Prieske, O, Negra, Y, and Granacher, U. Change of direction speed: Toward a strength training approach with accentuated eccentric muscle actions. *Sport Med* 48: 1773–1779, 2018.
- 37. Chang, YH and Kram, R. Limitations to maximum running speed on flat curves. *J Exp Biol* 210: 971–982, 2007.
- 38. Chelly, SM and Denis, C. Leg power and hopping stiffness: relationship with sprint running performance. *Med Sci Sport Exerc* 33: 326–333, 2001.
- 39. Churchill, SM, Salo, AIT, and Trewartha, G. The effect of the bend on technique and performance during maximal effort sprinting. *Sport Biomech* 14: 106–121, 2015.
- 40. Churchill, SM, Trewartha, G, Bezodis, IN, and Salo, AIT. Force production during maximal effort bend sprinting: Theory vs reality. *Scand J Med Sci Sport* 26: 1171–1179, 2016.
- 41. Clark, KP and Weyand, PG. Are running speeds maximized with simple-spring stance mechanics? *J Appl Physiol* 117: 604–615, 2014.
- 42. Clarke, R, Aspe, R, Sargent, D, and Hughes, J. Technical models for change of direction: Biomechanical principles. *Prof Strength Cond* 17–23, 2018.
- 43. Cloke, DJ, Spencer, S, Hodson, A, and Deehan, D. The epidemiology of ankle injuries occurring in English Football Association academies. *Br J Sports Med* 43: 1119–1125, 2009.
- 44. Dalen, T, Jørgen, I, Gertjan, E, Havard, HG, and Ulrik, W. Player load, acceleration, and deceleration during forty-five competitive matches of elite soccer. *J Strength Cond Res* 30: 351–359, 2016.
- 45. Daly, C, Mccarthy Persson, U, Twycross-Lewis, R, Woledge, RC, and Morrissey, D. The biomechanics of running in athletes with previous hamstring injury: A case-control study. *Scand J Med Sci Sport* 26: 413–420, 2016.
- 46. Daniel, DM, Stone, M Lou, Dobson, BE, Fithian, DC, Rossman, DJ, and Kaufman, KR. Fate of the ACL-injured patient: a prospective outcome study. *Am J Sports Med* 22: 632–644, 1994.
- 47. David, S, Mundt, M, Komnik, I, and Potthast, W. Understanding cutting maneuvers The mechanical consequence of preparatory strategies and foot strike pattern. *Hum Mov Sci* 62: 202–210, 2018.
- 48. De-Ste-Croix, M. Advances in paediatric strength assessment: Changing our perspective on strength development. *J Sport Sci Med* 6: 292, 2007.

- 49. Debaere, S, Delecluse, C, Aerenhouts, D, Hagman, F, and Jonkers, I. From block clearance to sprint running: Characteristics underlying an effective transition. *J Sports Sci* 31: 137–149, 2013.
- 50. Delgado-Bordonau, J and Mendez-Villanueva, A. Tactical periodization: Mourinho's best-kept secret. *Soccer Journal,* 57: 29–34, 2012.
- 51. Dempsey, AR, Elliott, BC, Munro, BJ, Steele, JR, and Lloyd, DG. Whole body kinematics and knee moments that occur during an overhead catch and landing task in sport. *Clin Biomech* 27: 466–474, 2012.
- 52. Dempsey, AR, Lloyd, DG, Elliott, BC, Steele, JR, and Munro, BJ. Changing sidestep cutting technique reduces knee valgus loading. *Am J Sports Med* 37: 2194–2200, 2009.
- 53. Di-Salvo, V, Baron, R, González-Haro, C, Gormasz, C, Pigozzi, F, and Bachl, N. Sprinting analysis of elite soccer players during European Champions League and UEFA Cup matches. *J Sports Sci* 28: 1489–1494, 2010.
- 54. Di-Salvo, V, Baron, R, Tschan, H, Calderon Montero, F, Bachl, N, and Pigozzi, F. Performance characteristics according to playing position in elite soccer. *Int J Sports Med* 28: 222–227, 2007.
- 55. Donnelly, CJ, Elliott, BC, Ackland, TR, Doyle, TLA, Beiser, TF, Finch, CF, et al. An anterior cruciate ligament injury prevention framework: Incorporating the recent evidence. *Res Sport Med* 20: 239–262, 2012.
- 56. Dorn, TW, Schache, AG, and Pandy, MG. Muscular strategy shift in human running: Dependence of running speed on hip and ankle muscle performance. *J Exp Biol* 215: 1944–1956, 2012.
- 57. Dos'Santos, T, Thomas, C, Comfort, P, and Jones, PA. The effect of angle and velocity on change of direction biomechanics: An angle-velocity trade-off. *Sport Med* 48: 2235–2253, 2018.
- Dos'Santos, T, Thomas, C, Comfort, P, and Jones, PA. The effect of training interventions on change of direction biomechanics associated with increased anterior cruciate ligament loading: a scoping review. *Sport Med* Sep 6: 1–23, 2019.
- 59. Dos'Santos, T, Thomas, C, Jones, PA, and Comfort, P. Mechanical determinants of faster change of direction speed performance in male athletes. *J Strength Cond Res* 31: 696–705, 2017.
- 60. Dos'Santos, T, McBurnie, A, Thomas, C, Comfort, P, and Jones, P. Biomechanical comparison of cutting techniques: A review and practical applications. *Strength Cond J* 41: 40–45, 2019.
- 61. Dos'Santos, T, McBurnie, A, Thomas, C, Comfort, P, and Jones, PA. Biomechanical determinants of the modified and traditional 505 change of direction speed test. *J strength Cond Res* 34: 1285–1296, 2020.
- 62. Dos'Santos, T, Thomas, C, Comfort, P, and Jones, PA. Role of the penultimate foot contact during change of direction: implications on performance and risk of injury. *Strength Cond J* 41: 87–104, 2019.
- 63. Faude, O, Koch, T, and Meyer, T. Straight sprinting is the most frequent action in goal situations in professional football. *J Sports Sci* 30: 625–631, 2012.

- 64. Ferguson, M and Patricios, J. What is the relationship between groin pain in athletes and femoroacetabular impingement? Br. J. Sports Med. 1074–1075, 2014.
- 65. Filter, A, Olivares-Jabalera, J, Santalla, A, Morente-Sánchez, J, Robles-Rodríguez, J, Requena, B, et al. Curve sprinting in soccer: Kinematic and neuromuscular analysis. *Int J Sports Med* 41: 1–7, 2020.
- 66. Fílter, A, Olivares, J, Santalla, A, Nakamura, FY, Loturco, I, and Requena, B. New curve sprint test for soccer players: Reliability and relationship with linear sprint. *J Sports Sci* 1–6, 2019.
- 67. Fitzpatrick, JF, Linsley, A, and Musham, C. Running the curve: a preliminary investigation into curved sprinting during football match-play. *Sport Perform and Sci Reports* 55: 1–3, 2019.
- 68. Folland, JP and Williams, AG. The adaptations to strength training: Morphological and neurological contributions to increased strength. *Sport Med* 37: 145–168, 2007.
- 69. Fuerst, P, Gollhofer, A, and Gehring, D. Preparation time influences ankle and knee joint control during dynamic change of direction movements. *J Sports Sci* 35: 762–768, 2017.
- 70. Le Gall, F, Carling, C, Reilly, T, Vandewalle, H, Church, J, and Rochcongar, P. Incidence of injuries in elite French youth soccer players: A 10-season study. *Am J Sports Med* 34: 928–938, 2006.
- 71. Gastin, PB, Bennett, G, and Cook, J. Biological maturity influences running performance in junior Australian football. *J Sci Med Sport* 16: 140–145, 2013.
- 72. Gleeson, N, Eston, R, Marginson, V, and McHugh, M. Effects of prior concentric training on eccentric exercise induced muscle damage. *Br J Sports Med* 37: 119–125, 2003.
- 73. Goto, H, Morris, JG, and Nevill, ME. Match analysis of U9 and U10 English Premier League Academy soccer players ising a Global Positioning System. *J Strength Cond Res* 29: 954–963, 2015.
- 74. Granero-Gil, P, Bastida-Castillo, A, Rojas-Valverde, D, Gómez-Carmona, CD, Sánchez, E de la C, and Pino-Ortega, J. Influence of contextual variables in the changes of direction and centripetal force generated during an elite-level soccer team season. *Int J Environ Res Public Health* 17: 967, 2020.
- 75. Guilhem, G, Doguet, V, Hauraix, H, Lacourpaille, L, Jubeau, M, Nordez, A, et al. Muscle force loss and soreness subsequent to maximal eccentric contractions depend on the amount of fascicle strain in vivo. *Acta Physiol* 217: 152–163, 2016.
- 76. Al Haddad, H, Simpson, BM, Buchheit, M, Di Salvo, V, and Mendez-Villanueva, A. Peak match speed and maximal sprinting speed in young soccer players: Effect of age and playing position. *Int J Sports Physiol Perform* 10: 888–896, 2015.
- 77. Hader, K, Mendez-Villanueva, A, Palazzi, D, Ahmaidi, S, and Buchheit, M. Metabolic power requirement of change of direction speed in young soccer players: Not all is what it seems. *PLoS One* 11: e0149839, 2016.
- 78. Hader, K, Palazzi, D, and Buchheit, M. Change of direction speed in soccer: How much braking is enough? *Kinesiol Int J Fundam Appl Kinesiol* 47: 67–74, 2015.
- 79. Hall, ECR, Larruskain, J, Gil, SM, Lekue, JA, Baumert, P, Rienzi, E, et al. An injury

audit in high-level male youth soccer players from English, Spanish, Uruguayan and Brazilian academies. *Phys Ther Sport* 44: 53–60, 2020.

- Harley, JA, Barnes, CA, Portas, M, Lovell, R, Barrett, S, Paul, D, et al. Motion analysis of match-play in elite U12 to U16 age-group soccer players. *J Sports Sci* 28: 1391– 1397, 2010.
- 81. Harper, DJ, Carling, C, and Kiely, J. High-intensity acceleration and deceleration demands in elite team sports competitive match play: A systematic review and metaanalysis of observational studies. *Sport Med* 49: 1923–1947, 2019.
- 82. Harper, DJ, Jordan, AR, and Kiely, J. Relationships between eccentric and concentric knee strength capacities and maximal linear deceleration ability in male academy soccer players. *J Strength Cond Res* Published ahead of print, 2018.
- 83. Harper, DJ and Kiely, J. Damaging nature of decelerations: Do we adequately prepare players? *BMJ Open Sport Exerc Med* 4, 2018.
- Haugen, TA, Tønnessen, E, and Seiler, S. Anaerobic performance testing of professional soccer players 1995-2010. *Int J Sports Physiol Perform* 8: 148–156, 2013.
- 85. Hewett, TE, Myer, GD, and Ford, KR. Decrease in neuromuscular control about the knee with maturation in female athletes. *J Bone Jt Surg* 86: 1601–1608, 2004.
- 86. Hewett, TE, Myer, GD, Ford, KR, Heidt, RS, Colosimo, AJ, McLean, SG, et al. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: A prospective study. *Am J Sports Med* 33: 492–501, 2005.
- 87. Hicks, DS, Schuster, JG, Samozino, P, and Morin, J-B. Improving mechanical effectiveness during sprint acceleration: Practical recommendations and guidelines. *Strength Cond J* 42: 45–62, 2020.
- De Hoyo, M, Cohen, DD, Sañudo, B, Carrasco, L, Álvarez-Mesa, A, del Ojo, JJ, et al. Influence of football match time–motion parameters on recovery time course of muscle damage and jump ability. *J Sports Sci* 34: 1363–1370, 2016.
- Hunter, JP, Marshall, RN, and McNair, PJ. Relationships between ground reaction force impulse and kinematics of sprint-running acceleration. *J Appl Biomech* 21: 31– 43, 2005.
- 90. Iga, J, George, K, Lees, A, and Reilly, T. Cross-sectional investigation of indices of isokinetic leg strength in youth soccer players and untrained individuals. *Scand J Med Sci Sport* 19: 714–719, 2009.
- 91. Ishimura, K and Sakurai, S. Comparison of inside contact phase and outside contact phase in curved sprinting. *Proc 28th Int Conf Biomech Sport* In: ISBS-C, 2010.
- 92. Jaspers, A, Kuyvenhoven, JP, Staes, F, Frencken, WGP, Helsen, WF, and Brink, MS. Examination of the external and internal load indicators' association with overuse injuries in professional soccer players. *J Sci Med Sport* 21: 579–585, 2018.
- 93. Jeffreys, I. A task-based approach to developing context-specific agility. *Strength Cond J* 33: 52–59, 2011.
- 94. Jeffreys, I. Agility training for team sports running the OODA loop. *Prof Strength Cond* 42: 15–21, 2016.

- 95. Jeffreys, I, Huggins, S, and Davies, N. Delivering a gamespeed-focused speed and agility development program in an english premier league soccer academy. *Strength Cond J* 40: 23–32, 2018.
- 96. Johnson, DM, Williams, S, Bradley, B, Sayer, S, Murray Fisher, J, and Cumming, S. Growing pains: Maturity associated variation in injury risk in academy football. *Eur J Sport Sci* 20: 544–552, 2019.
- 97. Jones, P, Bampouras, TM, and Marrin, K. An investigation into the physical determinants of change of direction speed. *J Sports Med Phys Fitness* 49: 97–104, 2009.
- 98. Jones, P, Thomas, C, Dos'Santos, T, McMahon, J, and Graham-Smith, P. The role of eccentric strength in 180 turns in female soccer players. *Sports* 5: 42, 2017.
- 99. Jones, PA, Dos'Santos, T, McMahon, JJ, and Graham-Smith, P. Contribution of eccentric strength to cutting performance in female soccer players. *J Strength Cond Res* Published ahead of print, 2019.
- 100. Jones, PA, Herrington, L, and Graham-Smith, P. Braking characteristics during cutting and pivoting in female soccer players. *J Electromyogr Kinesiol* 30: 46–54, 2016.
- 101. Jones, PA, Herrington, LC, and Graham-Smith, P. Technique determinants of knee joint loads during cutting in female soccer players. *Hum Mov Sci* 42: 203–211, 2015.
- 102. Kawamori, N, Nosaka, K, and Newton, RU. Relationships between ground reaction impulse and sprint acceleration performance in team sport athletes. *J Strength Cond Res* 27: 568–573, 2013.
- 103. Kemper, GLJ, Van Der Sluis, A, Brink, MS, Visscher, C, Frencken, WGP, and Elferink-Gemser, MT. Anthropometric injury risk factors in elite-standard youth soccer. *Int J Sports Med* 36: 1112–1117, 2015.
- 104. Kenneally-Dabrowski, CJB, Brown, NAT, Lai, AKM, Perriman, D, Spratford, W, and Serpell, BG. Late swing or early stance? A narrative review of hamstring injury mechanisms during high-speed running. *Scand J Med Sci Sport* 29: 1083–1091, 2019.
- 105. Khayambashi, K, Ghoddosi, N, Straub, RK, and Powers, CM. Hip muscle strength predicts noncontact anterior cruciate ligament injury in male and female athletes: a prospective study. *Am J Sports Med* 44: 355–361, 2016.
- 106. King, E, Franklyn-Miller, A, Richter, C, O'Reilly, E, Doolan, M, Moran, K, et al. Clinical and biomechanical outcomes of rehabilitation targeting intersegmental control in athletic groin pain: Prospective cohort of 205 patients. *Br J Sports Med* 52: 1054–1062, 2018.
- 107. Kugler, F and Janshen, L. Body position determines propulsive forces in accelerated running. *J Biomech* 43: 343–348, 2010.
- 108. Von Lieres Und Wilkau, HC, Bezodis, NE, Morin, J-B, Irwin, G, Simpson, S, and Bezodis, IN. The importance of duration and magnitude of force application to sprint performance during the initial acceleration, transition and maximal velocity phases. *J Sports Sci* Published ahead of print, 2020.
- 109. Lloyd, DG and Buchanan, TS. Strategies of muscular support of varus and valgus isometric loads at the human knee. *J Biomech* 34: 1257–1267, 2001.

- 110. Lloyd, RS, Cronin, JB, Faigenbaum, AD, Haff, GG, Howard, R, Kraemer, WJ, et al. National Strength and Conditioning Association position statement on long-term athletic development. *J Strength Cond Res* 30: 1491–1509, 2016.
- 111. Lloyd, RS, Meyers, RW, and Oliver, JL. The natural development and trainability of plyometric ability during childhood. *Strength Cond J* 33: 23–32, 2011.
- 112. Lloyd, RS and Oliver, JL. The youth physical development model: A new approach to long-term athletic development. *Strength Cond J* 34: 61–72, 2012.
- 113. Lloyd, RS, Radnor, JM, De Ste Croix, MBA, Cronin, JB, and Oliver, JL. Changes in sprint and jump performances after traditional, plyometric, and combined resistance training in male youth pre- and post-peak height velocity. *J Strength Cond Res* 30: 1239–1247, 2016.
- 114. Lloyd, RS, Read, P, Oliver, JL, Meyers, RW, Nimphius, S, and Jeffreys, I. Considerations for the development of agility during childhood and adolescence. *Strength Cond J* 35: 2–11, 2013.
- 115. Lockie, RG, Murphy, AJ, Schultz, AB, Knight, TJ, and De Jonge, XAKJ. The effects of different speed training protocols on sprint acceleration kinematics and muscle strength and power in field sport athletes. *J Strength Cond Res* 26: 1539–1550, 2012.
- 116. Lockie, RG, Schultz, AB, Jeffriess, MD, and Callaghan, SJ. The relationship between bilateral differences of knee flexor and extensor isokinetic strength and multidirectional speed. *Isokinet Exerc Sci* 20: 211–219, 2012.
- 117. Lovell, R, Fransen, J, Ryan, R, Massard, T, Cross, R, Eggers, T, et al. Biological maturation and match running performance: A national football (soccer) federation perspective. *J Sci Med Sport* 22: 1139–1145, 2019.
- 118. Luo, G and Stefanyshyn, D. Limb force and non-sagittal plane joint moments during maximum-effort curve sprint running in humans. *J Exp Biol* 215: 4314–4321, 2012.
- 119. Maćkała, K, Fostiak, M, and Kowalski, K. Selected determinants of acceleration in the 100m Sprint. *J Hum Kinet* 45: 135–148, 2015.
- 120. Malina, RM, Bouchard, C, and Bar-Or, O. Growth, maturation, and physical activity. 2nd ed. Champaign, Illinois, United States: Human Kinetics, 2004.
- 121. Malina, RM, Reyes, MEP, Eisenmann, JC, Horta, L, Rodrigues, J, and Miller, R. Height, mass and skeletal maturity of elite Portuguese soccer players aged 11–16 years. *J Sports Sci* 18: 685–693, 2000.
- 122. Malone, S, Owen, A, Mendes, B, Hughes, B, Collins, K, and Gabbett, TJ. High-speed running and sprinting as an injury risk factor in soccer: Can well-developed physical qualities reduce the risk? *J Sci Med Sport* 21: 257–262, 2018.
- 123. Malone, S, Roe, M, Doran, DA, Gabbett, TJ, and Collins, K. High chronic training loads and exposure to bouts of maximal velocity running reduce injury risk in elite Gaelic football. *J Sci Med Sport* 20: 250–254, 2017.
- 124. Maniar, N, Schache, AG, Cole, MH, and Opar, DA. Lower-limb muscle function during sidestep cutting. *J Biomech* 82: 186–192, 2019.
- 125. Maniar, N, Schache, AG, Sritharan, P, and Opar, DA. Non-knee-spanning muscles contribute to tibiofemoral shear as well as valgus and rotational joint reaction moments during unanticipated sidestep cutting. *Sci Rep* 8: 1–10, 2018.

- 126. Marshall, BM, Franklyn-Miller, AD, King, EA, Moran, KA, Strike, SC, and Falvey, ÉC. Biomechanical factors associated with time to complete a change of direction cutting maneuver. *J Strength Cond Res* 28: 2845–2851, 2014.
- 127. Materne, O, Farooq, A, Johnson, A, Greig, M, and McNaughton, L. Relationship between injuries and somatic maturation in highly trained youth soccer players. In: Proceedings of the World Congress on Science and Soccer.2015.
- 128. McBurnie, AJ, Dos'Santos, T, and Jones, PA. Biomechanical associates of performance and knee joint loads during a 70–90° cutting maneuver in sub-elite soccer players. *J Strength Cond Res* Published ahead of print, 2019.
- 129. McBurnie, AJ, Parr, J, Kelly, DM, and Dos'Santos, T. Multi-directional speed in youth soccer players: Practical applications and programming considerations. *Strength Cond J* Accepted for publication in SCJ, 2021.
- 130. McCall, A, Carling, C, Nedelec, M, Davison, M, Le Gall, F, Berthoin, S, et al. Risk factors, testing and preventative strategies for non-contact injuries in professional football: current perceptions and practices of 44 teams from various premier leagues. *Br J Sports Med* 48: 1352–1357, 2014.
- 131. McCall, A, Pruna, R, Van der Horst, N, Dupont, G, Buchheit, M, Coutts, AJ, et al. Exercise-based strategies to prevent muscle injury in male elite footballers: An expertled delphi survey of 21 practitioners belonging to 18 teams from the Big-5 European leagues. *Sport Med* Published online: 16 July, 2020.
- 132. McKay, D, Broderick, C, and Steinbeck, K. The adolescent athlete: A developmental approach to injury risk. *Pediatr Exerc Sci* 28: 488–500, 2016.
- McLean, SG, Lipfert, SW, and Van Den Bogert, AJ. Effect of gender and defensive opponent on the biomechanics of sidestep cutting. *Med Sci Sports Exerc* 36: 1008, 2004.
- 134. Mendiguchia, J, Conceição, F, Edouard, P, Fonseca, M, Pereira, R, Lopes, H, et al. Sprint versus isolated eccentric training: Comparative effects on hamstring architecture and performance in soccer players. *PLoS One* 15: e0228283, 2020.
- 135. Mendiguchia, J, Samozino, P, Martinez-Ruiz, E, Brughelli, M, Schmikli, S, Morin, JB, et al. Progression of mechanical properties during on-field sprint running after returning to sports from a hamstring muscle injury in soccer players. *Int J Sports Med* 35: 690–695, 2014.
- 136. Morin, JB, Edouard, P, and Samozino, P. Technical ability of force application as a determinant factor of sprint performance. *Med Sci Sports Exerc* 43: 1680–1688, 2011.
- 137. Morin, JB, Gimenez, P, Edouard, P, Arnal, P, Jiménez-Reyes, P, Samozino, P, et al. Sprint acceleration mechanics: The major role of hamstrings in horizontal force production. *Front Physiol* 6: 404, 2015.
- 138. Morin, JB, Slawinski, J, Dorel, S, de villareal, ES, Couturier, A, Samozino, P, et al. Acceleration capability in elite sprinters and ground impulse: Push more, brake less? *J Biomech* 48: 3149–3154, 2015.
- 139. Mornieux, G, Gehring, D, Fürst, P, and Gollhofer, A. Anticipatory postural adjustments during cutting manoeuvres in football and their consequences for knee injury risk. *J Sports Sci* 32: 1255–1262, 2014.
- 140. Myer, GD, Ford, KR, Di Stasi, SL, Barber Foss, KD, Micheli, LJ, and Hewett, TE. High

knee abduction moments are common risk factors for patellofemoral pain (PFP) and anterior cruciate ligament (ACL) injury in girls: Is PFP itself a predictor for subsequent ACL injury? *Br J Sports Med* 49: 118–122, 2015.

- 141. Nagahara, R, Kanehisa, H, and Fukunaga, T. Ground reaction force across the transition during sprint acceleration. *Scand J Med Sci Sport* 30: 450–461, 2019.
- 142. Nagahara, R, Matsubayashi, T, Matsuo, A, and Zushi, K. Kinematics of transition during human accelerated sprinting. *Biol Open* 3: 689–699, 2014.
- 143. Nagahara, R, Naito, H, Morin, JB, and Zushi, K. Association of acceleration with spatiotemporal variables in maximal sprinting. *Int J Sports Med* 35: 755–761, 2014.
- 144. Naughton, G, Farpour-Lambert, NJ, Carlson, J, Bradney, M, and Van Praagh, E. Physiological issues surrounding the performance of adolescent athletes. *Sport Med* 30: 309–325, 2000.
- Nedergaard, NJ, Kersting, U, and Lake, M. Using accelerometry to quantify deceleration during a high-intensity soccer turning manoeuvre. *J Sports Sci* 32: 1897– 1905, 2014.
- 146. Neptune, RR, Wright, IC, and Van Den Bogert, AJ. Muscle coordination and function during cutting movements. *Med Sci Sports Exerc* 31: 294–302, 1999.
- 147. Nikolaidis, PT, Ingebrigtsen, J, and Jeffreys, I. The effects of anthropometry and leg muscle power on drive and transition phase of acceleration: A longitudinal study on young soccer players. *J Sports Med Phys Fitness* 56: 1156–62, 2016.
- 148. Nimphius, S. Training change of direction and agility. In: Advanced Strenth and Conditioning: An Evidence-based Approach.Routledge, 2017. pp. 293–308
- 149. Nimphius, S, Callaghan, SJ, Bezodis, NE, and Lockie, RG. Change of direction and agility tests: Challenging our current measures of performance. *Strength Cond J* 40: 26–38, 2018.
- 150. Nimphius, S, McGuigan, MR, and Newton, RU. Relationship between strength, power, speed, and change of direction performance of female softball players. *J Strength Cond Res* 24: 885–895, 2010.
- 151. O'Kane, JW, Neradilek, M, Polissar, N, Sabado, L, Tencer, A, and Schiff, MA. Risk factors for lower extremity overuse injuries in female youth soccer players. *Orthop J Sport Med* 5: 2325967117733963, 2017.
- 152. Oliver, JL, Lloyd, RS, and Rumpf, MC. Developing speed throughout childhood and adolescence: The role of growth, maturation and training. *Strength Cond J* 35: 42–48, 2013.
- 153. Opar, DA, Williams, MD, Timmins, RG, Hickey, J, Duhig, SJ, and Shield, AJ. Eccentric hamstring strength and hamstring injury risk in Australian footballers. *Med Sci Sports Exerc* 47: 857–865, 2015.
- 154. Padua, DA, DiStefano, LJ, Hewett, TE, Garrett, WE, Marshall, SW, Golden, GM, et al. National athletic trainers' association position statement: Prevention of anterior cruciate ligament injury. *J Athl Train* 53: 5–19, 2018.
- 155. Palucci Vieira, LH, Carling, C, Barbieri, FA, Aquino, R, and Santiago, PRP. Match running performance in young soccer Players: A systematic review. *Sport Med* 49: 289–318, 2019.

- 156. Ploutz-Snyder, LL, Tesch, PA, and Dudley, GA. Increased vulnerability to eccentric exercise-induced dysfunction and muscle injury after concentric training. *Arch Phys Med Rehabil* 79: 58–61, 1998.
- 157. Potier, TG, Alexander, CM, and Seynnes, OR. Effects of eccentric strength training on biceps femoris muscle architecture and knee joint range of movement. *Eur J Appl Physiol* 105: 939–944, 2009.
- 158. Powell, JW and Barber-Foss, KD. Sex-related injury patterns among selected high school sports. *Am J Sports Med* 28: 385–391, 2000.
- 159. Price, R, Hawkins, R, Hulse, M, Hodson, A, and Gibson, M. The Football Association medical research programme: an audit of injuries in academy youth football. *Br J Sports Med* 38: 466–471, 2004.
- 160. Quatman-Yates, CC, Quatman, CE, Meszaros, AJ, Paterno, M V., and Hewett, TE. A systematic review of sensorimotor function during adolescence: A developmental stage of increased motor awkwardness? *Br J Sports Med* 46: 649–655, 2012.
- 161. Rabita, G, Dorel, S, Slawinski, J, Sàez-de-Villarreal, E, Couturier, A, Samozino, P, et al. Sprint mechanics in world-class athletes: A new insight into the limits of human locomotion. *Scand J Med Sci Sport* 25: 583–594, 2015.
- 162. Radnor, JM, Lloyd, RS, and Oliver, JL. Individual response to different forms of resistance training in school-aged boys. *J Strength Cond Res* 31: 787–797, 2017.
- 163. Read, PJ, Oliver, JL, De Ste Croix, MBA, Myer, GD, and Lloyd, RS. Neuromuscular risk factors for knee and ankle ligament injuries in male youth soccer players. *Sport Med* 46: 1059–1066, 2016.
- 164. Read, PJ, Oliver, JL, De Ste Croix, MBA, Myer, GD, and Lloyd, RS. The scientific foundations and associated injury risks of early soccer specialisation. *J Sports Sci* 34: 2295–2302, 2016.
- 165. Reilly, T, Bangsbo, J, and Franks, A. Anthropometric and physiological predispositions for elite soccer. *J Sports Sci* 18: 669–683, 2000.
- 166. Ritsche, P, Bernhard, T, Roth, R, Lichtenstein, E, Keller, M, Zingg, S, et al. M. biceps femoris architecture and sprint ability in youth soccer players: a cross-sectional analysis. *SportRxiv* Published ahead of print, 2020.
- 167. Robinson, G, O'Donoghue, P, and Nielson, P. Path changes and injury risk in English FA Premier League soccer. *Int J Perform Anal Sport* 11: 40–56, 2011.
- 168. Rosenbloom, CA, Loucks, AB, and Ekblom, B. Special populations: The female player and the youth player. *J Sports Sci* 24: 783–793, 2006.
- 169. Rumpf, MC and Cronin, J. Injury incidence, body site, and severity in soccer players aged 6-18 years: Implications for injury prevention. *Strength Cond J* 34: 20–31, 2012.
- 170. Rumpf, MC, Cronin, JB, Pinder, SD, Oliver, J, and Hughes, M. Effect of different training methods on running sprint times in male youth. *Pediatr Exerc Sci* 24: 170–186, 2012.
- 171. Rumpf, MC, Lockie, RG, Cronin, JB, and Jalilvand, F. Effect of different sprint training methods on sprint performance over various distances: a brief review. *J Strength Cond Res* 30: 1767–1785, 2016.

- 172. Samozino, P, Rabita, G, Dorel, S, Slawinski, J, Peyrot, N, Saez de Villarreal, E, et al. A simple method for measuring power, force, velocity properties, and mechanical effectiveness in sprint running. *Scand J Med Sci Sport* 26: 648–658, 2016.
- 173. Saward, C, Morris, JG, Nevill, ME, Nevill, AM, and Sunderland, C. Longitudinal development of match-running performance in elite male youth soccer players. *Scand J Med Sci Sports* 26: 933–942, 2016.
- 174. Schache, AG, Brown, NAT, and Pandy, MG. Modulation of work and power by the human lower-limb joints with increasing steady-state locomotion speed. *J Exp Biol* 218: 2472–2481, 2015.
- 175. Schache, AG, Dorn, TW, Williams, GP, Brown, NAT, and Pandy, MG. Lower-limb muscular strategies for increasing running speed. *J Orthop Sports Phys Ther* 44: 813–824, 2014.
- 176. Sheppard, J and Young, W. Agility literature review: Classifications, training and testing. *J Sports Sci* 24: 919–932, 2006.
- 177. Shin, CS, Chaudhari, AM, and Andriacchi, TP. Valgus plus internal rotation moments increase anterior cruciate ligament strain more than either alone. *Med Sci Sports Exerc* 43: 1484–1491, 2011.
- 178. Smirniotou, A, Katsikas, C, Paradisis, G, Argeitaki, P, Zacharogiannis, E, and Tziortzis, S. Strength-power parameters as predictors of sprinting performance. *J Sports Med Phys Fitness* 48: 447, 2008.
- 179. Smith, N, Dyson, R, Hale, T, and Janaway, L. Contributions of the inside and outside leg to maintenance of curvilinear motion on a natural turf surface. *Gait Posture* 24: 453–458, 2006.
- 180. Smith, NA, Dyson, RJ, and Hale, T. Lower extremity muscular adaptations to curvilinear motion in soccer. *J Hum Mov Stud* 33: 139–153, 1997.
- 181. Spiteri, T, Cochrane, JL, Hart, NH, Haff, GG, and Nimphius, S. Effect of strength on plant foot kinetics and kinematics during a change of direction task. *Eur J Sport Sci* 13: 646–652, 2013.
- 182. Spiteri, T, Newton, RU, Binetti, M, Hart, NH, Sheppard, JM, and Nimphius, S. Mechanical determinants of faster change of direction and agility performance in female basketball athletes. *J Strength Cond Res* 29: 2205–2214, 2015.
- 183. Spiteri, T, Nimphius, S, Hart, NH, Specos, C, Sheppard, JM, and Newton, RU. Contribution of strength characteristics to change of direction and agility performance in female basketball athletes. *J Strength Cond Res* 28: 2414–2423, 2014.
- 184. Stolen, T, Chamari, K, Castagna, C, and Wisloff, U. Physiology of soccer: An update. *Sport Med* 35: 501–536, 2005.
- 185. Stone, MH, Stone, M, and Sands, WA. Principles and Practice of Resistance Training. 2007.
- 186. Sugiura, Y, Saito, T, Sakuraba, K, Sakuma, K, and Suzuki, E. Strength deficits identified with concentric action of the hip extensors and eccentric action of the hamstrings predispose to hamstring injury in elite sprinters. *J Orthop Sports Phys Ther* 38: 457–464, 2008.
- 187. Sun, Y, Wei, S, Zhong, Y, Fu, W, Li, L, and Liu, Y. How joint torques affect hamstring

injury risk in sprinting swing-stance transition. Med Sci Sports Exerc 47: 373, 2015.

- 188. Tak, I, Weir, A, Langhout, R, Waarsing, JH, Stubbe, J, Kerkhoffs, G, et al. The relationship between the frequency of football practice during skeletal growth and the presence of a cam deformity in adult elite football players. *Br J Sports Med* 49: 630–634, 2015.
- 189. Taylor, KA, Terry, ME, Utturkar, GM, Spritzer, CE, Queen, RM, Irribarra, LA, et al. Measurement of in vivo anterior cruciate ligament strain during dynamic jump landing. *J Biomech* 44: 365–371, 2011.
- 190. Tillaar, R Van Den, Asmund, J, and Solheim, B. Comparison of hamstring muscle activation during high-speed running and various hamstring strengthening exercises. *Int J Sports Phys Ther* 12: 718, 2017.
- 191. Timmins, RG, Bourne, MN, Shield, AJ, Williams, MD, Lorenzen, C, and Opar, DA. Short biceps femoris fascicles and eccentric knee flexor weakness increase the risk of hamstring injury in elite football (soccer): A prospective cohort study. *Br J Sports Med* 50: 1524–1535, 2016.
- 192. Turner, A, Walker, S, Stembridge, M, Coneyworth, P, Reed, G, Birdsey, L, et al. A testing battery for the assessment of fitness in soccer players. *Strength Cond J* 33: 29–39, 2011.
- 193. Turner, AN and Stewart, PF. Strength and conditioning for soccer players. *Strength Cond J* 36: 1–13, 2014.
- 194. Van-der-Sluis, A, Elferink-Gemser, M, Coelho-e-Silva, M, Nijboer, J, Brink, M, and Visscher, C. Sport injuries aligned to peak height velocity in talented pubertal soccer players. *Int J Sports Med* 35: 351–355, 2013.
- 195. Van-Hooren, B, De, M, and Croix, S. Sensitive periods to train general motor abilities in children and adolescents: Do they exist? A critical appraisal. *Strength Cond J* Published ahead of print, 2020.
- 196. Vanrenterghem, J, Nedergaard, NJ, Robinson, MA, and Drust, B. Training load monitoring in team sports: a novel framework separating physiological and biomechanical load-adaptation pathways. *Sport Med* 47: 2135–2142, 2017.
- 197. Verheul, J, Warmenhoven, J, Lisboa, P, Gregson, W, Vanrenterghem, J, and Robinson, MA. Identifying generalised segmental acceleration patterns that contribute to ground reaction force features across different running tasks. *J Sci Med Sport* 22: 1355–1360, 2019.
- 198. Vescovi, JD. Sprint speed characteristics of high-level American female soccer players: Female Athletes in Motion (FAiM) Study. *J Sci Med Sport* 15: 474–478, 2012.
- 199. Vigne, G, Gaudino, C, Rogowski, I, Alloatti, G, and Hautier, C. Activity profile in an elite Italian soccer team. *Int J Sports Med* 31: 304–310, 2010.
- 200. Wang, Q, Ghasem-Zadeh, A, Wang, XF, Iuliano-Burns, S, and Seeman, E. Trabecular bone of growth plate origin influences both trabecular and cortical morphology in adulthood. *J Bone Miner Res* 26: 1577–1583, 2011.
- 201. Wang, Q, Wang, XF, Iuliano-Burns, S, Ghasem-Zadeh, A, Zebaze, R, and Seeman, E. Rapid growth produces transient cortical weakness: A risk factor for metaphyseal fractures during puberty. *J Bone Miner Res* 25: 1521–1526, 2010.

- 202. Weinhandl, JT, Earl-Boehm, JE, Ebersole, KT, Huddleston, WE, Armstrong, BSR, and O'Connor, KM. Reduced hamstring strength increases anterior cruciate ligament loading during anticipated sidestep cutting. *Clin Biomech* 29: 752–759, 2014.
- 203. Weinhandl, JT, Earl-Boehm, JE, Ebersole, KT, Huddleston, WE, Armstrong, BSR, and O'Connor, KM. Reduced hamstring strength increases anterior cruciate ligament loading during anticipated sidestep cutting. *Clin Biomech* 29: 752–759, 2014.
- 204. Welch, N, Richter, C, Moran, K, and Franklyn-Miller, A. Principal Component Analysis of the Associations Between Kinetic Variables in Cutting and Jumping, and Cutting Performance Outcome. *J Strength Cond Res* Published ahead of print, 2019.
- 205. Weyand, PG, Sternlight, DB, Bellizzi, MJ, and Wright, S. Faster top running speeds are achieved with greater ground forces not more rapid leg movements. *J Appl Physiol* 89: 1991–1999, 2000.
- 206. Wild, J, Bezodis, N, Blagrove, R, and Bezodis, I. A biomechanical comparison of accelerative and maximum velocity sprinting: Specific strength training considerations. *Prof Strength Cond* 21: 23–37, 2011.
- 207. Woods, C, Hawkins, RD, Maltby, S, Hulse, M, Thomas, A, and Hodson, A. The Football Association Medical Research Programme: An audit of injuries in professional football analysis of hamstring injuries. *Br J Sports Med* 38: 36–41, 2004.
- 208. Wormhoudt, R, Savelsbergh, GJP, Teunissen, AJW, and Davids, K. The athletic skills model: optimizing talent development through movement education. Routledge, 2017.
- 209. Wyatt, H, Weir, G, van Emmerik, R, Jewell, C, and Hamill, J. Whole-body control of anticipated and unanticipated sidestep manoeuvres in female and male team sport athletes. *J Sports Sci* 37: 2263–2269, 2019.
- 210. Young, W and Farrow, D. The importance of a sport-specific stimulus for training agility. *Strength Cond J* 35: 39–43, 2013.
- 211. Young, WB, Dawson, B, and Henry, GJ. Agility and change-of-direction speed are independent skills: Implications for training for agility in invasion sports. *Int J Sport Sci Coach* 10: 159–169, 2015.
- 212. Young, WB, James, R, and Montgomery, I. Is muscle power related to running speed with changes of direction? *J Sports Med Phys Fitness* 42: 282–288, 2002.
- Yu, B, Queen, RM, Abbey, AN, Liu, Y, Moorman, CT, and Garrett, WE. Hamstring muscle kinematics and activation during overground sprinting. *J Biomech* 41: 3121– 3126, 2008.
- 214. Yu, J, Sun, Y, Yang, C, Wang, D, Yin, K, Herzog, W, et al. Biomechanical insights into differences between the mid-acceleration and maximum velocity phases of sprinting. *J Strength Cond Res* 30: 1906–1916, 2016.
- 215. Zazulak, BT, Hewett, TE, Reeves, NP, Goldberg, B, and Cholewicki, J. Deficits in neuromuscular control of the trunk predict knee injury risk: A prospective biomechanical-epidemiologic study. *Am J Sports Med* 35: 1123–1130, 2007.