Manchester Metropolitan University

e-space

Manchester Metropolitan University's Research Repository

Please cite the Published Version

Leal Filho, W, Azeiteiro, UM, Balogun, AL, Setti, AFF, Mucova, SAR, Ayal, D, Totin, E, Lydia, AM, Kalaba, FK and Oguge, NO (2021) The influence of ecosystems services depletion to climate change adaptation efforts in Africa. Science of the Total Environment, 779. 146414 ISSN 0048-9697

DOI: https://doi.org/10.1016/j.scitotenv.2021.146414

Publisher: Elsevier

Version: Accepted Version

Downloaded from: https://e-space.mmu.ac.uk/628127/

Usage rights: CC BY-NC-ND Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Additional Information: Author accepted manuscript published by and copyright Elsevier.

Enquiries:
If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines)

The Influence of Ecosystems Services Depletion to Climate Change Adaptation Efforts in Africa

Science of the Total Environment 779 20 Jul 2021

https://www.sciencedirect.com/science/article/abs/pii/S0048969721014820?via%3Dihub

Walter Leal Filho[1], Ulisses Miranda Azeiteiro[2], Abdul-Lateef Balogun[3], Andréia Faraoni Freitas Setti[2]*, Serafino A.R. Mucova[2,9], Desalegn Ayal[4], Edmond Totin[5], Adeleke Mosunmola Lydia[6], Felix Kanungwe Kalaba[7], Nicholas Otienoh Oguge[8]

Corresponding author: *andreiasetti@ua.pt

1. Department of Natural Sciences, Manchester Metropolitan University, Chester Street, Manchester M1 5GD, United Kingdom. E-mail: w.leal@haw-hamburg.de.
2. Department of Biology & CESAM Centre for Environmental and Marine Studies, University of Aveiro, Portugal
3. Geospatial Analysis & Modelling Research (GAMR) Laboratory, Department of Civil & Environmental Engineering, University Teknologi PETRONAS (UTP), Malaysia
4. Centre for Food Security Studies (CFSS), College of Development Studies, Addis Ababa University, Ethiopia
5. Ecole de Foresterie Tropicale, Université Nationale d'Agriculture, Benin
6. Department of Fisheries and Aquaculture Technology, The Federal University of Technology Akure, Nigeria
7. School of Natural Resources, Copperbelt University, P.O Box 21692, Kitwe, Zambia.
8. Centre for Advanced Studies in Environmental Law & Policy (CASELAP), University of Nairobi, Kenya
9. Faculty of Natural Sciences, Lúrio University, Pemba, P.O. Box 958, Mozambique

Abstract

Climate change is one of the major challenges societies round the world face at present. Apart from efforts to achieve a reduction of emissions of greenhouse gases so as to mitigate the problem, there is a perceived need for adaptation initiatives urgently. Ecosystems are known to play an important role in climate change adaptation processes, since some of the services they provide, may reduce the impacts of extreme events and disturbance, such as wildfires, floods, and droughts. This role is especially important in regions vulnerable to climate change such as the African continent, whose adaptation capacity is limited by many geographic and socio-economic constraints. In Africa, interventions aimed at enhancing ecosystem services may play a key role in supporting climate change adaptation efforts. In order to shed some light on this aspect, this paper reviews the role of ecosystems services and investigates how they are being influenced by climate change in Africa. It contains a set of case studies from a sample of African countries, which serve the purpose to demonstrate the damages incurred, and how such damages disrupt ecosystem services. Based on the data gathered, some measures which may assist in fostering the cause of ecosystems services are listed, so as to cater for a better protection of some of the endangered Africa ecosystems, and the services they provide.

Key-words: Africa, ecosystem services, climate change, adaptation, best practices.

1 Introduction

Climate change is one of the major challenges societies round the world face at present. Apart from efforts to achieve a reduction of emissions of greenhouse gases so as to mitigate the problem, there is a perceived need for adaptation initiatives. Ecosystems are known to play an important role in climate change adaptation processes, since some of the services provide, may reduce the impacts of extreme events and disturbance, such as wildfires, floods, and

droughts. Ecosystem Services (ESs) have the potential to reduce vulnerability to climate change and enhance adaptation capacity (Pettinotti et al., 2018; Von Möllendorff and Hirschfeld, 2016; Hannah, 2015; Munang et al., 2013). Ecosystem-based adaptation (EbA), leverages nature's inherent capacity to protect communities from the severe effects of climate change. Capitalizing on ecosystems adaptation can increase the efficiency and effectiveness of adaptation strategies while promoting the resilience of social-ecological systems (Jones et al., 2012; Scarano, 2017, Pettinotti et al. 2018, Fedele, 2019). Productive EbA boosts biodiversity and thus ecosystem services which in turn eases adaptation to climate change impacts (CBD, 2009; Chong 2014; Zölch et al., 2018). It entails proper management of ESs to reduce the role of humans in altering ecosystems. Properly managed ecosystems can increase the guarantee of human safety from common natural hazards (Munang et al., 2013). For instance, ecosystems can be harnessed to minimise severe climatic impacts such as the rise of sea levels in coastal areas (Hannah, 2015). Generally, healthy ecosystems function as barriers to natural hazards thereby enhancing the resilience of socio-economic systems to climate change shocks (Abramovitz et al., 2002; Morecroft et al., 2012). Considering that ecosystem services also significantly support human well-being, it is imperative to preserve and enhance their integrity, by adopting appropriate management practices. In addition to protecting societies from disasters- bearing in mind that some are not strictly connected with climate change-, healthy ecosystems also improve societies' capacity to adjust to the impacts of such disasters and cope better (Sudmeier-Rieux et al., 2006; Gupta et al. 2017).

The wise use of ecosystems services in a climate change context focuses on utilizing various conservation strategies to safeguard these ecosystems -and the biodiversity they host- from climate change. This process also entails approaches aimed at the prevention of the occurrence of destructive factors like fire, coral bleaching, deforestation (Ojea et al., 2016; Hannah, 2015), and depletion of coral reefs, among others. The process of protection of

ecosystems services is also helpful in the sense that it prevents the deterioration of coastal and marine ecosystems (Liu and Stern, 2008), wetlands (Chaikumbung et al., 2016), mangroves (Brander et al., 2012), and peatlands (Pettinotti et al., 2018), among others. The safeguarding of such services has been documented as an effective strategy for climate change adaptation and to foster climate change resilience (Uy and Shaw, 2012, Leal Filho 2020).

For instance, mangroves' reforestation enhances coastal resilience by providing strong protection to shorelines, and flood plain management is a proven solution to flooding. Wind-sheltering and windbreaks to enhance the resilience of rangelands. Differentiating planting dates and crop types have a positive contribution to adaption in the agricultural sector (Munang et al., 2013). Tree planting forestalls erosion by enhancing soil stability. Mountain meadows, bushes, and forests serve as buffers from landslides and flash floods. Coral reefs serve as offshore breakwaters that minimize the effects of sea surges as well as storm waves on the shoreline (UNFCCC, 2008).

Multi-sectoral ecosystems services may also help to indirectly mitigate the effects of other phenomena such as landslides, floods, and erosion. Health ecosystems may also contribute to fight water and air pollution, to control diseases and pests, and to promote carbon sink (Nalau et al., 2018; Munang et al., 2013). A green environment created by a mix of trees, shrubs grass etc. helps decrease the negative impacts of the Urban Heat Island effect (UHI) (Chiabai et al., 2018). Collectively, the regulating services offered by ecosystems significantly influence climate change adaptation by reducing vulnerability and increasing overall resilience to extreme events. Similarly, ecosystems provide people with basic necessities such as food and water which increase their adaptive capacity (Uy and Shaw, 2012). The gradual depletion of ecosystem services seriously deprive people of their livelihoods. Low-income

levels, high rate of unemployment, and poverty are chronic stresses which negatively impact adaptation capacity in various regions, especially in Africa (Leal Filho et al., 2018).

The socio-economic, environmental, and cultural benefits emanating from ecosystems services conservation is a key component of sustainable adaptation. This is validated by the increasing body of evidence that highlights the crucial role of ecosystem services in building communities' resilience. The growing call for safeguarding ecosystems services, and mainstreaming this approach as part of climate change adaptation policies and implementation, illustrate the huge benefits and vast potentials offered (Uy and Shaw, 2012), which are often flexible, cost-effective, and have proven effective (Jones et al., 2012). This is also so in the context of urban planning through enhancing biodiversity and increasing recreation avenues (Doswald et al., 2014; Wamsler et al., 2014; Zölch et al., 2018), also harmonizing diverse interests of several stakeholders (Wamsler, 2017) and contributing to human security (Bourne et al., 2016).

Against this background, this paper reviews the role of ecosystems services and investigates how they are being influenced by climate change in Africa. It contains a set of case studies from a sample of African countries, which serve the purpose to demonstrate the damages incurred, and how such damages disrupt ecosystem services.

2 The importance of ecosystems services in an African context

Historically, ecosystem services had often been associated with tropical forests, freshwater, lakes, rivers, soil, minerals, and biodiversity (Holland et al., 2012; Green et al., 2013; MEA, 2014). Ecosystem services have been the bastion to livelihoods of most rural and urban communities in Africa. According to the IPBES (2018), more than 62% of the rural population depends directly on these services, while the urban and peri-urban population supplement their incomes, as well as their energy, medicine, and other essentials, from

98 ecosystem-based resources. Ecosystem services also significantly contributed to the
99 macroeconomy of many African countries (MEA, 2014; Ryan et al., 2016; IPBES, 2018).
100 The tourism sector is a major contributor to the national economies of East Africa (Kenya,
101 Uganda, Ethiopia, Rwanda, Democratic Republic of Congo, and Tanzania) with annual
102 average revenue of over USD 7 billion (World Bank, 2015).

103 Additionally, ecosystems have substantial spiritual value in traditional African belief systems.
104 Such belief systems have ecological roots or foundations (Schoffeleers, 1979; MEA, 2014;
105 IPBES, 2018). Also, these services function as gravesites and the dwelling places of powerful
106 ancestral spirits (Byers et al., 2001). Moreover, they are used as places to communicate with
107 the ancestors and for the annual rain-making ceremony (Wilson, 1989; Ryan et al., 2016).
108 The medicinal plants, generally classified as a provisioning service, also have a strong
109 cultural component, mainly for health problems (Ryan et al., 2016).

110 In Zambia, Mali, and Tanzania ecosystem services functioned as important assets to reduce
111 vulnerability during extreme climatic events (droughts and floods) (Robledo et al., 2012).
112 Additionally, the services provided by mangrove forests of the East Africa coast have been
113 extremely important in containing coastal erosion, reducing the intensity and frequency of
114 extreme climatic winds caused by tropical cyclones along the Indian Ocean coast (Ghermandi
115 et al., 2019).

116 The post-extreme event restoration of degraded areas in South Africa has carried out through
117 ecosystem-based adaptation which revived coastal and marine plant species (Mugwedi et al.,
118 2018). According to Ghermandi et al., (2019), coral reefs significantly prevented coastal
119 erosion in Comoros, Madagascar, Mayotte, Mozambique, Seychelles, and Tanzania with
120 100%, 55%, 98%, 16 %, 98%, 63% respectively. Services provided by seagrass were
121 responsible for 99% of the carbon sequestration in Comoros and Mayotte. Likewise,

122 mangroves contributed for to about 100% and 77% carbon sequestration in Seychelles and
123 Tanzania respectively. Finally, salt marshes were crucial for 75% carbon sequestration in
124 Mozambique (Ghermandi et al., 2019).

125 Besides, mangroves, sea grass, salt marshes, the woodlands play a crucial role in southern
126 Africa (Ryan et al., 2016). Besides sinking a large amount of carbon, the ecosystem service
127 contributed to biogeochemical regulation in the context or after extreme climatic events in
128 southern Africa (Ryan et al., 2016). Additionally, woodlands considerably reduce erosion
129 effects soil after climatic events by keeping the relatively nutrient and organic-rich surface
130 soil in place (Mkanda, 2002; Ryan et al., 2016). The water storage and regulation of the
131 hydrological cycle are attributable to the ecosystem service of woodlands in Southern Africa
132 (Bruijnzeel, 2004; Lopa et al., 2012; Ryan et al., 2016).

133 **3 Methodology: case studies on ecosystems services in Africa**

134 This work used three different methods. It initially consisted of an analysis of the literature,
135 aimed at identifying some key scientific works, which focused on ecosystem services in
136 Africa. Secondly, we deployed a set of case studies, which illustrate some of the pressures
137 that ecosystems are facing. These include:

138 a) Rangelands (Ethiopia, Kenya)
139 b) Mangroves (Benin, Mozambique)
140 c) Forests (Nigeria, Zambia)

141 To systematize data collection, a matrix was developed and used. The data was obtained by
142 the authors' teams of experts, all of whom live or have worked in the corresponding areas,
143 hence having first-hand information and experience with the sites. The case studies provided
144 by the experts were chosen based on the fact that there is sufficient evidence and literature

which supports them. They are also relevant since they provide examples from various African regions. Thirdly and finally, we used the *Driving Force – Pressure – State – Impact – Effect - Response (DPSIER)* (Gabrielsen and Bosch, 2003; Waheed et. al, 2009) model to: identify different pressures caused by human activities on ecosystem services, providing elements that predict the evolution of the problem concerning climate change and response actions carried out by the society, institutions, and governments to reduce or mitigate environmental degradation and expand capacity to adapt to climate change. The results obtained are described and discussed in the next section.

4 Results and Discussion

One of the first findings which characterize the case studies, is that the investigated ecosystems (i.e. rangelands, mangroves and forests) in the sampled countries (Figure 1). suffer from the interaction of multiple stressors, occurring at various levels such as endemic poverty, complex governance, and limited access to capital, including infrastructure and technology; ecosystem degradation; and complex conflicts (Boko et al., 2007). The countries studied already face serious environmental damages, which undermine ecosystem services and lead to land degradation, deforestation, and loss of biodiversity, among others. Such factors worsen vulnerability to climate change in the three ecosystems studied.



Figure 1. Case studies sites and ecosystems

Ecosystem services have on the one hand the potential to reduce vulnerability to climate change and to recover more easily from extreme weather events (Colls et al., 2009), but there are limits as to which what they can perform. The following case studies present an overview of the current situation.

4.1. Impacts of rangelands degradation on ecosystem services in Ethiopia and Kenya

Rangelands are defined as the land on which the potential native vegetation is predominately grasses, grass-like plants, forbs or shrubs. Rangelands produce a great variety of ecosystem services, including the provision of food and fiber, maintenance of biodiversity (conservation), regulating (carbon sequestration), and cultural services (Yahdjian et al., 2015).

However, rangelands are threatened ecosystems and are often undervalued as providers of ecosystem services. Table 1 lists climate-related pressures and impacts on rangelands in Ethiopia and Kenya, which were identified as part of the study.

Table 1. Rangelands in Ethiopia and Kenya

Ecosystem-service provided	Climate-related pressures	Impacts	References
Food (crop, livestock)	Temperature increase, increase in extreme events including droughts and floods, changes in the spatial and temporal variability in rainfall	Increased evapotranspiration and reduced soil moisture, soil degradation, invasive bush encroachment, reduction of crop yield, livestock sickness, and death, reduction in market value, increased food insecurity, increased resource-based conflicts	Shiferaw et al., 2018; Belay et al., 2018; Briske, 2017; Mussa et al., 2016; Lemma, 2011; Opiyo et al., 2015; RCRMD, 2016; Sala et al., 2017; Worden et al., 2009
Pastures	Drought, flood, reduction rainfall amount	Invasive bush encroachment, soil erosion, water depletion, and pollution, siltation of water points, reduction of the load capacity of rangeland, the decline in pasture availability and palatability, reduction of livestock production and productivity, reduce biodiversity, migration of wildlife, the proliferation of climate change sensitive livestock diseases, resource-based conflict, food insecurity	Assefa et al., 1986; Abule, 2009; Fenetahun et al., 2018; Mussa et al., 2016; Kassahun et al., 2008
Biodiversity	Drought, flood	Water pollution, siltation of water points, reduced biomass, the proliferation of climate change sensitive zoonotic diseases, soil erosion, reduced rangeland carrying capacity, the decline in graze availability, woody vegetation the encroachment, water scarcity, encroachment of migratory routes	Opiyo et al., 2015; Worden et al., 2009
Water	Increased extreme events, marked rainfall and, temperature variability, predicted temperature rise	Food insecurity, agrarian-pastoral communities' conflict, increased poverty, the proliferation of diseases	Kiggundu, 2002; Kubbinga, 2012; Malesu et al., 2006; Ngigi, 2003

In Ethiopia, the rangeland covers about 65% of the landmass (Bolo et al., 2019), supports more than two million people, contributes 19% of national DGP, and provides 80% of total annual milk production (Nyariki, 2017; Nyariki and Amwata, 2019). In Kenya, the rangeland covers over 70% of the country. It contributes to the country's GDP through livestock production and the tourism (RCRMD, 2016).

However, both countries' rangeland services are affected by recurrent drought and non-climate stressors. These include human and livestock pressure, overgrazing, erosion of customary resource management, deforestation, bush encroachment, and poor land use

186 policy, a trend which is consistent with what has been documented by the literature (e.g.
187 Fenetahun et al., 2018; Belay et al., 2018; Briske, 2017; Cerretelli et al., 2017; Abdu and
188 Robinson, 2017; Mussa et al., 2016; Hicks et al., 2015; Worden et al., 2009).

189 Rangelands degradation is associated with a decline of the carrying capacity of these
190 ecosystems, a decline of pasture availability and palatability, and high rate of soil erosion.
191 Rangelands degradation also triggers the migration of wildlife and is known to be associated
192 with resource-based conflicts, which have increased over the past decades (Bolo et al., 2019).
193 For instance, the Borana rangeland is affected by bush encroachment which claimed 40% of
194 the rangeland in the 1980s (Assefa et al., 1986). This increased to 63% in the 2000s (Abule,
195 2009). The cumulative effect of rangeland degradation altered the natural ecosystem service
196 thereby worsening poverty and food insecurity especially in the lowlands of Ethiopia
197 (Fenetahun et al., 2018; Mussa et al., 2016; Kassahun et al., 2008).

198 In Kenya, poverty is exacerbated by the harsh environment, poor infrastructure, and low
199 access to basic services. Pastoralism is the predominant source of livelihood accounting for
200 agropastoral and fisherfolks in the rural and peri-urban areas. Pastoralist households largely
201 bear the brunt of negative impacts of extreme climate events like drought, which include
202 increased poverty, water scarcity, resource-based conflicts, disease outbreaks, and food
203 insecurity (Opiyo et al., 2014).

204 The loss of key ecosystem services undermines the ability of the biophysical environment to
205 sustain human livelihoods, hence the need for adaptation efforts. Key areas for adaptation
206 include the livestock sector, water provision and access, biodiversity conservation, and
207 carbon sequestration (Sala et al., 2017).

208 Challenges to adaptation occur due to several socio-economic, political, and ecological
209 factors, affecting opportunities for long-term adaptation strategies to climate change (Opiyo
210 et al., 2015). In northern Kenya, these have been identified to include violent conflicts, lack
211 of affordable credit facilities and financial services, limited access to markets, land tenure
212 changes, and poor infrastructure (Opiyo et al., 2015).

213 In Ethiopia, cross-border and internal conflicts based on ethnic tensions are serious and
214 chronic problems that contributed to the prevalence of poverty and vulnerability to climate
215 change. As a survival strategy, most of the households are highly dependent on natural
216 resources for livestock-keeping, cropping, fishing, beekeeping, and hunting. However,
217 extended droughts and extreme flooding have increased the infestation of livestock diseases,
218 and conflicts. Differences in wealth, power, social values and natural resource bases within
219 the communities also determine their vulnerability to climate change and their capacity to
220 adapt to this also varies in time, and space (Mekonnen and Biruk, 2006; PROLINNOVA,
221 2009, IPCC 2012).

222 **4.2 Impacts of mangrove degradation in Benin and Mozambique**

223 Mangroves are important ecosystems for a wide range of various terrestrial and marine
224 organisms. Mangrove forests accumulate carbon in tree biomass, and most of this carbon is
225 lost by decomposition and export to adjacent ecosystems (Carugati et al., 2018; Igulu et al.,
226 2014). Table 2 presents an outline of the ecosystems services provided by mangroves in
227 Benin and Mozambique.

228 Mangroves' resources provide ecosystem services which influence the well-being and
229 livelihoods of local communities across the tropical and subtropical zones in Africa (Boateng,
230 2018; Chow, 2018). Over the last decades, it is widely reported that mangroves are facing a

high-pressure. For instance, due to both rapid demographic change in coastal mangrove areas and the effects of climate change, West-African mangrove diminished from 20,500 km^2 in 1980 to the current 9000 km^2 (Boateng, 2018; Feka and Morrison, 2017).

Table 2. Mangrove areas in Benin (B) and Mozambique (M)

Ecosystem-service provided	Climate-related pressures	Impacts	References
Physical barrier to floods	(B) Rapid demographic growth in coastal mangroves areas	(B) Increased vulnerability of local communities, mainly salt producers and Peda fishing communities	Boateng, 2018; Feka and Morrison, 2017; Ajonina et al., 2014; Teka et al., 2019
	(M) Floods, storms, and high wind speeds are caused by the tropical cyclones.	(M) Mangrove massive destruction: shifts in species composition; mangrove drowning; pronounced erosion; mangrove mortality	Charrua et al., 2020; Massuanganhe et al., 2015
Biodiversity protection	(B) Over-exploitation of mangroves resources	(B) Degradation of the fragile mangrove ecosystem	Teka et al., 2019
	(M) Prolonged submergence periods due to the combined effect of river flooding and heavy rains associated with the cyclone	(M) Variability in mangrove covers, changes in mangrove structure (tree mortality, canopy gaps, and a great amount of woody debris). Reduced capacity to feeds and protects juvenile fish, prawns, and endangered species, such as the dugong of Mozambique	Macamo et al., 2016; Chevallier, 2013
Food (fish, shrimps)	(B) Increased temperatures, a decrease in rainfall and the rapid sea-level rise	(B) Loss of biodiversity, especially the endemic species (Rhizophora racemosa Meyer and Avicennia germinans (L.) Stearn)	Boateng, 2018; Chow, 2018
	(M) Sea level rise: changes in inundation duration and frequency as well as salinity levels	(M) Reduction in species production, and fishing productivity	Makowski et al., 2018

In Benin, mangroves offer a wide range of goods, and services to communities, including fisheries, fuelwood, salt extraction and non-timber forest products with medicinal properties (Teka et al., 2019). But the over-exploitation of mangroves resources both for domestic use and for infrastructure building results in the degradation of the fragile mangrove ecosystem (Teka et al., 2019). The pressure on this ecosystem affects biodiversity conservation, especially the distribution of *Rhizophora racemosa* Meyer and *Avicennia germinans* (L.) species. It also increases the vulnerability of local communities, mainly salt producers and Peda fishing communities who make their livelihoods from the mangrove ecosystems (Ajonina et al., 2014; Teka et al., 2019).

Mozambique is often affected by natural disasters, including droughts and floods associated with the El Niño Southern Oscillation (USAID, 2012). Flooding, heat waves, cyclones and

247 drought are all getting more frequent and severe as the Earth gets hotter (Coumou and
248 Rahmstorf, 2012). Niquisse et al. (2017) assessed the changes of ecosystem services in
249 Mozambique between 2005 and 2009. The compound effects of climate change resulting
250 from increased temperatures, a decrease in rainfall, and the rapid sea-level rise are likely to
251 pose a significant threat to mangrove species adapted to a specific water salinity level
252 (Boateng, 2018; Chow, 2018). Some mangrove species have been described as highly
253 tolerant to salinity during the sea-level rise, which reduces the susceptibility of mangroves
254 due to salinity variation. This was particularity the case in Sofala province which has the
255 largest area of mangrove distribution and abundance in Mozambique. Mangroves of the
256 Zambezi River Delta are indicated as having a high capacity to store carbon (C) (Stringer et
257 al., 2015). The degradation of mangrove habitats will increase coastal vulnerability,
258 magnifying the effects of climate hazards and erosion, including the loss of lives (Cabral et
259 al., 2017).

260 **4.3 Deforestation and the dangers of climate change in Nigeria and Zambia**

261 The importance of forests to man's survival cannot be overemphasized due to the various
262 environmental, socia,l and economic benefits they provide. The wooded land areas have
263 decreased during the past two decades in Africa. Nigeria is among the five countries with the
264 biggest annual loss of (-3.7%) (Chakravarty et al., 2012) and Zambia has fast disappearing
265 forests (276, 000 hectares/annum) (Kalaba et al., 2013a).

266 With this trend, the potential of key ecosystem services (cycling of water, oxygen, carbon,
267 and nitrogen; stability to the hydrological system; supply of freshwater, prevention of flood
268 and soil erosion) is at risk (Oregon Forest Resources Institute, 2018; Adebayo, 2009). Table 3
269 presents the impacts on ecosystem services in forest areas in Nigeria and Zambia.

Table 3. Ecosystem-service provided climate-related pressures and impacts in forests in Nigeria (N) and Zambia (Z)

Ecosystem-service provided	Climate-related pressures	Impacts	References
Habitat for fauna and flora	(N) (Z) Drought, temperature rise, desertification	Washing away habitat for wildlife, loss of biodiversity, reduction in fish yields, food insecurity	Adeleke et al., 2019; Akande, 2017; Emeodilichi, 2018; Burivalova et al., 2015; Lembani et al., 2019; Kalaba et al., 2013; Ng'onga et al., 2019
Source of timber	(N) Land degradation, drought, increased desertification, soil erosion	Soil depletion, widespread flooding, loss of human settlements	Akande, 2017; Elisha et al., 2017; Ebele and Emodi, 2016; Olaniyi et al., 2013
Physical protection against extreme events	(N) (Z) Droughts, floods: landslides and bushfires	Soil erosion; loss of biodiversity and freshwater resources	Amanchukwu et al., 2015; Enete, 2014; BNRCC, 2011; Fashae and Onafeso, 2011
Balancing of hydrological regime	(N) Drought, sea-level rise, seawater temperature rise	Changes to ocean salinity, disturbance of marine life	Ifeanyi-obi et al., 2012; FAO, 2007; Nicholls and Cazenave, 2010

Despite the numerous advantages of the forest to man, fauna, and flora, a changing climate may increase and worsen the threats to forest and forest products. This in turn would intensify incidences of severe floods, drought, forest fires, landslides, temperature rise, rainfall variability, as well as insect and disease outbreaks (Adeleke et.al., 2019).

Nigeria is naturally endowed with a vast expanse of forest land the swamp forests in the extreme Southern part of the country, the tropical rainforest in the South-western axis, and the wooded savannah in the middle belt. Forests represent about 12% the total vegetation cover of the country and the forest resources account for 2.5% of the country's GDP (FAO, 2010).

In terms of Zambia, the majority of population depends on forest products for their livelihoods. Forest provisioning ecosystem services in the Miombo woodland, for instance, are an important adaptation strategy to reduce the impacts and to help communities cope with stresses and shocks (Kalaba et al., 2013a).

Habitats for fauna and flora are currently threatened by a reduction in forest cover driven by anthropogenic activities exacerbated by climate change. In Zambia the Miombo woodlands

288 are habitat for fauna and flora and are sources of timber. Besides carbon sequestration and
289 regulation of watersheds, woodlands play a crucial role in preventing extreme weather events
290 (Kalaba et al., 2013).

291 In freshwater ecosystems such as Lake Bangweulu, there is depletion of fish for fisheries-
292 based economy which may be attributed to the increase pressure of humans and increasing
293 temperatures (Ng'onga et al., 2019). Increased vulnerability of the rural people to climate
294 change further weakens their already low adaptive capacity. Fisheries-based livelihood
295 provides a full range of activities that involve the harvesting of fish, through the intermediary
296 phases of landing, processing, transporting, and delivery to final consumers. Temperature
297 increases have implications on fish yields.

298 **4.4 Conceptual models and ecosystem services**

299 Conceptual models summarize, produce a visualization of, and explain both actual or
300 predicted situations, as well as point out how they could be addressed, determining their
301 causes and consequences. In the last few years, the model *Driving Force – Pressure – State –*
302 *Impact – Effect - Response (DPSIER)* (Gabrielsen and Bosch, 2003; Waheed et. al, 2009)
303 have been crucial to the conceptualization of ecosystem issues, and to translate them to
304 shareholders, environmental managers and researchers (EC, 1999; Smith et al., 2016).

305 It is possible to observe how these issues are connected when we highlight cause-effect
306 relationships, although the product is a simplistic representation, given that not only it
307 suggests a linear relationship between human activities, the environment, and health, which,
308 generally is not true, but also hides underlying complexities in such interactions (OECD,
309 1993), which could be linked, and take place simultaneously (Gari et al., 2015). Moreover, it
310 does not stress differences in the nature, severeness, and time scales of changes in the state
311 concerning the intensity, frequency and duration of the pressure (Smith et al., 2016).

312 Although a single DPSIER cycle (Figures 2, 3 and 4) excessively simplify the "real world", it
313 could provide us with a conceptualization of the relations between environmental changes,
314 anthropic pressures and their effects (Svarstad et al., 2008; Smith et al., 2016) in the cases
315 studied.

316 **4.4.1 Identification of problems in rangelands areas, mangroves and forest**

317 Multiple environmental factors affect the health of the population and its ability to adapt to
318 climate change, thus forming complex cause and effect relations. When people are exposed to
319 a change in their environment, their health might be affected, let alone other socioeconomic
320 and cultural impacts possibly caused by the reduction or the loss of a few ecosystem services.
321 The review of the literature on pressures that rangelands, mangroves, and forests are facing
322 showed a vast range of reasons for changes in ecosystem services (Figures 2-4).

323 **Figure 2. Factors related to the degradation of rangelands areas in Ethiopia and Kenya**



324

We analyzed food production in the context of climate change, food insecurity and violence as an important example of the complexity of interactions in ecosystem services in rangeland areas in Ethiopia and Kenya.

We live in a global agri-food system guided by commercial interests and aimed at the consumption of ultra-processed items. Together with a lack of will of political leaders and a lack of effective action of the society as a whole, this led us into the current global syndemic – a synergic combination of the pandemics of obesity, malnutrition and climate change, which take place concurrently and have shared causes and effects, which feed into one another (Swinburn et al., 2019).

An example of that is what has been going on in Eastern Africa (Ethiopia, Kenya, Somalia, Tanzania and Uganda), which, for three consecutive years has been recording less than half the usual levels of seasonal precipitation, causing families to lose their crops and rangeland. The high cattle mortality rate, the low production of milk, the drop in market value of the animals who survive such adverse conditions, the lower agricultural production and the inflation in grain prices have increased hunger for approximately 20% of the population, and it is estimated that a million children under the age of five require treatment for acute malnutrition (FAO et al., 2018).

On the other hand, African obesity indexes have been growing faster than that of any other region in the world and causing a public health crisis. Eight of the twenty countries with the most drastic adult obesity rates are in Africa. The expanding economy and the presence of multinational companies in the food sector, which noticed the promising consumer market of developing countries, resulted in this high adult obesity prevalence (Jaffar and Gill, 2017; Gettleman, 2018; Martins, 2018). Factors such as a lack of access to nutritious foods and physiological adaptations to food deprivation explain how people that were subject to food

insecurity have an increased risk of gaining weight and a tendency towards obesity (FAO et al., 2018).

Climate variability, which affects rain patterns and agricultural seasonality, as well as extreme climate events (such as droughts and floods) are among the greatest drivers of food insecurity, together with wars and economic crises (FAO et al., 2018).

Water stress and the spread of climate change-related plagues – such as the swarm of locusts that emerged in Kenya, Ethiopia and Somalia in 2019 (FAO, 2020) – are devastating to these countries in which agriculture and cattle are crucial means of living, thus aggravating poverty because of a shortage of resources and income that can assure people sustainable means, and also due to the increase of hunger levels, malnutrition, the limited access to basic services, social exclusion and violence. In this case, social vulnerability is also linked to flooding events, which, in turn, generate even more poverty because of material loss and submit the population to an even greater risk of disease and death by water-borne diseases or other infectious diseases.

Figure 3. Factors related to mangrove degradation in Benin and Mozambique



365 The ability to recycle and retain nutrients is what makes mangroves an important ecosystem
366 for the survival of thousands of species and for the climate adaption and extreme weather
367 events. To understand and model the dynamic interaction between climate change-related
368 risks, ecosystem services, means of living and adaptation to climate change in coastal areas,
369 we have evaluated the context of the humanitarian crisis produced by natural disasters.

370 Mozambique and Benin are highly socially vulnerable to climate disasters. Besides human
371 loss, this type of disaster deeply affects the economy of these countries, which, in turn, have
372 poor indexes such as basic sanitation, drinking water, food safety and others concerning
373 income and literacy (UNISDR, 2018). On the other hand, these countries are developing very
374 rapidly because they are exploring natural resources such as wood, coal, gas and other
375 minerals. However, poor wealth and resource distribution continues to worsen poverty and
376 social inequality, as well as generate generalized pollution and the destruction of habitats,
377 including mangroves (Chevallier, 2013).

378 Therefore, the degradation of mangroves is a product of human activities related to survival,
379 the exploration of natural resources and urbanization, which impacts the environment (loss of
380 biodiversity, pollution, water shortage, increased salinity in the water, CO_2 release, soil
381 contamination.) The degradation of these ecosystems also causes a considerable
382 socioeconomic impact, increasing poverty, unemployment, conflicts, nutritional diseases,
383 water-borne diseases, vector-borne diseases, etc. Investing in disaster risk reduction strategies
384 based on ecosystem services – as well as in the adaptation to climate change to reduce the
385 social, environmental and economic impact of natural disasters – is fundamental.
386 Environmental management and the impact of climate change are interrelated, and the
387 management of their risks demand ecosystem approaches.

388 **Figure 4. Factors related to deforestation in the forest of Nigeria and Zambia**



389

390 An important example of the complexity of interactions between environmental damage,
391 ecosystem services and adaptation to climate change is the extraction of natural resources and
392 the march of agribusiness and anthropic activities threatening the forests of Nigeria and
393 Zambia. Deforestation in these countries takes place fundamentally for three reasons:
394 producing charcoal, their main source of fuel at national level, which is also exported to
395 countries in the European Community; exporting wood to China; and opening new
396 agricultural land, since mining is no longer profitable and the population started growing corn
397 to make ends meet (Chaves et al., 2020).

398 The population of Nigeria is growing at astounding rates, which drives a huge demand for
399 agricultural land, firewood, water and housing. Besides the effects of deforestation on climate
400 change, the excessive use of fertilizers and pesticides degrades the soil and pollutes the water.
401 The impact includes food safety, water scarcity, deforestation, desertification, forest fires,
402 loss of biodiversity, air and land pollution, rising sea levels and an excess of greenhouse gas
403 emissions.

The accelerated use of ecosystem services that are crucial to life (water, food, energy, medications, genetic resources among others) also drives climate change, which, in turn, affects food production, threaten livelihoods, increases poverty and is associated with the numbers of environmental refugees, especially in Africa (Leal Filho et al 2021). This social vulnerability increases the risk of nutritional diseases, water-borne diseases, vector-borne diseases and respiratory problems.

An increase in the risk of malaria, for instance, is pushed by the export of raw materials for rich countries. Chaves et al. (2020) showed that the global consumption and international trade of commodities related to deforestation in tropical countries – such as wood and agricultural products like tobacco, cocoa, coffee and cotton – increase the risk of contracting malaria in Africa, Southeast Asia, India and South America by 20%. The study includes a number of examples from developing countries where the exportation of commodities for the developed world is related to deforestation and an increase in malaria cases.

As far as adaptation approaches focusing on the protection of ecosystems services is concerned, a framework is necessary, and which may take into account four main components:

a) The adequacy of the adaptation initiatives to be pursued

b) The extent to which a particular ecosystem or set of ecosystems will benefit

c) The long-term prospects of the services to be protected

As Figure 5 shows, there are some means which may be deployed in order to maximise the contribution of ecosystems services to support climate change adaptation efforts in Africa.

Figure 5- Optimising the contribution of ecosystems services to climate change adaptation efforts.

But in order to yield the expected benefits, attempts to achieve such an optimization should not only be technically sound, but also financially viable, socially just and ethically acceptable. This is so in order to make sure that the many socio-economic problems which accompany climate change (e.g. inequalities, social justice, poverty) may also be addressed, also increasing the resilience of local populations, and helping them to better adapt to the manifold impacts of a changing climate.

5 Conclusions

Livelihoods and well-being are directly dependent on the presence of healthy ecosystems. Sustainable ecosystems may help to prevent the impact of extreme events and facilitate recovery processes. They may also assist people to adjust, reduce their vulnerability, and increase their resilience. However, as this paper has shown, many ecosystems are being

440 threatened by climate change and extreme weather events, which show that urgent action is
441 needed.

442 The diversity of causal factors confirms the complexity of the factors leading to the
443 degradation of the ecosystems studied. Previous works have documented the link between
444 the degradation of ecosystems and poverty. Echeverria (2000) for instance, relates poverty to
445 a trap (poverty-trap thesis), in which individuals, to survive, degrade the environment, and
446 that degradation, in turn, worsens poverty in a vicious cycle (Prakash, 1997; Stephens et al.,
447 2012). Meadows et al. (1973) connected the exacerbated consumption of natural resources to
448 economic and population growth, which, in turn, may lead to the depletion of resources and
449 an increase in social inequality (CMMAD, 1987; Drexhage and Murphy, 2010). In this
450 context, depressed economies lead to migration, food insecurity, exposure to the risk of
451 disasters, health risks and potential escalation of deeply-rooted socio-political tensions
452 (CEPAL, 2020).

453 The trends in Africa have been exacerbated by the COVID-19 pandemic, which has
454 unleashed the deepest global recession in the continent since the 1930s, leading to increased
455 poverty, food insecurity, unemployment, violence against women, inequality and
456 vulnerabilities to diseases (Sachs et al., 2020) and has been linked to environmental impact,
457 habitat destruction and indiscriminate exploration of natural (Afelt et al., 2018; Giatti et al.,
458 2020; Andersen et al., 2020).

459 Apart from social problems, the continuous degradation of African ecosystems also limits
460 their ability to respond to extreme events (e.g. droughts, floods), and to recover from them.
461 Thus there is an urgent need to address the threats posed to ecosystems, to stem the steady
462 loss of ecosystem services. To achieve this, some measures are needed. For instance:

a) a better monitoring of those ecosystem services which are being and likely to continue to be influenced by climate change, so that the steps to reduce these impacts are taken. This is especially needed in respect of aquatic ecosystems -especially rivers and lakes- which are currently under a double pressure: excessive use of the water resources and irregular rainfall due to changing climate conditions;

b) an improvement in the means to determine the climate sensitivity of ecosystem services, to avoid them reaching a tipping point. Here, aquatic ecosystems on the one hand, but terrestrial resources (e.g. forests) on the other, known to be already influenced by climate change need to be better understood, since there are signs of decline;

c) more efforts to optimize ecosystem services in a way that the resources and commodities deriving from them serve as an encouragement to actively pursue their protection. This is the case, for instance, of fishery in a lake or forests resources. If carefully planned, these resources and commodities may be available for a long period of time, avoiding the risks a sudden exhaustion could bring about;

d) to design, test and implement adaptive responses that are cost-effective, to upkeep the productivity of the ecosystems on the one hand, but without risk damaging them on the other. Here, the prevention of potential damages to ecosystems is an important component, so as to allow them to continue to function. Here, traditional knowledge may play a key role, since it is based on proven practices known to be successful.

This paper has some limitations. For instance, it has not covered all African countries and regions. Secondly, the fact that it aimed at providing an overview of the trends related to ecosystems depletion, means that it did not undertake an in-depth analysis of the many variables which characterize their vulnerability, nor it has looked at specific adaptation strategies. The paper is nonetheless a welcome addition to the literature, in the sense that it

487 offers a unique assessment of the extent to which large African ecosystems are being
488 negatively influenced by climate change. It also describes some of the measures which may
489 be deployed, so as to reduce the vulnerability of ecosystems and allow them to provide their
490 important contributions to climate change mitigation and adaptation efforts.

491 The drive towards protecting -or restoring- ecosystem services in an African context needs a
492 joint effort of all types of stakeholders. This includes government organizations, enterprises
493 and other representatives from the private sector on the one hand, but also the civil society
494 (e.g. citizens groups, non-governmental organizations, regional agencies) on the other, so as
495 to address the main constraints seen. Especially at the local level.

496 Moving forward, apart from the measures listed above, it is also important to assess how
497 impacts related to ecosystem services provision may affect the local populations, so that
498 suitable adaptation measures may be implemented, hence avoiding a situation where these
499 impacts may become irreversible.

500

References

Abdu N.H., Robinson L.W. (2017). Community-based rangeland management in Dirre rangeland unit: Taking Successes in Land Restoration to scale project. ILRI project report. Nairobi, Kenya: *International Livestock Research Institute (ILRI)*. https://hdl.handle.net/10568/89714

Abramovitz J.N., Banuri T., Girot P., Orlando B., Schneider N., Spanger-Siegfried E., Switzer J., Hammill A. (2002). Adapting to Climate Change: Natural Resource Management and Vulnerability Reduction. Background Paper to the Task Force on Climate Change, Adaptation and Vulnerable Communities. IISD – *International Institute for Sustainable Development*, Geneva, Switzerland.

Abule E. (2009). Rangeland suitability evaluation for livestock production in Laga-Wata/Laga Sura, Dawa and rift valley basins of the Borana Zone of Oromia, Southern Ethiopia Pp 441.

Adebayo A.A. (2009). Attaining sustainable rural development in Nigeria through effective management of forest based resources. *International Journal of Creativity and Technical Development* Vol. 1(3): 32-45. Available in: https://www.icidr.org/doc/ICIDR%20PDF%20contents/journal%20of%20research%20in%20education%20and%20society/IJCTD_v1no1to3/attaining%20sustainable%20rural.pdf

Adeleke M.L., Oluwatosin O.J., Fagbenro O.A., Amos T.T., Ajibefun I.A. (2019). Biodiversity, Ecosystem Degradation, and Climate Change Effects on Livelihoods in the Bitumen Area of Nigeria. *Springer Nature* Switzerland AG 2019 W. Leal Filho (ed.), Handbook of Climate Change Resilience. https://doi.org/10.1007/978-3-319-71025-9125-1

525 Andersen K.G., Rambaut A., Lipkin W.I, Holmes E.C., Garry R.F. (2020). The proximal
526 origin of SARS-CoV-2. *Nature Medicine* 26, 450–452. https://doi.org/10.1038/s41591-
527 020-0820-9

528 Afelt A., Frutos R., Devaux C. (2018). Bats, coronaviruses and deforestation: towards the
529 emergence of novel infectious diseases? *Front. Microbiol.*9:702. Doi:
530 10.3389/fmicb.2018.00702

531 Ajonina G.N., Ago E.E., Amoussou G., Mibog E.D., Akambi I.D., Dossa E. (2014).
532 Carbon budget as a tool for assessing mangrove forests degradation in the western,
533 coastal wetlands complex (Ramsar Site 1017) of Southern Benin, West Africa. The
534 Land/Ocean Interactions in the Coastal Zone of West and Central Africa. Springer, pp.
535 139-149. Doi: 10.1007/978-3-319-06388-1_12

536 Akande A. (2017). Geospatial Analysis of Extreme Weather Events in Nigeria (1985–
537 2015) using Self-Organizing Maps. Advances in Meteorology.
538 https://doi.org/10.1155/2017/8576150

539 Amanchukwu R.N., Amadi-Ali T.G., Ololube N.P. (2015). Climate change education in
540 Nigeria: The role of curriculum review. *Education*,5(3):71-79. Doi:
541 10.5923/j.edu.20150503.01

542 Assefa E., Bille J.C., Corra M. (1986). Ecological map of Southern Sidamo. JEPSS (Joint
543 Ethiopian Pastoral System Study). Research report. ILCA (International Livestock Centre
544 for Africa), Addis Ababa, Ethiopia. 44pp [ILCA library accession no. S9300188]

545 Belay Z., Bikila W., Demel T., Muluneh W. (2018). Rangeland Degradation and
546 Rehabilitation Efforts in the Somali National Regional State, Eastern Ethiopia: A Review.
547 *International Journal of Innovative Research & Development*, 7, 5, 1-18. Doi:
548 10.24940/ijird/2018/v7/i5/MAY18026

549 BNRCC - Building Nigeria's Response to Climate Change. (2011). National adaptation
550 strategy and plan of action on climate change for Nigeria (NASPA-CCN). Prepared for
551 the Federal Ministry of Environment Special Climate Change Unit. Available in:
552 http://csdevnet.org/wp-content/uploads/NATIONAL-ADAPTATION-STRATEGY-
553 AND-PLAN-OF-ACTION.pdf

554 Boateng I. (2018). An assessment of vulnerability and adaptation of coastal mangroves of
555 West Africa in the face of climate change. Threats to mangrove forests. Springer, pp. 141-
556 154. doi:10.1007/978-3-319-73016-5_7

557 Boko M., Niang I., Nyong A., Vogel C., Githeko A., Medany M., Osman-Elasha B., Tabo
558 R.. Yanda P. (2007). Africa. Climate Change 2007: Impacts, Adaptation and
559 Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the
560 Intergovernmental Panel on Climate Change (IPCC Report), Parry M.L., Canziani O.F.,
561 Palutikof J.P., van der Linden P.J., Hanson C.E., Eds., Cambridge University Press,
562 Cambridge UK, 433-467

563 Bolo P., Sommer R., Kihara J., Kinyua M., Nyawira S., Notenbaert A. (2019). Rangeland
564 degradation: Causes, consequences, monitoring techniques and remedies. Working Paper.
565 CIAT Publication No. 478. *International Center for Tropical Agriculture* (CIAT).
566 Nairobi, Kenya. Available in: https://hdl.handle.net/10568/102393

567 Bourne A., Holness S., Holden P., Scorgie S., Donatti C.I., Midgley G. (2016). A socio-
568 ecological approach for identifying and contextualising spatial ecosystem-based
569 adaptation priorities at the sub-national level. *PLoS One*, 11 (5).
570 https://doi.org/10.1371/journal.pone.0155235

571 Brander L.M., Wagtendonk A.J., Hussain S.S., McVittie A., Verburg P.H., de Groot R.S.,
572 van der Ploeg S. (2012). Ecosystem service values for mangroves in Southeast Asia: a

573 meta-analysis and value transfer application. *Ecosystem Services*., 1, 62-69.
574 https://doi.org/10.1016/j.ecoser.2012.06.003

575 Briske D. eds. (2017). Rangeland Systems Processes, Management and Challenges.
576 Springer, Cham, Switzerland. Doi: 10.1007/978-3-319-46709-2

577 Lemma B. (2011). The impact of climate change and population increase on Lakes
578 Haramaya and Hora-Kilole, Ethiopia (1986 – 2006). In: Lemma B., Getahun A. (eds).
579 Impacts of climate change and population on tropical aquatic resources, Proceedings of
580 the Third International Conference of the Ethiopian Fisheries and Aquatic Sciences
581 Association (EFASA), *AAU Printing Press*, Addis Ababa.

582 Bruijnzeel L.A. (2004). Hydrological functions of tropical forests: not seeing the soil for
583 the trees? *Agric. Ecosyst. Environ.* 104, 185–228.
584 https://doi.org/10.1016/j.agee.2004.01.015

585 Burivalova Z., Lee T.M, Giam X., Sekercioglu C.H., Wilcove D.S., Koh L.P. (2015).
586 Avian responses to selective logging shaped by species traits and logging practices.
587 *Proc.R.soc.lond.B*, 282, p. 20150164. https://doi.org/10.1098/rspb.2015.0164

588 Byers B.A., Cunliffe R.N., Hudak A.T. (2001). Linking the conservation of culture and
589 nature: a case study of sacred forests in Zimbabwe. *Human Ecology* 29, 187–218.
590 https://doi.org/10.1023/A:1011012014240

591 Cabral P., Augusto G., Akande A., Costa A., Amade N., Niquisse S., Atumane A., Cuna
592 A., Kazemi K., Mlucasse R., Santha R. (2017). Assessing Mozambique's exposure to
593 coastal climate hazards and erosion. International Journal of Disaster Risk Reduction, Vol
594 23, Pages 45-52, ISSN 2212-4209. https://doi.org/10.1016/j.ijdrr.2017.04.002

595 Carugati L., Gatto B., Rastelli E., Lo Martire M., Coral C., Greco S., Danovaro R. (2018).
596 Impact of mangrove forests degradation on biodiversity and ecosystem functioning.
597 Scientific Reports. Doi:10.1038/s41598-018-31683-0

598 CBD - Convention on Biological Diversity. (2009). Connecting Biodiversity and Climate
599 Change: Mitigation and Adaptation. Report of the Second Ad Hoc Technical Expert
600 Group on Biodiversity and Climate Change. *Technical Series* no. 41. Montreal, Canada:
601 Convention on Biological Diversity. ISBN: 92-9225-134-1

602 CEPAL – Comissão Econômica para a América Latina e Caribe. (2020). Economic
603 Survey of Latin America and the Caribbean 2020: Main conditioning factors of fiscal and
604 monetary policies in the post-COVID-19 era. *ECLAC* - Washington, D.C.
605 ISBN: 9789211220513

606 Cerretelli S., Poggio L., Gimona A., Peressotti A., Black H. (2017). The loss of
607 ecosystem services due to land degradation. Integration of mechanistic and probabilistic
608 models in an Ethiopian case study. Washington: *Journal of Geophysical Research*, 19,
609 8142. Doi: 10.13140/RG.2.2.20172.31361

610 Chaikumbung M., Doucouliagos H., Scarborough H. (2016). The economic value of
611 wetlands in developing countries: a meta-regression analysis. *Ecol. Econ*., 124,164-174.
612 ISSN 0921-8009. Doi: 10.1016/j.ecolecon.2016.01.022

613 Chakravarty S., Ghosh S.K., Suresh C.P., Dey A.N, Shukla. G. (2012). Deforestation:
614 Causes, Effects and Control Strategies.; In: Okia A. (Ed) Global Perspectives on
615 Sustainable Forest Management. UK: *IntechOpen*. Doi:10.5772/33342

616 Charrua A.B., Bandeira S.O., Catarino S., Cabral P., Romeiras, M.M. (2020). Assessment
617 of the vulnerability of coastal mangrove ecosystems in Mozambique. *Ocean & Coastal*
618 *Management*, 189, 105145. Doi: 10.1016/j.ocecoaman.2020.105145

619 Chaves L.S.M., Fry J., Malik A., Geschke A., Sallum M.A.M., Lenzen M. (2020). Global
620 consumption and international trade in deforestation-associated commodities could
621 influence malaria risk. *Nat Commun* 11, 1258. https://doi.org/10.1038/s41467-020-14954-
622 1

623 Chevallier R. (2013). Balancing Development and Coastal Conservation: Mangroves in
624 Mozambique. SAIIA Research Report 14. Governance of Africa's Resources Programme.
625 *SAIIA – South African Institute of International Issues*. ISBN: 978-1-919969-83-1

626 Chiabai A., Quiroga S., Martinez-Juarez P., Higgins S., Taylor T. (2018). The nexus
627 between climate change, ecosystem services and human health: Towards a conceptual
628 framework. *Science of The Total Environment*, 635, 1191-1204.
629 https://doi.org/10.1016/j.scitotenv.2018.03.323

630 Chong J. (2014). Ecosystem-based approaches to climate change adaptation: progress and
631 challenges. *Int Environ Agreements*. 14 391–405. Doi: 10.1007/s10784-014-9242-9

632 Chow J. (2018). Mangrove management for climate change adaptation and sustainable
633 development in coastal zones. *Journal of Sustainable Forestry.* 37:2, 139-156.
634 Doi: 10.1080/10549811.2017.1339615

635 Colls A., Ash N., Ikkala N. (2009). Ecosystem-based Adaptation: a natural response to
636 climate change. Gland, Switzerland: *IUCN*. 16pp. ISBN 978-2-8317-1215-4

637 CMMAD - Comisión Mundial del Medio Ambiente y del Desarroll. (1987). Nuestro
638 futuro común. Madri: *Alianza Editorial*. ISBN 978-84-206-9574-7

639 Coumou D., Rahmstorf S. (2012). A decade of weather extremes. *Nature Clim*
640 *Change* 2, 491–496. https://doi.org/10.1038/nclimate1452

641 Doswald N., Munroe R., Roe D., Giuliani A., Castelli I., Stephens J., Moller I., Spencer
642 T., Vira B., Reid H. (2014). Effectiveness of ecosystem-based approaches for adaptation:
643 review of the evidence-base. *Climate and Development,* 6:2 185-201,
644 https://doi.org/10.1080/17565529.2013.867247

645 Drexhage J., Murphy D. (2010). Sustainable Development: From Brundtland to Rio 2012.
646 Background Paper prepared for consideration by the High Level Panel on Global
647 Sustainability at its first meeting. New York: *United Nations Headquarters*.

648 Ebele N.E., Emodi N.V. (2016). Climate change and its impacts in the Nigerian economy.
649 *Journal of Scientific Research & Reports*, 10(6), 1-13. Doi: 10.9734/JSRR/2016/25162

650 EC – European Commission. (1999). Towards Environmental Pressure Indicators for the
651 EU. 1st Edition. Luxembourg: Office for Official Publications of the European
652 Communities. ISBN 92-828-4978-3

653 Echeverria R.G. (2000). Opciones para reducir la pobreza rural en América Latina y el
654 Caribe. Santiago: *Revista de la CEPAL 70*. ISBN: 9213215797

655 Elisha I., Sawa B.A., Lawrence E. (2017). Evidence of climate change and adaptation
656 strategies among grain farmers in Sokoto State, Nigeria. New York: *IOSR Journal of*
657 *Environmental Science, Toxicology and Food Technology (IOSR-JESTFT),* 11(3), 1-7.
658 Doi: 10.9790/2402-1103020107

659 Emeodilichi H.M. (2018). Assessment of Environmental impact of deforestation in
660 Enugu, Nigeria. *Resources and Environment*, vol. 8:4 207-215.
661 Doi:10.5923/j.re.20180804.03

662 Enete I.C. (2014). Impacts of climate change on agricultural production in Enugu State,
663 Nigeria. *Earth Sci Clin change* 5:234. Doi: 10.4172/2157-7617.1000234

664 FAO - Food Agricultural Organization of the United Nations. (2007). Adaptation to
665 climate change in agriculture, forestry and fisheries: Perspective, framework and
666 priorities. Rome: *Food and Agriculture Organization of the United Nations*. Available in:
667 http://www.fao.org/3/a-au030e.pdf

668 FAO - Food Agricultural Organization of the United Nations (2010). Global Forest
669 Resources Assessment 2010 (FRA 2010): Main Report. FAO Forestry Paper no. 163.
670 Available in: http://www.fao.org/3/i1757e/i1757e00.htm

671 FAO, IFAD, UNICEF, WFP and WHO. (2018). The State of Food Security and Nutrition
672 in the World 2018. Building climate resilience for food security and nutrition. Rome:
673 FAO. Licence: CC BY-NC-SA 3.0 IGO. Available in:
674 http://www.fao.org/3/I9553EN/i9553en.pdf

675 FAO - Food Agricultural Organization of the United Nations. (2020). Desert locust crisis.
676 FAO in emergencies. Available in:
677 http://www.fao.org/emergencies/crisis/desertlocust/intro/en/

678 Fashae O.A., Onafeso O.D. (2011). Impact of climate change on sea level rise in Lagos,
679 Nigeria. *International Journal of Remote Sensing*, 32(24), 9811-9819.
680 Doi:10.1080/01431161.2011.581709

681 Fedele G., Donatti C.I., Harvey C.A., Hannah L., Hole D.G. (2019). Transformative
682 adaptation to climate change for sustainable social-ecological systems. *Environmental*
683 *Science & Policy*. 101 116-125. Doi: 10.1016/j.envsci.2019.07.001

684 Feka Z.N., Morrison I. (2017). Managing mangroves for coastal ecosystems change: a
685 decade and beyond of conservation experiences and lessons for and from west-central
686 Africa. *Journal of Ecology and The Natural Environment* 9, 99-123.
687 https://doi.org/10.5897/JENE2017.0636

688 Fenetahun Y., Xu X., Wang Y. (2018). Assessment of Range Land Degradation, Major
689 Causes, Impacts, and Alternative Rehabilitation Techniques in Yabello Rangelands
690 Southern Ethiopia. *Preprints*, 2018070198. Doi: 10.20944/preprints201807.0198.v1

691 Gabrielsen P., Bosch P. (2003). Environmental Indicators: Typology and use in
692 Reporting. Copenhagen: *EEA - European Environmental Agency.* Internal Working
693 Paper, pp 1-20.

694 Gari S.R., Newton A., Icely J.D. (2015). A review of the application and evolution of the
695 DPSIR framework with an emphasis on coastal social-ecological systems. *Ocean Coast.*
696 *Manage*. 103, 63–77. Doi: 10.1016/j.ocecoaman.2014.11.013

697 Gettleman, J. (2018). In Kenya, and across Africa, un unexpected epidemic: Obesity. The
698 New York Times. Serial. Available in:
699 https://www.nytimes.com/2018/01/27/world/africa/kenya-obesity-diabetes.html

700 Ghermandi A., Obura D., Knudsen C., Nunes P.A.L.D. (2019). Marine ecosystem
701 services in the Northern Mozambique Channel: A geospatial and socio-economic analysis
702 for policy support. *Ecosystem Services*, 35, 1–12. Doi: 10.1016/j.ecoser.2018.10.009

703 Giatti L.L., da Paz M.G.A., Toledo R.F., Lauda-Rodriguez Z.L., Milz B. (2020). Editorial
704 - A emergência do coronavírus e os desafios socioambientais contemporâneos. São Paulo:
705 *Ambiente & Sociedade*, v. 23, p. 1-7, 2. https://doi.org/10.1590/1809-
706 4422asoceditorialvu2020l4ed

707 Green J.M.H., Larrosa C., Burgess N.D., Balmford A., Johnston A., Mbilinyi B.P., Coad
708 L. (2013). Deforestation in an African biodiversity hotspot: extent, variation and the
709 effectiveness of protected areas. *Biol. Conserv.*, 164, pp. 62-72.
710 https://doi.org/10.1016/j.biocon.2013.04.016

711 Gupta, A.K., Singh, S., Wajih, S.A., Mani, N., Singh, A.K. (2017). Urban Resilience and
712 Sustainability through Peri-Urban Ecosystems: Integrating Climate Change Adaptation
713 and Disaster Risk Reduction. Gorakhpur Environmental Action Group, *Gorakhpur* (U.P.)
714 India. ISBN: 9780124202184.

715 Hannah L. (2015). Ecosystem Services. In: Climate Change Biology (Second Edition),
716 Academic Press, 281-300. ISBN: 9780124202184

717 Hicks C.C., Cinner J.E., Stoeckl N., McClanahan T.R. (2015). Linking ecosystem
718 services and human-values theory: Ecosystem Services and Human Values. *Conservation*
719 *Biology*, 29(5), 1471–1480. https://doi.org/10.1111/cobi.12550

720 Holland R.A., Darwall W.R.T., Smith K.G. (2012). Conservation priorities for freshwater
721 biodiversity: the key biodiversity area approach refined and tested for continental Africa.
722 *Biol. Conserv.*, 148 (1), pp. 167-179. https://doi.org/10.1016/j.biocon.2012.01.016

723 Ifeanyi-obi C.C., Etuk U.R., Jike-wai O. (2012). Climate Change, Effects and Adaptation
724 Strategies: Implication for agricultural extension system in Nigeria. *Greener Journal of*
725 *Agricultural Sciences*, 2 (2): 053 -060.

726 Igulu M.M., Nagelkerken I., Dorenbosch M., Grol M.G.G., Harborne A.R., Kimirei I.A.,
727 Mumby P.J., Olds A.D., Mgaya Y.D. (2014). Mangrove Habitat Use by Juvenile Reef
728 Fish: Meta-Analysis Reveals that Tidal Regime Matters More than Biogeographic Region. *PLoS*
729 *ONE* 9(12), e114715. https://doi.org/10.1371/journal.pone.0114715

730 IPBES – Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem
731 Services. (2018). The regional assessment report on biodiversity and ecosystem services
732 for Africa. Archer E., Dziba L., Mulongoy K.J., Maoela M.A., Walters, M. (eds.). Bonn:
733 Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity and
734 Ecosystem Services. ISBN: 978-3-947851-05-8

735 IPCC (2012). Summary for Policymakers. In: Managing the Risks of Extreme Events and
736 Disasters to Advance Climate Change Adaptation [Field C.B., Barros V., Stocker T.F.,
737 Qin D., Dokken D.J., Ebi K.L., Mastrandrea M.D., Mach K.J., Plattner G.-K., Allen S.K.,
738 Tignor M., and Midgley P.M. (eds.)]. A Special Report of Working Groups I and II of the
739 Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge,
740 UK, and New York, NY, USA, pp. 3-21. Doi: 10.13140/2.1.3117.9529

741 Jaffar S., Gill G. (2017). The crisis of diabetes in sub-Saharan Africa. *The Lancet*
742 *Diabetes & Endocrinology*, Volume 5, Issue 8, 574 – 575. http://dx.doi.org/10.1016/
743 S2213-8587(17)30219-X

744 Jones H.P., Hole D.G., Zavaleta E.S. (2012). Harnessing nature to help people adapt to
745 climate change. *Nat. Clim. Chang.*, 2, 504-509. Doi: 10.1038/NCLIMATE1463

746 Kalaba F.K., Quinn C.H., Dougill A.J., Vinya R. (2013). Floristic composition, species
747 diversity and carbon storage in charcoal and agriculture fallows and management
748 implications in Miombo woodlands of Zambia. *Forest Ecology and Management,* 304 pp
749 99-109. https://doi.org/10.1016/j.foreco.2013.04.024

750 Kalaba F.K., Quinn C.H., Dougill A.J. (2013a). The role of forest provisioning ecosystem
751 services in coping with household stresses and shocks in Miombo woodlands, Zambia.
752 *Ecosystem Services*5, pp 143-148. https://doi.org/10.1016/j.ecoser.2013.07.008

753 Kassahun A., Snyman H.A., Smit G.N. (2008). Impact of rangeland degradation on the
754 pastoral production systems, livelihoods and perceptions of the Somali pastoralists in
755 Eastern Ethiopia. *Journal of Arid Environments*, 72, 7, 1265-1281.
756 https://doi.org/10.1016/j.jaridenv.2008.01.002

Kiggundu N., (2002). Evaluation of rainwater harvesting systems in Rakai and Mbarara District Uganda. GHARP case study report. *Greater Horn of Africa Rainwater Partnership (GHARP)*, Kenya Rainwater Association, Nairobi, Kenya.

Kubbinga B. (2012). Road runoff harvesting in the drylands of sub-Saharan Africa: Its potential for assisting smallholder farmers in coping with water scarcity and climate change, based on case studies in eastern Province, Kenya. *Unpublished MSc. Thesis.* Free University: Amsterdam, Netherlands.

Leal Filho W., Balogun A.L., Ayal D.Y., Bethurem E.M., Murambadoro M., Mambo J., Taddese H., Tefera G.W., Nagy G.J., Fudjumdjum H., Mugabe P. (2018). Strengthening climate change adaptation capacity in Africa- case studies from six major African cities and policy implications. *J. Environ. Sci. Policy*, 86, 29–37. https://doi.org/10.1016/j.envsci.2018.05.004

Leal Filho, W. et al (2020) Handbook of Climate Change Resilience. Springer, Cham.

Leal Filho, W. et al (2021) African Handbook of Climate Change Adaptation. Springer, Cham.

Lembani R., Knight J., Adam E., Kalaba F.K. (2019). Quantifying spatial-temporal changes of aboveground carbon stocks using Landsat time series data: A case study of Miombo woodlands. *International Journal of Environmental Studies*, pp.1-21. https://doi.org/10.1080/00207233.2019.1698870

Liu S., Stern D.I. (2008). A Meta-analysis of Contingent Valuation Studies in Coastal and Near-shore Marine Ecosystems. *MPRA Paper* 11608, University Library of Munich, Germany.

Lopa D., Mwanyoka I., Jambiya G., Massoud T., Harrison P., Ellis-Jones M., Blomley T.,

780 Leimona B., van Noordwijik M., Burgess N.D. (2012). Towards operational payments for
781 water ecosystem services in Tanzania: a case study from the Uluguru Mountains. *Oryx*,
782 46, 34–44. Doi: 10.1017/S0030605311001335

783 Macamo C.C.F., Massuanganhe E., Nicolau D.K., Bandeira S.O., Adams J.B. (2016).
784 Mangrove's response to cyclone Eline (2000): What is happening 14 years later. *Aquatic*
785 *Botany*, 134, 10–17. Doi: 10.1016/j.aquabot.2016.05.004

786 Makowski C., Finkl C.W. (2018). Threats to mangrove forests: Hazards, vulnerability,
787 and management. Coastal Research Library. *Springer International Publishing*. Doi:
788 10.1007/978-3-319-73016-5

789 Malesu M., Sang J.K., Odhiambo O.J., Oduor A.R., Nyabenge M. (2006). Hydrologic
790 impact of ponds on land cover change. RELMA-in-ICRAF, *Technical report no. 32.*
791 Netherlands Ministry of Foreign Affairs and Swedish International Development Agency
792 (SIDA).

793 Martins A.P.B. (2018). É preciso tratar a obesidade como um problema de saúde pública.
794 São Paulo: *Rev. Adm. Empresas*, FGV EAESP. https://doi.org/10.1590/s0034-
795 759020180312

796 Massuanganhe E.A., Macamo C., Westerberg L.O., Bandeira S., Mavume A., Ribeiro E.
797 (2015). Deltaic coasts under climate-related catastrophic events – Insights from the Save
798 River delta. *Mozambique. Ocean & Coastal Management*, 116, 331–340.
799 Doi:10.1016/j.ocecoaman.2015.08.008

800 Meadows D.H., Meadows D.L., Randers J., Behrens III W.W. (1973). Limites do
801 crescimento: um relatório para o projeto do Clube de Roma sobre o dilema da
802 humanidade. São Paulo: Perspectiva.

803 Mekonnen A., Biruk A. (2006). Flood hazard assessment in the Ghibe-Omo basin: the
804 case of 2006. In: Ensermu Kelbessa (ed.), Climate change and environmental
805 degradation: report on the 2006 flood disaster in Ethiopia (Addis Ababa: Forum for
806 Environment), pp19–35

807 Mkanda F.X. (2002). Contribution by farmers' survival strategies to soil erosion in the
808 Linthipe River catchment: implications for biodiversity conservation in Lake
809 Malawi/Nyasa. *Biodivers. Conserv*. 11, 1327–1359. Doi: 10.1023/A:1016265715267

810 Morecroft M.D., Crick H.Q.P., Duffield S.J. Macgregor N.A. (2012). Resilience to
811 climate change: translating principles into practice. *Journal of Applied Ecology*. 49 547–
812 551. Doi: 10.1111/j.1365-2664.2012.02136.x

813 Mugwedi L.F., Ray-Mukherjee J., Roy K.E., Egoh B.N., Pouzols F.M., Douwes E., Boon
814 R., O'Donoghue S., Slotow R., Di Minin E., Moilanen A., Rouget M. (2018). Restoration
815 planning for climate change mitigation and adaptation in the city of Durban, South
816 Africa. *International Journal of Biodiversity Science, Ecosystem Services &*
817 *Management,* 14(1), 132–144. Doi: 10.1080/21513732.2018.1483967

818 Munang R., Thiaw I., Alverson K., Liu J., Han Z. (2013). The role of ecosystem services
819 in climate change adaptation and disaster risk reduction. *Current Opinion in*
820 *Environmental Sustainability*, 5(1), 47-52. https://doi.org/10.1016/j.cosust.2013.02.002

821 Mussa M., Hashim H., Teha M. (2016). Rangeland degradation: Extent, impacts, and
822 alternative restoration techniques in the rangelands of Ethiopia. *Tropical and Subtropical*
823 *Agroecosystems*, 19, 305-318.

824 Nalau J., Becken S., Mackey B. (2018). Ecosystem-based Adaptation: A review of the
825 constraints. *Environmental Science & Policy*, 89, 357-364.
826 https://doi.org/10.1016/j.envsci.2018.08.014

827 Ng'onga M., Kalaba F.K., Mwitwa J., Bright N. (2019). The interactive effects of rainfall,
828 temperature and water level on fish yield in Lake Bangweulu fishery, Zambia. *Journal of*
829 *Thermal Biology*. 84: 45-52. https://doi.org/10.1016/j.jtherbio.2019.06.001

830 Ngigi S.N. (2003). Rainwater Harvesting for Improved Food Security: Promising
831 technologies in the Greater Horn of Africa. *Kenya Rainwater Association*, Nairobi,
832 Kenya. ISBN: 9966-9767-1-X

833 Nicholls R.J., Cazenave A. (2010). Sea-level rise and its impact on coastal zones. New
834 York: *Science*, 328 (5985): 1517-1520. Doi: 10.1126/science.1185782

835 Nyariki D.M. (2017). Regional Integration Support Programme (RISP III). Assessment of
836 the economic valuation of pastoralism in Kenya. A report for IGAD Centre for Pastoral
837 Areas and Livestock Development (ICPALD), Nairobi, Kenya.

838 Nyariki M., Amwata A. (2019). The value of pastoralism in Kenya: Application of total
839 economic value approach. *Pastoralism*: Research, Policy and Practice, 9:9.
840 https://doi.org/10.1186/s13570-019-0144-x

841 OECD – Organisation for Economic Co-operation and Development. (1993). OECD Core
842 Set of Indicators for Environmental Performance Reviews. A Synthesis Report by the
843 Group on the State of the Environment. Paris: *OECD*, 35 p.

844 Ojea E. (2015). Challenges for mainstreaming Ecosystem-based Adaptation into the
845 international climate agenda. *Curr. Opin. Environ. Sustain.*, 14, 41-48.
846 https://doi.org/10.1016/j.cosust.2015.03.006

847 Ojea E., Loureiro M.L., Alló M., Barrio M. (2016). Ecosystem services and REDD:
848 Estimating the benefits of non-carbon services in worldwide forests. *World Dev.*, 78, pp.
849 246-261. https://doi.org/10.1016/j.worlddev.2015.10.002

850 Olaniyi O.A., Ojekunle Z.O., Amujo B.T. (2013). Review of climate change and its effect
851 on Nigeria ecosystem. *International Journal of African and Asian Studies*, 1, 57.

852 Opiyo F.E., Wasonga O.V., Nyangito M.M. (2014). Measuring household vulnerability to
853 climate-induced stresses in pastoral rangelands of Kenya: Implications for resilience
854 programming. *Pastoralism*, 4, 10. https://doi.org/10.1186/s13570-014-0010-9

855 Opiyo F., Wasonga O., Nyangito M., Schilling J., Munang R. (2015). Drought Adaptation
856 and Coping Strategies Among the Turkana Pastoralists of Northern Kenya. *Int J Disaster*
857 *Risk Sci*, 6, 295–309. https://doi.org/10.1007/s13753-015-0063-4

858 OFRI - Oregon Forest Resources Institute. (2018). Oregon's Forest protection laws. An
859 illustrated Manual. Revised-Third Edition. Available in:
860 https://oregonforests.org/pub/oregons-forest-protection-laws-illustrated-manual

861 Pettinotti L., Ayala A., Ojea E. (2018). Benefits from Water Related Ecosystem Services
862 in Africa and Climate Change. Ecological Economics, 149, 294-305.
863 https://doi.org/10.1016/j.ecolecon.2018.03.021

864 Prakash S. (1997). Poverty and Environment: Linkages in mountains and uplands:
865 Reflections on the 'poverty trap' thesis. New York: CREED Working Paper 12.
866 ISBN: 978-1-84369-177-8

867 PROLINNOVA – Promoting Local Innovation in ecologically-oriented agriculture and
868 natural resource management. (2009). Local innovation in climate-change adaptation by
869 Ethiopian pastoralists. Ethiopia and Pastoralist Forum Ethiopia (PFE) Final report. Addis
870 Ababa, Ethiopia.

871 RCRMD – Regional Centre for Mapping of Resources for Development. (2016).
872 Rangeland Assessment and Monitoring Service. https://rcmrd.org/servir-rangeland-

assessment-and-monitoring-service (Accessed on 24 March 2020).

Robledo C., Clot N., Hammill A., Riché B. (2012). The role of forest ecosystems in community-based coping strategies to climate hazards: Three examples from rural areas in Africa. *Forest Policy and Economics*, 24, 20–28. Doi:10.1016/j.forpol.2011.04.006

Ryan C.M., Pritchard R., McNicol I., Owen M., Fisher J.A., Lehmann C. (2016). Ecosystem services from southern African woodlands and their future under global change. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 371(1703), 20150312. Doi: 10.1098/rstb.2015.0312

Sachs J., Schmidt-Traub G., Kroll C., Lafortune G., Fuller G., Woelm F. (2020). Sustainable Development Report 2020. The Sustainable Development Goals and COVID-19. Cambridge: *Cambridge University Press*.

Sala O.E., Yahdjian L., Havstad K., Aguiar M.R. (2017). Rangeland Ecosystem Services: Nature's Supply and Humans' Demand. In: Briske D. (eds) Rangeland Systems. Springer Series on Environmental Management. Springer, Cham. https://doi.org/10.1007/978-3-319-46709-2_14

Scarano F.R. (2017). Ecosystem-based adaptation to climate change: concept, scalability and a role for conservation Science. *Perspectives in Ecology and Conservation*. 15 65-73. Doi: 10.1016/j.pecon.2017.05.003

Schoffeleers J.M. (1979). Guardians of the land: essays on Central African territorial cults. Gwelo/Gweru, Zimbabwe: Mambo Press. ISBN: 0869227343 9780869227343

Shiferaw W., Demissew S., Bekele T. (2018). Invasive alien plant species in Ethiopia: ecological impacts on biodiversity a review paper. *Int J Mol Biol Open Access*, 3, 4, 171–178. Doi: 10.15406/ijmboa.2018.03.00072

Smith C.J., Papadopoulou K-N., Barnard S., Mazik K., Elliott M., Patrício J., Solaun O., Little S., Bhatia N., Borja A. (2016). Managing the Marine Environment, Conceptual Models and Assessment Considerations for the European Marine Strategy Framework Directive. *Front. Mar. Sci*. 3:144. Doi: 10.3389/fmars.2016.00144

Stephens E.C., Nicholson C.F., Brown D.R., Parsons D., Barret C.B., Lehmann J., Mbugua D., Ngoze S., Pell A.N., Riha S.J. (2012). Modeling the impact of natural resource-based poverty traps on food security in Kenya: the crops, livestock and soils in smallholder economic systems (CLASSES) model. *Food Sec*. 4, 423–439. https://doi.org/10.1007/s12571-012-0176-1

Stringer C.E., Trettin C.C., Zarnoch S.J., Tang W. (2015). Carbon stocks of mangroves within the Zambezi River Delta, Mozambique. *Forest Ecology and Management*, 354, 139–148. Doi:10.1016/j.foreco.2015.06.027

Sudmeier-Rieux K., Masundire H., Rizvi A., Rietbergen S. (2006). Ecosystems, livelihoods and disasters: An integrated approach to disaster risk management. Gland, Switzerland: *IUCN*. Doi:10.2305/IUCN.CH.2006.CEM.4.en

Swinburn B.A., Kraak V.I., Allender S., Atkins V., Baker P.I., Bogard J.R., Brinsden H., Calvillo A., de Schutter O., Devarajan R., Ezzati M., Friel S., Goenka S., Hammond R.A., Hastings G., Hawkes C., Herrero M., Hovmand P., Howden M., Jaacks L.M., Kapetanaki A.B., Kasman M., Kuhnhein H.V., Kumanyika S.K., Larijani B., Lobstein T., Long M.W., Matsudo V.K.R., Mills S.D.H., Morgan G., Morshed A., Nece P.M., Pan A., Patterson D.W., Sacks G., Shekar M., Simmons G.L., Smit W., Tootee A., Vandevijvere S., Waterlander W., Wolfenden L., Dietz W.H., (2019). The Global Syndemic of Obesity, Undernutrition, and Climate Change: The Lancet Commission report. *Lancet*. http://dx.doi.org/10.1016/S0140-6736(18)32822-8

920 Svarstad H., Petersen L.K., Rothman D., Siepel H., Wätzold F. (2008). Discursive biases
921 of the environmental research framework DPSIR. *Land Use Policy,* 25, 116–125. Doi:
922 10.1016/j.landusepol.2007.03.005

923 Teka O., Houessou L.G., Djossa B.A., Bachmann Y., Oumorou M., Sinsin B. (2019).
924 Mangroves in Benin, West Africa: threats, uses and conservation opportunities.
925 *Environment, Development and Sustainability* 21, 1153-1169.
926 https://doi.org/10.1007/s10668-017-0075-x

927 UNFCCC – United Nations Framework Convention on Climate Change. (2008). Report
928 of the Conference of the Parties on its thirteenth session, held in Bali from 3 to 15.
929 Geneva: *United Nations Digital Library.* FCCC/CP/2007/6/Add.1.

930 UNISDR – United Nations Office for Disaster Risk Reduction. (2018). Economic losses,
931 poverty & disasters 1998-2017. Centre for Research on the Epidemiology of Disasters.
932 [Internet]. Available from: https://www.undrr.org/publication/economic-losses-poverty-
933 disasters-1998-2017

934 Uy N.; Shaw R. (2012). The role of ecosystems in climate change adaptation and disaster
935 risk reduction. Uy N., Shaw R. (ed.) Ecosystem-Based Adaptation Community,
936 Environment and Disaster Risk Management. Bingley: *Emerald Group Publishing*
937 *Limited*, 12, 41–59. https://doi.org/10.1108/S2040-7262(2012)0000012009

938 Von Möllendorff C., Hirschfeld J. (2016). Measuring impacts of extreme weather events
939 using the life satisfaction approach. *Ecol. Econ.,* 121,108-116.
940 https://doi.org/10.1016/j.ecolecon.2015.11.013

941 Waheed, B.; Khan, F.; Veitch, B. (2009). Linkage-Based Frameworks for Sustainability
942 Assessment: Making a Case for Driving Force-Pressure-State-Exposure-Effect-Action
943 (DPSEEA) Frameworks. *Sustainability*, *1*, 441-463. https://doi.org/10.3390/su1030441

944 Wamsler C., Luederitz C., Brink E. (2014). Local levers for change: mainstreaming
945 ecosystem-based adaptation into municipal planning to foster sustainability transitions.
946 *Glob. Environ. Change*, 29, 189-201. https://doi.org/10.1016/j.gloenvcha.2014.09.008

947 Wamsler C. (2017). Stakeholder involvement in strategic adaptation planning:
948 transdisciplinarity and co-production at stake? *Environ. Sci. Policy*, 75, 148-157.
949 https://doi.org/10.1016/j.envsci.2017.03.016

950 Wilson K.B. (1989). Trees in fields in southern Zimbabwe. *J. South. Afr. Stud.* 15, 369–
951 383. Doi: 10.1080/03057078908708205

952 Worden J., Western D., Waruingi L. (2009). Exploring Potential Economic and
953 Livelihood Impacts of Climate Change and Possible Adaptation Mechanisms in the
954 Kenyan Rangelands. *Stockholm Environment Institute*.

955 World Bank (2015). Tourism in Africa: Harnessing Tourism for Growth and Improved
956 Livelihoods. Washington DC 20433. Available in: http://www.worldbank.org/afr/tourism.

957 Yahdjian L., Sala E.O., Havstad K.M. (2015). Rangeland ecosystem services: shifting
958 focus from supply to reconciling supply and demand. *The Ecological Society of America,*
959 doi: 10.1890/140156

960 Zölch T., Wamsler C., Pauleit S. (2018). Integrating the ecosystem-based approach into
961 municipal climate adaptation strategies: The case of Germany. *Journal of Cleaner*
962 *Production.* 180, 966-977. Doi: 10.1016/j.jclepro.2017.09.146