

Please cite the Published Version

Gebrye, Tadesse , Fatoye, Clara, Yusuf, Mohamed and Fatoye, Francis (2020) The Economic Burden of Musculoskeletal Disorders Resulting from Road Trafic Accident in Low and Middle Income Countries: A Systematic Review. In: ISPOR 2020, 16 May 2020 - 20 May 2020, Virtual.

DOI: https://doi.org/10.1016/j.jval.2020.04.727

Publisher: Elsevier on behalf of International Society for Pharmacoeconomics and Outcomes Research

Version: Published Version

Downloaded from: https://e-space.mmu.ac.uk/628294/

Usage rights: O In Copyright

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines)

THE ECONOMIC BURDEN OF MUSCULOSKELETAL DISORDERS RESULTING FROM ROAD TRAFIC ACCIDENT IN LOW AND MIDDLE INCOME COUNTRIES: A SYSTEMATIC REVIEW

- T. Gebrye
- C. Fatoye
- M. Yusuf
- F. Fatoye

Objectives

Approximately 20 - 50 million people sustain nonfatal injuries as a result of road traffic accidents (RTA). Treatment costs as well as lost wages associated with this nonfatal injury are substantial. The study reviewed the economic burden of musculoskeletal (MSK) disorders caused from RTA in low and middle-income countries (LMICs).

Methods

We performed a systematic search of MEDLINE, CINAHL, Cochrane Database of Systematic Reviews (CDSR), Health Economic Evaluations Database (HEED), and Web of Science Database from inception to September 2019. Studies that assessed the economic burden of MSK resulting from RTA in LMICs published in English language were included. The quality of the included studies was assessed using the Drummond and Jefferson criteria. A descriptive data synthesis was used to present the results.

Results

The search identified 456 studies, and nine of them met the inclusion criteria and were included in this review. The studies were conducted in Iran (n = 3), Nigeria (n = 2), Vietnam (n = 1), India (n = 1), Thailand (n = 1), and Nepal (n = 1). The quality score of the included studies were \geq 50%. The mean age of the patients in the included studies ranged between 24.6 to 40.3 years. Mean direct and indirect costs in the included studies ranged from \$11.04 to \$2494 and \$33.3 to \$189.2, respectively over 12 months. Hospitalisation and treatment were identified as the major cost drivers.

Conclusions

The findings of the review showed that the costs of injuries resulting from RTA were considerably high relative to health expenditures levels. Hence, it is important that prevention

strategies be in place to reduce the economic burden of MSK disorders caused by RTA in LMICs.