



Please cite the Published Version

McAuley, Scott, Dobbin, Nicholas , Morgan, Chris and Goodwin, Peter  (2022) Predictors of time to return to play and re-injury following hamstring injury with and without intramuscular tendon involvement in adult professional footballers : a retrospective cohort study. *Journal of Science and Medicine in Sport*, 25 (3). pp. 216-221. ISSN 1440-2440

DOI: <https://doi.org/10.1016/j.jsams.2021.10.005>

Publisher: Elsevier

Version: Accepted Version

Downloaded from: <https://e-space.mmu.ac.uk/628538/>

Usage rights:  [Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0](#)

Additional Information: This is an Author Accepted Manuscript of a paper which appeared in *Journal of Science and Medicine in Sport* published by and copyright Elsevier.

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

Manuscript Title: Predictors of time to return to play and re-injury following hamstring injury with and without intramuscular tendon involvement in adult professional footballers: A retrospective cohort study.

Short title: Time to return to play and re-injury rate following hamstring injury.

Submission Type: Original research

Authors: Scott McAuley^a, Nick Dobbin^b, Chris Morgan^a, Peter C Goodwin^b

Affiliation:

^aLiverpool Football Club, Liverpool, UK

^bDepartment of Health Professions, Manchester Metropolitan University, Manchester, UK

Corresponding Author: Scott McAuley, Liverpool Football Club, Liverpool, UK, L33 5XB.

Work Tel: 07886 838 414

Work Email: scott.mcauley@liverpoolfc.com

Abstract Word Count: 248

Manuscript Word Count: 3234

Number of References: 35

Tables: 2

Figures: 1

Abstract

Objectives: In one English Premier League football club over four seasons, 1) describe the number of hamstring strain injuries (HSI) sustained using the British Athletics Muscle Injury Classification (BAMIC); 2) determine if intramuscular tendon HSI influenced the time to return to play (TTRTP) and reinjury rate; 3) determine the predictors of TTRTP and reinjury.

Design: Retrospective cohort design.

Methods: All first team players who sustained a HSI between 2014 and 2018 were included. Players underwent an MRI scan that was graded by a Radiologist using the BAMIC (0a-4) criteria. TTRTP, reinjury rate and information on suspected predictors were recorded.

Results: Thirty-five HSI experienced by 24 players (age = 26 ± 4 years) were recorded over the 4 seasons. There was a difference in TTRTP between grades 1a and 2c ($P = 0.007$), but not between 2b and 2c ($P = 0.845$). Grade of HSI ($P = < 0.001$) and removal of the player ($P < 0.001$) were predictors of TTRTP, with each increase in grade resulting in an additional 3 days of TTRTP, and being removed, an additional 11 days. Grade and all other predictors did not influence reinjury rate, albeit higher odds were evident for previous HSI, experiencing the HSI during sprinting, passing a ball or stretching, and reported increase days of pain during walking.

Conclusion: HSIs extending into the intramuscular tendon (2b *cf.* 2c) do not influence TTRTP or re-injury, albeit TTRTP was affected by the BAMIC grade and if the player was removed from activity.

Keywords: Hamstring injury, tendon, professional football, rehabilitation, BAMIC.

Introduction

63 Hamstring strain injuries (HSI) account for ~12% of all injuries in professional football¹ and are
64 consistently the most prevalent time-loss injury reported during a competitive season.² Such is
65 the prevalence (~12%) and incidence rates (0.4-1.9/1000 exposure hours)³ of HSI, it's reported
66 that a 25-player squad typically experience 5-6 HSI per season, resulting in more than 80 days
67 lost due to injury.¹ It has also been reported that the average cost per injured player competing
68 in the UEFA Champions League is approximately €500,000 per month.⁴ For example, during
69 the 2016/2017 English Premier League (EPL) season, £177 million was spent in wages to
70 injured players,⁵ increasing to £217m in 2017/2018 and £221m in 2018/2019, with HSI
71 accounting for a large proportion of these injuries.⁵

72 Given the high prevalence and incidence rates of HSI in football players and the implications
73 this has for the player, club and medical team, return to play (RTP) research has become a
74 focus for many researchers and clinicians involved in football.⁶ Indeed, two Delphi surveys, an
75 expert led consensus statement, and a survey of 131 professional football teams have made
76 recommendations for reducing HSI and improving RTP in professional football.⁶⁻⁸ A worldwide
77 Delphi survey concluded that RTP following HSI should be defined as “the moment a player has
78 received criteria-based medical clearance and is mentally ready for full availability for
79 match/team training with RTP decisions based on shared decision-making, providing clarity of
80 when RTP is appropriate to reduce the risk of injury recurrence”.⁹ The RTP consensus
81 statement by Arden et al.⁶ recommended that: 1) RTP should be viewed as a continuum rather
82 than a single event at the conclusion of rehabilitation, and follow through to ‘return-to-
83 performance’; 2) objective markers should guide RTP and; 3) practitioners should follow a
84 shared decision-making process including key stakeholders.^{6, 7} In part, these guidelines aim to
85 improve time to return to play (TTRTP) and moderate the risk of HSI re-injury.

86 Predicting TTRTP and minimising injury recurrence is challenging as reflected in that HSI
87 recurrence rates are 15-20% (range 14% to 63%).^{10, 11} Further, there is weak evidence to
88 suggest that scanning with Magnetic Resonance Imaging (MRI) can aid prediction of TTRTP
89 and prognosis for RTP following HSI.¹² Time to RTP has been shown to vary greatly between

and within MRI classification systems following HSI,¹³ so, when used, the specific MRI classification should be reported to avoid miscommunication.¹³

Severity and site of tear is an important factor when considering RTP, as evidence suggests intramuscular tendon injuries require longer rehabilitation and TTRTP, with a higher risk of injury recurrence compared to HSI with no tendon involvement.^{10, 14, 15} Pollock et al.¹⁰ proposed the British Athletics Muscle Injury Classification (BAMIC), using a 0-4 scale, with grades 1-4 including a grade of a-c to specify injury site within the muscle; a = myofascial, b = musculotendinous junction, or c = intratendinous.¹⁰ This method has been shown to demonstrate 'substantial' intra-rater and 'almost perfect' inter-rater reliability when concerning the overall BAMIC classification.¹⁵ The first study to use the BAMIC in a sporting context looked at TTRTP and injury recurrence over a 4-year period in track and field athletics.¹⁰ The results showed injuries extending into the tendon [grade c] are more prone to reinjury and resulted in a greater number of days taken to RTP.¹⁰ Since then, the relevance of intramuscular tendon tears has been disputed and remains controversial, with evidence indicating complete resolution of these injuries on MRI is not necessary for successful RTP.¹⁶⁻¹⁹ As such, further research investigating if intramuscular tendon tears influence the duration of time-loss and reinjury rate is warranted in a range of populations including football players.

The aims of this study were, 1) To use the BAMIC to describe the number of HSI over four competitive football seasons; 2) Determine if HSI including an intratendinous tear results in extended TTRTP and higher reinjury; and 3). Explore possible predictors of TTRTP and reinjury in professional footballers.

Methods

This was a retrospective observational study of HSI over four consecutive English Premier League seasons 2014-2018, including domestic and European cup matches. Ethical approval was obtained from the Faculty of Health, Psychology and Social Care Research Ethics

Committee at Manchester Metropolitan University (Ref: 1055) with consent obtained from the athletes and club.

Participants included professional footballers contracted to a single football club in the English Premier League. All first team players registered at the club at the beginning of each season who sustained an HSI were included. Players who transferred to another club or whose contract had finished before the end of the season were included providing they played for the club in that season, provided prior consent and had return to play to allow TTRTP to be determined. Over four seasons, 52 players were eligible, receiving full-time medical care with all injury episodes recorded on an electronic medical record system by the physiotherapists at the club. Injury and rehabilitation data were subsequently collected and analysed by the lead researcher who was not directly involved in the key decisions related to the care of the players. Injury definitions and data collection procedures followed the consensus statement and data collection of football injuries,^{20, 21} thus standardising the reporting across all physiotherapists at the club and when completing the electronic medical record system. Included were HSI reported between June 2014 and May 2018 that occurred during training, matches or rehabilitation. These were defined as “*an acute injury to the posterior thigh muscle resulting in further assessment and referred for MRI investigation within 7 days of the injury*”.⁸ Excluded were HSI including contusions, haematomas, chronic tendinopathies, and players who experienced an exacerbation of the current injury during rehabilitation.^{10, 20}

Injury data were reviewed by one independent physiotherapist (musculoskeletal specialist with >10 years clinical experience) to determine all episodes of HSI that occurred between the 2014 and 2018 seasons. Over the study period, treatment was provided by different physiotherapists, all with >10 years clinical experience each. HSI-related RTP criteria included hamstring length; strength (isometric and isokinetic, <10% limb symmetry) and function (pitch-based running, acceleration, and high speed). Criteria were not prescriptively implemented, as clinicians acted autonomously providing individualised rehabilitation approaches according to clinical and pitch-based objective markers to determine RTP.

TTRTP was defined as *'the number of days until the player had resumed full team training and available for match selection'*.²¹ To establish TTRTP, the electronic medical record system was reviewed to calculate the number of days each player was unavailable based on the player's status changing from 'unavailable for play or training' to 'available'.

Injury recurrence was defined as *'an injury of the same type, at the same site as an index injury on return to full participation'* that was recorded during the subsequent 12-month period.²¹ To determine when players were most at risk of re-injury after RTP, follow-ups were documented routinely at 2, 6 and 12 months.

Other information and potential predictors included previous HSI, age, mechanism of injury, muscle injured and location, days taken to walk pain-free, and if the player was removed from play (match, training, or rehabilitation).^{10, 22-24} A decision to remove a player from play was based on clinical reasoning following on-pitch assessment including: muscle palpation, length and (isometric) strength testing, combined with the player's subjective feedback (ability to continue), often determined by the mechanism of injury (sudden traumatic onset of pain vs. gradual onset of pain).

Players were imaged on either 1.5T or 3T Siemens MRI platforms using dedicated surface coils. The imaging protocol consisted of coronal T1 and inversion recovery sequences with 4mm slice thickness, axial T1 and fat-saturated proton density and sagittal fat-saturated proton density sequences with 3mm slice thickness. All sequences used high-resolution matrix parameters. To minimise bias, all scans were read by a fellowship-trained musculoskeletal radiologist with 14 years' experience in sports imaging and were conducted 48 hours after the injury. The radiologist was not provided with any players' medical history - only a note on the suspect HSI - and provided a report on the athlete immediately after the scan to support the club's medical staff during rehabilitation. The radiologist provided a grade for severity and location of HSI that was in accordance with the BAMIC along with their clinical interpretation and recommendations.¹⁰

All statistics were analysed using the IBM SPSS package Version 25 (IBM SPSS Statistics for Windows, Version 25.0. Armonk, NY: IBM Corp). Descriptive statistics were used to describe player and HSI characteristics (Aim 1). The Shapiro-Wilk Test was used to determine if the data conformed to the assumptions of normality. BAMIC grades were compared according to TTRTP using Kruskal Wallis tests with Bonferroni corrected Mann Whitney post-hoc test (Aim 2). Differences in reinjury across BAMIC grades was assessed using Fisher's Exact test. Mean, SD and 95% confidence limits (CL) were calculated for descriptive purposes.

Linear and binary logistic regression using an 'Enter' method were used to determine the independent effects of grade of injury, age, previous history of HSI, mechanism of injury, muscle injured and location, injured during match/training/rehabilitation (TTRPT only), removed from play (player unable to continue due to HSI, and number of days to walk pain free on TTRTP and injury recurrence (Aim 3), respectively. Odds ratios were derived for predictors in the binary logistic regression. The compatibility of the data with the hypothesis was interpreted using the exact probability statistic (*P* value).

Results

Player and injury characteristics

Forty HSI were reported by twenty-six players over four English Premier League seasons (Table 1). Five injuries were excluded due to surgical intervention to address other chronic musculoskeletal injuries not related to each hamstring injury. Three HSI were excluded from the analysis for TTRTP as they experience a re-injury during rehabilitation and a return date was not available. However, these were included in the analysis for injury recurrence. Thus, the final analysis was based on 32 and 35 HSI's for TTRTP and recurrence, respectively, from 24 players.

******Table 1. Player and injury characteristics******

Grade 0, 1, 2 and 3 accounted for a total of 2, 12, 19 and 2 HSI's respectively. Of the 35 injuries, 9 injury recurrences were recorded after the players had returned to full team training. Six of these occurred <2 months of RTP and three recurrences occurring <12 months of RTP.

Intramuscular tendon injury and RTP and injury recurrence

There were differences in TTRTP according to BAMIC injury types 0b-3b ($H = 17.3$, $df = 6$, $P = 0.008$). TTRTP differences were observed between small myofascial (1a) and moderate musculotendinous junction tears (2b) (mean difference = 14 days, 95%CI = 4-24 days; $P = 0.016$) and small myofascial (1a) and moderate intratendinous tears (2c) (mean difference = 14 days; 95% CI = 5-23 days; $P = 0.007$). There were no differences (at $P = 0.05$) in TTRTP between injuries with and without tendon involvement (1a & 1b, or 2b & 2c; all $P > 0.05$) (Figure 1). There were no differences between injury recurrence according to the BAMIC grading, 0b-3b (Fishers Exact value = 3.897, $P = 0.844$), two reinjuries observed for grades 1b and 2c, and one observed for grade 0b, 1a, 2a, 2b and 3b.

******Figure 1******

Predictors of TTRTP and injury recurrence

Results from the linear regression indicated that all predictors combined explained 84.7% of the variance of the TTRTP ($P < 0.001$) (Table 2). Grade and whether the athlete was removed from the activity were significant predictors (Table 2). The unstandardised beta for grade indicated that for every 1 increase of grade of HSI, TTRTP increased by ~3 days (95% CI = 2 to 4). Being removed from the field of play was associated with an additional ~11 days (95% CI = 7 to 15) to RTP. All other predictors had confidence limits that overlapped the null effect, implying that association between the predictor (e.g., previous injury) and TTRTP is minimal (Table 2).

****Table 2. Predictors of TTRTP and injury recurrence****

Results from the binary logistical regression indicated that the predictors explained 49.7% of the variance in injury recurrence (Nagelkerke $R^2 = 0.497$). However, none of the predictors significantly altered the odds of experiencing a reinjury at $P = 0.05$. The odds ratio, whilst not significant (95%CI cross 1), did suggest that there are 'slightly' higher odds of a reinjury when players have had a previous HSI, experience the HSI during sprinting, passing a ball or stretching and experience increased days of pain during walking (Table 2).

Discussion

The aims of this study were to use the BAMIC to describe HSI over four English Premier League seasons; to determine if HSI including an intratendinous tear results in extended RTP and higher injury recurrence; and to determine the influence of some predictors of RTP and injury recurrence.

Injury characteristics

The current findings of 35 HSIs in 24 professional footballers over four seasons was higher than previous UEFA audits where 5-6 HSIs were reported per season.¹ Whilst this was undoubtedly a unique period for HSI in this club, comparisons to other clubs are beyond the scope of this study. Furthermore, it is difficult to account for differences in medical expertise, player characteristics, training loads and management strategies amongst many other factors between clubs. However, the number of HSI across multiple seasons allows further exploration beyond studies that are based over a single season or experience a limited number of HSI.

Findings in the current study highlight that most HSI occurred during match-play, which is noteworthy given this is 3.5 times more than training. Such findings might be explained by higher

overall intensity, greater physical loads and a greater degree of risk taking when compared to training. Bengtsson et al.²⁵ suggested that injury rates in the first match after RTP increase by 87% compared with a 'typical match', with the odds of injury dropping by 7% with each additional training session before the first match. However, training-related HSI rates have increased since 2001 (~4%), whilst match-related HSI rates have remained relatively stable.¹ In this study, the final data analysis of RTP and injury recurrence consisted of 35 HSI in 24 players with the moderate, (2a-c), injuries most common (54%) and with intratendinous tears (2c) producing the longest RTP of all grades.

The long head of biceps femoris was the most injured muscle, with the highest injury recurrence rate, which is in accordance with previous research.^{10, 23, 26} The long head of biceps femoris undergoes the greatest lengthening during high-speed running²⁷ and has a separate nerve supply to the short head, potentially with asynchronous contraction.²⁸ This might, in part, explain the greater recurrence rate, though we urge caution given the limited number of re-injuries. To confirm such findings, a multi-club, multi-league and/or multi-nation approach is required.

Intratendinous injuries

It has been suggested that injuries extending into the tendon require prolonged rehabilitation with longer RTP and higher risk of injury recurrence.^{10, 14, 16} Whilst the current study found differences in TTRTP between myofascial and intratendinous tears, they were also different sizes (small 1a vs. medium 2b, 2c) which needs consideration. No difference in TTRTP was found between tears with and without intratendinous involvement of the same size (2b vs. 2c). These findings suggest that HSI extending into the tendon do not always mean a greater time-loss for professional footballers, though further research with a larger sample from multiple clubs is needed to substantiate or refute this finding.

Recurrence

Player unavailability due to injury increases pressure on medical teams to return players to training as quickly as possible, whilst ensuring injury recurrence is no greater than a pre-injured

level.¹¹ There is no consensus on a single clinical or functional test or imaging investigation that provides a safe indication of RTP, but a multi-disciplinary approach is likely to be important with the use of subjective and objectives measures.^{8, 23}

Previous research indicates HSI injury recurrence is not related to tendon involvement.¹⁸ In the current study, the HSI reoccurrence rate of 26% was slightly higher than other, such as the 19% and 14% reported by Van der Made et al¹⁸ and Malliaropoulos et al.²⁹, respectively. Further comparisons cannot be made to other studies due to a lack of sufficient power to determine differences in the current study. However, 89% of reinjuries occurred in grade 0, 1 and 2 injuries, similar to Malliaropoulos et al.²⁹ who found low grade 1 & 2 HSI were more likely to recur than high grade 3 & 4, thus potential reflecting a short rehabilitation period and less monitoring.

The first month following RTP has the highest risk for HSI recurrence.^{16, 30} Of the 29 HSI in the current study, four re-injuries occurred within two months following RTP, with the highest recurrence rate seen in grade 1 HSI as opposed to higher grades of HSI. These findings suggest professional footballers are at greatest risk of injury recurrence within the first 2 months on RTP following a grade 1 HSI, with a key risk factor potentially due to incomplete healing times and ongoing muscle regeneration in lower grades of injury.³¹

Predictors

The finding that injury grade, determined using BAMIC, and whether a player was removed from play are significant predictors of TTRTP provide a useful insight with estimated number of days that medical personnel can use in future practice when giving estimates to coaching staff and the athlete. Our results indicated that for every increase in BAMIC grading, the number of days before the athletes return increased by 3, with an upper confidence limit of 4 days. A result that the severity of HSI increases the TTRTP is not surprising, but the qualification of an upper value is clinically and practically meaningful in the context of a professional football club where accurate estimates can be used. The results also indicated that an inability to continue with the activity (match, training, rehabilitation) was associated with an increased TTRTP of between 7 and 15 days. A sudden onset of symptoms in the posterior thigh (the player who “pulls up”) is a

good indicator that a significant muscle injury has taken place, and this can usually be confirmed with length, strength (isometric) and palpation testing on pitch.³² This contrasts with the lower grade injury where a player won't experience a significant sudden onset of pain and will often report the issue at the end of a game as the resultant on pitch symptoms haven't limited their ability to perform or they have continued with play regardless.³²

We also note how a range of other factors appeared to have limited predictive ability for TTRTP and reinjury following HSI. Indeed, our results, and those of others, indicate that an increased age, passive loss of range of movement and previous HSI, maximum pain at time of injury, delay in physiotherapy, time taken to walk pain free, change in pain during strength testing, duration of palpation pain, eccentric hamstring strength, hamstring fascicle length, maximal velocity running exposure and hamstring fatigue resistance require further investigation before being ruled in/out as clinical predictors.^{22, 33} Our results do, however, suggest a greater emphasis and responsibility should be placed on the club's medical department to monitoring player training and match loads closely in the first 2 months on RTP and for low-grade HSI due to the elevated risk of injury recurrence and natural healing likely to be incomplete.³¹ Furthermore, medical and sport science personnel should ensure injury prevention programmes are performed routinely and targeting high-risk players (e.g. those with previous HSIs).³⁴ Recommendations for HSI prevention suggest a holistic approach that translates existing knowledge of risk and applying this to a football context.³³

Strengths of study

This study was based over four English Premier League seasons with final data collection analysis consisting of 29 HSI in 24 professional footballers. To limit bias, injury definitions, two main outcome measures and data collection followed the consensus statement and data collection of football injuries, ensuring a clear and accurate methodological process.^{1, 20, 21} Rehabilitation adopted the same principals and recommendations advised by the 2016 RTP consensus statement with clear objective progressive markers used to guide RTP combined

with a shared decision-making process involving key stakeholders from the club's medical department.⁶

Limitations and future directions

The relatively small sample included in this study is a limitation and impacts on the statistical power, especially during the predictive analysis on re-injury. However, there needed to be a balance between the number of seasons to capture enough HSIs and the variances in coaches, managers, and medical personnel. As such, future work might consider a multi-club, multi-league and/or a multi-nation approach to ensure enough HSI are observed to provide additional statistical power, explore other predictors, and tailor programmes to moderate injury risk.

Conclusion

Following acute HSI in professional footballers, the BAMIC is a detailed and useful method of categorisation. Differences in TTRTP were observed between small myofascial (1a) and moderate musculotendinous tears (2b) and small myofascial (1a) and moderate intratendinous tears (2c). However, the effect of injuries with and without intramuscular tendon injury on TTRTP and injury recurrence remains inconclusive.

Practical Implications

- Medical personnel working in professional football can use the BAMIC classification for categorising hamstring strain injuries in a standardised manner, support future interpretation and comparison across other teams, age categories and/or sports.
- Indications are that hamstring strain injury's extending into the intramuscular tendon (2b *cf.* 2c) tend not to influence RTP or reinjury in professional footballers.
- Low grade (0b, 1a & 1b) hamstring strain injury reflected 89% of the reinjury rate, most occurring <2 months on return to play. Indications are that professional footballers are

at greatest risk of injury recurrence within the first 2 months of RTP following a low grade HSI.

- BAGIC grade of hamstring strain injury and removed from play were significant predictors of RTP and may prove a useful guide for medical personnel in professional football to support estimates for RTP.

Acknowledgements

We would like to acknowledge Liverpool Football Club for supporting this research and the players involved in the study.

Funding The authors declare no specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests: None declared.

References

1. Ekstrand J, Walden M, Hagglund M. Hamstring injuries have increased by 4% annually in men's professional football since 2001: a 13-year longitudinal analysis of the UEFA Elite Club injury study. *Br J Sports Med* 2016; 50:731-737.
2. Ekstrand J, Krutsch W, Spreco W, et al. Time before return to play for the most common injuries in professional football: a 16-year follow-up of the UEFA elite club injury study. *Br J Sports Med* 2020;54(7):421-426.
3. Diemer WM, Winters M, Tol JL, et al. Incidence of acute hamstring injuries in soccer: a systematic review of 13 studies involving more than 3800 athletes with 2 million sport exposure hours. *J Orthop Sports Phys Ther* 2021;51(1):27-36.
4. Ekstrand J. Keeping your top players on the pitch: the key to football medicine at a professional level. *Br J Sports Med* 2013; 47:723–724.
5. <https://www.jlt.com/industry/sports-media-and-entertainment/insurance/insights/football-injury> (accessed on 14th January 2020).
6. Ardern CL, Glasgow P, Schneiders A, et al. Consensus statement on return to sport from the First World Congress in sports physical therapy, Bern. *Br J Sports Med* 2016; 50:853-864.
7. Dunlop G, Ardern CL, Andersen TE, et al. Return-to-Play Practices Following Hamstring Injury: A Worldwide Survey of 131 Premier League Football Teams. *Sports Med* 2020; 50:829-840.
8. Zambaldi M, Beasley I, Rushton A. Return to play criteria after hamstring muscle injury in professional football: a Delphi consensus study. *Br J Sports Med* 2017; 51:1221-1226.
9. van der Horst N, Backx F, Goedhart EA, et al. Return to play after hamstring injuries in football (soccer): a worldwide Delphi procedure regarding definition, medical criteria, and decision-making. *Br J Sports Med* 2017 Nov;51(22):1583-1591.
10. Pollock N, Patel A, Chakraverty J, et al. Time to return to full training is delayed and recurrence rate is higher in intratendinous ('c') acute hamstring injury in elite track and

- field athletes: clinical application of the British Athletics Muscle Injury Classification. *Br J Sports Med* 2016; 50:305-310.
11. De Visser H, Reijman M, Heijboer M, et al. Risk factors of recurrent hamstring injuries: a systematic review. *Br J Sports Med* 2011; 46:124-130.
12. Reurink G, Brilman EG, de Vos RJ, et al. Magnetic resonance imaging in acute hamstring injury: can we provide a return to play prognosis? *Sports Med* 2015; 45: 133-146.
13. Wangensteen A, Guermazi A, Tol JL, Roemer FW, Hamilton B, Alonso JM, Whiteley R, Bahr R. New MRI muscle classification systems and associations with return to sport after acute hamstring injuries: a prospective study. *Eur Radiol* 2018; 28(8):3532-3541.
14. Comin J, Malliaras P, Baquie P, et al. Return to competitive play after hamstring injuries involving disruption of the central tendon. *Am J Sports Med* 2013; 41:111–115.
15. Wangensteen A, Tol JL, Roemer FW, et al. Intra-and interrater reliability of three different MRI grading and classification systems after acute hamstring injuries. *Eur J Radiol* 2017; 89:182-190.
16. Askling CM, Tengvar M, Saartok T, et al. Acute first-time hamstring strains during high-speed running: a longitudinal study including clinical and magnetic resonance imaging findings. *Am J Sports Med* 2007; 35:197-206.
17. Vermeulen R, Almusa E, Buckens S, et al. Complete resolution of a hamstring intramuscular tendon injury on MRI is not necessary for a clinically successful return to play. *Br J Sports Med* 2020. Doi: 10.1136/bjsports-2019-101808.
18. van der Made AD, Almusa E, Whiteley R, et al. Intramuscular tendon involvement on MRI has limited value for predicting time to return to play following acute hamstring injury. *Br J Sports Med* 2018; 52:83-88.
19. van der Made AD, Almusa E, Reurink G, et al. Intramuscular tendon injury is not associated with an increased hamstring reinjury rate within 12 months after return to play. *Br J Sports Med* 2018; 52:1261-1266.

20. Ekstrand J, Lee JC, Healy JC. MRI findings and return to play in football: a prospective analysis of 255 hamstring injuries in the UEFA Elite Club Injury Study. *Br J Sports Med* 2016; 50:738-743.
21. Fuller CW, Ekstrand J, Junge A, et al. Consensus statement on injury definitions and data collection procedures in studies of football (soccer) injuries. *Br J Sports Med* 2006; 40:83-92.
22. Green B, Bourne MN, van Dyk N, et al. Recalibrating the risk of hamstring strain injury (HSI): A 2020 systematic review and meta-analysis of risk factors for index and recurrent HSI in sport. *Br J Sports Med* 2020; 54:1081-1088.
23. van Heumen M, Tol JL, de Vos R, et al. The prognostic value of MRI in determining reinjury risk following acute hamstring injury: a systematic review. *Br J Sports Med* 2017; 51:1355-1363.
24. Jacobsen P, Witvrouw E, Muxart P, et al. A combination of initial and follow-up physiotherapist examination predicts physician-determined time to return to play after hamstring injury, with no added value of MRI. *Br J Sports Med* 2016; 50:431-439.
25. Bengtsson H, Ekstrand J, Walden M, et al. Few training sessions between return to play and first match appearance are associated with an increased propensity for injury: a prospective cohort study of male professional football players during 16 consecutive seasons. *Br J Sports Med* 2020; 54:427-432.
26. Wangensteen A, Tol JL, Witvrouw E, et al. Hamstring reinjuries occur at the same location and early after return to sport: A descriptive study of MRI-confirmed reinjuries. *Am J Sports Med* 2016; 44:2112-2121.
27. Heiderscheit BC, Hoerth DM, Chumanov ES, et al. Identifying the time of occurrence of a hamstring strain injury during treadmill running: a case study. *Clin Biomech* 2005; 20:1072-1078.
28. Woodley SJ, Mercer SR. Hamstring Muscles: Architecture and Innervation. *Cells, Tissues, Organs* 2005; 179:125-141.

29. Malliaropoulos N, Isinkaye T, Tsitas K, et al. Reinjury after acute posterior thigh muscle injuries in elite track and field athletes. *Am J Sports Med* 2011; 39: 304-310.
30. Brooks JHM, Fuller CW, Kemp SPT, et al. Incidence, Risk and Prevention of Hamstring Muscle Injuries in Professional Rugby Union. *Am J Sports Med* 2006; 34:1297-1306.
31. Pieters D, Wezenbeek E, Schuermans J, et al. Return to play after a hamstring strain injury: it is time to consider natural healing. *Sports Med* 2021; <https://doi.org/10.1007/s40279-021-01494-x>.
32. Verrall GM, Slavotinek JP, Barnes PG, et al. Diagnostic and prognostic value of clinical findings in 83 athletes with posterior thigh injury: comparison of clinical findings with magnetic resonance imaging documentation of hamstring muscle strain. *Am J Sports Med* 2003;31(6):969-973
33. Buckthorpe M, Wright S, Bruce-Low S, et al. Recommendations for hamstring injury prevention in elite football: translating research into practice. *Br J Sports Med* 2019; 53:449-456.
34. Van Dyk N, Behan FP, Whiteley R. Including the Nordic hamstring exercise in injury prevention programmes halves the rate of hamstring injuries: a systematic review and meta-analysis of 8459 athletes. *Br J Sports Med* 2019; 53:1362-1370.
35. Bahr R. Why screening tests to predict injury do not work – and probably never will...: a critical review. *Br J Sports Med* 2016; 50:776-780.

Table 1. Player and injury characteristics

Variable	N	% of total
Players available	52	-
All HSI		
Players with HSI between 2014-2018	26	-
HSI	40	-
2014-2015 season	8	-
2015-2016 season	22	-
2016-2017 season	4	-
2017-2018 season	6	-
Player age (years)	26 ± 4	-
Players with >2 separate HSI from analysed 24 players	8	-
HSI excluded due to additional rehabilitation/intervention	5	-
Analysed HSI		
Players with HSI analysed	24	
HSI - TTRTP	35	
HSI – Recurrence	32	
Injury occurrence		
Match injury	25	71%
Training injury	7	20%
Rehabilitation	3	9%
Mechanism of Injury		
Sprint	13	37%
Stretch	8	23%
Insidious (gradual onset)	12	34%
During a pass	2	6%
Location of injury		
Proximal	22	63%
Central (Mid 1/3)	1	3%
Distal	11	31%
Combined proximal & distal	1	3%
Muscle injured		
Biceps Femoris (Long Head)	17	48%
Biceps Femoris (Short Head)	1	3%
Semitendinosus	5	14.2%
Semimembranosus	3	8.6%
Multiple muscles (inc. tendon injury)	9	25.7%
Multiple Muscles		
Biceps Femoris	3	33.3%
Biceps Femoris & semitendinosus	2	22.2%
Semitendinosus	1	11.1%
Semimembranosus	2	22.2%
Sacroterous Ligament	1	11.1%
Intramuscular Tendon Injury (Multiple)		
No tendon disruption	26	74%
Free tendon disruption	3	9%
Intramuscular tendon disruption	6	17%
Tendon disruption (Tendon Injury Orientation)		
Longitudinal	8	89%
Transverse	1	11%
Recurrences / Exacerbations		
No	26	74%
Yes, after RTP	6	17%
Yes, before RTP (Injured during rehabilitation)	3	9%

HSI reinjury - muscle

Biceps Femoris (Long head)	6	67%
Biceps Femoris (Short head)	0	0
Semitendinosus	1	11%
Semimembranosus	1	11%
Multiple muscles (inc. tendon injury)	1	11%

HSI reinjury - location

Proximal	7	78%
Central	0	0
Distal	2	22%

476




Figure 1. Time taken to return to play (RTP) according to British Athletics Muscle Injury Classification (BAMIC) grade of injury.

Note: The whiskers box plots represent the 25th to 75th percentile of results inside the box, the median is indicated by the horizontal line across the box, and the mean by a solid black circle. The whiskers on each box represent the 5th to 95th percentile of results.

484 **Table 2.** Predictors of RTP and injury recurrence

Predictors of TTRTP	Unstandardized beta (95%CI)	Std. Error	t	P
Constant (days)	31.4 (7.6; 55.1)	11.5	2.737	0.012
Grade	3.3 (2.1; 4.4)	0.5	5.998	<0.001
Activity	1.3 (-1.9; 4.5)	1.5	0.868	0.395
Previous injury	-2.1 (-6.9; 2.7)	2.3	-0.918	0.368
Age	-0.4 (-0.9; 0.2)	0.3	-1.296	0.209
Mechanism	-2.6 (-6.9; 1.8)	2.1	-1.234	0.230
Removed from activity	-11.0 (-15.3; -6.7)	2.1	-5.298	<0.001
Muscle injured	-0.4 (-1.6; 0.8)	0.6	-0.677	0.505
Location of injury	0.6 (-1.7; 2.9)	1.0	0.561	0.580
No. days walking with pain	-0.6 (-1.9; 0.6)	0.6	-1.078	0.293

Predictors of a reinjury occurring	Unstandardized beta (95%CI)	Std. Error	Wald	P	Adjusted OR
Constant	-2.652	3.853	0.474	0.491	0.07
Grade	-0.493	0.529	0.871	0.351	0.61
Previous injury					
<i>No (reference)</i>	-	-	-	-	-
Yes	-0.260	1.462	0.032	0.859	0.77
Age	0.150	0.158	0.896	0.344	1.16
Mechanism					
<i>Insidious (reference)</i>	-	-	-	-	-
<i>Pass</i>	1.683	1.690	0.992	0.319	5.38
<i>Stretching</i>	0.672	1.900	0.125	0.723	1.96
<i>Sprinting</i>	3.435	2.271	2.290	0.130	31.1
Removed from activity					
<i>No (reference)</i>	-	-	-	-	-
Yes	0.713	1.569	0.207	0.649	2.04
Muscle injured	-0.229	0.504	0.207	0.649	0.80
Location	-1.473	0.816	3.257	0.071	0.23
No. days walking with pain	0.779	0.407	3.672	0.055	2.18

485 *Note: TTRTP = time to return to play, OR = odds ratio, std = standard, 95%CI = 95%*

486 *confidence intervals.*