

Please cite the Published Version

Kirwan, Guy M, Broughton, Richard K, Lees, Alexander C, Ottenburghs, Jente and Tobias, Joseph A (2021) The 'Meidum geese' revisited: Early historical art is not a suitable basis for taxonomic speculation. *Journal of Archaeological Science: Reports*, 41. p. 103322. ISSN 2352-409X

DOI: <https://doi.org/10.1016/j.jasrep.2021.103322>

Publisher: Elsevier BV

Version: Accepted Version

Downloaded from: <https://e-space.mmu.ac.uk/628890/>

Usage rights: © In Copyright

Additional Information: This is an Author Accepted Manuscript of an article published in *Journal of Archaeological Science: Reports*.

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

The ‘Meidum Geese’ revisited: early historical art is not a suitable basis for taxonomic speculation

Guy M. Kirwan^{1,2}, Richard Broughton³, Alexander C. Lees⁴, Jente Ottenburghs⁵, Joseph A. Tobias⁶

¹*Bird Group, Department of Life Sciences, Natural History Museum, Tring, Herts. HP23 6AP, UK*

²*Field Museum of Natural History, 1400 South Lakeshore Drive, Chicago, IL 60605 USA*

³*UK Centre for Ecology & Hydrology, Maclean Building, Benson Lane, Crowmarsh Gifford, Wallingford, Oxfordshire OX10 8BB, UK*

⁴*Department of Natural Sciences, Manchester Metropolitan University, All Saints Building, Manchester M15 6BH, UK*

⁵*Wildlife Ecology and Conservation & Forest Ecology and Forest Management, Wageningen University and Research, Droevendaalsesteeg 3a, 6708 PB Wageningen, the Netherlands*

⁶*Department of Life Sciences, Imperial College London, Silwood Park, Ascot, Surrey SL5 7PY, UK*

Keywords:

Ancient Egypt, Avian taxonomy, Branta ruficollis, Historical artwork, Tobias criteria

ABSTRACT

Romilio (2021) used a taxonomic scoring system to compare differences between three species of geese (Anseriformes) depicted in the Chapel of Itet, one of which he speculated might represent an undescribed (presumably now extinct) species. Despite some apparently distinctive features, the depiction has traditionally been associated with the well-known modern species, red-breasted goose *Branta ruficollis*. We discuss limitations in applying the Tobias et al. (2010) scoring system to cases such as this, for which it was not designed, and we outline the many pitfalls that must be considered when attempting to identify historical artwork of birds based on examples discussed recently in the ornithological literature. We conclude that the illustrations proposed by Romero to represent a new *Branta* goose species are within the range of known plumage variation and potential artistic licence for red-breasted goose, and that this very probably is the species upon which the artwork was based. More generally, we caution against applying the Tobias criteria to cases where a series of specimens cannot be measured, and highlight the pitfalls of using illustrations to inform taxonomy.

1. Introduction

One of the most widely known examples of Ancient Egyptian art depicting three species of geese (Anseriformes) was found in the Chapel of Itet and now housed in the Egyptian Museum in Tahrir Square, Cairo. Recently, in this journal, Romilio (2021) re-examined the taxonomic assignment of these species, including two strikingly patterned birds that have usually been identified as a well-known extant species, the red-breasted goose (*Branta ruficollis*) (Nicoll 1919, Raven 1947, Houlihan 1986, Goodman & Meininger 1989, Wyatt 2013), although this was not universally accepted (Weesie 1988). Analysing the depictions using the Tobias et al. (2010) criteria for species delimitation, Romilio (2021) concluded that these two birds represent “an extinct taxon that has no modern counterpart, or a partially accurate extant but locally extinct species, or a fabricated bird that includes elements of goose morphologies.”

Although Romilio (2021) accepted the potential validity of these three different interpretations, his paper clearly suggests that the initial option, an undescribed but extinct species, was the favoured hypothesis. This eye-catching conclusion attracted considerable international media attention (Hurrell 2021), often of the more sensationalist clickbait variety (“Extinct species of goose discovered in ancient Egyptian pyramid”: www.express.co.uk/news/science). In this commentary, we focus on two potential pitfalls in the Romilio paper: misapplication of the Tobias et al. criteria, and the difficulties associated with diagnosing taxa based solely on historical images, especially a unique iconography. We go on to highlight variation in red-breasted goose phenotypes which are similar to the birds illustrated from Itet, and thereby offer a more straightforward conclusion as to the latter’s identity.

2. Historical artwork for species delimitation

The gold standard for taxonomists is the complete specimen (e.g., Collar 1999, Raposo & Kirwan 2017). Taxonomic determination can be (and frequently is) based on a single representative of an organism, the type specimen or holotype. However, a range of material that encapsulates variation, especially in morphology, provides greater confidence in the validity of a diagnosis. For example, in describing *Scytalopus petrophilus* (rock tapaculo), a new species of Rhinocryptidae from southeast Brazil, Whitney et al. (2010) nominated an additional 19 specimens as paratypes. Irrespective of the designation of paratypes, it is standard in modern ornithology to base taxonomic decisions on evidence sampled from a larger series of specimens or, in the case of vocalisations, from recordings taken from many living individuals (Tobias et al. 2010).

The validity, or otherwise, of bird species known from single specimens remains of considerable interest to ornithologists (Kirwan & Schweizer 2020). Due to inferior methods of preservation available at the time, in addition to natural disasters (e.g., fires), poor curation and unsuitable maintenance of collections, type specimens of many taxa described in the initial post-Linnaeus wave of scientific discovery and global exploration no longer exist (Sharpe 1906, Jansen 2015), despite avian taxidermy being of interest among sections of Western society since at least the 13th century (Schulze-Hagen et al. 2003).

Extant pre-19th century bird specimens in museums were recently estimated to number just 1,500–3,000, the vast majority of them in just six, well-curated European institutions (Steinheimer 2005); Gouraud (2014) highlighted the relative importance of a seventh, otherwise relatively minor collection (Baillon, La Châtre, France) for such material. Overall, the world’s museums have been estimated to house c. 9,000,000 bird specimens (Goodman & Lanyon 1994), indicating just how few (as a percentage) from the ‘Linnean and pre-Linnean eras’ survive.

Given this temporal bias, our knowledge of most taxa is reliant on relatively modern specimen material. This is often abundant for widespread species, but can be extremely limited in the case of very rare or extinct species, despite some evidence of collection bias in favour of the latter (Gotelli et al. 2021). Between at least 80 and 100 described species are generally considered to have become extinct in relatively modern times (post-1600; an arbitrary date selected by several authors for such analyses) (e.g., Fuller 2002). The number is potentially higher considering that many species probably became extinct before they could be described by scientists (e.g., Lees & Pimm 2015). For a small subset of extinct species, no specimen exists, and our knowledge of them rests solely on illustrations or descriptions by early naturalists (e.g., Fuller 2002, Wiley & Kirwan 2013).

Examples of these include the drawings of zoological novelties commissioned by Sir Stamford Raffles (Wilson 2021). In addition, several members of Captain James Cook's circumnavigatory second (1772–75) and third expeditions (1776–80) produced natural history artwork (Sawyer 1949, Stresemann 1950). In some cases, the relevant accounts and illustrations represent virtually our entire knowledge of the species they 'discovered', with consequent uncertainty surrounding their taxonomic diagnosis. For example, the Polynesian genus *Prosobonia* (Scolopacidae) is currently believed to comprise five species, of which just one is extant (Tuamotu sandpiper *P. parvirostris*), and the four extinct species are represented by just a single known specimen (Cibois et al. 2012, De Pietri et al. 2020). However, variations in both the descriptions and artwork representing some of these other species recently led Jansen et al. (2021) to postulate that Moorea sandpiper *P. ellisi* should be considered a synonym of Tahiti sandpiper *P. leucoptera* and that the originally perceived differences between these two species can be attributed to artistic freedom.

Another example is the echo parakeet (*Psittacula eques*) (Psittaciformes), an endemic of Mauritius and Réunion (Juniper & Parr 1998). Clarifying the taxonomic status of the now extinct population on the latter island, of which specimens no longer exist, is especially complicated. Although there was just one original artistic representation of the unique but now lost specimen from Réunion (Cheke & Hume 2008, Cheke & Jansen 2016), by François-Nicolas Martinet, each copy of the work in which his drawing appeared (Daubenton 1770–83) had to be coloured separately. Jansen & Cheke (2020) determined, via analysis of five different copies of Buffon's work, that the presumably different colourists responsible, none of whom presumably had access to the specimen, introduced differences in the colour of the irides, bill and peri-ocular ring, the colour and pattern of the underparts, the shape and colour of the neck-ring, and the precise colour of the wings. None of the plates examined by Jansen & Cheke (2020) precisely matched the detailed description that Brisson (1760) made of the same specimen which served as Martinet's basis for the original illustration.

Similar problems exist in relation to other birds depicted in Daubenton's work, the *Planches enluminées* (see Hume 2007), with one such image of the so-called 'Oiseau de Paradis de la Nouvelle Guinée, dit le Superbe' (*Lophorina superba*) at the heart of recent controversy caused by the apparent loss of the physical type specimen on which it is based (Elliott et al. 2020, Schodde et al. 2021).

Problems of artistic license do not only affect our knowledge of extinct species without specimen material. The very distinctive white-eyed river martin (*Pseudochelidon sirintarae*) (Hirundinidae) was described on the basis of nine specimens from Thailand in 1968 (Thonglongya 1968), but has not been seen since 1980 (Tobias 2000). In the knowledge that all records of this species were from the winter period, Dickinson (1986) suggested that a Chinese scroll painting thought to date from sometime pre-1970 was a clear depiction of this ornithological enigma, and could provide an indication of where the species might breed. However, Parkes (1987) noted that the painting was not an accurate illustration of any known species, but given the degree of concordance between the observable features might equally be identified as an Oriental pratincole (*Glareola maldivarum*), a relatively abundant species of

shorebird, and a far more parsimonious interpretation. All of these examples offer serious counterweight to the hypothesis that it is possible or even advisable to diagnose species on the basis of illustrations alone, especially unique depictions.

As further evidence of the challenges involved, Romilio (2021) reported uncertainty over the species identification of the 'grey geese' in the Chapel of Itet illustration, which may have been either greylag geese (*Anser anser*) or bean geese sp. (*Anser fabalis/serrirostris*), although their features as depicted provide no more than an imperfect match for either. Such uncertainty among the depicted birds in the Chapel of Itet underlines the risks inherent to using these illustrations as a taxonomic voucher for a suggested new species.

3. Using the Tobias et al. criteria

Romilio (2021) applied the points-based scoring system published by Tobias et al. (2010) to assess the number and extent of differences between the Meidum artwork and known species of geese. The Tobias criteria provide a flexible system with a degree of objectivity in estimating such differences, offering fresh perspective on the identity of the illustrations. However, the system was designed to be applied to samples of individuals, typically museum specimens, with a minimum of 10 individuals recommended for calculating morphological differences in plumage, body parts and vocal traits. There is nothing to prevent an author from using a pared-down version of the Tobias et al. (2010) criteria, based on scores of visual differences, as attempted by Romilio (2021). Such an approach could provide a useful 'litmus test' of morphological divergence between related taxa, to assess whether the divergence is likely to reflect species status. The main difficulty is that such an approach relies heavily on the least quantitative, and consequently most subjective, aspect of the Tobias et al. criteria.

In their complete form, the criteria place more weight on the effect sizes of quantitative differences measured directly from specimens, rather than on visual impressions. For example, to remove the subjectivity of scoring bill size differences, a series of measurements permits a taxonomist to assess whether the differences between two taxa pass a quantitative threshold linked to the scoring system (1 for a minor difference, 2 for a medium difference, etc.). Estimating differences on a single measurement alone is risky because it is difficult to know whether a unique individual is especially large or small. This risk is elevated to extremes when the distinctions are estimated on the basis of an illustration, for all the many reasons that an artist may have depicted an inaccurately large or small bill.

Another aspect of the Tobias et al. (2010) criteria that does not appear to have been applied consistently is the need to avoid covariance of characters used in the scoring system. The fact that "covariance in colour-related traits, such as a whiter belly, broader white wing-bars and a larger white rump-patch, may be driven by the same genes underlying pigmentation" means that these differences need to be collapsed into a single character to avoid inflation of scores (Tobias et al. 2010). Incautious application of the Tobias criteria may have led to different traits of the Meidum Geese being scored independently, e.g., the pattern of white on the head, neck, and breast, whereas they are patently linked.

4. Variation in Red-breasted Geese

The notion that the 'Meidum Geese' represent a novel phenotype that can be recognized as an extinct taxon appears to overlook the issue of artistic licence, but also the degree of variation that exists in red-breasted geese, which can be seen in photos of individuals of captive and wild origin (Figs. 1–4). Romilio (2021) highlighted a number of differences, of varying levels of significance, between the Chapel of Itet depictions, and illustrations and online photos of undoubted red-breasted geese. The most salient of these differences were the predominately white faces and necks, with comparatively little red coloration on these feather tracts.

In addition, Romilio (2021) suggested that the slightly different characters of the two 'Meidum Geese' provided evidence of the species 'strongly dimorphic' nature, and that the frontal of the two individuals sports three greatly elongated flank-feathers (rather than wingbars as interpreted by others, including many ornithologists). It is difficult to be absolutely definitive either way on the latter point, but what Romilio considers to be dimorphism could clearly be interpreted as individual variation. For example, none of three captive birds in one photo online (https://www.animal.photos/bird3/goos-rb_files/rbg_young.jpg) has identical face, neck and upper breast patterns, with the birds showing varying degrees of white, black and red coloration. Indeed, plumage variation appears to be quite frequent in red-breasted geese, particularly among young birds; in addition to being generally duller than adults, and more weakly patterned overall, first-calendar-year birds can have variable head and neck patterns, with more extensive white around the ear-coverts and a duller or smaller rufous cheek patch, which may be lacking completely in some birds (Carboneras et al. 2021, Cramp & Simmons 1977, Reeber 2015). Young birds also typically show less white on the flanks than adults. This is shown in an image of two captive birds that display equal levels of variation, and show even less red on the face than the 'Meidum Geese' (<https://www.dudleyzoo.org.uk/wp-content/uploads/Red-Breasted-Goose-06092017-7-e1507721913279-180x150.jpg>), similar to the illustrations in Cramp & Simmons (1977).

Further variation in the face and neck pattern of red-breasted geese is visible among a flock of wild birds on the wintering grounds in Bulgaria (<https://www.neophron.com/the-first-red-breasted-geese-are-here/>). In this example, particularly note the younger individual centre, mid-distance, showing unusual amounts of white on the face, with reduced black and red, and the bird in the right foreground, with more black on the face and neck than is typical, and a different pattern of white as a result, which also partially obscures the red of the cheeks. Among other, presumed, wild birds, an adult in Germany also appears to lack any visible red on the face (https://macaulaylibrary.org/asset/197486241#_ga=2.170536672.1563421481.1613990720-2000280272.1583423099). Romilio (2021) does not appear to have considered this degree of age-related variation shown by red-breasted geese, leading to his suggestion that the differences between the two 'Meidum Geese' must reflect sex-related plumages, which would be a key distinction from red-breasted goose. In fact, sexual dimorphism is virtually unknown among true geese (Kear 2005), which would make such variation of clear taxonomic significance, but only if proven. For now, Romilio's contention that the apparent differences between the two birds are sex-related is pure supposition.

Even if captive geese were more likely to exhibit plumage anomalies than wild birds, due to the effects of inbreeding or hybridization, the Egyptians are known to have domesticated geese (Bailleul-LeSeur 2013), making it plausible that the 'Meidum Geese' were not even wild birds. However, it seems unlikely that inbreeding or hybridization would have resulted in the phenotypes depicted on the Chapel of Itet artwork. Red-breasted geese are known to hybridize with several species, such as greater white-fronted goose (*Anser albifrons*) and barnacle goose (*Branta leucopsis*), but the resulting hybrids tend to lack the bright markings of the red-breasted goose and develop 'drab' plumage colours

instead (Ottenburghs et al. 2016, Reeber 2015). Inbreeding in captivity could have contributed to the aberrant plumage patterns of the ‘Meidum Geese’. However, this explanation is not supported by red-breasted geese in modern waterfowl collections, in which this species displays the phenotypic variation that is also observed in wild populations.

Western Egypt’s paleolandscape was considerably wetter and more mesic prior to 2400 BC, so potentially more suitable for overwintering geese as a consequence (Wendorf et al. 1973, Butzer 1977), and red-breasted goose is a known constituent of the Egyptian avifauna, albeit only a vagrant in the modern era (Meinertzhagen 1930, Goodman & Meininger 1989). With these points in mind, our review of ornithological work pertaining to taxonomic enigmas suggests that none of the three hypotheses offered by Romilio (2021) is necessarily the most parsimonious explanation. The Chapel of Itet artwork represents only a partially accurate depiction, perhaps based on the artist’s memory or a second-hand description, of a well-known modern-day species whose regular winter range could have potentially reached Egypt in prehistory when the region’s paleo landscape was far less arid and more suitable for Palearctic geese (Yeakel et al. 2014).

CRedit authorship contribution statement

G. M. Kirwan: conceptualization and preparation of first draft; **all other authors:** review and editing.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We thank the photographers Viktor Vasilev, Mladen Vasilev, Chavdar Nikolov and Valentin Katrandzhiev for supplying their images for possible use here, and Dimiter Georgiev for facilitating contact with them.

References

- Bailleul-LeSeur, R., 2013. From kitchen to table: the practical role of birds in Ancient Egypt. In: Bailleul-LeSeur, R. (ed) *Between Heaven and Earth: Birds in Ancient Egypt*. Oriental Institute Museum Publications 35. Chicago: University of Chicago Press; pp. 23–32.
- Brisson, M.J., 1760. *Ornithologie, ou méthode contenant la division des oiseaux en ordres, sections, genres, espèces et leurs variétés à laquelle on a joint une description exacte de chaque espèce*. 6 vols. Paris: J.-B. Bauche.
- Butzer, K. 1977. Geographie. In: Heick, W., Westendorf, W. (eds) *Lexicon der Ägyptologie*. Wiesbaden: Otto Harrassowitz; pp. 526–530.

- Carboneras, C., Kirwan, G.M., Sharpe, C.J. 2020. Red-breasted Goose (*Branta ruficollis*), version 1.0. In Birds of the World (J. del Hoyo, A. Elliott, J. Sargatal, D. A. Christie, E. de Juana, Eds). Cornell Lab of Ornithology, Ithaca, NY, USA. <https://doi.org/10.2173/bow.rebgoo1.01>.
- Cheke, A.S., Hume, J.P., 2008. Lost land of the Dodo. An ecological history of Mauritius, Réunion and Rodrigues. T. & A.D. Poyser, London.
- Cheke, A.S., Jansen, J.J.F.J., 2016. An enigmatic parakeet – the disputed provenance of an Indian Ocean *Psittacula*. Ibis 158 (3), 439–443.
- Collar, N.J., 1999. New species, high standards and the case of *Laniarius liberatus*. Ibis 141 (3), 358–367.
- Cramp, S., Simmons, K.E.L. (Eds.), 1977. The birds of the Western Palearctic. Volume 1. Ostrich to Ducks. Oxford: Oxford University Press.
- De Pietri, V.L., Worthy, T.H., Scofield, R.P., Cole, T.L., Wood, J.R., Cibois, A., Jansen J.J.F.J., Zhang, G., Mitchell, K.J., Feng, S., Chen, W., Tennyson, A.J.D., Wragg, G.M., 2020. A new species of Polynesian sandpiper (Charadriiformes: Scolopacidae: *Prosobonia*) from Henderson Island, Pitcairn Group, and the scolopacid affinities of *Prosobonia*. Zoological Journal of the Linnean Society, zlaa115.
- Dickinson, E.C., 1986. Does the White-eyed River-Martin *Pseudochelidon sirintarae* breed in China? Forktail 2, 95–96.
- Elliott, A., Collar, N.J., Bruce, M.D., Kirwan, G.M., 2020. The nomenclature of *Lophorina* (Aves: Paradisaeidae), with remarks on the type and type locality of *L. superba*. Zootaxa 4732 (1), 57–78.
- Fuller, E., 2002. Foreword: Extinct Birds. In: del Hoyo, J., Elliott, A., Sargatal, J. (Eds.) Handbook of the birds of the world. Volume 7. Jacamars to Woodpeckers. Barcelona: Lynx Edicions; p. 11–72.
- Goodman, S.M., Lanyon, S.M., 1994. Scientific collecting. Conservation Biology 8, 314–315.
- Goodman, S.M., Meininger, P.L., 1989. Birds of Egypt. Oxford University Press.
- Gotelli, N.J., Booher, D.B., Urban, M.C., Ulrich, W., Suarez, A.V., Skelly, D.K., Russell, D.J., Rowe, R.J., Rothendler, M., Rios, N., Rehan, S.M., Ni, G., Moreau, C.S., Magurran, A.E., Jones, F.A.M., Graves, G.R., Fiera, C., Burkhardt, U., Primack, R.B., 2021. Estimating species relative abundances from museum records. Methods in Ecology and Evolution DOI: 10.1111/2041-210X.13705.
- Gouraud, C., 2014. Eighteenth century bird specimens in the Baillon Collection, La Châtre, France. Journal of the National Museum (Prague), Natural History Series 183 (2), 19–27.
- Houlihan, P.F., 1986. The birds of Ancient Egypt. Aris and Phillips Ltd, Warminster.
- Hume, J.P., 2007. Reappraisal of the parrots (Aves: Psittacidae) from the Mascarene Islands, with comments on their ecology, morphology and affinities. Zootaxa 1513, 1–76.
- Hurrell, S., 2021. When ancient art meets modern taxonomy. World Birdwatch 44 (1), 40–41.
- Jansen, J.J.F.J., 2015. The bird collection of the Muséum national d'Histoire naturelle, Paris, France: the first years (1793-1825). Journal of the National Museum (Prague), Natural History Series 184, 81–111.

- Jansen, J.J.F.J., Cheke, A., 2020. Martinet's engravings in Buffon (1770–83): variation in their hand-colouring and its implications for defining Echo Parakeet *Psittacula eques* (Boddaert, 1783). *Bulletin of the British Ornithologists' Club* 140 (3), 299–308.
- Jansen, J.J.F.J., Kamminga, P., Argeloo, M., 2021. Taxonomic implications of the original illustrations of Prosobonia from Tahiti and Moorea made during the second and third Cook expeditions. *Bulletin of the British Ornithologists' Club* 141 (2), 133–141.
- Juniper, T., Parr, M., 1998. *Parrots: a guide to the parrots of the world*. Pica Press, Robertsbridge.
- Kear, J., (ed.) 2005. *Ducks, geese and swans. Volume 1. General chapters and species accounts (Anhima to Salvadorina)*. Oxford: Oxford University Press.
- Kirwan, G.M., Schweizer, M., 2020. Trends in systematics. Resolving the mystery of Various Nightjar and problems posed by single-specimen species. *Dutch Birding* 42 (5), 355–360.
- Lees, A.C., Pimm, S.L. 2015. Species, extinct before we know them? *Current Biology* 25 (5), R177.
- Meinertzhagen, R., 1930. *Nicoll's birds of Egypt*. London: Hugh Rees.
- Nicoll, M.J., 1919. *Handlist of the birds of Egypt*. Cairo: Government Press.
- Ottenburghs, J., van Hooft, P., van Wieren, S.E., Ydenberg, R.C. & Prins, H.H.T., 2016. Hybridization in geese: a review. *Frontiers in Zoology* 13:20.
- Parkes, K.C., 1987. Was the 'Chinese' White-eyed River-Martin an Oriental Pratincole? *Forktail* 3, 68–69.
- Raposo, M.A., Kirwan, G.M., 2017. What lies beneath the controversy as to the necessity of physical types for describing new species? *Bionomina* 12, 52–56.
- Raven, C.E., 1947. *English naturalists from Neckam to Ray. A study of the making of the modern world*. The Macmillan Company, New York.
- Reeber, S., 2015. *Wildfowl of Europe, Asia and North America*. Bloomsbury Publishing, London.
- Romilio, A., 2021. Assessing 'Meidum Geese' species identification with the 'Tobias criteria'. *Journal of Archaeological Science: Reports* 36, 102834.
- Sawyer, F., 1949. Notes on some original drawings of birds used by Dr John Latham. *Journal of the Society for the Bibliography of Natural History* 2, 173–180.
- Schodde, R., Christidis, L., Batalha-Filho, H., Ericson, P.G.P., Irestedt, M., 2021. Why neotypification of *Lophorina superba* (Pennant, 1781) (Aves: Paradisaeidae) is justified—and necessary. *Zootaxa* 4951 (2), 304–320.
- Schulze-Hagen, K., Steinheimer, F.D., Kinzelbach, R., Gasser, C., 2003. Vogelpräparation in Europa im Mittelalter und in der Renaissance. *Journal für Ornithologie* 144 (2), 459–478.
- Sharpe, R.B., 1906. *The history of the collections contained in the natural history departments of the British Museum (Birds)*. Trustees of the British Museum, London.
- Steinheimer F.D., 2005. The whereabouts of pre-nineteenth century bird specimens. *Zoologische Mededelingen (Leiden)* 79, 45–67.

- Stresemann, E., 1950. Birds collected during Capt. James Cook's last expedition (1776–1780). *Auk* 67 (1), 66–88.
- Thonglongya, A., 1968. A new martin of the genus *Pseudochelidon* from Thailand. Thai National Science Papers, Fauna Series 1.
- Tobias, J.A., 2000. Critically Threatened: White-eyed River-martin. *Oriental Bird Club Bulletin* 31, 45–48.
- Tobias, J.A., Seddon, N., Spottiswoode, C.N., Pilgrim, J.D., Fishpool, L.D.C., Collar, N.J., 2010. Quantitative criteria for species delimitation. *Ibis* 152 (4), 724–746.
- Weesie, P.D.M., 1988. The Quaternary avifauna of Crete, Greece. *Palaeo Vertebrata* 18(1).
- Wendorf, F., Schild, R., Said, R., Haynes, C.V., Gautier, A., Kobusiewicz, M., 1976. The prehistory of the Egyptian Sahara. *Science* 193 (4248), 103–114.
- Whitney, B.M., Vasconcelos, M.F., Silveira, L.F., Pacheco, J.F., 2010. *Scytalopus petrophilus* (Rock Tapaculo): a new species from Minas Gerais, Brazil. *Revista Brasileira de Ornitologia*, 18 (2), 73–88.
- Wiley, J.W., Kirwan, G.M., 2013. The extinct macaws of the West Indies, with special reference to Cuban Macaw *Ara tricolor*. *Bulletin of the British Ornithologists' Club* 133 (2), 125–156.
- Wilson, J.-J., 2021. Birds from Sumatra given by Sir Stamford Raffles to Lord Stanley: links to names, types and drawings. *Bulletin of the British Ornithologists' Club* 141 (1), 39-49.
- Wyatt, J., 2013. Birds, Pharaonic Egypt. In: Bagnall, R.S., Brodersen, K., Champion, C.B., et al. (Eds.). *The Encyclopedia of Ancient History*. First Edition ed. Blackwell Publishing Ltd; p. 1125–1127.
- Yeakel, J.D., Pires, M.M., Rudolf, L., Dominy, N.J., Koch, P.L., Guimarães, P.R., Gross, T., 2014. Collapse of an ecological network in Ancient Egypt. *Proceedings of the National Academy of Sciences* 111(40), 14472–14477.

Figs. 1–2. Mixed-species flocks of geese, predominantly red-breasted geese *Branta ruficollis*, including varying numbers of young birds, with much smaller numbers of greater white-fronted geese *Anser albifrons* photographed in Bulgaria, showing variation, much of it age-related, in the first-named species in the colour of the red feathers on the head, neck and breast, and the shape and extent of the intervening and bordering white areas, e.g., the young bird in the foreground just right of centre in Fig. 1, but many more such individuals in Fig. 2. Figs. 1 (Viktor Vasilev) and 2 (Mladen Vasilev) . Colour online.

