

# Please cite the Published Version

Morse, Christopher I, Onambele-Pearson, Gladys, Edwards, Bryn, Wong, Sze Choong and Jacques, Matthew F (2022) Objective and subjective measures of sleep in men with Muscular Dystrophy. PLoS One, 17 (9). e0274970-e0274970. ISSN 1932-6203

DOI: https://doi.org/10.1371/journal.pone.0274970

Publisher: Public Library of Science (PLoS)

Version: Published Version

Downloaded from: https://e-space.mmu.ac.uk/630438/

Usage rights: (cc) BY

Creative Commons: Attribution 4.0

**Additional Information:** This is an Open Access article which appeared in PLoS One, published by Public Library of Science

**Data Access Statement:** Data are available from the university repository: https://e-space.mmu.ac.uk/630206/.

# **Enquiries:**

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines)



# G OPEN ACCESS

**Citation:** Morse CI, Onambele-Pearson G, Edwards B, Wong SC, Jacques MF (2022) Objective and subjective measures of sleep in men with Muscular Dystrophy. PLoS ONE 17(9): e0274970. <u>https://</u>doi.org/10.1371/journal.pone.0274970

**Editor:** Pei-Lin Lee, National Taiwan University Hospital, TAIWAN

Received: March 24, 2022

Accepted: September 7, 2022

Published: September 22, 2022

**Copyright:** © 2022 Morse et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Data are available from the university repository: https://e-space. mmu.ac.uk/630206/.

**Funding:** The author(s) received no specific funding for this work.

**Competing interests:** The authors have declared that no competing interests exist.

**RESEARCH ARTICLE** 

# Objective and subjective measures of sleep in men with Muscular Dystrophy

# Christopher I. Morse<sup>1</sup>\*, Gladys Onambele-Pearson<sup>1</sup>, Bryn Edwards<sup>2</sup>, Sze Choong Wong<sup>3</sup>, Matthew F. Jacques<sup>4</sup>

1 Department of Sport and Exercise Sciences, Manchester Metropolitan University, Manchester, United Kingdom, 2 The Neuromuscular Centre, Winsford, United Kingdom, 3 Developmental Endocrinology Research Group, Royal Hospital for Children, University of Glasgow, Glasgow, United Kingdom, 4 School of Life Sciences, University of Nottingham, Nottingham, United Kingdom

\* c.morse@mmu.ac.uk

# Abstract

# Purpose

Despite poor sleep quality being recognised in Duchenne Muscular Dystrophy, reports from milder forms of Muscular Dystrophy (MD), and accompanied associations with quality of life (QoL), pain and fatigue, remain limited however.

# Methods

Adult males (n = 15 Beckers MD (BMD), n = 12 Limb-Girdle MD (LGMD), n = 12 Fascioscapulohumeral (FSHD), n = 14 non-MD (CTRL)) completed assessments of body composition (Bio-electrical impedance), sleep (7-day 24-hour tri-axial accelerometer, Pittsburgh Sleep Quality Index (PSQI) and Insomnia Severity Index, QoL (SF36-v2), pain (Visual analogue scale), fatigue (Modified Fatigue Index Scale) and functional assessments (Brookes and Vignos).

# Results

FSHD and BMD reported worse sleep than CTRL on the PSQI. FSHD scored worse than CTRL on the Insomnia Severity Index (P<0.05). 25–63% and 50–81% of adults with MD reported poor sleep quality using the Insomnia Severity Index and PSQI, respectively. Accelerometery identified no difference in sleep quality between groups. Associations were identified between sleep measures (PSQI global and insomnia severity) with mental or physical QoL in LGMD, BMD and FSHD. Multiple regression identified associations between sleep impairment and fatigue severity (all MDs), body composition (BMD & LGMD), upper and lower limb function (LGMD, FSHD) and age (FSHD).

# Conclusions

25–81% of men with MD, depending on classification, experience sleep impairment, using self-report sleep measures. Whilst BMD and FSHD showed worse sleep outcomes than CTRL, no group difference was observed between LGMD and CTRL, however all groups showed associations with sleep impairment and higher levels of fatigue. These findings, and

associations with measures of health and wellbeing, highlight an area for further research which could impact QoL in adults with MD.

### Introduction

Muscular Dystrophy (MD) is an umbrella term for a set of progressive muscle weakness conditions, for which the focus of research and interventions has been genetics and clinical characteristics, in particular muscle strength and the maintenance of ambulation in Duchenne MD (DMD) [1–4]. More recently, an increasing volume of research has been developed in order to identify influences on quality of life (QoL) and health status in other forms of MD [5–10]. The Health Status in MD model by de Groot et al. [11] acknowledges the importance of sleep disturbances along with pain and fatigue as contributing factors to QoL and health in MD. Evidence surrounding prevalence and impact of pain and fatigue on QoL is relatively wellestablished [12–15], sleep disturbances on the other hand, and the broader concept of sleep quality, remains comparably under-reported in adults with MD.

The term 'Sleep Quality' is commonly used to encompass a variety of sleep measures, including sleep time, onset of sleep, sleep efficiency and sleep disruptions [16], with poor sleep quality a defining feature of insomnia [17]. Sleep quality is known to be reduced in conditions with chronic pain [18, 19], and has been shown to result in sensations of fatigue in other clinical conditions [18]. Both pain and fatigue have been shown to be symptomatic within MD [6, 14, 20–22], indeed, our research group recently presented associations between both pain and fatigue with QoL in adults with MD [5]. Despite the presentation of pain and fatigue in MD [5, 14], well established associations with sleep quality in other clinical conditions [19], and inclusion within the Health Status in MD model [11], direct evidence of associations between sleep quality with pain, fatigue and QoL remains limited in adults with MD, with the exception of DMD [23].

In adults with DMD, poor sleep quality is linked to a combination of supine body position, severely weakened respiratory system, and the use of a night-time ventilator (indeed, severe respiratory weakness can often result in the daytime use of a ventilator also [23–28]. In comparison, reports of sleep quality within milder conditions of MD, such as Beckers MD (BMD), Limb-Girdle MD (LGMD) and Fascioscapulohumeral MD (FSHD), are limited. In LGMD and FSHD, sleep disordered breathing has been identified previously [29–31] using recall questionnaires (LGMD and FSHD) and polysomnography (FSHD) [22]. Accelerometery as a technique has been used recently to identify impaired sleep quality in children with DMD [28], however this measurement technique has not yet been adopted in adults with milder forms of MD. While understanding of sleep quality in LGMD and FSHD is limited to these aforementioned studies [29–31], further research is required to understand the prevalence of poor sleep quality in adults with BMD, LGMD and FSHD, but also its implications on Health related QoL and associations with pain and fatigue.

Therefore this research study aimed to: 1) Assess sleep quality in adults with BMD, LGMD and FSHD using 3 methods, 2) Assess the relationship between sleep quality and QoL, 3) Identify relationships between Pain, Fatigue and Participant Characteristics, and sleep quality.

#### Material and methods

Fifty-Three adult males volunteered to participate in this study (n = 15 BMD, n = 12 LGMD, n = 12 FSHD, n = 14 non-MD controls (CTRL), <u>Table 1</u>). Participants were grouped by dystrophic condition which has been confirmed by gene analysis in their referral. All participants with MD were recruited from, and tested at, The Neuromuscular Centre (Winsford, UK),

	BMD	LGMD	FSHD	CTRL
N	15	12	11	14
Age (Years)	40.7 (12.3)	43.3 (11.3)	46.3 (12.4)	37.8 (12.8)
Mass (Kg)	79.5 (13.1)	97.6 (17.3)	87.2 (12.1)	84.2 (17.9)
Stature (cm)	178.3 (5.9)	179.6 (7.2)	180.2 (8.3)	179.5 (8.0)
BMI	25.1 (3.5)	29.4 (4.5)	27.2 (3.7)	26.0 (4.0)
Body Fat (%)	26.1 (7.6) <sup>C</sup>	33.8 (4.8) <sup>F,C,B</sup>	$26.4(7.8)^{\rm C}$	18.4 (5.0)
Lean Body Mass (Kg)	58.2 (8.5) <sup>C</sup>	64.4 (8.7)	62.6 (8.9)	68.4 (12.7)
Brooks	2 (1-3)	2 (2-4)	2 (1-3)	-
Vignos	$6 (2-9)^{\rm F}$ $9 (3-9)^{\rm F}$ $3 (1-9)^{\rm F}$		3 (1-9)	-
Respiratory Support				
Night Time Only	0/15	2/12	0/11	-
24-7	0/15	0/12	0/11	-

Table 1. Participant characteristics and body composition
---

Participant Characteristics. Data presented as Mean (SD), or median (range) where appropriate. Significant differences are denoted in those furthest away from CTRL first

<sup>C</sup> denotes significantly different from CTRL

<sup>B</sup> denotes significantly different from BMD

<sup>LG</sup> denotes significantly different for LGMD

<sup>F</sup> denotes significantly different from FSHD. Kg = Kilograms; cm = centimetres; BMI = body mass index;
BMD = Beckers Muscular Dystrophy; LGMD = Limb-Girdle Muscular Dystrophy; FSHD = Fascioscapulohumeral Muscular Dystrophy; CTRL = Control.

https://doi.org/10.1371/journal.pone.0274970.t001

where they habitually participate in monthly physiotherapy sessions (Average monthly attendance =  $2 \pm 0.34$  days). CTRL participants, were recruited from the general population and free from any health conditions. CTRL participants were tested at the local university campus, using identical methods and equipment as the MD participants (with the exception of height and body mass, see below). None of the MD or CTRL participants reported any change in their activity levels or physiotherapy provision in the 3-months prior to inclusion in this study. All participants arrived for their testing in a fasted state. Ethical approval was obtained through the Sports and Exercise Science Ethics Committee, and all participants provided informed written consent prior to participation. All procedures complied with the World Medical Association Declaration of Helsinki [32].

#### Procedures

Each participant was tested in a single testing session. The same equipment was used for all participants, with the exception of seated scales for body mass measures in non-ambulatory MD participants. Anthropometric measures were performed first, followed by questionnaires for sleep quality, insomnia, quality of life, fatigue and pain were completed independently. The principal investigator was present to aid with any questions, or in some cases, to tick the desired box for participants with limited upper-limb function. Upon completion of questionnaires, an accelerometer was strapped to the wrist of the self-reported dominant arm and worn for seven consecutive days and nights (GENEActiv, Cambridge, United Kingdom).

#### Anthropometry

Control participants' mass was measured whilst standing (unshod) using digital scales (Seca model 873, Seca, Germany). MD participants were weighed in digital seated scale (6875,

Detecto, Webb City, Mo, USA). Slings, shoes, splints etc. were weighed separately and subtracted from the gross weight. All participants' stature was calculated as point to point (index finger, elbow, shoulder and across midline) to replicate a previously used method on nonambulatory participants [33, 34]. A correction of 3.5% was applied to the raw data, consistent with regression data from Caucasian males in order to account for the known discrepancy between height and arm span measures [35].

Body Mass Index (BMI) was calculated using the following equation [36]:

$$BMI\left(\frac{Kg}{m^2}\right) = Body \ Mass \ (Kg) \div Height^2 \ (m^2)$$

#### **Body composition**

Body composition measures of lean body mass (LBM) and fat mass (FM) were measured using BIA in a fasted state. Two adhesive electrodes were placed on the dorsal surfaces of the metacarpals and metatarsals of the right foot and hand. Two proximal electrodes were placed between the medial and lateral malleoli of the right ankle, and between the medial and lateral malleoli of the right radius and right ulna. BIA is a promoted method of body composition assessment in MD, given its speed and ease of use within populations that may be non-ambulant. BIA has been promoted as a measure for change in FM and LBM over time in MD [37].

#### **Functional scales**

Lower and upper limb function was assessed using common functional scales in MD, namely Brooke [38, 39] and Vignos [1] scales, respectively. The Brooke scale ranges from 1–6, whereby 1 means the participant is able to "start with arms at the sides and abduct the arms in a full circle until they tough above the head" and 6 means "Cannot raise hands to the mouth and has no function of hands" [40]. The Vignos scale ranges from 1–10, with 1 "Walk and climb stairs without assistance, and 10 "Confined to a bed" [40]. All functional scales were performed by a chartered physiotherapist, and are reported in MD participants only [14, 41].

#### Sleep assessment

**Pittsburgh sleep quality index.** The Pittsburgh Sleep Quality Index (PSQI) is a reliable ( $\alpha = 0.73-0.81$ ) [42–44] and commonly used measure of sleep quality within clinical and research settings [23, 45, 46]. The PSQI is a 21-item questionnaire, determining a global score, representing the sum of its seven domain scores: Subjective Sleep Quality, Sleep Latency, Sleep Duration, Habitual Sleep Efficiency, Sleep Disturbances, Use of Medications for Sleep and Daytime Dysfunction. Individual domains are scored on a scale of 0–3, whereby 0 denotes no difficulty, and 3 denotes severe difficulty. "Poor sleep quality" is determined from a global score of 5 or greater.

**Insomnia severity index.** The Insomnia Severity Index is a self-report measure of Sleep that has been widely used in clinical conditions [47–50], consisting of 7 items. These 7 items are: Severity of sleep-onset, Sleep Maintenance, Early morning wakening, Satisfaction with sleep pattern, interference with daily functioning, impairment attributed to sleep and distress caused by sleep [51]. Questions are on a 5-point Likert scale, from "Not at all (0) to "extremely" (4), Total scored range from 0–28, with higher scores indicative of greater insomnia severity. Total scores up to 7 is considered 'Not Clinically Significant Insomnia', scores 8–14 are considered 'Subthreshold Insomnia', Scores 15–21 are considered 'Clinical Insomnia (Moderate)', and scores 22–28 are considered 'Clinical Insomnia (Severe) [52].

**Accelerometer.** Participants wore a wrist-watch triaxial accelerometer (GENEACTIV, Kimbolton, Cambs, United Kingdom) over a consecutive 7 day period [53], which has been

reported as reliable and valid previously [54, 55]. All participants wore accelerometers on the wrist to increase adherence and remove the potential discomfort of waist-worn accelerometers for non-ambulant participants [56]. Furthermore, the use of accelerometers to assess sleep has shown 83–89% agreement with polysomnography [57, 58], and has become a more common assessment method due to participant convenience compared to polysomnography [59–62]. Wrist-worn accelerometers were worn for 24h a day on the self-reported dominant wrist of the participant, with daily physical activity reported previously by the authors [63]. Monitors were initialised to collect data at 100Hz. Once the monitors were returned, post 7-day data collection, data was downloaded from monitors into.bin files and converted in 60s epoch.csv files using the GENEACTIV PC Software (Version 2.1). 60s epoch data files were entered into a validated open source Excel Macro (v2, Activinsights Ltd.) [64]. The Excel macro reports participant bed time, rise time, total elapsed sleep time (time between bed time and rise time), sleep time (time actually asleep), Sleep Efficiency (Sleep Time/ Total Elapsed Sleep time), number of activity periods and median activity Periods and Activity Period Length for analysis.

**Pain.** A Visual Analog Scale of pain (Pain VAS) was used to quantify the level of whole body pain felt by participants over the 7 days preceding assessment. VAS is a common method of pain assessment [65] and used in many conditions [66, 67]. Participants were given a 10cm straight line, with at one end "No Pain", and the other "Worst Possible Pain", and instructed to mark where, on average, they felt their pain over the preceding 7 days was on the scale. The mark was then measured and presented as distance (cm) from the "No Pain" end.

**Fatigue.** The Modified Fatigue Index Scale is a 21-item questionnaire, was completed by participants to provide a total fatigue score (MFIS Total) or a subscore of its domains, namely Physical (MFIS Physical), Cognitive (MFIS Cognitive) or Psychosocial (MFIS Psychosocial). Participants rate the 21 questions on a 5-point Likert scale, from "Never" (0) to "Always" (4). MFIS Total is out of a possible score of 84, with scores 38 or over indicative of Fatigue.

**Quality of life.** All participants completed the SF-36v2 questionnaire, a reliable and validated measure, with eight domains of quality of life (QoL) [68, 69], for which full description is available in our research group's previous work [5]. For the purpose of the present paper, data is presented as composite Total Mental (QoL Mental) and Total Physical (QoL Physical) component scores. All data was analysed using Health Outcomes Scoring Software 4.5 (QualityMetric Health Outcomes<sup>™</sup>, Lincoln, United Kingdom).

#### Statistical analysis

All analyses were performed using IBM Statistics 26 software. The critical level of statistical significance was set at 5%. All data, except for stature and LBM, were non-parametric as determined through Levene's and Shapiro-Wilk tests. Kruskal-Wallis test was used to compare between groups with post hoc Mann-Whitney U (least significant difference) pairwise comparisons used where appropriate. Stature and LBM were compared between groups using a one-way analysis of variance (ANOVA), with Tukey's test used for post hoc pairwise comparisons. Where appropriate, bias corrected accelerated confidence intervals were calculated with 1000 bootstrap replicates [70]. Linear regressions between Sleep Quality (PSQI—Global Score; Insomnia Severity Index—Total Score; and Accelerometer—Sleep Time and Sleep Efficiency) and QoL Measures (SF36-v2 Physical and SF36-v2 Mental) were conducted. Stepwise Multiple Linear Regressions were conducted between co-variates (Pain, Fatigue and Participant Characteristics of Age, Fat Mass, Fat Free Mass, Vignos and Brookes rating) and measures of Sleep Quality (PSQI—Global Score; Insomnia Severity Index—Total Score; and Accelerometer— Sleep Time and Sleep Efficiency). ANCOVA was conducted to determine whether there were group differences in QoL (Physical and Mental) while controlling for covariables of fatigue, pain, and sleep (PSQI global) with Bonferroni post-hoc. Significant differences in tables and figures are denoted in the MD group furthest from CTRL in comparison, and thereafter from most affected to least affected MD.

# Results

# Participant characteristics

No differences were identified between groups for Age, Stature, Mass or BMI (P>0.05, Table 1). BMD (42%, P = 0.10, 95% CI [1.5, 14.1]), LGMD (84%, P<0.001, 95% CI [8.9, 22.2]) and FSHD (44%, P = 0.036, 95% CI [0.4, 14.4]) had higher body fat % than CTRL (Table 1). In addition LGMD had higher body fat % than BMD (30%, P = 0.015, 95% CI [1.2, 14.3]) and FSHD (28%, P = 0.022, 95% CI [-15.4, -0.9]). LBM was 15% lower in BMD compared to CTRL (P = 0.04, 95% CI [-20.0, -0.5], Table 1). There were no other anthropometric differences (P>0.05).

#### **Functional scales**

FSHD participants scored lower on the Vignos Scale than BMD (33%, P = 0.044, Table 1) and LGMD (67%, P = 0.002, Table 1), while no differences were identified between BMD and LGMD participants. No differences were identified between MD groups for Brooks Scale (P > 0.05, Table 1).

#### Sleep

For clarification, higher domain and global scores for PSQI, as well as Insomnia Severity Index, are indicative of worse measures of sleep.

FSHD participants scored higher in the PSQI Sleep Quality Domain than LGMD (130%, P = 0.02, 95% CI [0.12, 1.81], Fig 1). FSHD participants scored higher in the PSQI Sleep Latency Domain than CTRL (240%, P = 0.006, 95% CI [0.35, 2.62], Fig 1). BMD participants scored higher in the PSQI Sleep Efficiency Domain than CTRL (88%, P = 0.008, 95% CI [2.18, 2.52], Fig 1). No other differences were identified between groups for PSQI Domains (P > .05), however all data are presented in Fig 1.

FSHD and BMD participants scored higher on the PSQI Global Score than CTRL (88%, P = 0.011, 95% CI [0.96, 9.50] and 123%, P = 0.032, 95% CI [0.26, 7.93], respectively, Table 2). No other difference were identified between groups.

FSHD participants scored higher on the Insomnia Severity Index Total Score than CTRL participants (186%, P = 0.015, 95% CI [1.06, 13.2], Table 2). No other differences were identified between groups for Insomnia Severity Index Total Score (P>0.05, Table 2).

No differences were found between groups for any of the accelerometer determined measures of sleep time, sleep efficiency, number of activity periods or activity period length (P>0.05).

#### Pain, fatigue and quality of life

Pain rating was higher in BMD (P = 0.002, 95% CI 0.92, 5.22), LGMD (P = 0.004, 95% CI [0.84, 5.39]) and FSHD (P<0.001, 95% CI [1.18, 5.97]) compared to CTRL (Table 3). Physical fatigue (MFIS Physical) was higher in BMD (P<0.001, 95% CI [8.9, 21.0]), LGMD (P<0.001, 95% CI [10.1, 22.9]) and FSHD (P<0.001, 95% CI [-30.4, -16.9]), compared to CTRL (Table 3). BMD participants had lower physical fatigue than FSHD (P = 0.006, 95% CI [-15.4, -2.1], Table 3). Cognitive fatigue (MFIS cognitive) was higher in FSHD than CTRL (P = 0.03, -2.1]).



**Fig 1. Group scores for PSQI domains.** For clarity, differences between groups are described in text but not displayed on this Figure. https://doi.org/10.1371/journal.pone.0274970.g001

95% CI [0.6, 15.4]. Psychosocial fatigue (MFIS Psychosocial) was higher in BMD (P<0.001, 95% CI [1.74, 4.84]), LGMD (P = 0.001, 95% CI [1.30, 4.58]) and FSHD (P<0.001, 95% CI [3.33, 6.78]) than CTRL (Table 3). LGMD participants had lower psychosocial fatigue than FSHD (P = 0.014, 95% CI [-3.90, -0.33], Table 3). Overall fatigue (MFIS total) was higher in BMD (P<001, 95% CI [11.3, 37.3]), LGMD (P = 0.001, 95% CI [9.33, 36.8]) and FSHD (P<0.001, 95% CI [22.2, 51.2]) compared to CTRL (Table 3). SF36-v2 Physical was lower in BMD (27%, P<0.001, 95% CI [-27.6, -16.1]), LGMD (33%, P<0.001, 95% CI [-32.3, -20.1]) and FSHD (32%, P<0.001, 95% CI [-31.9, -19.1]) participants compared to CTRL (Table 3). No other differences were identified between groups (P>0.05). ANCOVA revealed group differences in QoL (Physical) (P<0.001) when controlling for pain, fatigue (MFIS Total), and sleep (PSQI global), between all MD groups and CTRL: BMD (P<0.001, 95% CI [-21.7, -9.2]), LGMD (P<0.001, 95% CI [-27.6, -14.1]) and FSHD (P<0.001, 95% CI [-23.1, -8.0]). There was no group or covariance effect between QoL (Mental) and pain, fatigue, and sleep.

#### Associations with sleep measures

**QoL.** Linear regression identified significant relationships for PSQI with QoL Mental in LGMD ( $R^2 = 0.593$ , P = 0.003, Table 4), and QoL Physical in BMD ( $R^2 = 0.302$ , P = 0.034, Table 4). Linear regression identified significant relationships for Insomnia Severity with QoL Mental in LGMD ( $R^2 = 0.371$ , P = 0.036, Table 4), and Physical QoL in FSHD ( $R^2 = 0.382$ , P = 0.043, Table 4). No other significant relationships were identified between QoL and Sleep Quality (P > 0.05).

#### Table 2. Sleep measures in adults with MD.

	BMD	LGMD	FSHD	CTRL
PSQI		·		·
Global Score	8.7 (4.7) <sup>C</sup>	5.5 (3.3)	10.3 (5.0) <sup>C,L</sup>	4.6 (2.0)
Prevalence of Poor Sleep Quality	80%	50%	81%	43%
Insomnia Severity Index				
Total Score	8.40 (6.1)	5.25 (5.6)	11.0 (6.7) <sup>C,L</sup>	3.85 (2.9)
No Clinically Significant Insomnia	40%	75%	36%	86%
Subthreshold Insomnia	40%	17%	36%	14%
Moderate Severity Insomnia	20%	8%	27%	-
Severe Insomnia	-	-	-	-
Accelerometer				
Sleep Time (mins) 411 (14		426 (146)	428 (86)	392 (62)
Sleep Efficiency (%)	67% (15%)	70% (17%)	70% (10%)	77% (12%)
Activity Periods (n)	12.2 (5.0)	10.6 (5.6)	10.6 (5.4)	10.5 (3.6)
Activity Period (mins)	10.8 (3.0)	13.4 (11.0)	13.1 (8.8)	8.6 (2.9)

Measures of Sleep in Adults with MD. Data presented as Mean (SD). Significant differences are denoted in those furthest away from CTRL first

 $^{\rm C}$  denotes significantly different from CTRL

<sup>B</sup> denotes significantly different from BMD

 $^{\rm LG}$  denotes significantly different for LGMD

<sup>F</sup> denotes significantly different from FSHD. mins = minutes; PSQI = Pittsburgh Sleep Quality Index; BMD = Beckers Muscular Dystrophy; LGMD = Limb-Girdle Muscular Dystrophy; FSHD = Fascioscapulohumeral Muscular Dystrophy; CTRL = Control.

https://doi.org/10.1371/journal.pone.0274970.t002

**Multiple linear regressions.** In BMD, MFIS Physical was identified as the strongest relationship with PSQI ( $R^2 = 0.588$ , P = 0.001), with all other variables excluded. Body Fat (%) was identified as the strongest relationship with Severity Index ( $R^2 = 0.358$ , P = 0.018), with all other variables excluded. No models were identified including any variables for Sleep Time or Sleep Efficiency in BMD (P>0.05).

	BMD	LGMD	FSHD	CTRL
Pain	·		·	·
VAS Pain	3.50 (2.6) <sup>C</sup>	3.54 (2.7) <sup>C</sup>	3.73 (2.2) <sup>C</sup>	0.43 (0.8)
Fatigue				
MFIS Physical	19.6 (7.4) <sup>C</sup>	21.2 (6.6) <sup>C</sup>	27.9 (4.4) <sup>C,B</sup>	4.64 (5.1)
MFIS Cognitive	11.3 (8.1)	8.83 (6.0)	12.5 (6.9) <sup>C</sup>	5.21 (5.4)
MFIS Psychosocial	3.93 (2.0) <sup>C</sup>	3.58 (1.9) <sup>F,C</sup>	5.6 (1.1) <sup>C</sup>	0.64 (0.9)
MFIS Total	34.8 (16.6) <sup>C</sup>	33.6 (12.2) <sup>C</sup>	45.9 (11.3) <sup>C</sup>	10.5 (10.8)
Quality of Life			·	
SF36-v2 Physical Score	34.7 (7.3) <sup>C</sup>	30.4 (5.4) <sup>C</sup>	32.0 (6.5) <sup>C</sup>	56.3 (3.8)
SF36-v2 Mental Score	51.0 (8.6)	57.1 (9.0)	49.5 (10.9)	56.2 (3.8)

#### Table 3. Pain, fatigue and quality of life in adults with MD.

Pain, Fatigue and Quality of Life in Adults with MD. Data presented as Mean (SD). Significant differences are denoted in those furthest away from CTRL first <sup>C</sup> denotes significantly different from CTRL

<sup>B</sup> denotes significantly different from BMD

<sup>LG</sup> denotes significantly different for LGMD

<sup>F</sup> denotes significantly different from FSHD. MFIS = Modified Fatigue Index Scale; VAS = Visual Analogue Scale; BMD = Beckers Muscular Dystrophy; LGMD = Limb-Girdle Muscular Dystrophy; FSHD = Fascioscapulohumeral Muscular Dystrophy; CTRL = Control.

https://doi.org/10.1371/journal.pone.0274970.t003

	PSQI	Insomnia Severity Index	Accelerometer	
	Global Score	Total Score	Sleep Time	Sleep Efficiency
SF36-v2 Physical	0.302 <sup>B</sup>	$0.382^{\mathrm{F}}$	-	-
SF36-v2 Mental	0.593 <sup>LG</sup>	0.371 <sup>LG</sup>	-	-

#### Table 4. Associations between sleep measures and QoL.

Associations between sleep measures and QoL measures. Data presented as R values between corresponding variables.

<sup>B</sup> denotes significant association for BMD participants

 $^{\rm LG}$  denotes significant association for LGMD participants

<sup>F</sup> denotes significant association for FSHD participants. MFIS = Modified Fatigue Index Scale; VAS = Visual Analog Scale; BMI = Body Mass Index.

https://doi.org/10.1371/journal.pone.0274970.t004

In LGMD, Brooke, Vignos, Body Fat (%) and Age were identified as the strongest relationship with Sleep Efficiency ( $R^2 = 0.952$ , P<0.001), with all other variables excluded. MFIS Total was identified as the strongest relationship with PSQI ( $R^2 = 0.332$ , P = 0.049), while MFIS Cognitive was identified as the strongest relationship with Insomnia Severity Index ( $R^2 = 0.548$ , P = 0.006), with all other variables excluded. No model was identified including any variables however for Sleep Time in LGMD (P>0.05).

In FSHD, Vignos and MFIS Physical were identified as the strongest relationship with Insomnia Severity Index ( $R^2 = 0.770$ , P = 0.003), with all other variables excluded. MFIS Physical was identified as the strongest relationship with PSQI ( $R^2 = 0.408$ , P = 0.034), with all other variables excluded. Age and MFIS Cognitive were identified as the strongest relationship with Sleep Efficiency ( $R^2 = 0.652$ , P = 0.015). No model was identified including any variables however for Sleep Time in FSHD (P>0.05).

# Discussion

This study has assessed sleep in adults with BMD, LGMD and FSHD using three separate assessments. Between 25–81% of Adults with MD, depending on classification, experience some form of sleep impairment compared to 14–43% of CTRL, using self-report measures of sleep impairment. The severity of sleep impairment in adults with MD was associated with low physical function, higher body fat, higher levels of fatigue and lower QoL. These findings, although consistent with work in children with DMD [23, 26–28], could represent a new avenue for potentially improving QoL in adults with MD.

While using accelerometery, adults with MD showed comparable sleep to CTRL, in contrast, self-reported outcomes consistently identified impaired sleep quality in Adults with MD. Of particular note, using the PSQI, poor sleep quality was noted in 80% and 81% of adults with BMD and FSHD, respectively. In comparison, Della Marca, Frusciante (30) reported 59% of adults with FSHD had poor sleep quality using the same assessment, while more recently Crescimanno et al. [23] reported 56% adults with DMD using night time ventilators had poor sleep quality, with a PSQI Global Score of 6.1, within the present study PSQI Global Score ranged from 5.5 (LGMD) to 10.3 (FSHD). The increased prevalence of impaired sleep quality in the present study, despite participants not using night-time ventilators may be surprising. It is possible that physical limitations and an impaired ability to re-position during sleep may explain these findings, given that accelerometers did not identify differences in sleep quality. This is further evidenced by the associations identified between physical function, namely the Brooks (FSHD) and Vignos (LGMD, FSHD) scales, and either the Insomnia Severity Index or sleep efficiency. It is pertinent to note, that despite being a valid measure of sleep compared with polysomnography [57, 58], there is a lack of polysomnography data from adults with MD from which we can make a comparison. Regardless, in the present study, adults with MD had a 5081% prevalence of poor sleep quality (PSQI), and 8–27% had moderate insomnia (Insomnia Severity Index), emphasising the need for further investigation into impaired sleep quality in adults with MD.

Despite experiencing low QoL, high pain and high fatigue, those with LGMD did not present differences in sleep outcomes from CTRL, despite having a 50% prevalence of participants with poor sleep quality. Of interest however was the within group associations, showing that those with poor sleep tended to have higher body fat consistent with BMD, and higher levels of fatigue (MFIS total), consistent with BMD and FSHD. The present study has therefore, provided further evidence for the health model proposed previously by de Groot et al. [11], through associations between sleep in multiple classifications of MD and QoL. In addition, associations were consistently identified between measures of body composition, function scales and fatigue with Sleep Measures in adults with MD in the present study. Previous work has reported associations between sleep disruption and respiratory impairment in LGMD [29], these respiratory impairments are likely exacerbated by body fat% and BMI associations identified in the present study, resulting in apnoea and a likely eventual need for nocturnal ventilation [30, 31]. While relationships between visceral fat and sleep apnoea are recognised within healthy populations [71], further research is required to explore the relationships between absolute adiposity and fat distribution with sleep in adults with MD [72]. Associations between sleep impairment and functional measures, fatigue and QoL in the present study are similar to those reported previously in FSHD [22]. Fatigue has become a prevalent feature of MD research [13, 73], particularly within FSHD [21, 74], whereby the current authors reported its impact on QoL across adults with MD previously [5]. Therefore, the present authors propose an updated health model for adults with MD (Fig 2), whereby independence, body composition and sleep disturbances are included and evidenced to impact QoL across adults with



Fig 2. Proposed health model for adults with muscular dystrophy, adapting the previous work from de Groot et al. [11] (White Boxes), added new components (Grey Boxes), using the present study (Denoted by [\*]) and the authors previous findings [5, 63]. Increases/improvements or decreases/impairments are denoted by + or-signs, respectively.

https://doi.org/10.1371/journal.pone.0274970.g002

MD, through our current and previous work [5, 63], as well as work evidenced in the original de Groot model [11].

Using the updated health model, it could be suggested that physical activity may be a key area to improve QoL and health status in adults with MD. The effect of increased physical activity on improvements in body composition and reducing body fat is well established within the general population [75] and recognised in MD [33, 63, 76, 77]. This may therefore reduce the effect of increased body fat percentage on sleep disturbances in adults with MD. Furthermore, Voet et al. [74] has reported previously how increased aerobic exercise resulted in decreased experienced fatigue in adults with FSHD, however the applicability across MD conditions remains unexplored. The description of Physical behaviour (during waking hours from physical activity through to sedentary behaviours) is relatively limited in adults with MD however [63, 78], future studies should consider the potential benefits of physical activity interventions on fatigue, sleep quality and QoL in adults with MD.

#### Limitations

It should be recognised that the MD population recruited for this study voluntarily attend a centre for care and rehabilitative treatments. Participants must be referred to the centre via a clinician, therefore participants may be pre-disposed to higher levels of physical dysfunction, pain or poor sleep quality, resulting in their referral. In addition, this study has used a relatively novel method of accelerometery to assess sleep quality in a clinical population [28]. Acceler-ometers allow for an objective method to be used within a home setting, rather than the intrusive nature of polysomnography [79]. Outcome measures however are estimates of sleep quality based on movement, and as our results attest, such should be treated with some caution. The present study included two self-report methods of sleep quality to supplement the objective method of assessment, and consistent in previous work in children with DMD [61], all three found associations relative to the conclusions proposed.

# Conclusion

In conclusion, this study has identified a high prevalence of impaired self-reported sleep in adults with MD, along with a wide range of associations including fatigue and QoL. These findings have helped further develop an initial health model for adults with MD by de Groot et al. [11], resulting in a more comprehensive health model suggested by the present authors. This health models proposes future interventions that can improve sleep quality, will have benefits on QoL in adults with MD, and reduce some of the burden associated with these conditions.

# **Author Contributions**

**Conceptualization:** Christopher I. Morse, Bryn Edwards, Sze Choong Wong, Matthew F. Jacques.

Data curation: Matthew F. Jacques.

Formal analysis: Gladys Onambele-Pearson, Matthew F. Jacques.

Investigation: Christopher I. Morse, Matthew F. Jacques.

Methodology: Christopher I. Morse, Gladys Onambele-Pearson, Sze Choong Wong, Matthew F. Jacques.

Project administration: Christopher I. Morse.

Supervision: Christopher I. Morse, Gladys Onambele-Pearson, Sze Choong Wong.

Writing - original draft: Christopher I. Morse, Matthew F. Jacques.

Writing - review & editing: Gladys Onambele-Pearson, Bryn Edwards, Sze Choong Wong.

#### References

- Vignos PJ, Archibald KC. Maintenance of ambulation in childhood muscular dystrophy. Journal of chronic diseases. 1960; 12(2):273–90. <u>https://doi.org/10.1016/0021-9681(60)90105-3</u> PMID: 13842210
- Hilton-Brown P, Stålberg E. Motor unit size in muscular dystrophy, a macro EMG and scanning EMG study. Journal of Neurology, Neurosurgery & Psychiatry. 1983; 46(11):996–1005. https://doi.org/10. 1136/jnnp.46.11.996 PMID: 6655485
- Williams EA, Read L, Ellis A, Morris P, Galasko CS. The management of equinus deformity in Duchenne muscular dystrophy. The Journal of bone and joint surgery British volume. 1984; 66(4):546– 50. https://doi.org/10.1302/0301-620X.66B4.6746690 PMID: 6746690
- 4. Cros D, Harnden P, Pellissier JF, Serratrice G. Muscle hypertrophy in Duchenne muscular dystrophy. Journal of neurology. 1989; 236(1):43–7. https://doi.org/10.1007/BF00314217.
- Jacques MF, Stockley RC, Onambele-Pearson GL, Reeves ND, Stebbings GK, Dawson EA, et al. Quality of life in adults with muscular dystrophy. Health and quality of life outcomes. 2019; 17(1). https:// doi.org/10.1186/s12955-019-1177-y PMID: 31307472
- Morís G, Wood L, FernáNdez-Torrón R, González Coraspe JA, Turner C, Hilton-Jones D, et al. Chronic pain has a strong impact on quality of life in facioscapulohumeral muscular dystrophy. Muscle & Nerve. 2017; 57(3):380–7. https://doi.org/10.1002/mus.25991.
- Jeronimo G, Nozoe KT, Polesel DN, Moreira GA, Tufik S, Andersen ML. Impact of corticotherapy, nutrition, and sleep disorder on quality of life of patients with Duchenne muscular dystrophy. Nutrition. 2016; 32(3):391–3. https://doi.org/10.1016/j.nut.2015.09.004 PMID: 26701140
- Wei Y, Speechley KN, Zou G, Campbell C. Factors associated with Health-Related Quality of Life in children with Duchenne muscular dystrophy. Journal of child neurology. 2016; 31(7):879–86. https:// doi.org/10.1177/0883073815627879 PMID: 26863997
- Landfeldt E, Lindgren P, Bell CF, Guglieri M, Straub V, Lochmüller H, et al. Health-related quality of life in patients with Duchenne muscular dystrophy: a multinational, cross-sectional study. Developmental Medicine & Child Neurology. 2016; 58(5):508–15. https://doi.org/10.1111/dmcn.12938.
- Messina S, Vita GL, Sframeli M, Mondello S, Mazzone E, D'Amico A, et al. Health-related quality of life and functional changes in DMD: A 12-month longitudinal cohort study. Neuromuscular Disorders. 2016; 26(3):189–96. https://doi.org/10.1016/j.nmd.2016.01.003 PMID: 26916554
- de Groot IJM, Voet NBM, Jansen M, van den Engel-Hoek L. Rehabilitation in Muscular Dystrophies: Changing Approach. The Netherlands: Radboud University Nijmegen Medical Centre. 2012; 12:235– 46.
- Abresch RT, Carter GT, Jensen MP, Kilmer DD. Assessment of pain and health-related quality of life in slowly progressive neuromuscular disease. American Journal of Hospice and Palliative Medicine®. 2002; 19(1):39–48. https://doi.org/10.1177/104990910201900109 PMID: 12173612
- Alschuler KN, Jensen MP, Goetz MC, Smith AE, Verrall AM, Molton IR. Effects of pain and fatigue on physical functioning and depression in persons with muscular dystrophy. Disability and health journal. 2012; 5(4):277–83. https://doi.org/10.1016/j.dhjo.2012.07.002 PMID: 23021739
- Jacques MF, Stockley RC, Bostock EI, Smith J, DeGoede CG, Morse CI. Frequency of reported pain in adult males with muscular dystrophy. PloS one. 2019; 14(2):e0212437. <u>https://doi.org/10.1371/journal.pone.0212437</u> PMID: 30763387
- Pangalila RF, Van Den Bos GA, Bartels B, Bergen M, Stam HJ, Roebroeck ME. Prevalence of fatigue, pain, and affective disorders in adults with Duchenne muscular dystrophy and their associations with quality of life. Archives of physical medicine and rehabilitation. 2015; 96(7):1242–7. https://doi.org/10. 1016/j.apmr.2015.02.012 PMID: 25731937
- Krystal AD, Edinger JD. Measuring sleep quality. Sleep medicine. 2008; 9:S10–S7. <u>https://doi.org/10.1016/S1389-9457(08)70011-X PMID: 18929313</u>
- Edinger JD, Bonnet MH, Bootzin RR, Doghramji K, Dorsey CM, Espie CA, et al. Derivation of research diagnostic criteria for insomnia: report of an American Academy of Sleep Medicine Work Group. Sleep. 2004; 27(8):1567–96. https://doi.org/10.1093/sleep/27.8.1567 PMID: 15683149

- Nicassio PM, Moxham EG, Schuman CE, Gevirtz RN. The contribution of pain, reported sleep quality, and depressive symptoms to fatigue in fibromyalgia. Pain. 2002; 100(3):271–9. https://doi.org/10.1016/ S0304-3959(02)00300-7 PMID: 12467998
- Sayar K, Arikan M, Yontem T. Sleep quality in chronic pain patients. The Canadian Journal of Psychiatry. 2002; 47(9):844–8. https://doi.org/10.1177/070674370204700905 PMID: 12500754
- Hoffman AJ, Jensen MP, Abresch RT, Carter GT. Chronic pain in persons with neuromuscular disease. Physical Medicine and Rehabilitation Clinics. 2005; 16(4):1099–112. <u>https://doi.org/10.1016/j.pmr</u>. 2005.08.015 PMID: 16214063
- Kalkman JS, Schillings ML, Van Der Werf SP, Padberg GW, Zwarts MJ, van Engelen BGM, et al. Experienced fatigue in facioscapulohumeral dystrophy, myotonic dystrophy, and HMSN-I. Journal of Neurology, Neurosurgery & Psychiatry. 2005; 76(10):1406–9. http://dx.doi.org/10.1136/jnnp.2004.050005.
- Kalkman JS, Schillings ML, Zwarts MJ, van Engelen BGM, Bleijenberg G. The development of a model of fatigue in neuromuscular disorders: a longitudinal study. Journal of psychosomatic research. 2007; 62(5):571–9. https://doi.org/10.1016/j.jpsychores.2006.11.014 PMID: 17467412
- Crescimanno G, Greco F, D'Alia R, Messina L, Marrone O. Quality of life in long term ventilated adult patients with Duchenne muscular dystrophy. Neuromuscular Disorders. 2019; 29(8):569–75. <a href="https://doi.org/10.1016/j.nmd.2019.06.599">https://doi.org/10.1016/j.nmd.2019.06.599</a> PMID: 31395305
- Barbe F, Quera-Salva MA, McCann C, Gajdos P, Raphael JC, De Lattre J, et al. Sleep-related respiratory disturbances in patients with Duchenne muscular dystrophy. European respiratory journal. 1994; 7 (8):1403–8. https://doi.org/10.1183/09031936.94.07081403 PMID: 7957826
- Smith PEM, Calverley PMA, Edwards RHT. Hypoxemia during sleep in Duchenne muscular dystrophy. American Journal of Respiratory and Critical Care Medicine. 1988; 137(4):884–8. https://doi.org/10. 1164/ajrccm/137.4.884 PMID: 3354996
- Suresh S, Wales P, Dakin C, Harris MA, Cooper DM. Sleep-related breathing disorder in Duchenne muscular dystrophy: disease spectrum in the paediatric population. Journal of paediatrics and child health. 2005; 41(9-10):500–3. https://doi.org/10.1111/j.1440-1754.2005.00691.x PMID: 16150067
- Smith PEM, Edwards RHT, Calverley PMA. Ventilation and breathing pattern during sleep in Duchenne muscular dystrophy. Chest. 1989; 96(6):1346–51. https://doi.org/10.1378/chest.96.6.1346 PMID: 2582842
- Siegel BI, Cakmak A, Reinertsen E, Benoit M, Figueroa J, Clifford GD, et al. Use of a wearable device to assess sleep and motor function in Duchenne muscular dystrophy. Muscle & nerve. 2020; 61 (2):198–204. https://doi.org/10.1002/mus.26759 PMID: 31725915
- Dohna-Schwake C, Ragette R, Mellies U, Straub V, Teschler H, Voit T. Respiratory function in congenital muscular dystrophy and limb girdle muscular dystrophy 2I. Neurology. 2004; 62(3):513–4. <u>https://</u> doi.org/10.1212/01.wnl.0000106937.14703.d6 PMID: 14872048
- Della Marca G, Frusciante R, Vollono C, Dittoni S, Galluzzi G, Buccarella C, et al. Sleep quality in facioscapulohumeral muscular dystrophy. Journal of the neurological sciences. 2007; 263(1–2):49–53. https://doi.org/10.1016/j.jns.2007.05.028 PMID: 17597162
- Della Marca G, Frusciante R, Dittoni S, Vollono C, Buccarella C, Iannaccone E, et al. Sleep disordered breathing in facioscapulohumeral muscular dystrophy. Journal of the neurological sciences. 2009; 285 (1–2):54–8. https://doi.org/10.1016/j.jns.2009.05.014 PMID: 19501370
- World Medical A. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. Jama. 2013; 310(20):2191. <u>https://doi.org/10.1001/jama.2013</u>. 281053 PMID: 24141714
- Jacques MF, Orme P, Smith J, Morse CI. Resting Energy Expenditure in Adults with Becker's Muscular Dystrophy. PloS one. 2017; 12(1):e0169848. https://doi.org/10.1371/journal.pone.0169848 PMID: 28060911
- Morse CI, Bostock EL, Twiss HM, Kapp LH, Orme P, Jacques MF. The cardiorespiratory response and physiological determinants of the assisted 6-minute handbike cycle test in adult males with muscular dystrophy. Muscle & nerve. 2018; 58(3):427–33. https://doi.org/10.1002/mus.26146 PMID: 29669172
- Reeves SL, Varakamin C, Henry CJ. The relationship between arm-span measurement and height with special reference to gender and ethnicity. European Journal of Clinical Nutrition. 1996; 50(6):398–400. PMID: 8793422
- McCabe MP, Ricciardelli LA, Parent P. BODY MASS INDEX. Eating Disorders: An Encyclopedia of Causes, Treatment, and Prevention. 2013; 34:90.
- Mok E, Letellier G, Cuisset J-M, Denjean A, Gottrand F, Hankard R. Assessing change in body composition in children with Duchenne muscular dystrophy: anthropometry and bioelectrical impedance analysis versus dual-energy X-ray absorptiometry. Clinical nutrition. 2010; 29(5):633–8. https://doi.org/10. 1016/j.clnu.2010.03.011 PMID: 20427103

- Brooke MH, Fenichel GM, Griggs RC, Mendell JR, Moxley R, Florence J, et al. Duchenne muscular dystrophy Patterns of clinical progression and effects of supportive therapy. Neurology. 1989; 39(4):475–. https://doi.org/10.1212/wnl.39.4.475 PMID: 2927672
- Brooke MH, Griggs RC, Mendell JR, Fenichel GM, Shumate JB, Pellegrino RJ. Clinical trial in Duchenne dystrophy. I. The design of the protocol. Muscle & Nerve: Official Journal of the American Association of Electrodiagnostic Medicine. 1981; 4(3):186–97. <u>https://doi.org/10.1002/mus.880040304</u> PMID: 7017401
- 40. Lue Y-J, Lin R-F, Chen S-S, Lu Y-M. Measurement of the functional status of patients with different types of muscular dystrophy. The Kaohsiung journal of medical sciences. 2009; 25(6):325–33. <u>https://doi.org/10.1016/S1607-551X(09)70523-6 PMID: 19560997</u>
- Archibald KC, Vignos PJ Jr. A study of contractures in muscular dystrophy. Archives of physical medicine and rehabilitation. 1959; 40(4):150–7. PMID: 13638069
- Nicassio PM, Ormseth SR, Custodio MK, Olmstead R, Weisman MH, Irwin MR. Confirmatory factor analysis of the Pittsburgh Sleep Quality Index in rheumatoid arthritis patients. Behavioral sleep medicine. 2014; 12(1):1–12. https://doi.org/10.1080/15402002.2012.720315 PMID: 23390921
- Beaudreau SA, Spira AP, Stewart A, Kezirian EJ, Lui L-Y, Ensrud K, et al. Validation of the Pittsburgh Sleep Quality Index and the Epworth Sleepiness Scale in older black and white women. Sleep medicine. 2012; 13(1):36–42. https://doi.org/10.1016/j.sleep.2011.04.005 PMID: 22033120
- 44. Beck SL, Schwartz AL, Towsley G, Dudley W, Barsevick A. Psychometric evaluation of the Pittsburgh Sleep Quality Index in cancer patients. Journal of pain and symptom management. 2004; 27(2):140–8. https://doi.org/10.1016/j.jpainsymman.2003.12.002 PMID: 15157038
- 45. Grandner MA, Kripke DF, Yoon IY, Youngstedt SD. Criterion validity of the Pittsburgh Sleep Quality Index: Investigation in a non-clinical sample. Sleep and biological rhythms. 2006; 4(2):129–36. https:// doi.org/10.1111/j.1479-8425.2006.00207.x PMID: 22822303
- Løppenthin K, Esbensen BA, Jennum P, Østergaard M, Tolver A, Thomsen T, et al. Sleep quality and correlates of poor sleep in patients with rheumatoid arthritis. Clinical rheumatology. 2015; 34(12):2029– 39. https://doi.org/10.1007/s10067-015-2875-4 PMID: 25620673
- Bahouq H, Allali F, Rkain H, Hmamouchi I, Hajjaj-Hassouni N. Prevalence and severity of insomnia in chronic low back pain patients. Rheumatology international. 2013; 33(5):1277–81. <u>https://doi.org/10.1007/s00296-012-2550-x PMID: 23124732</u>
- Savard MH, Savard J, Simard S, Ivers H. Empirical validation of the Insomnia Severity Index in cancer patients. Psycho-Oncology: Journal of the Psychological, Social and Behavioral Dimensions of Cancer. 2005; 14(6):429–41. https://doi.org/10.1002/pon.860 PMID: 15376284
- Kim J, Kim Y, Yang KI, Kim D-e, Kim SA. The relationship between sleep disturbance and functional status in mild stroke patients. Annals of rehabilitation medicine. 2015; 39(4):545. <u>https://doi.org/10.5535/arm.2015.39.4.545</u> PMID: 26361590
- Asih S, Neblett R, Mayer TG, Brede E, Gatchel RJ. Insomnia in a chronic musculoskeletal pain with disability population is independent of pain and depression. The Spine Journal. 2014; 14(9):2000–7. https://doi.org/10.1016/j.spinee.2013.11.052 PMID: 24333458
- Bastien CH, Vallières A, Morin CM. Validation of the Insomnia Severity Index as an outcome measure for insomnia research. Sleep medicine. 2001; 2(4):297–307. <u>https://doi.org/10.1016/s1389-9457(00)</u> 00065-4 PMID: 11438246
- Gagnon C, Bélanger L, Ivers H, Morin CM. Validation of the Insomnia Severity Index in primary care. The Journal of the American Board of Family Medicine. 2013; 26(6):701–10. https://doi.org/10.3122/ jabfm.2013.06.130064 PMID: 24204066
- Dillon CB, Fitzgerald AP, Kearney PM, Perry IJ, Rennie KL, Kozarski R, et al. Number of days required to estimate habitual activity using wrist-worn GENEActiv accelerometer: A cross-sectional study. PloS one. 2016; 11(5):e0109913. https://doi.org/10.1371/journal.pone.0109913 PMID: 27149674
- Rosenberger ME, Buman MP, Haskell WL, McConnell MV, Carstensen LL. 24 hours of sleep, sedentary behavior, and physical activity with nine wearable devices. Medicine and science in sports and exercise. 2016; 48(3):457. https://dx.doi.org/10.1249%2FMSS.000000000000778.
- 55. Phillips LRS, Parfitt G, Rowlands AV. Calibration of the GENEA accelerometer for assessment of physical activity intensity in children. Journal of science and medicine in sport. 2013; 16(2):124–8. https://doi.org/10.1016/j.jsams.2012.05.013 PMID: 22770768
- Arteaga D, Donnelly T, Crum K, Markham L, Killian M, Burnette WB, et al. Assessing Physical Activity Using Accelerometers in Youth with Duchenne Muscular Dystrophy. Journal of neuromuscular diseases. 2020; (Preprint):1–12.
- Cole RJ, Kripke DF, Gruen W, Mullaney DJ, Gillin JC. Automatic sleep/wake identification from wrist activity. Sleep. 1992; 15(5):461–9. https://doi.org/10.1093/sleep/15.5.461 PMID: 1455130

- Van Hees VT, Sabia S, Anderson KN, Denton SJ, Oliver J, Catt M, et al. A novel, open access method to assess sleep duration using a wrist-worn accelerometer. PLoS One. 2015; 10(11):e0142533. <a href="https://doi.org/10.1371/journal.pone.0142533">https://doi.org/10.1371/journal.pone.0142533</a> PMID: 26569414
- Verhoog S, Braun KVE, Bano A, van Rooij FJA, Franco OH, Koolhaas CM, et al. Associations of activity and sleep with quality of life: a compositional data analysis. American journal of preventive medicine. 2020; 59(3):412–9. https://doi.org/10.1016/j.amepre.2020.03.029 PMID: 32713616
- Lorenz RA, Auerbach S, Nisbet P, Sessanna L, Alanazi N, Lach H, et al. Improving sleep among adults with multiple sclerosis using mindfulness plus sleep education. Western Journal of Nursing Research. 2021; 43(3):273–83. https://doi.org/10.1177/0193945920947409 PMID: 32744186
- Crescimanno G, Greco F, Abbate A, Canino M, Bertini M, Marrone O. Subjective sleep quality in adult patients affected by Duchenne muscular dystrophy. Beyond nocturnal hypoventilation. Sleep medicine. 2020; 69:168–71. https://doi.org/10.1016/j.sleep.2020.01.025 PMID: 32092475
- van Hees VT, Sabia S, Jones SE, Wood AR, Anderson K, Trenell MI, et al. Assessment of sleep parameters from raw accelerometry data. Sleep Medicine. 2017; 40:e333.
- Jacques MF, Onambele-Pearson GL, Reeves ND, Stebbings GK, Smith J, Morse CI. Relationships between muscle size, strength, and physical activity in adults with muscular dystrophy. Journal of Cachexia, Sarcopenia and Muscle. 2018; 9(6):1042–52. <u>https://doi.org/10.1002/jcsm.12347</u> PMID: 30338901
- Esliger DW, Rowlands AV, Hurst TL, Catt M, Murray P, Eston RG. Validation of the GENEA Accelerometer. Medicine and Science in Sports and Exercise. 2011; 43(6):1085–93. https://doi.org/10.1249/MSS. 0b013e31820513be PMID: 21088628
- Price DD, McGrath PA, Rafii A, Buckingham B. The validation of visual analogue scales as ratio scale measures for chronic and experimental pain. Pain. 1983; 17(1):45–56. <u>https://doi.org/10.1016/0304-3959(83)90126-4</u> PMID: 6226917
- Moulin DE, Hagen N, Feasby TE, Amireh R, Hahn A. Pain in Guillain-Barré syndrome. Neurology. 1997; 48(2):328–31. https://doi.org/10.1212/wnl.48.2.328 PMID: 9040715
- Padua L, Aprile I, Frusciante R, Iannaccone E, Rossi M, Renna R, et al. Quality of life and pain in patients with facioscapulohumeral muscular dystrophy. Muscle & Nerve: Official Journal of the American Association of Electrodiagnostic Medicine. 2009; 40(2):200–5. <u>https://doi.org/10.1002/mus.21308</u> PMID: 19609906
- Ware JE Jr, Sherbourne CD. The MOS 36-item short-form health survey (SF-36): I. Conceptual framework and item selection. Medical care. 1992:473–83.
- Jenkinson C, Stewart-Brown S, Petersen S, Paice C. Assessment of the SF-36 version 2 in the United Kingdom. Journal of Epidemiology & Community Health. 1999; 53(1):46–50. <u>https://doi.org/10.1136/jech.53.1.46</u> PMID: 10326053
- Efron B. Better bootstrap confidence intervals. Journal of the American statistical Association. 1987; 82 (397):171–85.
- Sweatt SK, Gower BA, Chieh AY, Liu Y, Li L. Sleep quality is differentially related to adiposity in adults. Psychoneuroendocrinology. 2018; 98:46–51. https://doi.org/10.1016/j.psyneuen.2018.07.024 PMID: 30098512
- 72. Resta O, Foschino Barbaro MP, Bonfitto P, Giliberti T, Depalo A, Pannacciulli N, et al. Low sleep quality and daytime sleepiness in obese patients without obstructive sleep apnoea syndrome. Journal of internal medicine. 2003; 253(5):536–43. <u>https://doi.org/10.1046/j.1365-2796.2003.01133.x</u> PMID: 12702031
- Angelini C, Tasca E. Fatigue in muscular dystrophies. Neuromuscular Disorders. 2012; 22:S214–S20. https://doi.org/10.1016/j.nmd.2012.10.010 PMID: 23182642
- 74. Voet N, Bleijenberg G, Hendriks J, de Groot I, Padberg G, van Engelen B, et al. Both aerobic exercise and cognitive-behavioral therapy reduce chronic fatigue in FSHD An RCT. Neurology. 2014; 83 (21):1914–22. https://doi.org/10.1212/WNL.000000000001008 PMID: 25339206
- 75. Elward K, Larson EB. Benefits of exercise for older adults: a review of existing evidence and current recommendations for the general population. Clinics in geriatric medicine. 1992; 8(1):35–50. <u>https://doi.org/10.1016/S0749-0690(18)30496-8</u>.
- 76. McDonald CM, Widman LM, Walsh DD, Walsh SA, Abresch RT. Use of step activity monitoring for continuous physical activity assessment in boys with Duchenne muscular dystrophy. Archives of physical medicine and rehabilitation. 2005; 86(4):802–8. <u>https://doi.org/10.1016/j.apmr.2004.10.012</u> PMID: 15827935
- McDonald CM. Physical activity, health impairments, and disability in neuromuscular disease. American journal of physical medicine & rehabilitation. 2002; 81(11):S108–S20.

- 78. Jimenez-Moreno AC, Newman J, Charman SJ, Catt M, Trenell MI, Gorman GS, et al. Measuring Habitual Physical Activity in Neuromuscular Disorders: A Systematic Review. Journal of Neuromuscular Diseases. 2017; 4(1):25–52. https://doi.org/10.3233/JND-160195 PMID: 28269791
- 79. Jean-Louis G, Kripke DF, Cole RJ, Assmus JD, Langer RD. Sleep detection with an accelerometer actigraph: comparisons with polysomnography. Physiology & behavior. 2001; 72(1–2):21–8. https://doi. org/10.1016/s0031-9384(00)00355-3 PMID: 11239977