


**Please cite the Published Version**

Jackson, Thea, Bostock, Emma L, Hassan, Amal, Greeves, Julie P, Sale, Craig  and Elliott-Sale, Kirsty J (2022) The legacy of pregnancy: elite athletes and women in arduous occupations. *Exercise and Sport Sciences Reviews*, 50 (1). pp. 14-24. ISSN 0091-6331

**DOI:** <https://doi.org/10.1249/JES.0000000000000274>

**Publisher:** American College of Sports Medicine

**Version:** Accepted Version

**Downloaded from:** <https://e-space.mmu.ac.uk/630597/>

**Usage rights:**  [Creative Commons: Attribution-Noncommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/)

**Additional Information:** This is an Accepted Manuscript of an article which was published in *Exercise and Sport Sciences Reviews*.

**Enquiries:**

If you have questions about this document, contact [openresearch@mmu.ac.uk](mailto:openresearch@mmu.ac.uk). Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

# The Legacy of Pregnancy: Elite Athletes and Women in Arduous Occupations

Jackson T, Bostock EL, Hassan A, Greeves JP, Sale C, Elliott-Sale, KJ

## ABSTRACT

Best-practice guidance and management of pregnant and postpartum elite athletes and women in arduous occupations is limited by the lack of high-quality evidence available within these populations. We have summarized the adaptations and implications of pregnancy and childbirth, proposed a novel integrative concept to address these changes, and made recommendations to progress research in this area.

## INTRODUCTION

Motherhood has been considered incompatible with an athletic career; traditionally, sportswomen have been encouraged to end their careers to have children (1), although there are now increasing numbers of elite female athletes conceiving, giving birth, and successfully returning to sport (2). Similarly, there are many women employed in arduous occupations, such as servicewomen (11% of the UK Regular Forces are female (3)), police officers (31% of police officers in England and Wales are female (4)), firefighters (7% of all firefighters employed by Fire and Rescue Authorities England are female (5)), and manual laborers (15% of construction jobs are held by women (6)), who have returned to their physically demanding occupations after childbirth. Women in elite sport and arduous occupations must adapt training, competition, and occupational activities during and after pregnancy to minimize adverse effects on maternal and fetal health outcomes and reduce their risk of injury. Elite female athletes face many challenges when considering having a baby, planning a pregnancy within an Olympic/Paralympic cycle, time away from training and competition, profound pregnancy- and postpartum-related physiological and anatomical changes, and returning to previous levels of performance within a limited time frame (7). In 2016, the International Olympic Committee produced a five-part series providing expert opinion on these matters and highlights the need for more high-quality evidence investigating the impact of high-intensity exercise on pregnancy, childbirth, and the return to competitive sport in elite athletes (8–12). Subsequent evidence has shown that postpartum elite athletes are still unsatisfied with the training and exercise advice given to them (13), which may present practical challenges when rehabilitating in preparation for returning to sport. Since 2016, servicewomen have been fully integrated into all UK Armed Forces roles, including ground close combat, and are required to be physically and mentally fit and ready for intense physical training and deployment. Upon declaring pregnancy, servicewomen are medically downgraded, meaning they are not deployable for the duration of their pregnancy. UK servicewomen can return to physically and mentally demanding roles as early as 2 wk postpartum. As of October 1, 2020, 99% of servicewomen who went on maternity leave returned to work within 27 to 40 wk after giving birth (3). Recent data suggest that postpartum UK servicewomen are at greater risk of illness and injury in the year after giving birth compared with prepregnancy (14). The number of working days lost due to illness and injury per week was twofold higher postpartum than prepregnancy (14). Postnatal depression and gynecological disorders were the most common causes of workdays lost due to illness (14). Similarly, several studies have shown that U.S. military

servicewomen are returning to work unprepared for the demands of their role because of reduced physical fitness compared with prepregnancy, and with symptoms of depression and anxiety (15,16). Little is known about pregnancy and postpartum fitness in other arduous occupations, including police officers, firefighters, and manual laborers. The characteristics of these occupations require women to be able to perform physically strenuous activities, such as sustained heavy lifting and running. There is a dearth of evidence regarding the implications of pregnancy on the risk of injury and illness when returning to work, although physically strenuous jobs are a risk factor for pelvic floor dysfunction in the general female population (17). At 12 months postpartum, women with physically strenuous occupations experience more symptoms of pelvic organ prolapse (POP) compared with women who did not participate in heavy lifting and spent <50% of their day seated (18). Moreover, reduced abdominal muscle function (19) and incomplete recovery of the structural integrity of the linea alba observed after childbirth (20) (decreased strength (21,22), increased fatigability (22), and decreased steadiness of muscle contraction (22)) expose high-functioning postpartum women to increased injury risk, especially those who return to occupations that demand repetitive, sustained, or strenuous physical activity (19). Despite evidence to suggest that women in arduous occupations are at an increased risk of pelvic floor dysfunction and musculoskeletal injury postpartum, there are no data or guidelines on the most appropriate management of pregnant or postpartum women in arduous and manual occupations. The aim of this opinion article is to detail the changes that occur because of pregnancy, provide opinion on the current management of pregnancy in elite athletes and women employed in arduous occupations, propose a novel integrative concept to improve the legacy of pregnancy in these groups, and recommend future high-quality, impactful studies.

PREGNANCY AND POSTPARTUM ADAPTATIONS AND IMPLICATIONS

Pregnancy and childbirth cause unique physiological, musculoskeletal, and psychological adaptations that have potential implications for pregnancy and postpartum health, function, and physical activity (Table). Many of these adaptations resolve naturally after childbirth, as physiological homeostasis is restored, but some adaptations can persist and affect maternal health, posing significant challenge on physical activity participation. It is understood now that the effects of pregnancy and childbirth can extend beyond 12 months postpartum (130). The long-term health implications of returning to high-impact and intense physical activity too soon after childbirth are unknown (10).

TABLE

Adaptations and implications during pregnancy and after childbirth.

Full Size Table

Pregnancy Adaptations			Pregnancy Implications	Postpartum Adaptations
Trimester 1	Trimester 2	Trimester 3		
Hormonal				

TABLE

Adaptations and implications during pregnancy and after childbirth.

Full Size Table

	Pregnancy Adaptations			Pregnancy Implications	Postpartum Adaptations
	Trimester 1	Trimester 2	Trimester 3		
Human chorionic gonadotropin	↑ (23)	Peaks at 8–11 wk, then ↓ and levels off (23)		Stimulates the production of estrogen and progesterone within the ovary (23)	Appears to rapidly ↓ after delivery (24)
Relaxin	↑ to peak (23)	↓ (23)	Peaks again at delivery (23)	Plays a role in: ↑ cardiac output, renal blood flow, and arterial compliance ↑ joint laxity (25)	Appears to ↓ to nonpregnant values within the first week postpartum (26)
Progesterone	Gradual ↑ (23)			Maintains the uterus lining and supportive environment for the developing fetus (23)	Rapid ↓ after childbirth (27)
Estrogen	Gradual ↑ (23)			Plays a role in (23): • Maintaining uterus lining and regulating hormones essential for fetal growth • Preparing the body for breastfeeding • Enabling the uterus to respond to oxytocin in labor • ↑ venous capacitance	Rapid ↓ after childbirth (27)
Prolactin	↑ throughout and by term levels are ~10 to 20 times higher (23,28)			Preparation for lactation Progesterone blocks prolactin from exerting its effect on milk secretion (28)	Signals for milk glands to start milk production but ↓ as lactation is established (28)

TABLE

Adaptations and implications during pregnancy and after childbirth.

Full Size Table

	Pregnancy Adaptations			Pregnancy Implications	Postpartum Adaptations
	Trimester 1	Trimester 2	Trimester 3		
Oxytocin	Gradual ↑ (23)		↑ in labor (23)	↑ length, strength, and frequency of contraction in labor Keeps uterine contraction going after birth to shrink the uterus	↓ (27) Released in response to breastfeeding (29)
Physiological					
Cardiovascular	Substantial, rapid, and progressive cardiovascular system changes from ~5th week gestation (30) which ensure blood supply to the fetus: • Gradual ↑ in resting cardiac output during trimesters 1 and 2, plateaus ~20 wk gestation, and remains elevated until term (~50% over nonpregnant values by 16–20 wk gestation) (31) • ↑ resting HR throughout pregnancy (~15 bpm ↑ by term) (32) • ↑ stroke volume (~10% end of trimester 1), plateaus ~20 wk (31) Mean arterial pressure ↓ up to mid-pregnancy (~10 mm HG ↓), during trimester 3 blood pressure gradually ↑ to prepregnancy levels (33) RPE scale does not strongly correlate with HR (34) Swelling (edema) is common, often in the legs, ankles, feet, and fingers			Gestational hypertension can lead to preeclampsia in severe cases If monitoring HR during exercise, RPE should not be used as the only measure of exercise intensity	Blood pressure often ↓ straight after delivery, then ↑ to peak by 3–6 d (33) Dramatic stroke volume ↑ after delivery, ↑ cardiac output and HR remaining elevated for 24 h after birth, returning to baseline by 12 wk (31,35) Other reports that changes gradually return toward baseline but remain different from prepregnancy 1 year after birth (32)

TABLE

Adaptations and implications during pregnancy and after childbirth.

Full Size Table

	Pregnancy Adaptations			Pregnancy Implications	Postpartum Adaptations
	Trimester 1	Trimester 2	Trimester 3		
Respiratory	Elevation of the diaphragm and altered thoracic configuration (36,37) = ↓ expiratory reserve volume during 2nd half of pregnancy (8%–40% ↓ at term) = ↓ functional residual capacity (9.5%–25% at term) and ↑ inspiratory capacity to maintain constant total lung capacity (36,38,39) Sensitivity to carbon dioxide ↑ in early pregnancy (40) ↑ tidal volume + ↔ respiratory rate = ↑ minute ventilation during trimester 1 (up to 48% ↑), maintained through pregnancy (36–39,41) ↓ arterial carbon dioxide tension and ↑ arterial oxygen tension (38) Respiratory discomfort (dyspnea) often reported, trimester 3 (42)			Perception of respiratory effort and dyspnea ↓ during submaximal steady-state exercise (43)	↓ in intra-abdominal pressure allows ↑ expansion of diaphragm Respiratory changes during pregnancy return to prepregnancy levels within a few months (44)
Thermoregulation	Fetal temperature regulation is dependent upon maternal temperature, fetal metabolism, and uterine blood flow Enhanced thermoregulatory capacity as pregnancy progresses (45,46): • ↑ plasma volume • ↑ heat dissipation ↓ body temperature thresholds for sweating (46)			Moderate intensity exercise (~60%–70% V'O <sub>2</sub> max) tolerated with no significant core temperature changes in most women (47) Hyperthermia should be avoided, especially in trimester 1 (48)	Transient chill or shivering is often experienced ~15 min after birth Transient maternal temperature ↑ (up to 38°C) can be experienced in the first 24 h after delivery (49)
Glucose metabolism	Maternal blood glucose = major energy substrate for fetoplacental unit = maternal metabolism adapts to supply adequate glucose (51) Hormonal events cause: • ↑ maternal blood glucose, ↑ maternal insulin levels and liver glucose release			GDM is glucose intolerance with onset or first recognition during pregnancy (58) and can result in adverse perinatal	Women with GDM have ↑ risk for developing diabetes (59,60)

TABLE

Adaptations and implications during pregnancy and after childbirth.

Full Size Table

	Pregnancy Adaptations			Pregnancy Implications	Postpartum Adaptations
	Trimester 1	Trimester 2	Trimester 3		
	(52) • ↓ liver glycogen storage (53) = ↑ insulin resistance in skeletal muscle (54) = ↓ maternal utilization of glucose in peripheral tissues = more maternal glucose for fetal use (55) Fetoplacental unit can use up to 30%–50% of maternal glucose pool in late gestation (56) ↑ maternal insulin levels in trimester 1 = ↑ stored maternal body fat (57)			outcomes Elite athletes and those regularly exercising at a high level are less likely to have the risk factors associated with GDM	
	Recommended GWG (61):				
GWG	Prepregnancy BMI ( $\text{kg}\cdot\text{m}^{-2}$ )	Total weight gain (kg)	Weekly weight gain range in trimesters 2 and 3 (kg)	GWG is an indicator for sufficient energy intake for fetal growth and development (61) Excessive or inadequate GWG can lead to adverse pregnancy outcomes (62–66) Intensive exercise during pregnancy could cause inadequate GWG and energy intake should be adjusted accordingly	0.5 to 4 kg average weight retention 1 year after pregnancy in the general population (67,68)
	<18.5	12.5–18.0	0.44–0.58		
	18.5–24.9	11.5–16.0	0.35–0.50		
	25.0–29.9	7.0–11.5	0.23–0.33		
Energy intake	Recommended additional calorie intake across trimesters (71):				Additional ~330 kcal $\text{d}^{-1}$ in the first 6 months of lactation, if exclusively breastfeeding (72)
	Underweight: 150 kcal $\text{d}^{-1}$	200 kcal $\text{d}^{-1}$	300 kcal $\text{d}^{-1}$		

TABLE

Adaptations and implications during pregnancy and after childbirth.

Full Size Table

	Pregnancy Adaptations			Pregnancy Implications	Postpartum Adaptations
	Trimester 1	Trimester 2	Trimester 3		
Nutrition (73)	Normal weight: 0 kcal d <sup>-1</sup>	350 kcal d <sup>-1</sup>	500 kcal d <sup>-1</sup>		
	Overweight: 0 kcal d <sup>-1</sup>	450 kcal d <sup>-1</sup>	350 kcal d <sup>-1</sup>		
	Growing fetus needs to receive sufficient energy in the form of glucose During trimesters 2 and 3, protein is deposited in maternal, fetal, and placental tissues, and therefore, the recommended protein intake is increased A gestational diet should contain essential fatty acids, choline, sterols, phospholipids, and triglycerides to support fetal growth (74)			Recommendations during pregnancy (75): Carbohydrate intake: 175 g d <sup>-1</sup> Protein intake: 71 g d <sup>-1</sup> Fat intake: does not change (~20%–35% of total calories)	Carbohydrate intake important to maintain milk supply to meet baby's nutritional needs Protein needs are ↑ for breastfeeding women Dietary fats are important for baby's growth and development
Bone and calcium	In some rare cases women develop osteoporosis during pregnancy In most cases bone lost during pregnancy is recovered after childbirth or after breastfeeding ends (76)			Ensuring adequate calcium intake is important, especially if energy expenditure is high through training	Women often lose 3–5% of their bone mass breastfeeding but this usually recovers within a few months after breastfeeding ends (76)
Fatigue	Fatigue (an overwhelming sustained sense of exhaustion and decreased capacity for physical and mental work (77)) is common in pregnancy (78), particularly in trimester 1 and again in trimester 3 Severe fatigue or fatigue that lasts the entire pregnancy			Lack of rest and recovery could ↑ symptoms of fatigue Important to rule out	General lack of energy, tiredness, and irregular sleep are common Severe postpartum fatigue may be a symptom of



TABLE

Adaptations and implications during pregnancy and after childbirth.

Full Size Table

	Pregnancy Adaptations			Pregnancy Implications	Postpartum Adaptations
	Trimester 1	Trimester 2	Trimester 3		
			could be sign of a more serious condition, medical advice should be sought	hyperthyroidism or anemia	a significant medical or psychiatric illness (79,80)
Hyperemesis gravidarum			Nausea affects ~70% and vomiting ~60% of pregnant women (81) Symptoms usually begin between 4 and 7 wk gestation and disappear by 16 wk gestation (81,82) Excessive and persistent nausea and vomiting = hyperemesis gravidarum Reported incidence of 1 in 200 (82), often needs hospital treatment	Can result in: • Dehydration • Weight loss • Electrolyte imbalances • Low blood pressure when standing • Fetal growth restriction and prematurity in severe cases	None
Pelvic floor					
Anorectal		↑ progesterone levels relax smooth muscle including the digestive tract, slowing the digestion of food, may cause constipation (83)	Engagement of the fetus head in the pelvis ↑s pressure on the rectum anal sphincter descent (84)	Potential pathologies: • Fecal incontinence (85) • Defecation dysfunction including constipation (83)	Childbirth can result in: • Neuromuscular trauma (86) • Anal sphincter injury (87) • Perineal tears • Levator ani muscle Injury (88) • Pelvic floor muscle weakness (89)
Urinary		↑ bladder neck mobility and ↑ urethral mobility (90) ↑ pelvic floor distensibility (91) Uterine weight exerts pressure on and irritates the bladder (92) Reduced bladder volume from trimesters 1 to 3 (93)		Potential pathologies: • Urinary incontinence (94) • Nocturia (excessive urination at night) (95) • ↑ micturition	Childbirth can result in: • Perineal tears • Neuromuscular trauma (86) • ↑ bladder neck mobility (97) • ↑ levator hiatus distensibility (97) • Pelvic floor muscle

TABLE

Adaptations and implications during pregnancy and after childbirth.

Full Size Table

	Pregnancy Adaptations			Pregnancy Implications	Postpartum Adaptations
	Trimester 1	Trimester 2	Trimester 3		
POP		Downward movement of the anterior and posterior vaginal walls leads to descent of one or more of the pelvic organs. This is associated with: • ↑ straining associated with defecation dysfunction • ↑ peroneal body length and genital hiatus (components of the POP quantification assessment) (98)		(urination) frequency (96)  Symptoms include vaginal: • Vaginal heaviness (99) • Vaginal dragging (99) POP can: • Be asymptomatic • Impact bladder (100) and bowel function (101) • Impact quality of life (102)	weakness (89) Instrumental delivery can ↑ risk of developing pathologies (89)  Childbirth can result in: • Perineal tears • Neuromuscular trauma • ↑ levator hiatus distensibility (97) • Levator ani muscle Injury • Pelvic floor muscle weakness (89) Instrumental delivery and episiotomy can ↑ risk: 50% of postpartum women have some degree of symptomatic or asymptomatic prolapse (103)
Sexual function		Reduced sexual function is often caused by or results in (104,105): • ↑ physical discomfort as the fetus grows • ↑ fear of injury to baby • Perceived lack of attractiveness and physical awkwardness • Dyspareunia (painful intercourse) • ↓ sexual desire and sexual arousal • Orgasmic disorders		Sexual dysfunction leads to (106): • ↓ quality of life • Dissatisfaction with others • Changes in sexual and marital relationships	Factors affecting sexual function (105,107): • Lactation • Vaginal lubrication • ↓ sexual arousal and desire • Postnatal depression (108) • Dyspareunia • Orgasmic disorders

TABLE

Adaptations and implications during pregnancy and after childbirth.

Full Size Table

	Pregnancy Adaptations			Pregnancy Implications	Postpartum Adaptations
	Trimester 1	Trimester 2	Trimester 3		
Musculoskeletal					
Pelvic girdle pain	Experienced by 1 in 5 pregnant women and can be caused by (109): • A history of falls or trauma to the pelvis and/or falls during pregnancy (110) • A history of lower back pain before pregnancy (110)			May reduce function and movement, with potential implications on: • Performance • Emotional well-being • Quality of life	Often resolved postpartum but can be persistent (111) Caesarean section ↑ risk of persistent pelvic girdle pain (112)
Postural balance	None	Shift in the center of gravity causes ↓ postural balance and ↑ risk of falls (114)		2 to 3 times more likely to be injured because of falls than nonpregnant women (114)	↓ postural stability is persistent 6–8 wk after delivery (115)
Rectus diastasis	Rectus diastasis +/- protrusion or hernia (occurs in 100% of women from 37 wk onward to varying degrees) as a result of (116): • ↑ mechanical stresses placed on the abdominal wall and displacement of the abdominal organs by the growing fetus • Thinning and widening of the linea alba			Reduced function of the abdominal wall, particularly rectus abdominus function (19,20,22)	Rectus diastasis +/- protrusion or hernia may naturally resolve after childbirth (117) (39% of women 6 months postpartum have a rectus diastasis of >2 cm (116))
Breast enlargement	Can result in pain and ↑ sensitivity			Pain and discomfort may affect training and performance	Breast enlargement with lactation

TABLE

Adaptations and implications during pregnancy and after childbirth.

Full Size Table

	Pregnancy Adaptations			Pregnancy Implications	Postpartum Adaptations
	Trimester 1	Trimester 2	Trimester 3		
Gait biomechanics	None	Expanding uterus and breast enlargement: • Displaces the center of gravity • ↑ lumbar lordosis • ↑ anterior rotation of the pelvis on the femur Gait cycle changes: • ↓ length of gait cycle (120–123) • ↓ step length (120–123) • ↑ double support time (122,123) • ↑ width of step and stance (124)		Postural and biomechanical adaptations could impact performance specific tasks	Gait cycle changes early postpartum (123): • ↓ length of gait cycle • ↓ step length • ↑ double support time • ↑ width of step • ↑ width of stance
Psychological					
Depression	Symptoms: low mood, insomnia, feelings of irritability, sadness, guilt, and aversion to activity affecting thoughts, feelings, and well-being			↑ the risk of poorer birth outcomes including (125): • ↑ risk of preterm birth • Low birth weight • Preeclampsia • Intrauterine growth restriction	Postpartum depression = ↑ depressive symptoms from 0–12 months after birth (126)

TABLE

Adaptations and implications during pregnancy and after childbirth.

Full Size Table

	Pregnancy Adaptations			Pregnancy Implications	Postpartum Adaptations
	Trimester 1	Trimester 2	Trimester 3		
Anxiety	1 in 3 women experience anxiety during pregnancy (129)			Associated depressive feelings ↑ risk of poor birth outcomes (125)	Anxiety can persist after childbirth and correlates with depressive symptoms (129)

BMI, body mass index; GDM, gestational diabetes; GWG, gestational weight gain; HR, heart rate; RPE, rating of perceived exertion. [\[Context Link\]](#)

In the absence of contraindications, exercise can be beneficial during pregnancy (131), including fewer newborn complications and improved maternal health outcomes (132). The American College of Obstetricians and Gynecologists committee proposed that the type and volume of training modality should be considered between trimesters because of the normal anatomical and physiological changes ([Table](#)) that occur (133). Observational studies have shown that both elite athletes and physically active controls reduced their training volume between the first, second, and third trimesters, and none participated in high-intensity training in the final trimester (13). Although elite athletes and women in arduous roles are more likely than the general population to be physically active throughout pregnancy, they still undergo significant anatomical and physiological changes. Emerging evidence suggests that physically demanding occupations (heavy lifting, prolonged standing, prolonged walking, and prolonged bending) during pregnancy are associated with increased risk of adverse pregnancy outcomes including preterm delivery, low birth weight, small for gestational weight, and preeclampsia (134). Furthermore, the changes that occur during pregnancy may affect exercise capacity (8); however, the full extent of these implications on exercise performance is not fully understood (for a review on the state of the art, see (135)). The time taken to recover from pregnancy and childbirth can vary, dependent on gestational complications, mode of delivery (vaginal, instrumental, or cesarean section), pelvic floor dysfunction, pain, and depression. After an uncomplicated pregnancy, it is possible for women to return to low-impact physical activity in less than 12 wk after childbirth (136), and elite athletes have returned to physical activity less than 6 wk postpartum (13,137,138). Women who return to high-impact physical activity in the early postpartum period risk becoming injured without adequate rehabilitation and recovery after childbirth (13,137,139,140). Twelve weeks or more are required for the musculoskeletal soft tissues to heal after an injury, which is accounted for by the time taken to progress through the defined stages of inflammation, proliferation, and remodeling (141). Recent self-published guidelines for returning to running after childbirth suggest that symptomatic women or asymptomatic women who are without the guidance of a health care professional may benefit from waiting until 12 wk postpartum to resume high-impact activities (142). Without adequate healing and rehabilitation of the pelvic floor (143) and abdominal muscles (19–22), women are at risk of

dysfunction, which may present with (symptomatic) or without (asymptomatic) physical symptoms. If asymptomatic, it is unknown whether pelvic floor dysfunction, such as POP, could worsen because of lack of detection and intervention and become a barrier to returning to physical activity, competition, or occupational performance. Moreover, it is important to understand that the degree to which symptoms relate to function can vary between individuals, and symptomatic women will be better placed to seek help to manage these symptoms. Women may return to work or training/competition prematurely after pregnancy because of external pressure regardless of physical health or symptoms. Elite athletes are forced to consider the timing of major sporting events including the Olympic/Paralympic cycle. Taking time away from sport to have a baby may impact on ranking, team selection, or qualification to major sporting events. In addition to considering qualification or selection, elite athletes face financial worries, because paid maternity leave is uncommon. Most sports do not offer financial protection for pregnant athletes, and often, funding is withheld. Women in arduous occupations also face financial pressures as maternity pay decreases over time; for example, UK servicewomen are entitled to 26 wk' full pay, 13 wk of additional maternity leave paid at the statutory pay rate, followed by 13 wk of unpaid leave. The reductions in paid maternity leave coincide with when UK servicewomen typically return to work after childbirth (3). Moreover, women in elite sport are faced with social and psychological challenges driven by cultural beliefs and traditions regarding their responsibility to prioritize their role as a new mother over their career as an athlete. Elite female athletes often strive to meet the expectation that they can “do it all” but may experience psychological distress when they do not feel they meet the expectation of being a mother and able to perform at the highest level (7).

#### CURRENT AND PROPOSED MANAGEMENT

There is a lack of evidence available to develop best-practice guidelines for the care of pregnant and postpartum elite athletes and women employed in arduous occupations. Consequently, beyond the standard care provided by the UK National Health Service (NHS), there are disparities in the guidance and care available. For uncomplicated pregnancies, standard NHS care provides a midwife and general practitioner-led model of care, and women are offered additional care provided by specialist teams, if problems are identified (144). The management of pregnant and postpartum elite athletes is guided by the experiences of support staff and any medical professionals that form part of the wider support team (*e.g.*, sports physiotherapists) rather than by scientific evidence or clinical guidelines. As such, there is no standardized approach to managing a pregnant/postpartum elite athlete, and organizational policies make little to no reference to the support of pregnant or nursing athletes at competitive events or even for day-to-day training. Women in arduous occupations are subject to unique organizational policies, for example, there are differences in the UK approach to returning women to physical training after childbirth between the Royal Air Force, Royal Navy, and the British Army. The British Army provides women with a graduated return to fitness program for 6–12 wk after their return to work, whereas the Royal Air Force and Royal Navy do not provide a specific phased return to physical activity, unless requested by the individual. Herein, we propose an approach that could be implemented to address inconsistencies in the management of pregnant and postpartum elite athletes and women in arduous occupations using a multidisciplinary team (MDT). An MDT approach has been successfully adopted in many sporting contexts including both team and individual sports. For instance, Dijkstra *et al.* (145) describes an MDT and integrated approach to athlete health that underpinned the success of the UK track and field team at the London 2012 Olympic/Paralympic Games. An MDT approach ([Fig. 1](#)) provides a range of skills and expertise for holistic athlete health management and, consequently, achievement of performance goals. It also ensures specialist care is provided and the unique implications of pregnancy and childbirth are addressed (136). In this model, different members of an MDT are required at different stages

throughout an athlete's journey, for instance, coaches, physiotherapists, and nutritionists are required to adapt physical training, performance, and dietary needs for a pregnant athlete (Fig. 1), and the priority of care is adapted to an individual's needs and stages of pregnancy.




Figure 1

Novel integrative concept to return elite female athletes and women in arduous occupations to high-level performance after pregnancy. CV, cardiovascular; HR, heart rate; RED-S, relative energy deficiency in sport; RER, respiratory exchange ratio; RPE, rating of perceived exertion. Women in arduous occupations face similar physical and psychological demands as elite athletes, and on this basis also warrant an MDT approach to care. At present, UK servicewomen, policewomen, and firefighters access standard NHS care throughout pregnancy and after childbirth. Access to an MDT, such as specialist pelvic health physiotherapists, is not routine practice. Only in the presence of symptoms will they be able to access additional services. It is, therefore, vital that women, coaches, employers, and members of the wider MDT (Fig. 1) develop a sound understanding of the adaptations that occur during each trimester of pregnancy and after childbirth and how these changes may impact upon exercise physiology. Women also should be encouraged to evaluate their own risk for exercise participation during pregnancy, using tools such as the Canadian Society of Exercise Physiology “Get Active Questionnaire for Pregnancy” (146). Using a questionnaire such as this in combination with better awareness of pregnancy adaptations (Table), women and MDT are better placed to make informed decisions regarding training, performance, or occupational duties through pregnancy to avoid injury and complications and to facilitate postpartum recovery (Fig. 1). Contrary to traditional approaches and beliefs, it is possible to prevent common complications associated with pregnancy, such as urinary incontinence (144) and lower back pain (147). Often, women wait until after childbirth to address common complaints, meaning that they have become more established conditions and are harder to treat (148). Evidence suggests that if the intervention is intensive and early enough, pelvic floor muscle training throughout pregnancy in healthy women (with no prior pelvic floor dysfunction) may reduce the incidence of urinary incontinence and pelvic floor dysfunction in late pregnancy and the postpartum period (10,143,149). Adverse pregnancy outcomes, such as preterm birth, preeclampsia, and intrauterine growth restriction associated with physically demanding occupations, could be prevented through careful consideration of workplace guidance and strategies that reduce activities such as prolonged standing (>4 h), prolonged walking (>4 h), and heavy lifting (>100 kg·d<sup>-1</sup>) (134). The use of similar preventative approaches in elite athletes and women in arduous occupations may help limit pelvic floor dysfunction throughout pregnancy and after childbirth and reduce the risk of adverse pregnancy outcomes, although more research is needed to investigate preventative strategies within these cohorts. After an uncomplicated pregnancy, labor, and birth, individualized rehabilitation can start as early as the first

day after birth (142). Using a phased approach, such as the one proposed by Bø *et al.* (10), and updated by Deering *et al.* (136), elite athletes and women in arduous occupations should progress through three distinct postpartum phases: active recovery, training, and performance, under the supervision of an MDT (Fig. 1). The “active recovery” phase should support the transition into motherhood and address pregnancy- and birth-related concerns. To enhance continuity of care, a pelvic health physiotherapist should be employed throughout pregnancy to address the musculoskeletal complaints and pelvic floor dysfunctions identified in Table. The “training” phase should prioritize functional or sport-specific exercises in preparation for returning to competition. During the “performance” phase, the primary goal is to develop athletic performance that requires continuing communication between all members of the MDT involved. The transition between each phase is a dynamic process, whereby women can reenter phases dependent upon injury or after competition in an off-season. We suggest that a similar approach should be adopted by women in arduous occupations to ensure graded return to exercise including a progression from low- to high-impact exercise and an increase in training volume before intensity. The initial active recovery phase would not need to be altered; however, the training phase should place emphasis on occupation-specific exercise and functional tasks replicating occupational demands. The final phase should encourage occupational performance, enhancing maternal well-being and improving physical and psychological health on returning to work. The unique adaptations of pregnancy and childbirth pose a significant challenge to elite athletes and women in arduous occupations, requiring person-centered, MDT-based management. It is important to raise awareness of the changes that occur during pregnancy and the possible implications on health and performance in elite athletes and women in arduous occupations and to educate the MDT. Furthermore, preventative approaches during pregnancy aimed at improving maternal health after childbirth require further investigation within the context of elite sport and arduous occupations. Women in arduous occupations urgently need improved provision and accessibility to services to reduce injury and illness concerns upon returning to work. Overall, it should be emphasized that recovery after childbirth is not necessarily linear and women should be prepared to transition back and forth between training and performance phases.

## FUTURE PERSPECTIVES

Major gaps in our understanding exist because of the lack of studies investigating pregnancy, postpartum, and return to exercise in elite athletes and arduous occupations. It is important to acknowledge the similarities between the populations discussed in this article and use this collaborative approach to broaden the evidence base. Bø *et al.* (11) outlined important research questions that still need answering to advance the care of elite female athletes (spanning fertility; medical conditions, physiological and anatomical changes, exercise testing, athletic training, and exercise interventions) and recommended investigating the impact of strenuous exercise and the prevalence and risk factors associated with pregnancy complications. With reference to the postpartum period, they called for research to consider athletic training and exercise, breastfeeding, physiological and anatomical changes, and medical conditions. In addition to these recommendations, we suggest several other considerations to progress research and, thus, the care of elite female athletes and women in arduous occupations.

- **Point 1:** A wider MDT should be used throughout pregnancy and after childbirth to maximize recovery. Seeking and reflecting upon qualitative data from elite female athletes who have previously trained and competed through and beyond pregnancy might help shape future MDT configurations as well as timelines of care.



- **Point 2:** Pelvic health physiotherapists are well placed to evaluate specific pregnancy-related and postpartum concerns and to implement preventative strategies during, and even before, pregnancy. Further research is required to (i) assess the effectiveness of pelvic health physiotherapy interventions, such as pelvic floor muscle training during pregnancy; (ii) establish whether these interventions are required in the absence of dysfunction to improve overall performance outcomes after childbirth; and (iii) investigate whether implementation of interventions in asymptomatic women improves long-term health outcomes. A 5- to 10-year follow-up of women after these interventions would provide useful information on the long-term effectiveness of such interventions.
- **Point 3:** Swift *et al.* () showed that “early pregnancy” is a period of significant and heterogeneous behavior change, which is prejudiced by women's lived experience and their perceptions of risk. As such, the time scale for pregnancy-based interventions needs to be established to maximize the advantages of an active pregnancy.
- **Point 4:** The most effective exercise-based interventions for the active recovery phase of postpartum care needs to be investigated. We need to establish whether a combination of pelvic floor, abdominal, and general strengthening and conditioning exercises is more effective than each component in isolation.
- **Point 5:** Six weeks postpartum is an arbitrary time point at which commencing exercise is often advised and many interventions start. For active women, where time to return to activity is often critical, research that investigates rehabilitation and exercise interventions from day 1 postpartum, rather than 6 wk, is needed. A similar approach is already adopted in the immediate postoperative period for many conditions.
- **Point 6:** Observational studies of the interventions used by active women throughout pregnancy and postpartum would provide useful retrospective data for analysis to establish the most effective methods used to return to performance and competition. Furthermore, a repository of this case study-type information that is made accessible to researchers and medical professionals would improve the current state of the art.
- **Point 7:** Continued surveillance of injuries and illness incurred in the first year after childbirth would provide useful information and direction for future intervention-based studies. Similarly, documentation of training, competition, and working days lost due to injury and illness during pregnancy and after childbirth may provide useful information of the significance of the injuries and illnesses experienced.
- **Point 8:** Research is needed to demonstrate the effectiveness of an MDT approach with a structured exercise program to return women in arduous occupations to postpartum occupational fitness. Such studies should include an active recovery phase followed by a gradual increase in loading through both resistance and endurance training. Research should explore rehabilitative postpartum interventions with a focus on long-term health outcomes of UK servicewomen and incidences of illness and injury after return to work.
- **Point 9:** Sociological implications of pregnancy and childbirth need to be assessed in active women. It is imperative that we improve our understanding of the impact of pregnancy and childbirth, including maternity pay, sponsorship, pressures to secure position on a team, and pressures to be fit and ready for major sporting events or deployment.

- **Point 10:** Anecdotally, there are more mothers in elite para-sport than able-bodied sport, perhaps due, in part, to the longevity of their careers. It would be pertinent to scrutinize how para-athletes deal with their synchronous roles as mothers and Paralympians.

## Conclusion

More research is required to advance our knowledge, understanding, and quality of care of pregnant and postpartum elite athletes and women in arduous occupations. Such advancement will facilitate the safest and most timely return to employment.

## Acknowledgments

TJ and ELB are joint first authors. No funding was received for the preparation of this manuscript. CS and KJES have received funding from the UK Ministry of Defence (MOD: ASC\CMRCL\RFQ\00694, task 0157; NTU: 349408) for a study on the efficacy of a postpartum physical development programme on occupational physical performance and musculoskeletal health in UK Servicewomen. The salaries for TJ and ELB are also supported by this grant. KJES is a member of the scientific advisory board for the Active Pregnancy Foundation (Registered Charity Number – 1190780) and is an Expert for biamother.

## References

- 1

Palmer FR, Leberman SI. Elite athletes as mothers: managing multiple identities. *Sport. Manag. Rev.* 2009; 12(4):241–54. [\[Context Link\]](#)

- 2

McGannon KR, Curtin K, Schinke RJ, Schweinbenz AN. (De)constructing Paula Radcliffe: exploring media representations of elite running, pregnancy and motherhood through cultural sport psychology. *Psychol. Sport Exerc.* 2012; 13(6):820–9. [\[Context Link\]](#)

- 3

Ministry of Defence. UK Armed Forces biannual diversity statistics 1 October 2020 [Internet]. 2020 [cited 2021 November 31]. Available from: <https://www.gov.uk/government/statistics/uk-armed-forces-biannual-diversity-statistics-2020/html>. [\[Context Link\]](#)

- 4

Home Office. Police workforce: England and Wales [Internet]. 2021 [cited 2021 November 31]. Available from: [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/955182/police-workforce-mar20-hosb2020.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/955182/police-workforce-mar20-hosb2020.pdf). [\[Context Link\]](#)

- 5

Home Office. Fire and rescue workforce and pensions statistics, England, April 2019 to March 2020 [Internet]. 2020 [cited 2021 November 31]. Available from: [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/928167/fire-rescue-workforce-pensions-1920-hosb3020.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/928167/fire-rescue-workforce-pensions-1920-hosb3020.pdf). [\[Context Link\]](#)

- 6

Francis-Devine B, Ward M, Foley N. Briefing paper: women and the economy. *House Commons Libr.* 2020; CBP06838:2–18.[\[Context Link\]](#)

- 7

Roberts C-M, Kenttä G. Motherhood in the exercising female. In: Forsyth J, Roberts C-M, editors. *The Exercising Female*. New York: Taylor and Francis Group; 2018. p. 224–35.[\[Context Link\]](#)

- 8

Bø K, Artal R, Barakat R, et al. Exercise and pregnancy in recreational and elite athletes: 2016 evidence summary from the IOC expert group meeting, Lausanne. Part 1—exercise in women planning pregnancy and those who are pregnant. *Br. J. Sports Med.* 2016; 50(10):571–89.[\[Context Link\]](#)

- 9

Bø K, Artal R, Barakat R, et al. Exercise and pregnancy in recreational and elite athletes: 2016 evidence summary from the IOC expert group meeting, Lausanne. Part 2—the effect of exercise on the fetus, labour and birth. *Br. J. Sports Med.* 2016; 50(21):1297–305.[\[Context Link\]](#)

- 10

Bø K, Artal R, Barakat R, et al. Exercise and pregnancy in recreational and elite athletes: 2016/17 evidence summary from the IOC expert group meeting, Lausanne. Part 3—exercise in the postpartum period. *Br. J. Sports Med.* 2017; 51(21):1516–25.[\[Context Link\]](#)

- 11

Bø K, Artal R, Barakat R, et al. Exercise and pregnancy in recreational and elite athletes: 2016/17 evidence summary from the IOC expert group meeting, Lausanne. Part 4—recommendations for future research. *Br. J. Sports Med.* 2017; 51(24):1724–6.[\[Context Link\]](#)

- 12

Bø K, Artal R, Barakat R, et al. Exercise and pregnancy in recreational and elite athletes: 2016/2017 evidence summary from the IOC expert group meeting, Lausanne. Part 5. Recommendations for health professionals and active women. *Br. J. Sports Med.* 2018; 52(17):1080–5.[\[Context Link\]](#)

- 13

Sundgot-Borgen J, Sundgot-Borgen C, Myklebust G, Sjølvberg N, Torstveit MK. Elite athletes get pregnant, have healthy babies and return to sport early postpartum. *BMJ Open Sport Exerc. Med.* 2019; 5:e000652.[\[Context Link\]](#)

- 14

Cordell RF, Wickes CK, Casey A, Greeves JP. Female UK Army Service personnel are at greater risk of work-related morbidity on return to duty postpartum. *BMJ Mil. Heal.* 2020;jramc-2019-001282.[\[Context Link\]](#)

- 15

Weina SU. Effects of pregnancy on the army physical fitness test. *Mil. Med.* 2006; 171(6):534–7.[\[Context Link\]](#)

- 16

Rychnovsky JD. Postpartum fatigue in the active-duty military woman. *J. Obstet. Gynecol. Neonatal Nurs.* 2007; 36(1):38–46.[\[Context Link\]](#)

- 17

Hunskaar S, Burgio KL, Clark A, et al. Epidemiology of urinary (UI) and faecal (FI) incontinence and pelvic organ prolapse (POP). *ICS.* 2009; 35–112.[\[Context Link\]](#)

- 18

Kolberg Tennfjord M, Hilde G, Stær-Jensen J, Siafarikas F, Engh ME, Bø K. Effect of postpartum pelvic floor muscle training on vaginal symptoms and sexual dysfunction—secondary analysis of a randomised trial. *BJOG.* 2016; 123(4):634–42.[\[Context Link\]](#)

- 19

Deering RE, Senefeld J, Pashibin T, Neumann DA, Cruz M, Hunter SK. Fatigability of the lumbopelvic stabilizing muscles in women 8 and 26 weeks postpartum. *J. Womens Health Phys. Therap.* 2018; 42(3):128–38.[\[Context Link\]](#)

- 20

Liaw LJ, Hsu MJ, Liao CF, Liu MF, Hsu AT. The relationships between inter-recti distance measured by ultrasound imaging and abdominal muscle function in postpartum women: a 6-month follow-up study. *J. Orthop. Sports Phys. Ther.* 2011; 41(6):435–43.[\[Context Link\]](#)

- 21

Hills NF, Graham RB, McLean L. Comparison of trunk muscle function between women with and without diastasis recti abdominis at 1 year postpartum. *Phys. Ther.* 2018; 98(10):891–901.[\[Context Link\]](#)

- 22

Deering RE, Cruz M, Senefeld JW, Pashibin T, Eickmeyer S, Hunter SK. Impaired trunk flexor strength, fatigability, and steadiness in postpartum women. *Med. Sci. Sports Exerc.* 2018; 50(8):1558–69.[\[Context Link\]](#)

- 23

Tal R, Taylor HS. Endocrinology of pregnancy [Internet]. *Endotext.* 2000 [cited 2021 November 31]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25905197>.[\[Context Link\]](#)

- 24

Faiman C, Ryan RJ, Zwirek SJ, Rubin ME. Serum FSH and HCG during human pregnancy and puerperium. *J. Clin. Endocrinol. Metab.* 1968; 28(9):1323–9.[\[Context Link\]](#)

- 25

Cherni Y, Desseauve D, Decatoire A, et al. Evaluation of ligament laxity during pregnancy. *J. Gynecol. Obstet. Hum. Reprod.* 2019; 48(5):351–7.[\[Context Link\]](#)

- 26

Lafayette RA, Hladunewich MA, Derby G, Blouch K, Druzin ML, Myers BD. Serum relaxin levels and kidney function in late pregnancy with or without preeclampsia. *Clin.Nephrol.* 2011; 75(3):226–32.[\[Context Link\]](#)

- 27

Stock S, Bremme K, Uvnäs-Moberg K. Plasma levels of oxytocin during the menstrual cycle, pregnancy and following treatment with HMG. *Hum. Reprod.* 1991; 6(8):1056–62.[\[Context Link\]](#)

- 28

Ostrom KM. A review of the hormone prolactin during lactation. *Prog. Food vNutr. Sci.* 1990; 14(1):1–44.[\[Context Link\]](#)

- 29

Moberg KU, Ekström-Bergström A, Buckley S, et al. Maternal plasma levels of oxytocin during breastfeeding—a systematic review. *PLoS One.* 2020; 15(8):e0235806.[\[Context Link\]](#)

- 30

Duvekot JJ, Cheriex EC, Pieters FA, Menheere PP, Peeters LH. Early pregnancy changes in hemodynamics and volume homeostasis are consecutive adjustments triggered by a primary fall in systemic vascular tone. *Am. J. Obstet. Gynecol.* 1993; 169(6):1382–92.[\[Context Link\]](#)

- 31

Hunter S, Robson SC. Adaptation of the maternal heart in pregnancy. *Br. Heart J.* 1992; 68(6):540–3.[\[Context Link\]](#)

- 32

Clapp JF 3rd, Capeless E. Cardiovascular Function before, during, and after the first and subsequent pregnancies. *Am. J. Cardiol.* 1997; 80(11):1469–73.[\[Context Link\]](#)

- 33

Bramham K, Nelson-Piercy C, Brown MJ, Chappell LC. Postpartum management of hypertension. *BMJ.* 2013; 346:f894.[\[Context Link\]](#)

- 34

O'Neill ME, Cooper KA, Mills CM, Boyce ES, Hunyor SN. Accuracy of Borg's ratings of perceived exertion in the prediction of heart rates during pregnancy. *Br. J. Sports Med.* 1992; 26(2):121–4.[\[Context Link\]](#)

- 35

Atkins AFJ, Watt JM, Milan P, Davies P, Crawford JS. A longitudinal study of cardiovascular dynamic changes throughout pregnancy. *Eur. J. Obstet. Gynecol. Reprod. Biol.* 1981; 12(4):215–24.[\[Context Link\]](#)

- 36

Gilroy RJ, Mangura BT, Laviates MH. Rib Cage and Abdominal Volume Displacements during Breathing in Pregnancy]. *Am. Rev. Respir. Dis.* 1988; 137(3):668–72.[\[Context Link\]](#)

- 37

Contreras G, Gutiérrez M, Beroiza T, et al. Ventilatory Drive and Respiratory Muscle Function in Pregnancy. *Am. Rev. Respir. Dis.* 1991; 144(4):837–41. [\[Context Link\]](#)

- 38

Weinberger SE, Weiss ST, Cohen WR, Weiss JW, Johnson TS. Pregnancy and The Lung. *Am. Rev. Respir. Dis.* 1980; 121(3):559–81. [\[Context Link\]](#)

- 39

Jensen D, Webb KA, Davies GAL, O'Donnell DE. Mechanical ventilatory constraints during incremental cycle exercise in human pregnancy: implications for respiratory sensation. *J. Physiol.* 2008; 586(19):4735–50. [\[Context Link\]](#)

- 40

Weissgerber TL, Wolfe LA, Hopkins WG, Davies GAL. Serial respiratory adaptations and an alternate hypothesis of respiratory control in human pregnancy. *Respir. Physiol. Neurobiol.* 2006; 153(1):39–53. [\[Context Link\]](#)

- 41

Kolarzyk E, Szot WM, Lyszczarz J. Lung function and breathing regulation parameters during pregnancy. *Arch. Gynecol. Obstet.* 2005; 272(1):53–8. [\[Context Link\]](#)

- 42

Milne JA, Howie AD, Pack AI. Dyspnoea during normal pregnancy. *BJOG An. Int. J. Obstet. Gynaecol.* 1978; 85(4):260–3. [\[Context Link\]](#)

- 43

Field SK, Bell SG, Cenaiko DF, Whitelaw WA. Relationship between inspiratory effort and breathlessness in pregnancy. *J. Appl. Physiol.* 1991; 71(5):1897–902. [\[Context Link\]](#)

- 44

Datta S, Kodali B, Segal S. Maternal Physiological Changes During Pregnancy, Labor, and the Postpartum Period. In: *Obstetric Anesthesia Handbook*. 5th Edition. 2010. p. 1–14. [\[Context Link\]](#)

- 45

Lindqvist PG, Marsal K, Merlo J, Pirhonen JP. Thermal response to submaximal exercise before, during and after pregnancy: a longitudinal study. *J. Matern. Neonatal. Med.* 2003; 13(3):152–6. [\[Context Link\]](#)

- 46

Clapp JF. The changing thermal response to endurance exercise during pregnancy. *Am. J. Obstet. Gynecol.* 1991; 165(6 Pt 1):1684–9. [\[Context Link\]](#)

- 47

Soultanakis HN, Artal R, Wiswell RA. Prolonged exercise in pregnancy: glucose homeostasis, ventilatory and cardiovascular responses. *Semin. Perinatol.* 1996; 20(4):315–27. [\[Context Link\]](#)

- 48

Chambers CD. Risks of hyperthermia associated with hot tub or spa use by pregnant women. ***Birth. Defects Res. A. Clin. Mol. Teratol.*** 2006; 76(8):569–73.[\[Context Link\]](#)

- 49

Ely JW, Dawson JD, Townsend AS, Rijhsinghani A, Bowdler NC. Benign fever following vaginal delivery. ***J. Fam. Pract.*** 1996; 43(2):146–51.[\[Context Link\]](#)

- 50

World Health Organisation. International Classification of Diseases (ICD) Information Sheet. ***WHO [Internet]***. 2015 [cited 2021 November 31]; (May 1990):10–1. Available from: <http://www.who.int/classifications/icd/factsheet/en/%0Ahttps://www.who.int/classifications/icd/factsheet/en/%0Ahttp://www.who.int/classifications/icd/factsheet/en/>.[\[Context Link\]](#)

- 51

Bauer MK, Harding JE, Bassett NS, et al. Fetal growth and placental function. ***Mol. Cell. Endocrinol.*** 1998; 140(1–2):115–20.[\[Context Link\]](#)

- 52

Mottola MF, Christopher PD. Effects of maternal exercise on liver and skeletal muscle glycogen storage in pregnant rats. ***J. Appl. Physiol.*** 1991; 71(3):1015–9.[\[Context Link\]](#)

- 53

Catalano PM, Tyzbir ED, Roman NM, Amini SB, Sims EA. Longitudinal changes in insulin release and insulin resistance in nonobese pregnant women. ***Am. J. Obstet. Gynecol.*** 1991; 165(6 Pt 1):1667–72.[\[Context Link\]](#)

- 54

Buchanan TA, Metzger BE, Freinkel N, Bergman RN. Insulin sensitivity and B-cell responsiveness to glucose during late pregnancy in lean and moderately obese women with normal glucose tolerance or mild gestational diabetes. ***Am. J. Obstet. Gynecol.*** 1990; 162(4):1008–14.[\[Context Link\]](#)

- 55

Lesser KB, Carpenter MW. Metabolic changes associated with normal pregnancy and pregnancy complicated by diabetes mellitus. ***Semin. Perinatol.*** 1994; 18(5):399–406.[\[Context Link\]](#)

- 56

Clapp JF III. Maternal carbohydrate intake and pregnancy outcome. ***Proc. Nutr. Soc.*** 2002; 61(1):45–50.[\[Context Link\]](#)

- 57

Boden G. Fuel metabolism in pregnancy and in gestational diabetes mellitus. ***Obstet. Gynecol. Clin. North Am.*** 1996; 23(1):1–10.[\[Context Link\]](#)

- 58

Metzger BE, Coustan DR. Summary and recommendations of the Fourth International Workshop-Conference on Gestational Diabetes Mellitus. The Organizing Committee. *Diabetes Care*. 1998; 21(Suppl. 2):B161–7.[\[Context Link\]](#)

- 59

Kaaja R, Rönnemaa T. Gestational diabetes: pathogenesis and consequences to mother and offspring. *Rev. Diabet. Stud.* 2008; 5(4):194–202.[\[Context Link\]](#)

- 60

Lauenborg J, Hansen T, Jensen DM, et al. Increasing incidence of diabetes after gestational diabetes: a long-term follow-up in a Danish population. *Diabetes Care*. 2004; 27(5):1194–9.[\[Context Link\]](#)

- 61

Institute of Medicine (US) and National Research Council (US) Committee to Reexamine IOM Pregnancy Weight Guidelines, Rasmussen KM, Yaktine AL, editors. In: *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC: The National Academies Press; 2009.[\[Context Link\]](#)

- 62

Siege-Riz AM, Viswanathan M, Moos M-K, et al. A systematic review of outcomes of maternal weight gain according to the Institute of Medicine recommendations: birthweight, fetal growth, and postpartum weight retention. *Am. J. Obstet. Gynecol.* 2009; 201(4):339.e1–339.e14.[\[Context Link\]](#)

- 63

Alavi N, Haley S, Chow K, McDonald SD. Comparison of national gestational weight gain guidelines and energy intake recommendations. *Obes. Rev.* 2013; 14(1):68–85.[\[Context Link\]](#)

- 64

Artal R, Lockwood CJ, Brown HL. Weight gain recommendations in pregnancy and the obesity epidemic. *Obstet. Gynecol.* 2010; 115(1):152–5.[\[Context Link\]](#)

- 65

Crane JMG, White J, Murphy P, Burrage L, Hutchens D. The effect of gestational weight gain by body mass index on maternal and neonatal outcomes. *J. Obstet. Gynaecol. Can.* 2009; 31(1):28–35.[\[Context Link\]](#)

- 66

Macdonald-Wallis C, Tilling K, Fraser A, Nelson SM, Lawlor DA. Gestational weight gain as a risk factor for hypertensive disorders of pregnancy. *Am. J. Obstet. Gynecol.* 2013; 209(4):327.e1–17.[\[Context Link\]](#)

- 67

Linné Y, Barkeling B, Rössner S. Long-term weight development after pregnancy. *Obes. Rev.* 2002; 3(2):75–83.[\[Context Link\]](#)

- 68



Olson CM, Strawderman MS, Hinton PS, Pearson TA. Gestational weight gain and postpartum behaviors associated with weight change from early pregnancy to 1 y postpartum. *Int. J. Obes.* 2003; 27(1):117–27. [\[Context Link\]](#)

- 69

Lim CC, Mahmood T. Obesity in pregnancy. *Best Pract. Res. Clin. Obstet. Gynaecol.* 2015; 29(3):309–19. [\[Context Link\]](#)

- 70

Rooney BL, Schauburger CW, Mathiason MA. Impact of perinatal weight change on long-term obesity and obesity-related illnesses. *Obstet. Gynecol.* 2005; 106(6):1349–56. [\[Context Link\]](#)

- 71

Butte NF, Wong WW, Treuth MS, Ellis KJ, O'Brian Smith E. Energy requirements during pregnancy based on total energy expenditure and energy deposition. *Am. J. Clin. Nutr.* 2004; 79(6):1078–87. [\[Context Link\]](#)

- 72

Reader D, Franz MJ. Lactation, diabetes, and nutrition recommendations [Internet]. *Curr. Diab. Rep.* 2004; 4(5):370–6. [\[Context Link\]](#)

- 73

Elliott-Sale KJ, Graham A, Hanley SJ, Blumenthal S, Sale C. Modern dietary guidelines for healthy pregnancy; maximising maternal and foetal outcomes and limiting excessive gestational weight gain. *Eur. J. Sport Sci.* 2019; 19(1):62–70. [\[Context Link\]](#)

- 74

Brown LS. Nutrition requirements during pregnancy. In: *Life Cycle Nutrition An Evidence–Based Approach*. Burlington, MA: Jones & Bartlett Learning; 2011. p. 1–24. [\[Context Link\]](#)

- 75

Institute of Medicine. *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids*. Washington, DC: The National Academies Press; 2005. [\[Context Link\]](#)

- 76

Grizzo FMF, Alarcão ACJ, Dell' Agnolo CM, et al. How does women's bone health recover after lactation? A systematic review and meta-analysis [Internet]. *Osteoporos. Int.* 2020; 31(3):413–27. [\[Context Link\]](#)

- 77

Pugh LC, Milligan R. A framework for the study of childbearing fatigue. *ANS Adv. Nurs. Sci.* 1993; 15(4):60–70. [\[Context Link\]](#)

- 78

Bialobok KM, Monga M. Fatigue and work in pregnancy. *Curr. Opin. Obstet. Gynecol.* 2000; 12(6):497–500. [\[Context Link\]](#)

- 79

Corwin EJ, Arbour M. Postpartum fatigue and evidence-based interventions. *MCN Am. J. Matern. Child Nurs.* 2007; 32(4):215–20. [\[Context Link\]](#)

- 80

Henderson J, Alderdice F, Redshaw M. Factors associated with maternal postpartum fatigue: an observational study. *BMJ Open.* 2019; 9(7):e025927. [\[Context Link\]](#)

- 81

Nelson-Piercy C. Treatment of nausea and vomiting in pregnancy. When should it be treated and what can be safely taken? *Drug. Saf.* 1998; 19(2):155–64. [\[Context Link\]](#)

- 82

Eliakim R, Abulafia O, Sherer DM. Hyperemesis gravidarum: a current review. *Am. J. Perinatol.* 2000; 17(4):207–18. [\[Context Link\]](#)

- 83

Ferdinande K, Dorreman Y, Roelens K, Ceelen W, De Looze D. Anorectal symptoms during pregnancy and postpartum: a prospective cohort study. *Colorectal Dis.* 2018; 20(12):1109–16. [\[Context Link\]](#)

- 84

Routzong MR, Rostaminia G, Moalli PA, Abramowitch SD. Pelvic floor shape variations during pregnancy and after vaginal delivery. *Comput. Methods Programs Biomed.* 2020; 194:105516. [\[Context Link\]](#)

- 85

Johannessen H, Wibe A, Stordahl A, Sandvik L, Backe B, Mørkved S. Prevalence and predictors of anal incontinence during pregnancy and 1 year after delivery: a prospective cohort study. *BJOG An. Int. J. Obstet. Gynaecol.* 2014; 121(3):269–80. [\[Context Link\]](#)

- 86

South MMT, Stinnett SS, Sanders DB, Weidner AC. Levator ani denervation and reinnervation 6 months after childbirth. *Am. J. Obstet. Gynecol.* 2009; 200(5):519.e1–7. [\[Context Link\]](#)

- 87

Oberwalder M, Connor J, Wexner SD. Meta-analysis to determine the incidence of obstetric anal sphincter damage. *Br. J. Surg.* 2003; 90(11):1333–7. [\[Context Link\]](#)

- 88

Schwertner-Tiepelmann N, Thakar R, Sultan AH, Tunn R. Obstetric levator ani muscle injuries: current status [Internet]. *Ultrasound Obstet. Gynecol.* 2012; 39(4):372–83. [\[Context Link\]](#)

- 89

Friedman S, Blomquist JL, Nugent JM, McDermott KC, Muñoz A, Handa VL. Pelvic muscle strength after childbirth. *Obstet. Gynecol.* 2012; 120(5):1021–8. [\[Context Link\]](#)

- 90

Dietz HP, Benness CJ. Voiding function in pregnancy and puerperium. *Int. Urogynecol. J.* 2005; 16(2):151–4.[\[Context Link\]](#)

- 91

van Veelen GA, Schweitzer KJ, van der Vaart CH. Ultrasound imaging of the pelvic floor: changes in anatomy during and after first pregnancy. *Ultrasound Obstet. Gynecol.* 2014; 44(4):476–80.[\[Context Link\]](#)

- 92

Francis WJ. Disturbances of bladder function in relation to pregnancy. *J. Obstet. Gynaecol. Br. Emp.* 1960; 67(3):353–66.[\[Context Link\]](#)

- 93

Bulchandani S, Coats AC, Gallos ID, Tooze-Hobson P, Parsons M. Normative bladder diary measurements in pregnant women. *Eur. J. Obstet. Gynecol. Reprod. Biol.* 2017; 210:319–24.[\[Context Link\]](#)

- 94

Granese R, Adile B. Urinary incontinence in pregnancy and in puerperium: 3 months follow-up after delivery. *Minerva Ginecol.* 2008; 60(1):15–21.[\[Context Link\]](#)

- 95

Li Z, Xu T, Zhang L, Zhu L. Prevalence, potential risk factors, and symptomatic bother of lower urinary tract symptoms during and after pregnancy [Internet]. *LUTS Low Urin Tract Symptoms*. 2019; 11(4):217–23.[\[Context Link\]](#)

- 96

Hanif S. Frequency and pattern of urinary complaints among pregnant women. *J. Coll. Physicians. Surg. Pak.* 2006; 16(8):514–7.[\[Context Link\]](#)

- 97

Stær-Jensen J, Siafarikas F, Hilde G, Benth JS, Bø K, Engh ME. Postpartum recovery of levator hiatus and bladder neck mobility in relation to pregnancy. *Obstet. Gynecol.* 2015; 125(3):531–9.[\[Context Link\]](#)

- 98

O'Boyle AL, O'Boyle JD, Calhoun B, Davis GD. Pelvic organ support in pregnancy and postpartum. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* 2005; 16(1):69–72.[\[Context Link\]](#)

- 99

Barber MD. Symptoms and outcome measures of pelvic organ prolapse. *Clin. Obstet. Gynecol.* 2005; 48(3):648–61.[\[Context Link\]](#)

- 100

Dietz HP, Haylen BT, Vancaillie TG. Female pelvic organ prolapse and voiding function. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* 2002; 13(5):284–8.[\[Context Link\]](#)

- 101

Dietz HP, Cartmill J. Imaging in patients with obstructed defecation [Internet]. *Tech. Coloproctol.* 2013; 17(5):473–4. [\[Context Link\]](#)

- 102

Jelovsek JE, Barber MD. Women seeking treatment for advanced pelvic organ prolapse have decreased body image and quality of life. *Am. J. Obstet. Gynecol.* 2006; 194(5):1455–61. [\[Context Link\]](#)

- 103

Hagen S, Stark D. Conservative prevention and management of pelvic organ prolapse in women. *Cochrane Database Syst. Rev.* 2011;(12):CD003882. [\[Context Link\]](#)

- 104

Aslan G, Aslan D, Kizilyar A, İspahi Ç, Esen A. A prospective analysis of sexual functions during pregnancy. *Int. J. Impot. Res.* 2005; 17(2):154–7. [\[Context Link\]](#)

- 105

Tennfjord MK, Hilde G, Stær-Jensen J, Ellström Engh M, Bø K. Dyspareunia and pelvic floor muscle function before and during pregnancy and after childbirth. *Int. Urogynecol. J.* 2014; 25(9):1227–35. [\[Context Link\]](#)

- 106

Connolly A, Thorp J, Pahel L. Effects of pregnancy and childbirth on postpartum sexual function: a longitudinal prospective study [Internet]. *Int. Urogynecol. J. Pelvic Floor Dysfunct.* 2005; 16(4):263–7. [\[Context Link\]](#)

- 107

Barrett G, Pendry E, Peacock J, Victor C, Thakar R, Manyonda I. Women's sexual health after childbirth [Internet]. *BJOG.* 2000; 107(2):186–95. [\[Context Link\]](#)

- 108

Galbally M, Watson SJ, Permezel M, Lewis AJ. Depression across pregnancy and the postpartum, antidepressant use and the association with female sexual function. *Psychol. Med.* 2019; 49(9):1490–9. [\[Context Link\]](#)

- 109

Vleeming A, Albert HB, Östgaard HC, Sturesson B, Stuge B. European guidelines for the diagnosis and treatment of pelvic girdle pain. *Eur. Spine J.* 2008; 17(6):794–819. [\[Context Link\]](#)

- 110

Albert HB, Godskesen<sup>†</sup> M, Korsholm L, Westergaard JG. Risk factors in developing pregnancy-related pelvic girdle pain. *Acta. Obstet. Gynecol. Scand.* 2006; 85(5):539–44. [\[Context Link\]](#)

- 111

Bjelland E, Stuge B, Engdahl B, Eberhard-Gran M. The effect of emotional distress on persistent pelvic girdle pain after delivery: a longitudinal population study. *BJOG*. 2013; 120(1):32–40.[\[Context Link\]](#)

- 112

Bjelland EK, Stuge B, Vangen S, Stray-Pedersen B, Eberhard-Gran M. Mode of delivery and persistence of pelvic girdle syndrome 6 months postpartum. *Am. J. Obstet. Gynecol.* 2013; 208(4):298.e1–7.[\[Context Link\]](#)

- 113

Mackenzie J, Murray E, Lusher J. Women's experiences of pregnancy related pelvic girdle pain: a systematic review. *Midwifery*. 2018; 56:102–11.[\[Context Link\]](#)

- 114

Cakmak B, Ribeiro AP, Inanir A. Postural balance and the risk of falling during pregnancy. *J. Matern. Fetal. Neonatal. Med.* 2016; 29(10):1623–5.[\[Context Link\]](#)

- 115

Butler EE, Colón I, Druzin ML, Rose J. Postural equilibrium during pregnancy: decreased stability with an increased reliance on visual cues. *Am. J. Obstet. Gynecol.* 2006; 195(4):1104–8.[\[Context Link\]](#)

- 116

Gonçalves Fernandes da Mota P, Brites Andrade Pascoal AG, Andrade Dinis Carita AI, Bø K. Prevalence and risk factors of diastasis recti abdominis from late pregnancy to 6 months postpartum, and relationship with lumbo-pelvic pain. *Man. Ther.* 2015; 20(1):200–5.[\[Context Link\]](#)

- 117

Boissonnault JS, Blaschak MJ. Incidence of diastasis recti abdominis during the childbearing year. *Phys. Ther.* 1988; 68(7):1082–6.[\[Context Link\]](#)

- 118

Benjamin DR, Frawley HC, Shields N, van de Water ATM, Taylor NF. Relationship between diastasis of the rectus abdominis muscle (DRAM) and musculoskeletal dysfunctions, pain and quality of life: a systematic review. *Physiotherapy*. 2019; 105(1):24–34.[\[Context Link\]](#)

- 119

ACOG Committee opinion no. 650: physical activity and exercise during pregnancy and the postpartum period. *Obstet. Gynecol.* 2015; 126(6):e135–42.[\[Context Link\]](#)

- 120

Forczek W, Staszkiwicz R. Changes of kinematic gait parameters due to pregnancy. *Acta. Bioeng. Biomech.* 2012; 14(4):113–9.[\[Context Link\]](#)

- 121

Hagan L, Wong CK. Gait in pregnant women. *J. Womens Health Phys. Therap.* 2010; 34(2):46–56.[\[Context Link\]](#)

- 122

Branco M, Santos-Rocha R, Aguiar L, Vieira F, Veloso A. Kinematic analysis of gait in the second and third trimesters of pregnancy. *J. Pregnancy*. 2013; 2013:1–9.[\[Context Link\]](#)

- 123

Carpes FP, Griebeler D, Kleinpaul JF, et al. Women able-bodied gait kinematics during and post pregnancy period. *Brazilian J. Biomech*. 2008; 9(16):33–40.[\[Context Link\]](#)

- 124

Gilleard WL. Trunk motion and gait characteristics of pregnant women when walking: report of a longitudinal study with a control group. *BMC Pregnancy Childbirth*. 2013; 13:71.[\[Context Link\]](#)

- 125

Kim DR, Sockol LE, Sammel MD, Kelly C, Moseley M, Epperson CN. Elevated risk of adverse obstetric outcomes in pregnant women with depression. *Arch. Womens Ment. Health*. 2013; 16(6):475–82.[\[Context Link\]](#)

- 126

Woody CA, Ferrari AJ, Siskind DJ, Whiteford HA, Harris MG. A systematic review and meta-regression of the prevalence and incidence of perinatal depression. *J. Affect Disord*. 2017; 219:86–92.[\[Context Link\]](#)

- 127

Logsdon MC, Wisner KL, Pinto-Foltz MD. The impact of postpartum depression on mothering. *J. Obstet. Gynecol. Neonatal. Nurs*. 2006; 35(5):652–8.[\[Context Link\]](#)

- 128

Hay DF, Pawlby S, Waters CS, Sharp D. Antepartum and postpartum exposure to maternal depression: different effects on different adolescent outcomes. *J. Child Psychol. Psychiatry*. 2008; 49(10):1079–88.[\[Context Link\]](#)

- 129

Sandra NR, Tadinac M, Herman R. Anxiety during pregnancy and postpartum: course, predictors and comorbidity with postpartum depression. *Acta. Clin. Croat*. 2018; 57(1):39–51.[\[Context Link\]](#)

- 130

Thein-Nissenbaum J. The postpartum triathlete. *Phys. Ther. Sport*. 2016; 21:95–106.[\[Context Link\]](#)

- 131

Erdener U, Budgett R. Exercise and pregnancy: focus on advice for the competitive and elite athlete. *Br. J. Sports Med*. 2016; 50:567.[\[Context Link\]](#)

- 132

Mottola MF, Davenport MH, Ruchat SM, et al. No. 367-2019 Canadian guideline for physical activity throughout pregnancy. *J. Obstet. Gynaecol. Can*. 2018; 40(11):1528–37.[\[Context Link\]](#)

- 133

Kader M, Naim-Shuchana S. Physical activity and exercise during pregnancy. *Eur. J. Physiother.* 2014; 16(1):2–9.[\[Context Link\]](#)

- 134

Cai C, Vandermeer B, Khurana R, et al. The impact of occupational activities during pregnancy on pregnancy outcomes: a systematic review and metaanalysis. *Am. J. Obstet. Gynecol.* 2020; 222(3):224–38.[\[Context Link\]](#)

- 135

Pivarnik JM, Connolly CP, Marshall MR, Schlaff RA. Can pregnancy be an ergogenic aid to athletic performance? A review. *Women Sport Phys. Act. J.* 2017; 25(2):111–7.[\[Context Link\]](#)

- 136

Deering RE, Christopher SM, Heiderscheit BC. From childbirth to the starting blocks: are we providing the best care to our postpartum athletes? *J. Orthop. Sports Phys. Ther.* 2020; 50(6):281–4.[\[Context Link\]](#)

- 137

Tenforde AS, Toth KES, Langen E, Fredericson M, Sainani KL. Running habits of competitive runners during pregnancy and breastfeeding. *Sports Health.* 2015; 7(2):172–6.[\[Context Link\]](#)

- 138

Potteiger JA, Welch JC, Byrne JC. From parturition to marathon: a 16-wk study of an elite runner. *Med. Sci. Sports Exerc.* 1993; 25(6):673–7.[\[Context Link\]](#)

- 139

Kimber ML, Meyer S, McHugh T-L, et al. Health outcomes following pregnancy in elite athletes: a systematic review and meta-analysis. *Med. Sci. Sport Exerc.* 2021; 53(8):1739–47.[\[Context Link\]](#)

- 140

Solli GS, Sandbakk Ø. Training characteristics during pregnancy and postpartum in the world's most successful cross country skier. *Front. Physiol.* 2018; 9:595.[\[Context Link\]](#)

- 141

Houglum PA. Soft tissue healing and its impact on rehabilitation. *J. Sport Rehabil.* 1992; 1(1):19–39.[\[Context Link\]](#)

- 142

Goom T, Donnelly G, Brockwell E. Returning to running postnatal—guidelines for medical, health and fitness professionals managing this population [Internet]. 2019 [cited 2021 November 31]. doi:10.13140/RG.2.2.35256.90880/2.[\[Context Link\]](#)

- 143

Woodley SJ, Lawrenson P, Boyle R, et al. Pelvic floor muscle training for preventing and treating urinary and faecal incontinence in antenatal and postnatal women. *Cochrane Database Syst. Rev.* 2020; 5(5):CD007471.[\[Context Link\]](#)

- 144

National Institute for Health and Care Excellence. Antenatal care for uncomplicated pregnancies. *Clin. Guidel.* 2008; (March):1–55.[\[Context Link\]](#)

- 145

Dijkstra HP, Pollock N, Chakraverty R, Alonso JM. Managing the health of the elite athlete: a new integrated performance health management and coaching model. *Br. J. Sports Med.* 2014; 48(7):523–31.[\[Context Link\]](#)

- 146

Canadian Society for Exercise Physiology. *Get active questionnaire for pregnancy [Internet]*. [cited 2021 November 31]. Available from: <https://csep.ca/index.php/2021/05/27/get-active-questionnaire-for-pregnancy/>.[\[Context Link\]](#)

- 147

Shiri R, Coggon D, Falah-Hassani K. Exercise for the prevention of low back and pelvic girdle pain in pregnancy: a meta-analysis of randomized controlled trials. *Eur. J. Pain.* 2018; 22(1):19–27.[\[Context Link\]](#)

- 148

National Institute for Health and Clinical Excellence (NICE). Postnatal care up to 8 weeks after birth clinical guideline. *NICE Guidel.* 2006; (July 2006):1–63.[\[Context Link\]](#)

- 149

Schreiner L, Crivelatti I, de Oliveira JM, Nygaard CC, dos Santos TG. Systematic review of pelvic floor interventions during pregnancy. *Int. J. Gynecol. Obstet.* 2018; 143(1):10–8.[\[Context Link\]](#)

- 150

Swift JA, Langley-Evans SC, Pearce J, et al. Antenatal weight management: diet, physical activity, and gestational weight gain in early pregnancy. *Midwifery.* 2017; 49:40–6.[\[Context Link\]](#)