

# Please cite the Published Version

Field, Adam , Corr, Liam David, Sarmento, Hugo, Naughton, Robert, Clifford, Tom, Haines, Matthew, Page, Richard Michael and Harper, Liam David (2023) The impact of 120 minutes of soccer-specific exercise on recovery. Research Quarterly for Exercise and Sport, 94 (1). pp. 237-245. ISSN 0270-1367

DOI: https://doi.org/10.1080/02701367.2021.1964697

Publisher: Taylor & Francis (Routledge)

(cc)

Version: Accepted Version

Downloaded from: https://e-space.mmu.ac.uk/630649/

**BY-NC** 

Creative Commons: Attribution-Noncommercial 4.0

**Additional Information:** This is an Accepted Manuscript of an article published by Taylor & Francis in Research Quarterly for Exercise and Sport on 28th March 2022, available at: http://www.tandfonline.com/10.1080/02701367.2021.1964697. It is deposited under the terms of the Creative Commons Attribution-NonCommercial License (http://creativecommons.org/licenses/by-nc/4.0/), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

## **Enquiries:**

Usage rights:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines) 1 Manuscript Title: The impact of 120 minutes of soccer-specific exercise on recovery

3	Authors: Adam Field <sup>1</sup> • Liam David Corr <sup>1</sup> • Hugo Sarmento <sup>2</sup> • Robert Naughton <sup>1</sup> • Tom Clifford <sup>3</sup> • Matthew
4	Haines <sup>1</sup> • Richard Michael Page <sup>4</sup> • Liam David Harper <sup>1</sup> .
5	
6	
_	
/	<sup>1</sup> School of Human and Health Sciences, University of Huddersfield, Huddersfield, HD1 3DH, United Kingdom.
8	<sup>2</sup> Research Unit for Sport and Physical Activity (CIDAF), Faculty of Sport Sciences and Physical Education,
9	University of Coimbra, Coimbra, Portugal.
10	<sup>3</sup> School of Sport, Exercise and Health Sciences, Loughborough University, Epinal Way, Loughborough, LE11
11	3TU, United Kingdom.
12	<sup>4</sup> Department of Sport & Physical Activity, Edge Hill University, St. Helens Road, Ormskirk, Lancashire, L39
13	4QP, United Kingdom.
14	
15	Trial registration
16	The study was pre-registered on the Open Science Framework (DOI: 10.17605/OSF.IO/VGU6T_Date:
17	10/06/2019).
18	
19	
20	Author responsible for correspondence:
21	Mr Adam Field
22	Address as above
23	+44 (0) 1484 471157
24	Email: <u>Adam.Field@hud.ac.uk</u>
25	ORCID: 0000-0002-2600-6182

### 26 Abstract

Purpose: The extra-time (ET) period of soccer is competed during fixture congested schedules with often limited recovery time between matches. The aim of this study was to assess muscle damage recovery following 90- and 120-min (i.e., incorporation of ET) of simulated soccer match-play. Methods: Twelve semi-professional soccer players completed 90 and 120-min treadmill-based soccer-specific exercise in a counterbalanced order. Creatine kinase (CK), creatinine, urea, aspartate aminotransferase, perceived muscle soreness, pain pressure threshold, reactive strength index, countermovement jump height, and isokinetic strength assessments of eccentric knee flexors at 60, 180 and 270 deg·s<sup>-1</sup> were taken at baseline and immediately-, 24, 48 and 72-hr post-exercise to assess recovery. Results: No significant between-trial interactions except for CK were found. Pairwise comparisons detected a 53% increase in CK at 24-hr ( $455 \pm 29 \ \mu \cdot L^{-1}$ ) following 120-min of simulated match-play vs. the corresponding post 90-min time-point (299  $\pm$  29  $\mu$ ·L<sup>-1</sup>; p<0.01). The 120-min trial caused a 58% higher CK response at 72-hr ( $244 \pm 25 \ \mu \cdot L^{-1}$ ) vs. post 90-min comparisons ( $154 \pm 29 \ \mu \cdot L^{-1}$ ; p = 0.02). No interaction effects were detected for any other recovery variables. Creatine kinase and perceived muscle soreness remained elevated up to 72-hr in both trials (p < 0.01). Conclusions: These data indicate that 120 min of simulated soccer match-play delays the time-course of CK recovery up to 72-hr post-match. However, 120 min of simulated soccer has no additional impact on functional recovery and perceived muscle soreness vs. 90 min. Recovery should be investigated following 90- and 120-min of actual match-play.

Keywords Extra-time • Football • Soccer simulation • Muscle damage

- \_

## 55 Introduction

56 The physical demands of soccer matches involve a myriad of eccentric muscular efforts including sprints, jumps, 57 rapid accelerations, decelerations and changes of direction that induce fatigue (a debilitating symptom), perceptual 58 stress, deplete endogenous substrates and impose structural damage within skeletal muscle (Harper et al., 2016; 59 Mohr, Krustrup, & Bangsbo, 2005; Stevenson et al., 2017). Several mechanisms explain these detrimental effects 60 at the local level including an increased inflammatory response, disturbance to calcium homeostasis and ultra-61 structural damage to muscle fibres and the surrounding connective tissues (Mohr et al., 2005; Nédélec et al., 2013; 62 Proske & Morgan, 2001). Following these disturbances, recovery, defined as restoring homeostasis (i.e., returning 63 players to pre-performance values), is paramount post-match (Nédélec et al., 2012b). Traditionally, soccer is 64 contested over 90 min, and as such, there is a wealth of literature documenting fatigue and subsequent recovery 65 post matches of this duration (Brownstein et al., 2017; Nedelec et al., 2014b; Nédélec et al., 2012b). However, 66 when matches are tied, and an outright winner is required during the knockout phase of certain major tournaments 67 and domestic cup competitions, an additional 30-min period is played, known as extra-time (ET) (Harper et al., 68 2016).

69 The prevelance of matches that progress to ET have become increasingly common over recent years, with 36% 70 of the previous four FIFA World Cup knockout phase matches proceeding to this additional 30-min period (Field 71 et al., 2020). For the first time in the tournament's history, a team competed in three consecutive ET matches 72 (round 1/16, quarter- and semi-finals) in the knockout phase of the 2018 FIFA World Cup (Kołodziejczyk et al., 73 2021). Interestingly, players covered less distance across the 90 min duration in the final match compared to the 74 previous knockout-phase matches, which could be related to fatigue from the accumulation of ET matches. Further 75 evidence suggests that ET results in fatigue-induced decreases in relative (m·min<sup>-1</sup>) total distance and high-speed 76 distance covered, as well as the number of sprints, accelerations and decelerations performed (Peñas, Dellal, 77 Owen, & Gómez-Ruano, 2015; Russell et al., 2015a). Tournament matches are typically competed amid fixture 78 congested scheduling, with an insufficient recovery time between matches (Julian, Page, & Harper, 2020), 79 potentially predisposing players to injury/illness and impacting upon player availability and performance (Carling, 80 McCall, Le Gall, & Dupont, 2015). Previous empirical research examining five English Premier League reserve 81 team players found explosive power (i.e., countermovement jump [CMJ]) reduced and creatine kinase (CK) 82 activity increased at 24 and 48 hr post 120 min matches (Russell, Sparkes, et al., 2015a). As players only competed 83 in a single match that required ET and there were no comparisons to 90 min, it is difficult to determine the extent 84 to which the additional 30-min duration further impacted recovery. Winder, Russell, Naughton, and Harper (2018)

85 monitored four professional players competing in a micro cycle, with the second of three matches progressing to 86 ET (i.e., 90, 120, 90-min). The ET match negatively affected subjective measures of fatigue, muscle soreness, 87 mood and CMJ height vs. the initial 90-min match. Although these findings are indicative of impeded recovery 88 following ET matches, and notwithstanding the high ecological validity of these studies, limitations such as small 89 sample sizes, insufficient control of confounders (Drust et al., 2013) and high inter-match variability associated 90 with match-play reduce experimental rigour (Gregson, Drust, Atkinson, & Salvo, 2010). Therefore, there is a 91 paucity of research attention afforded to investigating the impact of ET on recovery under controlled experimental 92 conditions, utilising adequately powered trials and directly comparing the same group of subjects across both 93 experimental conditions (i.e., 90 and 120-min exercise durations).

94 Researchers often use soccer-specific exercise protocols to replicate the demands of soccer match-play, while 95 negating the influence of confounding factors to assess performance and recovery outcomes (Field et al., 2020). 96 Free-running protocols are appealing for their increased ecological validity (implementation of technical actions, 97 utility/multidirectional movements and attainment of maximum speeds) (Russell, Benton, & Kingsley, 2011; 98 Uddin et al., 2020). However, these simulations appear to lack reproducibility/experimental control, fail to omit 99 subconscious pacing approaches and are not reflective of the neuromechanical loading patterns of match-play 100 (Field et al., 2020; Page, Marrin, Brogden, & Greig, 2019). Therefore, motorised treadmill protocols are preferred 101 such that the activity profile is mechanistically valid and external running output can be standardised; thus, 102 changes in a given variable are likely a fatigue-impaired physical capacity as opposed to player pacing and 103 motivation (Field et al., 2020; Page et al., 2015). Therefore, utilising a treadmill-based model that elicits 104 experimental control, integrating fixed bouts of activity appears appropriate since the magnitude of the physical 105 output and the fatigue experienced whilst exercising, corresponds closely with the degree of muscle damage and 106 the subsequent nature of the recovery response (Nedelec et al., 2014a).

107 The purpose of this study was to assess recovery following 90- and 120-min of treadmill-based simulated soccer.
108 It was hypothesised that recovery of the primary outcome measure CMJ height, along with the other variables,
109 would be impaired following 120 min vs. 90 min of simulated soccer.

#### 110 Materials and Methods

111 Study design

112 Ten visits to the laboratory over a three-week period were required. The study was a single-blinded, within113 subjects, crossover design with trials completed in a counterbalanced order. The order was determined by a

number generator that randomly allocated participants to a sequence of trials. The two trials were performed 9 ± 2 days apart to ensure recovery. To eliminate the influence of self-pacing on running technique (Waldron & Highton, 2014), participants were told that both trials could last 120 min; however, the protocol was either terminated at 90 min or ET commenced. Recovery measures were measured in the following order: capillary blood sampling, perceived muscle soreness, pain pressure threshold (PPT), reactive strength index (RSI), CMJ and isokinetic peak torque, at baseline, immediately-, 24, 48 and 72 hr post-trial.

120 Participants

121 Following institutional ethical approval, 12 male semi-professional players (mass:  $74 \pm 8$  kg; stature:  $1.79 \pm 0.3$ 122 m; age:  $22 \pm 3$  years; estimated  $\dot{V}O2_{max}$ :  $59 \pm 7$  ml·kg·min<sup>-1</sup>) with  $13 \pm 3$  years soccer experience provided informed 123 consent prior to data collection. An apriori difference between two dependant means (matched pairs) power 124 calculation was undertaken (GPower v3.1; Germany) which deemed a sample size of 12 sufficient based on 80% power  $(1 - \beta)$ , an alpha ( $\alpha$ ) of 0.05, and a large effect size (Cohen's d = 0.8) to detect differences in the primary 125 outcome variable, CMJ. Participants were included on the basis that they attained  $\dot{V}O_{2max} \ge 48.5$  ml·min·kg<sup>-1</sup> as 126 per previous ET work (Field et al., 2020; Stevenson et al., 2017) and had no medical contraindications to exercise. 127 128 Participants were asked to refrain from strenuous activity and prohibited from alcohol consumption, non-steroidal 129 anti-inflammatory drugs and foods rich in polyphenols 72-hr prior and throughout testing. Players were also 130 instructed to avoid any non-nutritional recovery interventions (massage, foam rolling, cryotherapy etc). Dietary 131 intake was recorded throughout the five testing days (24-hr pre-trial to 72-hr post-trial) via weighed food diaries 132 and later analysed to assess macronutrient composition and caloric intake. Dietary intake was replicated as closely 133 as possible for the subsequent trial. In order to assess compliance and food wastage, pictures were sent to the lead 134 researcher on two days (pre-selected) during data collection (-24 and +48-hr of trial) using a free smartphone 135 messaging application (WhatsApp). This 'snap and send' method has been used in previous research (Costello, 136 Deighton, Dyson, Mckenna, & Jones, 2017; Zhang et al., 2015).

137 Preliminary visits and main trials

Mass (SECA 875 electronic flat scale, SECA, Germany) and stature (SECA 213 portable stadiometer, SECA, Germany) were taken during the preliminary visit and  $\dot{V}O_{2max}$  was estimated *via* an incremental treadmill protocol. Participants initially completed a standardised warm-up of intermittent speed changes and a dynamic stretching sequence. The incremental test then ensued, starting at a running speed of 10 km·h<sup>-1</sup> and continued to increase by 1 km·h<sup>-1</sup> every 30-s until 17 km·h<sup>-1</sup>, whereby the protocol inclined by 0.5° every 30-s until volitional exhaustion (Field et al., 2020). A second session was used to fully habituate participants with trial day procedures and the 144 completion of a 120-min familiarisation trial. Upon arrival for the main trials, participants provided a mid-flow 145 urine sample for measurement of urine osmolality (Osmocheck, Vitech Scientific, West Sussex, UK). Water 146 and/or a carbohydrate electrolyte beverage was consumed *ad libitum*, recorded by the research team and replicated 147 for the subsequent trial. Post-trial recovery measures were collected, and participants vacated the lab.

148 Soccer simulation

149 Participants completed the soccer-specific exercise protocol on a motorised treadmill (h/p/ cosmos pulsar® 3p: 150 h/p/cosmos sports & medical GmBH, Germany). The protocol has previously been validated (Page et al., 2015), 151 replicating the frequency, duration and speed of discrete locomotive phases, the distances covered, sprint 152 quantities and low-to-high speed running ratios observed during 90-min of a professional soccer match (Mohr, 153 Krustrup, & Bangsbo, 2003). Similar to ET match-play, participants perform 56 sprints and cover a distance of 154 16.26 km during the 120-min trial (Russell et al., 2015a; Winder et al., 2018). The protocol has also demonstrated 155 moderate-to-very strong reliability (Field et al., 2020) and has shown to elicit a peripheral and central fatigue 156 response comparable with soccer match-play (Field et al., 2020). Applying a 1% inclination at lower speeds and 157 a 2% gradient at higher speeds, better reflects the energy cost of outdoor running (Jones & Doust, 1996); thus, 158 these varying degrees of gradient were applied to account for the lack of air resistance indoors. The protocol was 159 structured as two x 45-min halves (separated by a 15-min half-time break) and either terminated or continued 160 following a 5-min rest period, with two additional 15-min periods (interspersed by a 2-min break). The same 161 activity profile was repeated every 15-min (Figure 1). Maximum sprint velocities reached 25 km h<sup>-1</sup> with changes in velocity set at the treadmill threshold (1.39  $\text{m}\cdot\text{s}^{-2}$ ). Participants were asked to provide differential ratings of 162 163 perceived exertion (d-RPE) in a counterbalanced order, for legs (RPE-L), breathlessness (RPE-B) and overall 164 (RPE-O), through use of the 6-20 Borg scale (Borg, 1998). The treadmill paused briefly at the cessation of each 165 15-min bout, and d-RPE, and a blood capillary sample was taken and later analysed for blood lactate (BLa) 166 (Biosen C-Line; EKF-diagnostic GmBH, Cardiff, Wales; CV both 1.5%). PlayerLoad™ (Catapult Innovations, 167 Australia) data were continuously recorded throughout exercise and defined across each 15-min bout.

168 \*\*\*INSERT FIGURE 1\*\*\*

169 Recovery measures

170 Blood sampling procedures and biochemical analyses

Finger-prick capillary blood samples (200 μl) were taken once the foremost droplet of blood was discarded. This
was collected in a lithium heparin coated microvette<sup>®</sup> and analysed for CK, creatinine, urea and aspartate
aminotransferase (AST) activity using a colorimetric assay procedure (Reflotron<sup>®</sup> plus, Roch Diagnostics,

174 Switzerland). All samples were analysed by a single researcher in duplicate to eliminate inter-assay variation.

175 Muscle soreness

A subjective assessment of perceived muscle soreness was undertaken using a visual analogue scale (VAS), which
has previously demonstrated good reliability (intra-class correlation [ICC] = 0.65) (Rampinini et al., 2011). Once
in a squat position (angle ~90°) with knees shoulder width apart, participants marked on a 100 mm horizontal line
which comprised of "no pain whatsoever" (0 mm) and "maximum pain imaginable" (100 mm) (Chen, Lin, Chen,
Lin, & Nosaka, 2011).

181 A PPT test was also performed at the biceps femoris (PPTBF) on the dominant leg using a handheld Baseline 182 Algometer (27.22 kg/ 60 lbs, Fabrication, Enterprises Inc., USA). The biceps femoris were identified as the most 183 common site of injury in a recent epidemiological systematic review and meta-analysis (López-Valenciano et al., 184 2019). A mark was positioned central to the muscle belly and re-applied to the skin daily using indelible ink to 185 ensure between-test consistency (Hermens et al., 2000). The lead researcher applied progressively increasing 186 pressure with the circular flat tip (1-cm in diameter) perpendicular to the pre-marked site. Participants were 187 instructed to signal at the point of shift from pressure-to-pain with the value recorded in pounds (lbs). Recordings 188 were taken twice, separated by ~60-s, and the mean of both scores were presented for analyses. The use of PPT 189 has previously demonstrated high inter- and intra-rater reliability (ICC = 0.63-0.97) (Binderup, Arendt-Nielsen, 190 & Madeleine, 2010; Walton et al., 2011).

191 Muscle function

For RSI, participants were instructed to drop from a 0.3 m platform and upon landing, jump maximally, whilst minimising ground contact time and maximising vertical jump height. The RSI values were calculated automatically using manufacturer software (Optojump, Microgate, Italy) through the sum of jump height (cm) divided by contact time (ms). Following a 30-s rest period, participants performed a CMJ with hands on hips, feet shoulder width apart and when prompted, descended into a squat (~60°) and jumped vertically with maximal effort. Jump efforts were measured using a portable optical measurement system (Optojump, Microgate, Italy) and were separated by a 30-s passive recovery period with the mean of three jump being used for analyses (Fieldet al., 2020).

200 Isokinetic testing

201 Unilateral peak torque of the eccentric knee flexors (eccKF) were measured using a Cybex HUMAC Norm 202 isokinetic dynamometer with HUMAC2009 software version 0.8.4 (CSMI, USA). The eccKF were selected, given 203 that fatigue-induced strength deficits are commonly observed in the eccentric hamstrings, increasing stretch injury 204 susceptibility due to an impaired capacity to resist over lengthening whilst fatigued (Small, McNaughton, Greig, 205 & Lovell, 2010). Each of three sets were performed at respective angular velocities of 180, 270 and 60 deg·s<sup>-1</sup>; 206 order chosen to attenuate the possible fatiguing effect of slower speeds (Greig, 2008). The dynamometer was 207 setup specific to the participant for each session as per manufacturers guidelines. Whilst seated, the passive limb 208 was weighed at anatomical zero (defined as full knee extension) and the effects of gravity were applied to torque 209 data. Five reps of each speed were performed and each of the three sets were interspersed by a 30-s passive rest 210 period. The preferred kicking leg was assessed through a range of  $0-90^{\circ}$  (0° representing full extension and 90° 211 full flexion). Similar soccer research has reported ICCs of 0.78, 0.76 and 0.78 for eccentric knee flexion at 180, 300 and 60 deg  $\cdot$  s<sup>-1</sup>, respectively (Greig, 2008). 212

213 Dietary analysis

214 Participant food diaries were inputted into dietary analysis software (Nutrimen.co.uk, Dark Green Media Ltd, ©2016). The estimated measurement error of inputting participant's dietary intake was calculated using the 215 216 standard error of measurement (SEM) with 95% confidence intervals (CI). The SEM was derived from the square 217 root of the mean square error from an ANOVA and expressed in the units of each given variable (Stratford & 218 Goldsmith, 1997). To examine intra-rater reliability, three participants were selected at random and a single 219 researcher inserted the same three food diaries into the analysis software on six separate occasions during a six-220 week period. To assess inter-rater reliability, the same diaries were entered by three separate researchers twice in 221 two-weeks.

222 Statistical analysis

Exploratory data analysis was undertaken to evaluate the assumptions associated with the linear mixed model (LMM). This statistical method was chosen as an appropriate test for a repeated measures design that involves random and fixed level factors with missing data; assuming data are missing at random (Di Salvo, Gregson, 226 Atkinson, Tordoff, & Drust, 2009). A visual inspection of q-q plots, histograms and boxplots was undertaken to 227 assess normality of residuals. Residuals > 3.0 SD from the mean values were removed prior to analyses in line 228 with the assumptions of the LMM. A basic variance components assessment revealed the model Akaike's 229 information criterion (AIC) was best fit for each recovery variable. Models were initially regarded as null and 230 subsequently developed to more stringent models. A basic variance components model was utilised to calculate 231 the intraclass correlation (ICC) of the random factors (i.e., participant) to establish if a significant variance 232 contributed to the recovery variables. Wald Z statistics were employed to assess the null hypothesis (i.e., that zero 233 variance existed between participants); if rejected, the random factor of participant ID was included in the 234 successive hierarchical models. The covariance structure of the random factors was set to variance components in 235 all models. The auto-regressive (AR-1) was established as the model most suitable for each recovery variable for 236 the repeated measures of time. The fixed effects and their interactions included were trial and time for each model. 237 All models estimated parameters using the maximum likelihood method. Least significant corrections were 238 applied post-hoc with 95% CI of the difference reported. A paired samples T-test assessed differences between 239 participant dietary intake across the five experimental days. Data are expressed as mean and  $\pm$  SE unless otherwise 240 stated and were processed using IBM SPSS Statistics 26 for windows (SPSS Inc., Chicago, IL, USA). Alpha was 241 accepted at < 0.05 prior to analyses.

### 242 Results

- 243 Between trials measures
- No differences were detected between trials for energy or macronutrient intake (p > 0.35; Table 1), urine osmolality, environmental conditions, cumulative PlayerLoad<sup>TM</sup>, BLa and d-RPE (all p > 0.33; Table 2).
- 246 \*\*\*INSERT TABLE 1\*\*\*
- **247** \*\*\*INSERT TABLE 2\*\*\*
- 248 Variance calculations
- Table 3 provides the ICC's (%) of the random factors accounted for in each of the LMMs. All recovery measures,
- 250 except for CK, creatinine and AST, contributed significant variance to the dependant variables and were included
- as a random factor in the larger hierarchical models.

252 \*\*\*INSERT TABLE 3\*\*\*

253 Recovery responses following 90 and 120 minutes of simulated soccer

values at 24-hr post 120-min (455.1  $\pm$  29.4 U·L<sup>-1</sup>; 95% CI = 396.7 to 513.7 U·L<sup>-1</sup>) compared to 24-hr following 90-min (298.8  $\pm$  29.3 U·L<sup>-1</sup>; 95% CI = 240.6 to 356.9 U·L<sup>-1</sup>; 95% CI for diff = 75.1 to 237.8 U·L<sup>-1</sup>; p < 0.01). Furthermore, trial interaction effects were observed at 72-hr post 120-min (244.1  $\pm$  24.7 µ·L<sup>-1</sup>; 95% CI = 194.7 to 293.4 U·L<sup>-1</sup>) vs. 72-hr following the 90-min simulation (154.1  $\pm$  29.3U·L<sup>-1</sup>; 95% CI = 240.6 to 356.9 U·L<sup>-1</sup>; 95% CI for diff = 17.2 to 176.9 U·L<sup>-1</sup>; p = 0.02). No interactions were identified for creatinine (p = 0.24), urea (p =0.59), AST (p = 0.83), VAS (p = 0.22), PPT<sub>RF</sub> (p = 0.99), PPT<sub>BF</sub> (p = 0.78), RSI (p = 0.35), CMJ (p = 0.21), eccKF<sub>180</sub> (p = 0.75), eccKF<sub>270</sub> (p = 0.67) and eccKF<sub>60</sub> (p = 0.42).

Interaction effects for trial and time were evident for CK (p = 0.01), with post-hoc comparisons revealing higher

262 Main effects for time were observed for CK, urea, VAS, PPT<sub>BF</sub>, RSI, CMJ, eccKF<sub>180</sub>, eccKF<sub>270</sub> and eccKF<sub>60</sub> (all 263 p < 0.01; Table 4). No main effects for time were evident for creatinine (p = 0.24), AST (p = 0.23) and PPT<sub>RF</sub> (p264 = 0.46).

#### 265 \*\*\*INSERT TABLE 4\*\*\*

### 266 Discussion

254

The purpose of this study was to assess recovery in response to a 90- and 120-min (i.e., inclusion of an ET period) simulated soccer match. No differences were established between trials for all variables aside from CK. Higher CK activity was observed at 24 and 72 hr post 120 min of simulated soccer compared with a typical 90 min duration. Contrary to the study hypothesis, little evidence that recovery is prolonged after 120 min vs. 90 min of simulated soccer match-play was found.

272 Despite most variables demonstrating that recovery is not further impeded following ET, the current findings 273 demonstrate that compared to the 90-min trial, CK activity increased by a further 53 and 58% at 24 and 72-hr 274 following 120 min, respectively. The magnitude of change in CK far exceeds the intra- and inter-assay 275 measurement error (i.e., coefficient of variation = 1.4-2.1% and 1.5-4.2%, respectively), indicating the 276 variation is low enough not to obscure a true experimental effect. Considering the CK response to 120 min alone, 277 a 317% increase at 24-hr post-trial was observed vs. baseline. Previous ET match-play investigations observed a 278 236% increase in CK at 24-hr post 120 min of soccer match-play vs. baseline in professional players (Russell et 279 al., 2015a). The higher magnitude of change in the current study compared to previous research is potentially 280 explained by a wealth of factors such as standard of player and innate blood marker variability (Baird et al., 2012), 281 though we speculate most likely as a consequence of the differences in exercise modality (i.e., simulation vs 282 match-play). For instance, the match-play study identified reductions in total (-12%) and high-speed distance (-283 37.5%), as well as number of accelerations (-14%) and decelerations (-13%) during ET, relative to the preceding 284 90 min (Russell et al., 2015a). However, the activity was standardised for the current protocol, with players 285 performing at the same intensity throughout the entire 120 min duration, unlike the players in the match-play 286 study (Russell et al., 2015a). This also supports that the current treadmill protocol was sufficiently intense to 287 induce a muscle damage response comparable with match-play and suggests recovery strategies may need to be 288 modulated to restore the physiological perturbations of players competing for 120 min. Notwithstanding, it is 289 important to note that CK has been criticised as a marker of muscle damage recovery (Twist & Highton, 2013), 290 and therefore this finding should not be interpreted independent of the functional measures discussed below.

291 No between-trial interactions were present for eccKF peak torque in the current research, although the current 292 soccer-specific exercise protocol reduced hamstring strength following both exercise durations. Due to the 293 logistical complexities associated with collecting isokinetic peak torque data during match-play, comparable data 294 are not available following 90 min or ET matches. However, a similar investigation also demonstrates that 295 completing an ET match simulation on a treadmill, significantly reduces eccKF peak torque, at the same speeds 296 to the present study, immediately post 120 min of exercise (Field et al., 2020). Therefore, the current research 297 supports previous data and contributes new infromation that indicates the deficits in eccKF persist up to 24 hr and 298 return to baseline at 72 hr following 120-min of soccer. It is important to note, however, that matches are 299 occasionally interpersed by ~48 hr (Ranchordas, Dawson, & Russell, 2017), and thus, practitioners are advised to 300 closely monitor recovery markers, before returning players to training and matches. The recovery of maximal 301 strength should largely be considered on an indvidual basis, depsite the current research reporting an average 302 response, since eccKF strength characteristics are susceptible to large individual variation following ET (Field et 303 al., 2020). It is advised that replacements are used strategically during or before the additional 30 min ET period 304 to attenuate the effects of fatigue, and maintain fitness and freshness since a fourth substitution is now permitted 305 during ET (Hills et al., 2020). Furthermore, to support player welfare during the COVID-19 pandemic, the 306 International Football Association Board approved a temporary amendment to the laws of the game, allowing 307 teams the option of using five substitutions per match, with an additional replacement permitted during ET. This 308 is especially important considering that the residual fatigue-induced decrements in eccKF in the intervening post 309 match period, and may explain the increased injury incidence during ET (Aoki, O'Hata, Kohno, Morikawa, & 310 Seki, 2012), especially if players are not entirely recovered prior to the next competitive encounter.

311 Epidemiological research, therefore, appears warranted to determine the extent to which ET matches exacerbate312 injury incidence in subsequent training and matches.

313 Soccer-specific activity involves repetitive eccentric to concentric actions, that can induce neuromuscular fatigue 314 and reduce peak power output (Nédélec et al., 2012b). As such, given the functional relevance of such markers 315 and their ability to detect perturbations in muscle function and elastic energy usage (Oliver, Armstrong, & Williams, 2008), RSI and CMJ were used as indicators of exercise-induced fatigue. Time-dependant reductions 316 317 in RSI and CMJ were evident for up to 48-hr following both trials in the current research, although no between-318 trial differences were identified for these measures. Several studies have shown that the recovery of these markers 319 are prolonged following 90 and 120 min of simulated and actual match-play (Abbott, Brashill, Brett, & Clifford, 320 2019; Russell, Northeast, et al., 2015b; Thomas et al., 2017; Winder et al., 2018). Therefore, although the muscle 321 function data limit inference concerning how player recovery should be managed following ET, the results add to 322 an expanding body of literature that suggests recovery may need to be considered alongside the congested fixture 323 periods associated with major tournaments (Julian et al., 2020; Page et al., 2019). However, given that ET may 324 not further prolong recovery of muscle function, there appears to be little need to adapt practice and provide 325 additional measures to restore muscle function capacity irrespective of match duration.

326 An increased presence of blood urea was evident immediately post-exercise in both trials, and although differences 327 did not reach statistical significance, these values were further ( $\sim 10\%$ ) increased immediately following the 120-328 min exercise trial. Given macronutrient intake was similar between trials (Table 1), this finding, although 329 speculative, is indicative of an increased protein degradation following ET. Urea is strongly associated with 330 training volume potentially owing to the upregulation of gluconeogenesis during exercise that results in the 331 breakdown of structural proteins (Haralambie et al., 1976; Meyer & Meister, 2011). The rate of gluconeogenesis 332 is increased when glycogen stores are depleted and is stimulated by the secretion of glucagon, which can be linked 333 to an upregulation in lipolysis (de Sousa et al., 2019; Shephard, 1999). This finding may corroborate previous 334 research that suggests glycogen depletion and an increased rate of lipolysis is apparent during ET (Field et al., 335 2020; Harper et al., 2016; Stevenson et al., 2017). This could potentially highlight the importance of modulating 336 fuelling strategies, including carbohydrate intake prior and during, as well as adapting protein intake post ET 337 matches. However, this research provides a foundation for future lines of exploration that are warranted to assess 338 substrate metabolism using invasive techniques such as a muscle biopsies following 120 min of soccer-specific 339 exercise.

340 Limitations are present within the current study that should be acknowledged. The cohort used within this study 341 may not be representative of the population against which the soccer-specific protocol is based (Mohr et al., 2003), 342 although semi-professional players were recruited with a  $VO_{2max}$  of  $59 \pm 7 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ , which is similar to that 343 of professional soccer players (Tønnessen, Hem, Leirstein, Haugen, & Seiler, 2013). Furthermore, the responses 344 observed in the current study are likely conservative estimates; indeed, actual match-play involves contacts with 345 other players and additional movements such as cutting manoeuvres that likely place a greater deal of stress on 346 the musculoskeletal system, and therefore evoke a muscle damage response of a higher magnitude (Castellano, 347 Blanco-Villaseñor, & Alvarez, 2011). It is well accepted that disparate physical loads, such as those experienced 348 during match-play, can impact upon the muscle damage recovery response (Nédélec et al., 2012b). Therefore, a 349 decision was taken to use a valid treadmill design that employs fixed running velocities of an equal exercise 350 volume to ensure that differences in the recovery response occurred as a result of the additional 30-min duration 351 rather than player pacing strategies to offset fatigue. Another limitation of the study was that the protocol elicited 352 a physiological response (blood lactate:  $2.4 \pm 1.5$  mmol·l<sup>-1</sup>) near the lower end of the ranges previously reported during match-play (2-10 mmol·l<sup>-1</sup>) (Bangsbo, Iaia, & Krustrup, 2007). This is a common observation with 353 354 treadmill variants (Greig, Mc Naughton, & Lovell, 2006; Page et al., 2015), although, the mechanical strain 355 emulates a match more closely, with the current protocol eliciting a comparable CK response to elite soccer match-356 play (Malone et al., 2018).

## 357 Practical applications

358 The findings demonstrate that although CK indices are higher following 120 min of soccer-specific exercise, 359 functional measures of recovery are not exacerbated by the ET period versus 90-min markers. Therefore, 360 according to the current data, it appears that recovery practices do not need to be adapted, and training can resume 361 as normal following ET matches like traditional 90 min approaches. This finding conflicts with empirical research, 362 which suggests that 82% of the professional soccer practitioners surveyed believe that practices should be adapted 363 following ET compared with a typical 90 min match (Field et al., 2021). However, inter-individual variances in 364 recovery potential are apparent (Nédélec et al., 2012), and as such, recovery protocols following ET matches 365 should be individualised as opposed to generic for the entire team. Therefore, monitoring players following ET 366 using physical, physiological, and subjective assessments may be required to identify individuals experiencing 367 residual fatigue. If players are identified as needing an additional 'rest', then practitioners may need to consult the 368 head coach to ensure training loads are sufficient to maintain optimal conditioning. Careful periodisation of training is particularly key during fixture dense tournaments to ensure players are adequately conditioned foroptimal performance and a reduced injury incidence during ET.

#### 371 Conclusion

372 In summary, this study highlights that simulated soccer match-play of 120 min in duration is associated with 373 increased CK compared with a typical 90-min duration. However, there were no other trial and time interactions, 374 suggesting ET has no further effect on the muscle damage recovery response following simulated soccer. This 375 suggests a disassociation between the CK response and functional variables. Collectively, these data might have 376 implications for practitioners responsible for managing training and match loads during fixture dense tournaments 377 that require ET matches. There appears to be little need (according to the current data) for adapting recovery 378 practices in response to an ET match during fixture congested micro cycles and training can proceed as normal. 379 Moving forward, recovery and subsequent performance should be investigated following 90- and 120-min of 380 actual match-play.

381

#### 382 Acknowledgments

383 The authors gratefully appreciate the participants that willingly took part in the research.

## 384 Disclosure of interest

385 The authors report no conflict of interest

#### 386 Figure captions

- **387** Figure 1. A schematic of an individual 15-min bout of the soccer-specific exercise simulation
- 388
- 389
- 390
- 391

392

393

394

## 395 References

- Abbott, W., Brashill, C., Brett, A., & Clifford, T. (2020). Tart cherry juice: no effect on muscle function
   loss or muscle soreness in professional soccer players after a match. *International journal of sports physiology and performance*, 15(2), 249-254.
- Aoki, H., O'Hata, N., Kohno, T., Morikawa, T., & Seki, J. (2012). A 15-year prospective epidemiological
  account of acute traumatic injuries during official professional soccer league matches in Japan. *The American journal of sports medicine*, 40(5), 1006-1014.
- Baird, M. F., Graham, S. M., Baker, J. S., & Bickerstaff, G. F. (2012). Creatine-kinase-and exerciserelated muscle damage implications for muscle performance and recovery. *Journal of nutrition and metabolism*, 2012.
- 405 4. Bangsbo, J., Iaia, F. M., & Krustrup, P. (2007). Metabolic response and fatigue in soccer. *International journal of sports physiology and performance*, 2(2), 111-127.
- 407 5. Binderup, A. T., Arendt-Nielsen, L., & Madeleine, P. (2010). Pressure pain sensitivity maps of the neck408 shoulder and the low back regions in men and women. *BMC musculoskeletal disorders*, *11*(1), 1-7.

409 6. Borg, G. (1998). *Borg's perceived exertion and pain scales*: Human kinetics.

- 410 7. Brownstein, C. G., Dent, J. P., Parker, P., Hicks, K. M., Howatson, G., Goodall, S., & Thomas, K. (2017).
  411 Etiology and recovery of neuromuscular fatigue following competitive soccer match-play. *Frontiers in physiology*, *8*, 831.
- 8. Carling, C., McCall, A., Le Gall, F., & Dupont, G. (2015). The impact of in-season national team soccer
  play on injury and player availability in a professional club. *Journal of sports sciences*, 33(17), 17511757.
- 416 9. Castellano, J., Blanco-Villaseñor, A., & Alvarez, D. (2011). Contextual variables and time-motion
  417 analysis in soccer. *International journal of sports medicine*, *32*(06), 415-421.
- 418 10. Chen, T. C., Lin, K. Y., Chen, H. L., Lin, M. J., & Nosaka, K. (2011). Comparison in eccentric exercise419 induced muscle damage among four limb muscles. *European journal of applied physiology*, *111*(2), 211420 223.
- 421 11. Costello, N., Deighton, K., Dyson, J., Mckenna, J., & Jones, B. (2017). Snap-N-Send: A valid and reliable
  422 method for assessing the energy intake of elite adolescent athletes. *European journal of sport*423 science, 17(8), 1044-1055.
- 424 12. de Sousa, M. V., Fukui, R., Dagogo-Jack, S., Krustrup, P., Zouhal, H., & da Silva, M. E. R. (2019).
  425 Biomarkers of insulin action during single soccer sessions before and after a 12-week training period in
  426 type 2 diabetes patients on a caloric-restricted diet. *Physiology & behavior*, 209, 112618.
- 427 13. Di Salvo, V., Gregson, W., Atkinson, G., Tordoff, P., & Drust, B. (2009). Analysis of high intensity
  428 activity in Premier League soccer. *International journal of sports medicine*, *30*(03), 205-212.
- 429 14. Reilly, T., Drust, B., & Cable, N. (1999). Metabolic and physiological responses to a laboratory-based
  430 soccer-specific intermitent protocol on a non-monitorized treadmill. *Journal of Sports Sciences*, *18*, 811431 813.
- 432 15. Field, A., Naughton, R. J., Haines, M., Lui, S., Corr, L. D., Russell, M., Harper, L. D. (2020). The
  433 Demands of the Extra-Time Period of Soccer: A Systematic Review. *Journal of Sport and Health*434 *Science*.

- 435 16. Field, A., Corr, L. D., Haines, M., Lui, S., Naughton, R., Page, R. M., & Harper, L. D. (2020a).
  436 Biomechanical and Physiological Responses to 120 min of Soccer-Specific Exercise. *Research Quarterly*437 *for Exercise and Sport*, 1-13.
- 438 17. Field, A., Page, R. M., Corr, L., Naughton, R., Haines, M., Harper, L. D., & Hudson, S. (2020b). Lower439 Limb Muscle Excitation, Peak Torque, and External Load Responses to a 120-Minute Treadmill-Based
  440 Soccer-Specific Simulation. *Research Quarterly for Exercise and Sport*, 1-11.
- 441 18. Field, A., Corr, L., Thompson, C., Gonzalez Lucena, J. C., Sarmento, H., Naughton, R., ... & Harper, L.
  442 (2021). Recovery following the extra-time period of soccer: Practitioner perspectives and applied
  443 practices. *Biology of Sport*. DOI: https://doi.org/10.5114/biolsport.2022.104066
- 444 19. Gregson, W., Drust, B., Atkinson, G., & Salvo, V. D. (2010). Match-to-match variability of high-speed
  445 activities in premier league soccer. *International journal of sports medicine*, *31*(04), 237-242.
- 446 20. Greig, M. (2008). The influence of soccer-specific fatigue on peak isokinetic torque production of the
  447 knee flexors and extensors. *The American journal of sports medicine*, *36*(7), 1403-1409.
- Greig, M. P., Mc Naughton, L. R., & Lovell, R. J. (2006). Physiological and mechanical response to
  soccer-specific intermittent activity and steady-state activity. *Research in Sports Medicine*, 14(1), 29-52.
- 450 22. Haralambie, G., & Berg, A. (1976). Serum urea and amino nitrogen changes with exercise
  451 duration. *European journal of applied physiology and occupational physiology*, *36*(1), 39-48.
- 452 23. Harper, L. D., Hunter, R., Parker, P., Goodall, S., Thomas, K., Howatson, G., & Russell, M. (2016). Test453 retest reliability of physiological and performance responses to 120 minutes of simulated soccer match
  454 play. *Journal of strength and conditioning research*, 30(11), 3178-3186.
- 455 24. Hermens, H. J., Freriks, B., Disselhorst-Klug, C., & Rau, G. (2000). Development of recommendations
  456 for SEMG sensors and sensor placement procedures. *Journal of electromyography and*457 *Kinesiology*, 10(5), 361-374.
- 458 25. Hills, S. P., Radcliffe, J. N., Barwood, M. J., Arent, S. M., Cooke, C. B., & Russell, M. (2020).
  459 Practitioner perceptions regarding the practices of soccer substitutes. *PloS one*, *15*(2), e0228790.
- 460 26. Jones, A. M., & Doust, J. H. (1996). A 1% treadmill grade most accurately reflects the energetic cost of
  461 outdoor running. *Journal of sports sciences*, 14(4), 321-327.
- 462 27. Julian, R., Page, R. M., & Harper, L. D. (2020). The Effect of Fixture Congestion on Performance During
  463 Professional Male Soccer Match-Play: A Systematic Critical Review with Meta-Analysis. Sports
  464 Medicine, 1-19.
- 465 28. Kołodziejczyk, M., Chmura, P., Milanovic, L., Konefał, M., Chmura, J., Rokita, A., & Andrzejewski,
  466 M. (2021). How did three consecutive matches with extra time affect physical performance? A case study
  467 of the 2018 football Men's World Cup. *Biology of Sport, 38*(1).
- 468 29. López-Valenciano, A., Ruiz-Pérez, I., Garcia-Gómez, A., Vera-Garcia, F. J., Croix, M. D. S., Myer, G.
  469 D., & Ayala, F. (2020). Epidemiology of injuries in professional football: a systematic review and meta470 analysis. *British journal of sports medicine*, 54(12), 711-718.
- 30. Malone, S., Mendes, B., Hughes, B., Roe, M., Devenney, S., Collins, K., & Owen, A. (2018). Decrements
  in neuromuscular performance and increases in creatine kinase impact training outputs in elite soccer
  players. *The Journal of Strength & Conditioning Research*, 32(5), 1342-1351.

- 474 31. Meyer, T., & Meister, S. (2011). Routine blood parameters in elite soccer players. *International journal*475 *of sports medicine*, 32(11), 875-881.
- 476 32. Mohr, M., Krustrup, P., & Bangsbo, J. (2003). Match performance of high-standard soccer players with
  477 special reference to development of fatigue. *Journal of sports sciences*, *21*(7), 519-528.
- 478 33. Mohr, M., Krustrup, P., & Bangsbo, J. (2005). Fatigue in soccer: a brief review. *Journal of sports*479 sciences, 23(6), 593-599.
- 480 34. Nedelec, M., McCall, A., Carling, C., Legall, F., Berthoin, S., & Dupont, G. (2014). The influence of
  481 soccer playing actions on the recovery kinetics after a soccer match. *The Journal of Strength & Conditioning Research*, 28(6), 1517-1523.
- 483 35. Nédélec, M., McCall, A., Carling, C., Legall, F., Berthoin, S., & Dupont, G. (2012). Recovery in
  484 soccer. *Sports medicine*, 42(12), 997-1015.
- 36. Nédélec, M., McCall, A., Carling, C., Legall, F., Berthoin, S., & Dupont, G. (2013). Recovery in
  soccer. *Sports medicine*, 43(1), 9-22.
- 487 37. Oliver, J., Armstrong, N., & Williams, C. (2008). Changes in jump performance and muscle activity
  488 following soccer-specific exercise. *Journal of sports sciences*, *26*(2), 141-148.
- 489 38. Page, R. M., Marrin, K., Brogden, C. M., & Greig, M. (2015). Biomechanical and physiological response
  490 to a contemporary soccer match-play simulation. *The Journal of Strength & Conditioning Research*,
  491 29(10), 2860-2866.
- 492 39. Page, R. M., Marrin, K., Brogden, C. M., & Greig, M. (2019). Physical Response to a Simulated Period
  493 of Soccer-Specific Fixture Congestion. *The Journal of Strength & Conditioning Research*, 33(4), 1075494 1085.
- 495 40. Peñas, C. L., Dellal, A., Owen, A. L., & Gómez-Ruano, M. Á. (2015). The influence of the extra-time
  496 period on physical performance in elite soccer. *International Journal of Performance Analysis in Sport*,
  497 15(3), 830-839. doi:10.1080/24748668.2015.11868834
- 498 41. Proske, U., & Morgan, D. L. (2001). Muscle damage from eccentric exercise: mechanism, mechanical
  499 signs, adaptation and clinical applications. *The Journal of physiology*, *537*(2), 333-345.
- 42. Rampinini, E., Bosio, A., Ferraresi, I., Petruolo, A., Morelli, A., & Sassi, A. (2011). Match-related fatigue
  in soccer players. *Medicine & Science in Sports & Exercise*, 43(11), 2161-2170.
- 43. Ranchordas, M. K., Dawson, J. T., & Russell, M. (2017). Practical nutritional recovery strategies for elite
   soccer players when limited time separates repeated matches. *Journal of the International Society of Sports Nutrition, 14*(1), 35.
- 44. Russell, M., Benton, D., & Kingsley, M. (2011). The effects of fatigue on soccer skills performed during
  a soccer match simulation. *International journal of sports physiology and performance*, 6(2), 221-233.
- 507 45. Russell, M., Sparkes, W., Northeast, J., & Kilduff, L. P. (2015a). Responses to a 120 min reserve team
  508 soccer match: a case study focusing on the demands of extra time. *Journal of sports sciences*, 33(20),
  509 2133-2139.
- 46. Russell, M., Northeast, J., Atkinson, G., Shearer, D. A., Sparkes, W., Cook, C. J., & Kilduff, L. P.
  (2015b). Between-match variability of peak power output and creatine kinase responses to soccer matchplay. *The Journal of Strength & Conditioning Research*, 29(8), 2079-2085.

- 513 47. Shephard, R. J. (1999). Biology and medicine of soccer: an update. *Journal of Sports Sciences*, 17(10),
  514 757-786.
- 515 48. Small, K., McNaughton, L., Greig, M., & Lovell, R. (2010). The effects of multidirectional soccer516 specific fatigue on markers of hamstring injury risk. *Journal of Science and Medicine in Sport, 13*(1),
  517 120-125.
- 518 49. Stevenson, E. J., Watson, A., Theis, S., Holz, A., Harper, L. D., & Russell, M. (2017). A comparison of
  519 isomaltulose versus maltodextrin ingestion during soccer-specific exercise. *European Journal of Applied*520 *Physiology*, *117*(11), 2321-2333. doi:10.1007/s00421-017-3719-5
- 50. Stratford, P. W., & Goldsmith, C. H. (1997). Use of the standard error as a reliability index of interest:
  an applied example using elbow flexor strength data. *Physical therapy*, 77(7), 745-750.
- 51. Thomas, K., Dent, J., Howatson, G., & Goodall, S. (2017). Etiology and recovery of neuromuscular
  fatigue following simulated soccer match-play. *Medicine & Science in Sports & Exercise*, 49(5), 955964.
- 526 52. Tønnessen, E., Hem, E., Leirstein, S., Haugen, T., & Seiler, S. (2013). Maximal aerobic power
  527 characteristics of male professional soccer players, 1989–2012. *International journal of sports*528 *physiology and performance*, 8(3), 323-329.
- 529 53. Twist, C., & Highton, J. (2013). Monitoring fatigue and recovery in rugby league players. *International journal of sports physiology and performance*, 8(5), 467-474.
- 54. Uddin, N., Jeffries, O., Read, P., Howe, L., Patterson, S., & Waldron, M. (2020). Physiological responses
  to linear and nonlinear soccer-specific match simulations and their effects on lower-limb muscle fatigue. *The Journal of Strength & Conditioning Research*, 34(11), 3232-3240.
- 534 55. Waldron, M., & Highton, J. (2014). Fatigue and pacing in high-intensity intermittent team sport: an
  535 update. *Sports medicine*, 44(12), 1645-1658.
- 56. Walton, D., MacDermid, J., Nielson, W., Teasell, R., Chiasson, M., & Brown, L. (2011). Reliability,
  standard error, and minimum detectable change of clinical pressure pain threshold testing in people with
  and without acute neck pain. *journal of orthopaedic & sports physical therapy*, 41(9), 644-650.
- 539 57. Winder, N., Russell, M., Naughton, R. J., & Harper, L. D. (2018). The impact of 120 minutes of match540 play on recovery and subsequent match performance: a case report in professional soccer
  541 players. *Sports*, 6(1), 22.
- 542 58. Zhang, W., Yu, Q., Siddiquie, B., Divakaran, A., & Sawhney, H. (2015). "snap-n-eat" food recognition
  543 and nutrition estimation on a smartphone. *Journal of diabetes science and technology*, 9(3), 525-533.

Mean dietary intake across 5 days			Inter-rater reliability		Intra-rater reliability		
Variable	90 min	120 min	<i>p</i> value	SEM	95% CI	SEM	95% CI
Energy (Kcal)	$2008\pm577$	$1985\pm486$	0.68	95	186	53	104
Carbohydrate (g)	$237\pm93$	$220\pm67$	0.35	12	23	4	7
Protein (g)	$109\pm40$	$109\pm47$	0.93	2	4	1	2
Fat (g)	$75\pm18$	$77 \pm 18$	0.76	3	6	4	8

Table 1. Mean energy and macro nutrient composition of participants diet across five days of testing for each trial and reliability of data input

Data are reported as mean  $\pm SD$ 

**Table 2.** Environmental conditions and responses across both trials.

	1			
Variable	90 min		120 min	<i>p</i> value
Urine osmolality (pre; mOsm·kg <sup>-1</sup> )	$536\pm120$		$556\pm167$	0.79
Urine osmolality (post; mOsm·kg <sup>-1</sup> )	$516\pm104$		$530\pm160$	0.83
Humidity (%)	$33.8\pm3.5$		$33.3 \pm 3.1$	0.66
Barometric pressure (%)	$29.2\pm0.4$		$29.4\pm0.6$	0.79
Ambient temperature (°C)	$20.1\pm0.9$		19. $7 \pm 1.1$	0.33
Variable	90 min	90 min (120 min)	<i>p</i> value	120 min
Variable PlayerLoad™ (a.u)	$\frac{90 \text{ min}}{1359 \pm 21}$	$\frac{90 \min{(_{120 \min})}}{1347 \pm 25}$	<i>p</i> value 0.82	$\frac{120 \text{ min}}{1817 \pm 25}$
Variable PlayerLoad <sup>™</sup> (a.u) Blood lactate (mmol · L <sup>-1</sup> )	$     \begin{array}{r}       90 \text{ min} \\       1359 \pm 21 \\       2.4 \pm 1.5     \end{array} $	$\frac{90 \min (_{120 \min})}{1347 \pm 25} \\ 2.4 \pm 1.5$	<i>p</i> value 0.82 0.98	$\frac{120 \text{ min}}{1817 \pm 25} \\ 2.5 \pm 1.5$
VariablePlayerLoad <sup>TM</sup> (a.u)Blood lactate (mmol $\cdot L^{-1}$ )RPE-L	$     90 min     1359 \pm 21     2.4 \pm 1.5     12 \pm 2     $	$\begin{array}{r} 90 \text{ min } (_{120 \text{ min}}) \\ 1347 \pm 25 \\ 2.4 \pm 1.5 \\ 11 \pm 2 \end{array}$	<i>p</i> value 0.82 0.98 0.65	$     \begin{array}{r}       120 \text{ min} \\       1817 \pm 25 \\       2.5 \pm 1.5 \\       12 \pm 2     \end{array} $
Variable PlayerLoad <sup>TM</sup> (a.u) Blood lactate (mmol · L <sup>-1</sup> ) RPE-L RPE-B	$     \begin{array}{r} 90 \text{ min} \\     1359 \pm 21 \\     2.4 \pm 1.5 \\     12 \pm 2 \\     11 \pm   \end{array} $	$\begin{array}{r} 90 \text{ min } (_{120 \text{ min}}) \\ 1347 \pm 25 \\ 2.4 \pm 1.5 \\ 11 \pm 2 \\ 11 \pm 2 \end{array}$	<i>p</i> value 0.82 0.98 0.65 0.95	$     \begin{array}{r}       120 \text{ min} \\       1817 \pm 25 \\       2.5 \pm 1.5 \\       12 \pm 2 \\       12 \pm 2     \end{array} $
Variable PlayerLoad <sup>™</sup> (a.u) Blood lactate (mmol · L <sup>-1</sup> ) RPE-L RPE-B RPE-O	$90 \text{ min} \\ 1359 \pm 21 \\ 2.4 \pm 1.5 \\ 12 \pm 2 \\ 11 \pm \\ 12 \pm 2$	$\begin{array}{r} 90 \text{ min } (_{120 \text{ min}}) \\ 1347 \pm 25 \\ 2.4 \pm 1.5 \\ 11 \pm 2 \\ 11 \pm 2 \\ 12 \pm 2 \end{array}$	<i>p</i> value 0.82 0.98 0.65 0.95 0.83	$     \begin{array}{r}       120 \text{ min} \\       1817 \pm 25 \\       2.5 \pm 1.5 \\       12 \pm 2 \\       12 \pm 2 \\       12 \pm 2     \end{array} $

*RPE-L* = *RPE legs; RPE-B* = *RPE breathlessness; RPE-O* = *RPE overall.* 

Urine and environmental significance values are reported as differences between both 90- and 120-min trials. 90 min  $(_{120 \text{ min}}) = 90$  min duration for the 120 min trial.

Responses to exercise significance values are reported as differences between the 90 min duration of both trials. PlayerLoad<sup>TM</sup> values are reported as the cumulative response across the specified duration.

Blood lactate and RPE values are reported as the mean response per epoch of exercise.

Data are reported as mean  $\pm$  SD.

Variable	ICC (%)
СК	14.73
Creatinine	20.57
Urea	41.49*
AST	11.47
VAS	46.32*
PPT <sub>RF</sub>	85.03*
PPT <sub>BF</sub>	63.65*
RSI	86.09*
СМЈ	95.14*
$eccKF_{180}$	57.77*
eccKF <sub>270</sub>	64.27*
eccKF <sub>60</sub>	66.24*

**Table 3.** The ICC's (%) of the random factor of participant ID for all recovery variables. Where significance was found, participant ID was included in the linear mixed model.

\* Represents significant determinant of variance within the linear mixed model (p < 0.05).

<b>Tuble 1.</b> Musele dumage receivery responses remembers former 120 mm of simulated second
---

Time								
Variable	Baseline	Post	24 hr	48 hr	72 hr			
$CK (U \cdot L^{-1})$								
90 min	$102.7\pm24.7$	$229.6\pm25.6^{\rm a}$	$298.8\pm29.3^{ab}$	$258.6\pm28.9^{\text{ac}}$	$154.1\pm29.3^{acd}$			
120 min	$109.4\pm25.7$	$233.8\pm26.8^{\rm a}$	$455.1\pm29.4^{*ab}$	$301.6\pm30.8^{\text{ac}}$	$244.1\pm24.7^{\text{*acd}}$			
Creatinine (µmol·L <sup>-1</sup> )								
90 min	$77.6\pm4.7$	$80.8\pm4.9$	$79.9\pm4.7$	$72.3\pm4.7$	$74.2 \pm 5.1$			
120 min	$68.1\pm4.9$	$89.6 \pm 5.1$	$80.2\pm5.1$	$77.1 \pm 4.7$	$72.1\pm4.9$			
Urea (mmol·L <sup>-1</sup> )								
90 min	$6.64\pm0.53$	$7.97\pm0.53^{\rm a}$	$8.05\pm0.53^{\rm a}$	$6.84\pm0.52^{\mathrm{bc}}$	$6.52\pm0.55^{\rm bc}$			
120 min	$6.91\pm0.52$	$9.01\pm0.52^{\rm a}$	$8.27\pm0.54^{\rm a}$	$7.48\pm0.52^{\mathrm{bc}}$	$7.51\pm0.53^{\rm bc}$			
$AST (U \cdot L^{-1})$								
90 min	$41.4\pm3.2$	$44.01\pm3.3$	$45.5\pm3.2$	$39.6\pm3.8$	$41.0 \pm 3.4$			
120 min	$38.0\pm3.1$	$40.07\pm3.2$	$44.5\pm3.3$	$41.7\pm3.4$	$37.9\pm3.1$			
VAS								
90 min	$2\pm 2$	$20\pm2^{\mathrm{a}}$	$22\pm2^{\mathrm{a}}$	$14\pm3^{\mathrm{ac}}$	$6\pm3^{abcd}$			
120 min	$1\pm 2$	$26\pm3^{\mathrm{a}}$	$23\pm3^{\mathrm{a}}$	$18\pm2^{\mathrm{ac}}$	$5\pm2^{abcd}$			
$PPT_{BF}(lbs)$								
90 min	$25 \pm 1$	25±1	$24 \pm 1$	$24 \pm 1$	$26 \pm 1^{cd}$			
120 min	$22 \pm 1$	$22 \pm 1$	$22 \pm 1$	$22 \pm 1$	$24 \pm 1^{cd}$			
RSI (a.u)								
90 min	$1.41\pm0.12$	$1.31\pm0.12^{\rm a}$	$1.30\pm0.11^{\rm a}$	$1.31\pm0.11^{\rm a}$	$1.37\pm0.12^{bcd}$			
120 min	$1.33\pm0.12$	$1.26\pm0.12^{\rm a}$	$1.23\pm0.12^{\rm a}$	$1.25\pm0.12^{\rm a}$	$1.40\pm0.12^{bcd}$			
CMJ (cm)								
90 min	$36.6\pm2.2$	$35.4\pm2.2$	$34.9\pm2.2^{ab}$	$35.3\pm2.2^{ac}$	$36.6\pm2.2^{\text{ce}}$			
120 min	$36.4\pm2.2$	$36.3\pm2.2$	$34.9\pm2.2^{ab}$	$36.0\pm2.2^{\rm\ ac}$	$36.5\pm2.2^{\text{ce}}$			
$eccKF_{180}$ (Nm)								
90 min	$160.4\pm8.7$	$153.4\pm8.7^{\rm a}$	$146.0\pm8.7^{\rm a}$	$154.3\pm8.8$	$163.8\pm8.9^{bcd}$			
120 min	$162.3\pm8.7$	$145.2\pm8.7^{\rm a}$	$145.9\pm8.9^{\rm a}$	$153.8\pm8.7$	$163.7\pm8.8^{bcd}$			
$eccKF_{270}$ (Nm)								
90 min	$159.4\pm8.8$	$153.7\pm8.8^{\rm a}$	$152.8\pm8.8^{\rm a}$	$160.4\pm8.9^{\mathrm{bc}}$	$164.5\pm8.9^{bc}$			
120 min	$159.0\pm8.8$	$147.1\pm8.8^{\rm a}$	$145.5\pm8.9^{\rm a}$	$161.8\pm8.8^{\rm bc}$	$167.5\pm8.8^{bc}$			
$eccKF_{60}$ (Nm)								
90 min	$164.1\pm7.6$	$148.9\pm7.6^{\rm a}$	$142.5\pm7.5^{ab}$	$148.0\pm7.5^{\rm ac}$	$159.0\pm7.5^{\circ}$			
120 min	$159.5 \pm 7.6$	$147.2\pm7.5^{\rm a}$	$137.9\pm7.7^{ab}$	$149.3\pm7.5^{\rm ac}$	$160.7 \pm 7.5^{\circ}$			

Note. CK = creatine kinase = AST = aminotransferase, PPT = pain pressure threshold,  $_{BF} = biceps$  femoris, RSI = reactive strength index, CMJ = countermovement jump height, eccKF = eccentric knee flexion,  $_{180, 270, 60}$  denote isokinetic angular velocities (deg·s<sup>-1</sup>), VAS = Visual analogue scale. \*Represents significantly higher value for specified time point between trials. <sup>a-e</sup> Represents significant difference for time from baseline, post, 24, 48 and 72 hr, respectively.

Data are reported as mean  $\pm$  SE.

