


**Please cite the Published Version**

Filho, Renato Guilherme Trede, Rodrigues, Thamires Cristina Perdigão, Brant, Alícia Correa, Rodrigues, Nara Lourdes Moreno, Mazuquin, Bruno  and Richard, Jim (2023) Classification of foot type from podography: correlation of results between six quantitative assessment methods. *Journal of the Foot & Ankle*, 17 (1). pp. 29-33.

**DOI:** <https://doi.org/10.30795/jfootankle.2023.v17.1690>

**Publisher:** Associação Brasileira de Medicina e Cirurgia do Tornozelo e Pé - ABTPé

**Version:** Published Version

**Downloaded from:** <https://e-space.mmu.ac.uk/631923/>

**Usage rights:**  [Creative Commons: Attribution-Noncommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/)

**Additional Information:** This is an Open Access article published in *Journal of the Foot & Ankle*.

**Enquiries:**

If you have questions about this document, contact [openresearch@mmu.ac.uk](mailto:openresearch@mmu.ac.uk). Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

## Original Article

# Classification of foot type from podography: correlation of results between six quantitative assessment methods

Renato Guilherme Trede Filho<sup>1</sup>, Thamires Cristina Perdigão Rodrigues<sup>1</sup>, Alícia Correa Brant<sup>1</sup>,  
Nara Lourdes Moreno Rodrigues<sup>1</sup>, Bruno Fles Mazuquim<sup>2</sup>, Jim Richard<sup>3</sup>

1. Universidade Federal dos Vales do Jequitinhonha e Mucuri - UFVJM, Campus JK, Diamantina, MG, Brazil.

2. Manchester Metropolitan University, Manchester, United Kingdom.

3. University of Central Lancashire - UCLan, Preston, United Kingdom.

## Abstract

**Objective:** Apply the different methods available in the literature to classify a sample of podography and evaluate the level of agreement between the results.

**Methods:** Six quantitative and one qualitative method to classify foot type from podography were recorded on 30 feet. The podography indexes were calculated, and the level of agreement between methods was explored.

**Results:** Correlation values were above  $r = 0.84$  except for the test arch footprint angle. The highest correlation values were found between the truncated arch index and footprint index (0.99), arch index and footprint index (-0.94), and arch index and truncated arch index (-0.94), and the lowest was the arch footprint angle with the other parameters. However, there was a difference in the classification between the foot types, indicating a lack of agreement of thresholds between foot types. Qualitative visual inspection was the faster method to classify foot type.

**Conclusion:** The visual inspection was the fastest test to apply, followed by the quantitative arch footprint angle test. High correlation values were found between tests, especially the arch index and the footprint index, arch-length index, truncated arch index, and Chippaux-Smirak index tests.

**Level of Evidence IV; Therapeutic Studies; Case Series.**

**Keywords:** Evaluation; Foot; Podography.

## Introduction

Feet are the base of support to the human body, allowing functional tasks such as standing, gait, and running<sup>(1)</sup>. Foot alignment can be classified as normal, flat, or high-arch<sup>(2)</sup>. Misalignments of the foot in stance position or during movement can overload interrelated muscles and joints, increasing the risk of injuries<sup>(3)</sup>. Evidence has shown high correlations between flat and high-arch foot with stress fractures<sup>(4)</sup>. Flat foot is associated with a greater number of

injuries reported in midfoot and knee in a cadet population<sup>(5)</sup>. Another study found that runners with high-arched foot present a greater incidence of ankle and bone injuries and injuries on the lateral anatomical structures of lower limbs<sup>(6)</sup>; runners with low-arched foot are more likely to develop soft tissue injuries on the medial side of the lower extremity, and knee pain<sup>(7)</sup>.

Foot alignment classification can be performed by qualitative techniques, such as visual inspection (VI), or quantitative

Study performed at the Universidade Federal dos Vales do Jequitinhonha e Mucuri - UFVJM, Diamantina, MG, Brazil.

**Correspondence:** Renato Guilherme Trede Filho. Universidade Federal dos Vales do Jequitinhonha e Mucuri - UFVJM, Campus JK - Rodovia MGT 367 - Km 583, N° 5000, Bairro Alto da Jacuba, 39100-000, Diamantina, MG, Brazil. **E-mail:** [renato.trede@gmail.com](mailto:renato.trede@gmail.com) **Conflicts of interest:** none. **Source of funding:** none.

**Date received:** February 13, 2023. **Date accepted:** March 30, 2023. **Online:** April 30, 2023.



methods using baropodometry<sup>(8)</sup> or podography<sup>(9)</sup>. Visual inspection is a method to assess the arch and foot alignment and is widely used by physicians; however, their classification is subjective and has high inter-rater variability<sup>(10)</sup>. Regarding quantitative methods, podographs are low-cost and easier to apply compared to baropodometry<sup>(11)</sup>; podography has lower variability than VI<sup>(12)</sup>. The interpretation of podography can be based on different methods, such as VI, arch index (AI), arch footprint angle (AFA), footprint index (FI), arch-length index (ALI), truncated arch index (TAI), and Chippaux-Smirak index (CSI)<sup>(12-17)</sup>. However, each technique uses different parameters to classify foot posture, and some do not present cut-off thresholds between classifications. Furthermore, the parameters used to classify the foot for each podography method are different<sup>(12-17)</sup>; it is important to clarify whether the agreement between techniques is satisfactory to allow physicians to use their preferred choice. Thus, the objective of this study is to compare the efficiency of different parameters used in the literature to classify foot types from podographic images and determine the levels of agreement between them.

## Methods

A sample of 15 volunteers (11 women and four men) was recruited by convenience through posters fixed at the university physiotherapy clinic. Both feet were assessed, giving a sample of 30 feet. The inclusion criteria of the volunteers were: A) age between 18-25 years, B) capability of staying in a uni-pedal stance, and C) ability to walk without assistance. Exclusion criteria were: A) existing foot deformities or amputations, B) pain in the lower limb during stance, and C) balance deficit or postural instability when in a uni-pedal stance.

Podography was obtained using an APEX Harris Mat Set (Apex Foot Products Corporation, Englewood, NJ, USA). This instrument consists of two rectangular plastic plates mediated by textured rubber. To obtain a podography with homogeneous color, an equivalent smooth rubber replaced the textured rubber. A Deskjet 2050 scanner (Hewlett-Packard Development Company, LP, Hanover Street, California, CA, USA) was used to digitalize the podography. In addition, a universal goniometer (Carci, São Paulo, SP, Brazil) was used to measure angles for the AFA test.


The study was approved by the Human Research Ethics Committee at UFVJM, and informed consent was obtained from all participants. Demographic data collected were age, gender, body mass index (BMI), height, shoe size, history of foot and ankle injuries, level of physical activity, and limb dominance. After preparing the foot printer with a layer of black ink, a white paper was positioned inside the mat, and the volunteer was asked to keep one foot centralized on the APEX Harris Mat Set. Then, the volunteer was asked to elevate the opposite foot and sustain a uni-pedal position for 30 seconds. Subsequently, the podography was re-inked, and the same procedure was undertaken with the contralateral leg. During the test, the examiner ensured the knee alignment

remained in a neutral position. Volunteers were allowed to rest their hands on a table to maintain stability during a uni-pedal stance.

The same researcher, previously trained, obtained and analyzed all podography. Podography order was randomized before applying qualitative and quantitative tests, described below, to classify the foot as flat, neutral, or high-arched. In addition, podographies were scanned with a resolution of 200 dots per inch and transferred to software developed in MatLab (MathWorks, Massachusetts, MA, USA). This software calculates the number of black and white pixels inside the scanned image and the foot area (Figure 1).

The podography was divided into three equal sections, proportional to the length of the foot, by two parallel lines excluding the toes to calculate the AI. Then, the printed area of each section was calculated, and the index was calculated as the ratio between the midfoot area B and the area of the whole foot (A + B + C), as shown in Figure 1(I).

The AFA was calculated from the intersection of two lines; the first was a line connecting the prominent points of the calcaneus and the first metatarsal. The second line was drawn towards the concavity of the foot arch following the medial foot border. The angle between the two lines was obtained by a goniometer, as shown in Figure 1(II). The FI was calculated by the formula A/B, where A was the arch area, delimited by the medial line connecting the most prominent point in the



Arch index =  $B/(A+B+C)$ ; (II) Arch footprint angle; (III) Footprint index =  $A/B$ ; (IV) Arch-length index =  $PQ/r$ ; (V) Truncated arch index =  $A/B$ ; (VI) Chippaux-Smirak index =  $C/B$ ; (VII) Visual inspection =  $A \geq 1/2$  of B or  $A \leq 1/2$  of C.

**Figure 1.** Methods to assess podography.

medial aspect of the forefoot and the rearfoot, and B was the area of the toeless podography as shown in Figure 1(III).

To calculate the ALI, the length of a medial line (PQ), which starts at the most prominent point on the medial aspect of the forefoot and ends at the most prominent point on the medial aspect of the rearfoot, was measured. The length of the “r” line was obtained starting from the same point defined on the forefoot, following the contour of the arch, and ending at the same medial point on the rearfoot. The ALI was obtained by calculating the ratio of the length of the medial line and the arc length (PQ/r), as shown in Figure 1(IV).

The TAI was calculated as the FI; however, the foot area (B) was limited to the truncated area. The truncated area is restricted between two horizontal lines through the most medial points of the metatarsal and the heel region, as shown in Figure 1(V) and defined as detailed in the ALI test.

The CSI was drawn as a lateral line connecting the prominent points of the calcaneus and the fifth metatarsal. In addition, two perpendicular lines with a 90° angle to the lateral line were drawn in the podography. One line in the wider zone of the forefoot (B) and another line in the narrower zone of the midfoot (C). The result was obtained by calculating C/B as shown in Figure 1(VI).

Visual inspection of podographs was performed according to the criteria presented by Viladot (1987). According to the Viladot method (Figure 1(VII)), the flat foot is present when the midfoot width is equal to or greater than half the width of the forefoot. High-arched foot is present when the width of the midfoot impression is equal to or less than half of the rearfoot or not visible. Podography not classified as flat or high-arched foot was considered neutral<sup>(18)</sup>.

## Data analysis

Data analysis was performed using SPSS version 11.0 (SPSS Inc. Chicago, IL, USA). Initially, the normality of the variables was tested using the Shapiro-Wilk test. The quantitative scores tests were analyzed using Spearman’s correlation coefficient, considering a significance level of 5%.

## Results

Eleven women and four men were recruited, and the mean age was 23 years (SD = 2.7). According to their BMI, ten participants were classified as normal, two as pre-obese, one as mild thinness, one as moderate thinness, and one as severe thinness. Regarding regular physical activity, ten were physically inactive, while five stated to practice exercise.

Table 1 describes the frequency of different foot categories when using VI, AI, AFA, and CSI. Categorization among the four tests was not similar. The AFA test categorized the sample into the high-arch classification. Conversely, the CSI categorized most of the sample as neutral and flat arched foot.

Table 2 shows the results of Spearman’s correlation coefficient among the six different categorization methods. Correlation values were above  $r = 0.84$  except for the AFA test. The highest correlation values were found between TAI and FI (0.99), AI and FI (-0.94), and AI and TAI (-0.94), and the lowest was the AFA with the other parameters.

Table 3 shows the mean time taken to classify individuals’ feet using different methods. The VI was the quickest method compared to all other approaches.

**Table 1.** Feet categorization according to the different indexes

	High-arch (n)	High-arch (%)	Neutral (n)	Neutral (%)	Flat arch (n)	Flat arch (%)
Visual inspection	9	30%	12	40%	9	30%
Arch index	15	50%	11	37%	4	13%
Arch footprint angle	27	90%	2	7%	1	3%
Chippaux-Smirak index	1	3%	16	53%	13	44%

**Table 2.** Spearman’s correlation coefficient among six different scores

	AI	FI	ALI	TAI	AFA	CSI
AI	1.0000					
FI	- 0.9431	1.0000				
ALI	0.8413	- 0.8853	1.0000			
TAI	- 0.9400	0.9933	- 0.8780	1.0000		
AFA	- 0.4883	0.5469	- 0.6276	0.5218	1.0000	
CSI	0.8836	- 0.7751	0.6703	- 0.7794	- 0.3844	1.0000

AI: Arch index; FI: Footprint index; ALI: Arch-length index; TAI: Truncated arch index; AFA: Arch footprint angle; CSI: Chippaux-Smirak Index.

**Table 3.** Mean time taken in seconds for feet classification using different methods

VI	AI	AFA	FI	ALI	TAI	CSI
1.03	646	19	814	52	832	33

VI: Visual inspection; AI: Arch index; AFA: Arch footprint angle; FI: Footprint index; ALI: Arch-length index; TAI: Truncated arch index; CSI: Chippaux-Smirak index.

## Discussion

The aim of our study was to assess the agreement between different methods to classify a sample of podography. The hypothesis was that the quantitative tests would show high correlation scores; therefore, regardless of the method used by physicians, patients would be categorized under the same group. Moreover, because the time taken to perform each test was recorded, it would be possible to show which test is faster to use, although producing similar results.

Many different methods for podography assessment are available in the literature; however, many have been published before 1987, making access to the full text challenging<sup>(12,13,19-24)</sup>. In addition, many methods have been adapted and do not follow the original procedure, which can be misleading and result in systematic errors<sup>(25-27)</sup>.

The findings of this study demonstrated that the quantitative and qualitative methods might not classify the podography under the same category. According to Forriol and Pascual<sup>(17)</sup>, the classification is sensitive to the method used. Moreover, the cut-off points differed, which may be the main reason for the low correlation scores<sup>(21)</sup>. Another important point is that because of soft tissue malleability, feet with similar structural shapes can have different podography<sup>(25)</sup>. According to Razeghi and Batt<sup>(28)</sup>, quantitative methods are discouraged when assessing high-arch feet because of the medial area discontinuity.

The different quantitative indexes can be categorized as those that use total area, angle values, or length measurements in their formula. These methods require additional time to perform the calculations needed for final results. As shown in Table 3, the VI, which does not need extra time, was the quickest way to assess the foot. However, the results from VI may vary according to the assessor’s experience, making their results less reliable than more systematic methods<sup>(24)</sup>. However, Dahle et al.<sup>(21)</sup> demonstrated that experienced physiotherapists have high agreement rates while using VI.

One way of improving VI reliability could be by using more systematic methods, which in this study was the criteria proposed by Viladot<sup>(18)</sup>. Xiong et al.<sup>(22)</sup> demonstrated a high correlation between AI and VI. Therefore, in the clinical

setting, the VI is of great value as it is quick and does not require any equipment, which may not always be available in the clinical setting.

After VI, the methods that used the angle or length of the podography were those that required less time to classify the podography. Among them, the AFA was the quickest. These tests require only drawing lines dividing the podography into areas. They are measured in millimeters with a simple ruler, or in the case of angles, lines are traced and then measured with a goniometer. However, the ALI does not have cut-off points, which are only available for the AFA and the CSI in this category of tests.

Regarding AFA and the CSI, a recent study demonstrated the advantage of a 5-level stratification system (high, normal, intermediary, lowered, and flat medial longitudinal arch). However, Sacco et al.<sup>(27)</sup> study, comparing the AI, CSI, and AFA, demonstrated that only the AI and the CSI are reliable for anthropometric assessment as the AFA showed strong disagreement with the other indexes.


The tests using the area for podography classification were more time-consuming. The AI is the most commonly described in the literature and was the fastest among the indexes in this group<sup>(29,30)</sup>. The extra time needed for this index is due to the additional steps, such as digital processing of the image or other equipment, such as planimeters, to obtain the figure area.

Only the AI has cut-off points established among the three methods in the group of indexes using area for classification. Therefore, although the FI and the TAI have similar scores to the AI, it is not possible to determine if the same cut-point from the AI can be used for the other two<sup>(31-33)</sup>.

Regarding the correlation scores, the high correlation scores among tests found were opposite to Hawes et al.<sup>(17)</sup>, where the authors performed a regression analysis in relation to the plantar arch height with five different podography indexes: AFA, FI, AI, ALI, and TAI. In our findings, most of the correlation scores were over 0.84 with the AI, which may suggest that the cut-off points from this index can potentially be adopted for the others, but more studies are needed. The test with the lowest correlation values was the ALI; however, because it was the one with the shortest time needed, it might also be useful in a clinical setting as it has cut-off points that enable the classification of flat, normal, or high-arched foot.

## Conclusion

The visual inspection was the fastest test to apply, followed by the quantitative arch footprint angle test. High correlation values were found between tests, especially the arch index and the footprint index, arch-length index, truncated arch index, and Chippaux-Smirak index tests.

**Authors' contributions:** Each author contributed individually and significantly to the development of this article: RGTF \*(<https://orcid.org/0000-0001-6118-1181>) Data collection and clinical examination; TCP \*(<https://orcid.org/0000-0002-5245-1298>) Survey of the medical records and data collection; ACB \*(<https://orcid.org/0000-0003-3670-3616>) Participated in the review process, formatting of the article and approved the final version; NLMR \*(<https://orcid.org/0000-0003-1981-8606>) Bibliographic review and formatting of the article; BFM \*(<https://orcid.org/0000-0003-1566-9551>) Interpreted the results of the study and statistical analysis; JR \*(<https://orcid.org/0000-0002-4004-3115>) Conceived and planned the activities that led to the study and approved the final version. All authors read and approved the final manuscript.\*ORCID (Open Researcher and Contributor ID) 

## References

- Souza TR, Pinto RZ, Trede RG, Kirkwood RN, Fonseca ST. Temporal couplings between rearfoot-shank complex and hip joint during walking. *Clin Biomech (Bristol, Avon)*. 2010;25(7):745-8.
- Razeghi M, Batt ME. Foot type classification: a critical review of current methods. *Gait Posture*. 2002;15(3):282-91.
- Kalender H, Uzuner K, Şimşek D, Bayram İ. Comparison of ankle force, mobility, flexibility, and plantar pressure values in athletes according to foot posture index. *Turk J Phys Med Rehabil*. 2022;68(1):91-9.
- Chuckpaiwong B, Nunley JA, Mall NA, Queen RM. The effect of foot type on in-shoe plantar pressure during walking and running. *Gait Posture*. 2008;28(3):405-11.
- Buldt AK, Forghany S, Landorf KB, Murley GS, Levinger P, Menz HB. Centre of pressure characteristics in normal, planus and cavus feet. *J Foot Ankle Res*. 2018;11:3.
- Chow TH, Chen YS, Wang JC. Characteristics of plantar pressures and related pain profiles in elite sprinters and recreational runners. *J Am Podiatr Med Assoc*. 2018;108(1):33-44.
- Iijima H, Ohi H, Isho T, Aoyama T, Fukutani N, Kaneda E, et al. Association of bilateral flat feet with knee pain and disability in patients with knee osteoarthritis: A cross-sectional study. *J Orthop Res*. 2017;35(11):2490-8.
- Fullin A, Caravaggi P, Picerno P, Mosca M, Caravelli S, De Luca A, et al. Variability of postural stability and plantar pressure parameters in healthy subjects evaluated by a novel pressure plate. *Int J Environ Res Public Health*. 2022;19(5):2913.
- Matušík J, Kočí V. What is a Footprint? A conceptual analysis of environmental Footprint indicators. *J Clean Prod*. 2021;285:124833.
- Swedler DI, Knapik JJ, Grier T, Jones BH. Validity of plantar surface visual assessment as an estimate of foot arch height. *Med Sci Sports Exerc*. 2010;42(2):375-80.
- Dugourd A, Saez-Rodriguez J. Footprint-based functional analysis of multiomic data. *Curr Opin Syst Biol*. 2019;15:82-90.
- Zuil-Escobar JC, Martínez-Cepa CB, Martín-Urrialde JA, Gómez-Conesa A. reliability and accuracy of static parameters obtained from ink and pressure platform footprints. *J Manipulative Physiol Ther*. 2016;39(7):510-7.
- Chu WC, Lee SH, Chu W, Wang TJ, Lee MC. The use of arch index to characterize arch height: a digital image processing approach. *IEEE Trans Biomed Eng*. 1995;42(11):1088-93.
- Clarke HH. An objective method of measuring the height of the longitudinal arch in foot examinations. *Res Quarterly Am Phys Educ Assoc*. 1933;4(3):99-107.
- Irwin LW. A study of the tendency of school children to develop flat-footedness. *Res Quarterly Am Phys Educ Assoc*. 1937;8(1):46-53.
- Hawes MR, Nachbauer W, Sovak D, Nigg BM. Footprint parameters as a measure of arch height. *Foot Ankle*. 1992;13(1):22-6.
- Forriol F, Pascual J. Footprint analysis between three and seventeen years of age. *Foot Ankle*. 1990;11(2):101-4.
- Viladot A. *Dez lições de patologia do pé*. 3ª ed. São Paulo: Roca;1987.
- Cantalino JLR, Mattos HM. Comparison of foot types classified by certain forms of clinical evaluation. *Ter Manual*. 2006;4(16):76-81.
- Ramos GM, Pereira SRF, Nucci A. Computational evaluation of the footprint: reference values of the plantar arch index in a sample of the Brazilian population. *Acta Fisiatr*. 2007;14(1):7-10.
- Dahle LK, Mueller M, Delitto A, Diamond JE. Visual assessment of foot type and relationship of foot type to lower extremity injury. *J Orthop Sports Phy Ther*. 1991;14(2):70-4.
- Xiong S, Goonetilleke RS, Witana CP, Weerasinghe TW, Au EY. Foot arch characterization: a review, a new metric, and a comparison. *J Am Podiatr Med Assoc*. 2010;100(1):14-24.
- Filoni E, Martins JF, Fukuchi RK, Gondo RM. Comparison between different plantar Arch Motriz Rio Claro. 2009;15(4):850-60.
- Cobey JC, Sella E. Standardizing methods of measurement of foot shape by including the effects of subtalar rotation. *Foot Ankle*. 1981;2(1):30-6.
- Nikolaidou ME, Boudolos KD. A Footprint -based approach for the rational classification of foot types in young schoolchildren. *Foot*. 2006;16(2):82-90.
- Souza PS, João SMA, Sacco ICN. Characterization of the longitudinal plantar arch of obese children using plantar print indexes. *J Human Growth Dev*. 2007;17(1):76-83.
- Sacco ICN, Nogueira GC, Bacarin TA, Casarotto R, Tozzi FL. Medial longitudinal arch change in diabetic peripheral neuropathy. *Acta Ortop Bras*. 2009;17(1):13-6.
- Razeghi M, Batt ME. Foot type classification: a critical review of current methods. *Gait Posture*. 2002;15(3):282-91.
- Balsdon ME, Khan M, Richards D, Dombroski CE. Arch height index, arch rigidity index, and arch stiffness values in a symptomatic population. *J Am Podiatr Med Assoc*. 2022;112(1):19-154.
- Chen B, Ma X, Xiao F, Chen P, Wang Y. Arch index measurement method based on plantar distributed force. *J Biomech*. 2022;144:111326.
- Zuil-Escobar JC, Martínez-Cepa CB, Martín-Urrialde JA, Gómez-Conesa A. Medial longitudinal arch: Accuracy, reliability, and correlation between navicular drop test and footprint parameters. *J Manipulative Physiol Ther*. 2018;41(8):672-9.
- Rosende-Bautista C, Munuera-Martínez PV, Seoane-Pillado T, Reina-Bueno M, Alonso-Tajes F, Pérez-García S, et al. Relationship of body mass index and footprint morphology to the actual height of the medial longitudinal arch of the foot. *Int J Environ Res Public Health*. 2021;18(18):9815.
- Kramer PA, Lautzenheiser SG. Foot morphology influences the change in arch index between standing and walking conditions. *Anat Rec (Hoboken)*. 2022;305(11):3254-62.