


Please cite the Published Version

Armstrong, Paul C, Allan, Harriet E, Kirkby, Nicholas S, Gutmann, Clemens, Joshi, Abhishek, Crescente, Marilena , Mitchell, Jane A, Mayr, Manuel and Warner, Timothy D (2022) Temporal in vivo platelet labeling in mice reveals age-dependent receptor expression and conservation of specific mRNAs. *Blood Advances*, 6 (23). pp. 6028-6038. ISSN 2473-9529

DOI: <https://doi.org/10.1182/bloodadvances.2022007099>

Publisher: American Society of Hematology

Version: Published Version

Downloaded from: <https://e-space.mmu.ac.uk/632489/>

Usage rights:  [Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0](#)

Additional Information: This is an Open Access article which appeared in *Blood Advances*, published by American Society of Hematology

Data Access Statement: Data are available on request from the corresponding author, Paul C. Armstrong (p.c.armstrong@qmul.ac.uk).

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

Temporal in vivo platelet labeling in mice reveals age-dependent receptor expression and conservation of specific mRNAs

Paul C. Armstrong,¹ Harriet E. Allan,¹ Nicholas S. Kirkby,² Clemens Gutmann,³ Abhishek Joshi,³ Marilena Crescente,^{1,4} Jane A. Mitchell,² Manuel Mayr,³ and Timothy D. Warner¹

¹Centre for Immunobiology, Blizard Institute, Faculty of Medicine and Dentistry, Queen Mary University of London, London, United Kingdom; ²National Heart and Lung Institute, Faculty of Medicine, Imperial College London, London, United Kingdom; ³King's British Heart Foundation Centre, Faculty of Life Sciences and Medicine, King's College London, London, United Kingdom; and ⁴Department of Life Sciences, Manchester Metropolitan University, Manchester, United Kingdom

Key Points

- Age-specific analysis of young platelets confirm higher thrombotic potential but independent of platelet size.
- Apparent refined control of individual mRNAs with those encoding granule proteins are greatly conserved across platelet lifespan.

The proportion of young platelets, also known as newly formed or reticulated, within the overall platelet population has been clinically correlated with adverse cardiovascular outcomes. However, our understanding of this is incomplete because of limitations in the technical approaches available to study platelets of different ages. In this study, we have developed and validated an in vivo temporal labeling approach using injectable fluorescent antiplatelet antibodies to subdivide platelets by age and assess differences in functional and molecular characteristics. With this approach, we found that young platelets (<24 hours old) in comparison with older platelets respond to stimuli with greater calcium flux and degranulation and contribute more to the formation of thrombi in vitro and in vivo. Sequential sampling confirmed this altered functionality to be independent of platelet size, with distribution of sizes of tracked platelets commensurate with the global platelet population throughout their 5-day lifespan in the circulation. The age-associated decrease in thrombotic function was accompanied by significant decreases in the surface expression of GPVI and CD31 (PECAM-1) and an increase in CD9. Platelet messenger RNA (mRNA) content also decreased with age but at different rates for individual mRNAs indicating apparent conservation of those encoding granule proteins. Our pulse-chase-type approach to define circulating platelet age has allowed timely reexamination of commonly held beliefs regarding size and reactivity of young platelets while providing novel insights into the temporal regulation of receptor and protein expression. Overall, future application of this validated tool will inform age-based platelet heterogeneity in physiology and disease.

Introduction

Understanding the fundamental changes that occur during a platelet's lifespan are central to many avenues of current research in both cardiovascular and hematologic disease. However, research has been severely limited because of a lack of methodologies to identify circulating platelets of different ages. Platelets inherit a finite complement of messenger RNA (mRNA) from the parent megakaryocyte,¹ which is rapidly lost once they are released into the circulation. As platelets are anucleate, they are unable to replenish these levels² and so nucleic acid dyes such as thiazole orange or SYTO13 have been widely used to differentiate

Submitted 19 January 2022; accepted 18 August 2022; prepublished online on *Blood Advances* First Edition 29 August 2022. <https://doi.org/10.1182/bloodadvances.2022007099>.

Data are available on request from the corresponding author, Paul C. Armstrong (p.c.armstrong@qmul.ac.uk).

The full-text version of this article contains a data supplement.

© 2022 by The American Society of Hematology. Licensed under [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International \(CC BY-NC-ND 4.0\)](#), permitting only noncommercial, nonderivative use with attribution. All other rights reserved.

young platelets based on their higher mRNA content (sometimes inferred to reflect an immature platelet fraction).³⁻⁸

Increased platelet turnover in a number of diseases, including diabetes mellitus, chronic kidney disease, and essential thrombocythemia^{9,10} and the consequent increases in the immature platelet fraction have been associated with a higher incidence of acute coronary syndromes¹¹ and a reduced effectiveness of antiplatelet drugs.^{12,13} Similarly, several groups have used nucleic acid dyes and cell sorting to establish that young platelets defined in this way are hyperreactive and of increased thrombotic potential.⁴⁻⁶ We have previously demonstrated that young platelets identified by higher mRNA content localize to the core of aggregates and drive aggregate formation even in the presence of standard antiplatelet therapies, potentially reducing drug effectiveness in high platelet turnover conditions.¹⁴

Although useful as basic methodologies, there are limitations with mRNA dye-based approaches to fractionate platelets. First, staining levels do not accurately advise on platelet age as fluorescence has not been temporally benchmarked. Second, following the rapid early loss of mRNA from young platelets, the fluorescent signal can no longer discriminate platelets by age. Moreover, confounding factors including platelet size and nucleotide-rich granule content can cause undue bias or interference with counts or estimated fractions of immature young platelets.^{15,16}

As a result of these shortcomings, there continues to be a search for alternative techniques to differentiate the newly formed platelet fraction. A common strategy in this regard has been to cause rapid thrombocytopenia in mice by injection of antiplatelet antibodies.^{17,18} This acute depletion is followed by a burst of platelet production such that the entire circulating platelet population is taken as being reticulated or newly formed. However, such an approach involves a pathological insult to drive rapid platelet production and makes the questionable assumption that platelets released under such conditions are representative of platelets newly formed under normal circumstances.

We therefore sought to develop an alternative *in vivo* labeling approach in mice using antiplatelet fluorescent antibodies to provide a tool for the age-related tracking and isolation of platelet subpopulations. Here, we demonstrate the application of this novel approach in revisiting widely held ideas regarding the phenotype of young platelets, identifying new age-related alterations in surface marker expression and revealing preferential retentions of mRNAs.

Methods

Ethical statement

Animal procedures were conducted in accordance with the United Kingdom Home Office approval (Project Licenses 70-7013 and PP1576048) under the United Kingdom Animals (Scientific Procedures) Act Amendment 2013 and were subject to local approval from the Queen Mary University of London (London, United Kingdom) and Imperial College London (London, United Kingdom) ethical review panels.

Temporal labeling of platelets *in vivo*

In vivo grade fluorescently conjugated anti-CD42c antibodies (DyLight-x488 or x649, Emfret) were administered (0.1 µg/g, IV) to the tail vein of C57BL/6 mice at specific time points as per

supplier guidance. In establishing the model, labeling efficacy was confirmed by comparative *in vitro* platelet labeling using anti-CD41-BV421 (clone MWReg30, BioLegend). Labeled platelets were identified and tracked using flow cytometry (Novocyte, Agilent Technologies) after microsampling from the tail vein.

Blood collection and isolation of platelets

Mice were anesthetized with intraperitoneal ketamine (Narketan, 100 mg/kg; Vetoquinol) and xylazine (Rompun, 10 mg/kg; Bayer), and blood was collected from the inferior vena cava into lepirudin (Refludan, 25 µg/mL; Celgene, Windsor) or trisodium citrate (3.2%; Sigma). Platelet-rich plasma (PRP) was isolated as previously published.¹⁹ Briefly, whole blood was diluted 1:1 with the modified Tyrode *N*-2-hydroxyethylpiperazine-*N'*-2-ethanesulfonic acid (HEPES) buffer (134 mmol/L sodium chloride, 2.9 mM potassium chloride, 0.34 mmol/L disodium phosphate, 12 mmol/L sodium bicarbonate, 20 mmol/L HEPES, and 1 mmol/L magnesium chloride; pH 7.4; Sigma) and centrifuged at 100 × *g* for 8 minutes.

Immunofluorescence of temporally labeled platelets

Platelets fixed in paraformaldehyde (4%; VWR) were centrifuged onto poly-L-lysine coverslips (VWR; 600 × *g*, 5 minutes). Samples were mounted with Prolong Diamond antifade mount (Thermo Fisher Scientific). Confocal microscopy was performed using an inverted Zeiss LSM880 with Airyscan confocal microscope; 63× lens objective, 1.4 Oil DICII (Zeiss), collecting 10 individual fields of view per mouse. Analysis was conducted using Zen Software (2.3 SP1, Zeiss) and ImageJ (National Institutes of Health).

Quantification of P-selectin expression and active confirmation of GPIIb/IIIa

PRP was diluted in phosphate-buffered saline (PBS; 1:40; Sigma) and incubated with the protease-activated receptor-4-activating peptide, AYPGKF amide (PAR4-amide; 200 µM; Bachem) or PBS for 20 minutes at 37°C in presence of anti-CD62P-BV421 (clone RB40.34; BD Bioscience) or anti-CD41/61-PE (clone JON/A, Emfret). Samples were subsequently fixed with 1% formalin (Sigma) and samples acquired and analyzed by flow cytometry (Novocyte) using NovoExpress software (Agilent Technologies).

Quantification of calcium dynamics

Platelets were isolated from mice 24 hours after injection with anti-CD42c-DyLight-x649 and were subsequently incubated with Fluo-4 AM (5 µM; Thermo Fisher Scientific) for 30 minutes at 37°C, followed by counterstaining with anti-CD41-BV421 (1:100; BD Biosciences) for 15 minutes and supplementation with calcium chloride (2 mM; Sigma). Baseline Fluo-4 fluorescence was recorded for 30 seconds, followed by challenge with PAR4-amide (25 µM), a collagen-related peptide (CRP-XL, 1 µg/mL; CAMBCOL), the stable thromboxane A₂ mimetic U46619 (10 µM; Enzo), or ionomycin (10 µM; Invitrogen) and subsequent recording for 2 minutes. Samples were acquired on a BD LSRII (BD Bioscience) using FACSDiva acquisition software and analyzed using FlowJo software, version 10.

Ex vivo whole-blood aggregation

Platelet aggregation was performed in whole blood as previously described.¹⁹ Briefly, in half-area 96-well microtiter plates (Greiner Bio-One), whole blood anticoagulated with lepirudin was incubated with PBS, arachidonic acid (0.05-0.5 mM; Sigma), the Horm

collagen (0.1-10 $\mu\text{g/mL}$; Takeda), PAR4-amide, (50-100 μM), or U46619 (0.1-10 μM) under mixing conditions (1200 revolutions per minute, Bioshake iQ, Q instruments). Samples were subsequently diluted (1:20 acid citrate dextrose) and analyzed by flow cytometry (Novocyte) to identify and determine proportions of the single-platelet population.

PE injury model

Pulmonary embolism (PE) was induced as previously described.²⁰ Briefly, the Horm collagen (50 $\mu\text{g/kg}$, Takeda) and U46619 (210 $\mu\text{g/kg}$, Enzo) were administered IV. Blood was micro-sampled before and after (4 minutes) administration of platelet activators for determination of circulating labeled platelets.

Surface marker expression

PRP obtained from temporally labeled mice was diluted 1:40 in modified Tyrode HEPES buffer and incubated for 20 minutes at room temperature with one of the following PE-conjugated antibodies: anti-CD9 (clone MZ3), anti-CD49b (clone DX5), anti-CD31 (clone 390), anti-H2 (clone 1/42; all BioLegend), or anti-GPVI (clone 784808, R&D systems). PE-conjugated isotype antibodies of Rat immunoglobulin M (IgM) (clone RTK2118), IgG1 (clone 2071), or IgG2a (clone RTK2758; all BioLegend) were used as corresponding nonspecific-binding controls and used to correct data values for both tracked and global populations before comparison. After staining, samples were fixed with 1% formalin, acquired on Novocyte flow cytometry, and analyzed using FlowJo software, version 10.

RNA extraction, complementary DNA synthesis, and quantitative real-time polymerase chain reaction of sorted, temporally labeled platelets

Full details of mRNA analysis of platelet populations are available in supplemental Methods. RNA was extracted from fluorescence-activated cell-sorted platelets (2.5 million per population) using the miRNeasy mini kit (Qiagen, 217004), following the manufacturer's instructions. Samples were spiked with Cel-miR-39-3p (Qiagen, 219600) and MS2 carrier RNA (Roche, 10165948001) after the first step of RNA isolation (ie, addition of Qiazol to sample). Complementary DNA was generated using VILO RT Superscript (Thermo Fisher, 11755-250) from 8 μL of RNA, as per the manufacturer's instructions. Transcripts were quantified using SYBR Select Master Mix (Applied Biosystems, 4472908) with custom-designed primers (Integrated DNA Technologies), in a ViiA 7 real-time polymerase chain reaction system (Applied Biosystems) using ViiA 7 software (Applied Biosystems).

Statistics and data analysis

Data are presented as mean \pm standard error of the mean (SEM) and data sets analyzed, including principal component analysis, using Prism, version 9 (GraphPad software). As required, the single-platelet population was gated based on forward scatter and antiplatelet immunoreactivity (fluorescence intensity). In addition, global platelet populations were defined as all platelets present in the sample, incorporating all platelet ages, and used to provide a baseline comparison for temporally labeled populations. Statistical significance was determined by paired *t* test unless otherwise stated, and data sets were considered different if $P < .05$. Each *n* value represents a data point from a separate animal.

Results

Temporal fluorescence labeling of platelets in vivo to identify age-defined populations

IV injections of short-lived fluorescence-labeled anti-CD42c antibody (DyLight-x488) produced $\sim 95\%$ labeling of circulating platelets as determined after in vitro counterstaining with anti-CD41-BV421 (supplemental Figure 1). Platelets remained fluorescently tagged for the duration of their life. After 24 hours, an unlabeled platelet subpopulation, representing $\sim 20\%$ of the entire population, was seen because of normal platelet removal and replacement. At this time, an injection of the same anti-CD42c antibody conjugated to an alternative fluorophore (DyLight-x649) created 2 differentially fluorescently labeled platelet subpopulations (Figure 1A). A minority single-labeled ($x649^+/x488^-$) population corresponding to platelets <24 hours old and a majority dual-labeled




Figure 1. Methodological diagram of in vivo temporal labeling of platelets in mice using fluorescently tagged antibodies. (A) Mice are injected with DyLight-x488-labeled anti-CD42c to achieve near universal labeling ($>95\%$), followed by a second injection after 24 hours with DyLight-x649-labeled anti-CD42c results in a dual-labeled platelet population and a minority x649 single-label population corresponding to platelets generated in intervening 24-hour period since the first injection. (B) Representative pseudocolor dot plot of differentially labeled platelets from blood sample as detected by flow cytometry.

(x649⁺/x488⁺; Figure 1B) population corresponding to platelets >24 hours old.


Younger platelets demonstrate increased reactivity, contributing disproportionately to thrombotic response

No significant differences were observed in basal P-selectin expression or JON-A binding between platelets <24 hours old and >24 hours old (Figure 2A-B). After incubation with a PAR-4 peptide agonist, the younger platelets demonstrated higher P-selectin expression ($83\,147 \pm 858$ mean fluorescence index [MFI]) than older platelets ($80\,880 \pm 1257$ MFI; $P < .005$; Figure 2A) but lower JON-A binding (864 ± 18 vs 884 ± 17 ; $P < .01$; Figure 2B). Younger platelets also displayed significantly stronger increases in intracellular calcium levels after exposure to the PAR4 peptide (Figure 2C), ionomycin (Figure 2D), and CRP-XL or U46619 (supplemental Figure 2).

Next, to assess the relative aggregatory potential of young and older platelets, the proportions of each subpopulation were

determined in whole-blood samples after stimulation. In such experiments, equal aggregatory potential and contribution would result in no differences between the proportions found in the unaggregated single-platelet population before and after stimulation. However, in response to multiple agonists, the proportions diverged significantly. In response to the PAR4 peptide, the proportion of younger platelets remaining unaggregated significantly decreased ($24.7\% \pm 0.3\%$ to $18.6\% \pm 0.6\%$; $P < .05$), whereas the older population significantly increased ($75.3\% \pm 0.3\%$ to $81.4\% \pm 0.6\%$; $P < .05$; Figure 3A). Similar patterns of response were seen when aggregation was stimulated by CRP-XL or U46619 (supplemental Figure 3). To confirm these in vitro observations in vivo, we used a murine model of PE and assessed circulating proportions of the labeled platelet populations before and after embolism. Compared with before injury, a significant decrease in the circulating proportion of young platelets ($19.0\% \pm 1.7\%$ to $17.0\% \pm 1.8\%$; $P < .05$) and a corresponding significant increase in the proportion of older platelets remaining unaggregated ($80.2\% \pm 1.8\%$ to $82.9\% \pm 1.7\%$; $P < .05$; Figure 3B) was observed.

Figure 2. Differential degranulation, integrin activation, and calcium flux of young platelets compared with older platelets. Younger (<24 hours old) platelets express higher levels of P-selectin (A) but lower levels of active GPIIb/IIIa (B) in response to PAR4-specific stimulation compared with corresponding older platelets in the same sample. Younger platelets have a greater calcium flux (measured as area under the curve [AUC]) in response to PAR4-specific stimulation (C) and ionomycin (D) ($n = 5$). (C) Data are individual replicates or mean \pm SEM. ** $P < .01$, *** $P < .001$ by paired t test as indicated. ns, not significant.






Figure 3. Younger platelets disproportionately contribute to thrombotic response. (A) Population tracking in anticoagulated whole blood after stimulation with a range of concentrations (30–100 μ M) of PAR4-specific stimulation to drive ex vivo aggregation resulted in decreasing proportions of younger platelets and corresponding increased proportions of older platelets remaining in the nonaggregated single-platelet portion ($n = 4$). Data are mean \pm SEM or individual replicates. (B) Circulating proportions of labeled younger platelets in vivo decreased following PE challenge with corresponding increase in proportions of circulating older platelets. * $P < .05$, ** $P < .01$ by 2-way analysis of variance with the Dunnett multiple comparison to their respective basal level or paired t test as appropriate.

Newly formed platelets are heterogenous in size and do not change size distribution with age

Platelets identified as <24 hours old were heterogenous in size with no discernable difference in size distribution to the global platelet population as determined using cytometry-derived forward scatter (FSC) and side scatter. The same tracked population remained heterogenous in size and commensurate relative to the global population in samples obtained from the same mice across days 2 to 5 (Figure 4A; supplemental Figure 4). Comparison of the calculated geometric mean of FSC values relative to the global population at each time point, indicates that the tracked population decreases in size with age, (ratio 1.2:1 at day 1 vs 0.9:1 at day 4; $P < .05$; Figure 4B; supplemental Figure 4). However, immunofluorescence performed on fixed samples did not discern a difference in platelet cross-sectional area between young (<24 hours) and older (1–5 days) platelets ($5.28 \pm 0.13 \mu\text{m}$ vs $5.24 \pm 0.25 \mu\text{m}$, respectively; $P > .05$; Figure 4C–D).

MHC-I expression decreases with platelet age

The expression of major histocompatibility class I (MHC-I) on platelets aged <24 hours was higher than the global median and decreased significantly with age (Figure 5A), equating to a loss of $30\% \pm 6\%$ across the platelet lifespan.

Platelets lose GPVI and PECAM-1 but increase expression of CD9 with age

GPVI expression was significantly higher on platelets <24 hours old and significantly lower on day 5 when compared with the global medians, a total loss equating to $39\% \pm 2\%$ ($P < .01$; Figure 5B).

Expression of PECAM-1 (CD31) significantly decreased between days 1 and 5 ($20\% \pm 3\%$ loss; $P < .001$; Figure 5C). By contrast, CD9 expression was significantly higher than the global median on day 5 ($P < .01$; Figure 5F) equating to an increase of $13\% \pm 4\%$

relative to day 1. No significant changes in expression of integrins $\alpha 2$ (CD49b) or $\alpha 2b$ (CD41) were found (Figure 5D–E).

Platelets favor retention of granule protein encoding mRNA

Full data set of mRNA measures are reported in supplemental Table 1 and subject to the following analysis. Principal component analysis of mRNA content demonstrated 3 analyzed populations, younger (<24 hours), older (1–5 days old), and global, clustered individually (Figure 6A). The loading plot of individual mRNAs demonstrated a unidirectional pattern for PC1 indicating that a decreasing abundance of mRNAs likely predominates this algorithm. Interestingly, however, mRNAs also appeared to cluster separately according to PC2, indicating potential for individual patterns of decreased abundance (Figure 6B). Further investigation grouping mRNAs by function, namely signaling, structural, receptors, and granules, suggested a reduction in abundance associated with age, with platelets <24 hours old having the highest and platelets 1 to 5 days old having the lowest amounts of each mRNA (Figure 6C). However, the amount of loss varied by specific mRNA (median Δ Cq range, -2.31 to -5.85). These changes were not contingent on starting levels (Figure 6C), but there was some association with functional categorization. In particular, mRNAs classified as associated with signaling proteins demonstrated the greatest reduction in abundance, whereas mRNAs associated with granule proteins demonstrated the greatest retention (median Cq change, -4.87 vs -2.99 , respectively).

Discussion

We have developed an in vivo labeling approach, which allows for age-related subpopulations of platelets to be identified and tracked. This technique can be coupled with a range of functional and molecular analyses to provide unique insights into platelet age-associated physiology. Here, we have used it to address the




Figure 4. Platelets are heterogeneous in size when formed and do not change size with age in the circulation. Following temporal labeling, serial blood samples were obtained each day for 5 days. (A) Representative data of age-tracked platelets (blue) in an individual mouse. Tracked platelets were detectable in circulation for the full 5 days, and distribution of size was comparable with the counterstained global platelet population (red) as determined by flow cytometric FSC and side scatter (SSC) properties. (B) Decrease in ratio of geometric mean FSC of tracked platelet relative to global populations across circulating lifespan ($n = 4$). (C) Confocal microscopy of fixed platelets in




Figure 5. Differential surface receptor expressions of aging platelets in circulation. Isotype control–corrected expression of receptors on age-tracked platelets, relative to global population expression (represented as 0 on y-axis for normalization), as measured by flow cytometry in samples obtained from individual mice on day 1 (green) and 5 (cyan) after labeling. Between days 1 and 5 of age, significant decreases in expression of MHC-1 (A), GPVI (B), and CD31/PECAM-1 (C), but no difference in expression of CD49a/integrin $\alpha 2$ (D) or CD41/integrin $\alpha 2b$ (E) was detected, whereas significant increase in CD9 expression (F) was detected. ** $P < .01$, *** $P < .001$ as analyzed by paired t test.

2 fundamental assumptions that newly formed platelets are larger and more reactive. Furthermore, we have used this new approach to gain insights into changes in the expression of platelet receptors and the retention of mRNA during aging that may underpin alterations in their functionality.

The nature of the relationship among platelet age, size, and reactivity has remained contentious for many years. Perhaps the point of least contention is that mRNA-rich “young” platelets are more reactive, with a raft of literature over the years using *in vitro*, *ex vivo*, and *in vivo* studies supporting this hypothesis.^{14,21} Indeed, our recent studies identifying platelet age through RNA labeling of healthy human platelets are consistent with a decline in hemostatic function as platelets age. We demonstrated that young platelets

express higher levels of P-selectin and have an enhanced calcium signaling capacity.^{4,6} Supporting our work in healthy human blood, our temporal labeling approach confirmed that the youngest platelets have enhanced degranulation and calcium signaling in response to a range of agonists. Conversely, after stimulation, older platelets have increased levels of the activated GPIIb/IIIa receptor, indicating a higher propensity for fibrinogen binding; however, the functional significance of this remains to be determined. Despite this, *ex vivo* aggregation studies and an *in vivo* PE thrombosis model demonstrate that under healthy turnover conditions, young platelets contribute highly and disproportionately to forming thrombi.

A factor that continues to confound our understanding of why young platelets are more reactive or have higher thrombotic

Figure 4 (continued) PRP revealed no difference in area of younger and older platelets. (D) Representative image of younger (blue) and older (blue and red) present in analyzed platelet samples isolated from labeled mouse. Data are individual replicates from 4 mice with overlaid mean \pm SEM with * $P < .05$ as analyzed by Friedman test with the Dunn multiple comparison test or mean \pm SEM with paired t test.




Figure 6. Platelet mRNA content decreases with age but differential loss according to function of encoded protein. Younger, older, and total platelets (2.5 million per population) were isolated by fluorescence-activated cell sorting and mRNA extracted for comparative analysis of 25 highly abundant transcripts. (A) Principal component analysis of samples resulted in individual clustering of samples according to age. (B) Loading plot analysis depict unidirectional relationship for principal component 1 (PC1) but differential PC2 relationship. (C) Grouping of measured mRNAs by downstream function and mRNA level (quantification cycle [C_q]) analysis illustrates variable content of individual mRNAs (green) but with universal loss from highest levels in younger (blue) to lowest levels in older (red) platelets. (D) Calculated change (ΔC_q) from younger to older populations reveals differential scales of loss within groupings with mRNAs associated with encoding granule proteins experiencing greatest retention. Data are mean \pm SEM ($n = 4$).

potential is the enduring dogma that young platelets are larger. Originally proposed in the 1960s,²² this hypothesis posits that younger platelets have greater volume resulting in greater granule content and greater surface area resulting in greater surface receptor number. In support of this hypothesis, *in vivo* platelet studies in a range of species after pathological insult such as platelet depletion have observed that the subsequently produced platelets are larger in size.²³ Similarly, associations have been found between mean platelet volume and immature platelet fraction measurements in patient populations. However, a series of reports in the mid-1980s reported no correlation between platelet age and size.²⁴⁻²⁶ Similarly, *in vivo* biotinylation studies in healthy rabbits have shown that younger platelets are no larger than older platelets.²⁷ We therefore coupled our temporal labeling technique to flow cytometry and immunofluorescence to assess the sizes of platelets of different ages. Our data clearly show that in healthy mice, younger and older platelets exist in the same range of sizes throughout their 5-day lifespan in the circulation and therefore cannot be differentiated on the basis of size alone. Notably, the aforementioned *in vivo* studies that associate younger platelets with larger size are predicated on a pathological insult and the subsequent acute response to restore platelet number. It seems reasonable to conclude that the processes that underpin such an acute response differ from those supporting normal platelet production and release and could produce a different population of platelets. Interestingly, comparison of the average sizes of our tracked and global populations indicates, as measured by flow cytometry, that platelets may decrease to a degree in size during their aging process in the circulation. In our recent studies of human platelets, we observed an association between platelet age and decreased cytoskeletal protein content that may be consistent with small alterations in the FSC measures detected in this study.⁶ However, such potential alterations in size will require further investigation by other modalities to validate this observation.

Similar research has also been applied to identifying other markers of age. The observation that platelet size does not change substantially with age, coupled with our recent observation of loss of protein content in aging human platelets,⁶ might support proposals that platelet density is a good marker of platelet age.²⁸ However, the calculation of platelet density is technically challenging and laborious, whereas the measurement of surface molecular markers or receptor expression is much more easily accomplished. Angenieux et al²⁹ recently identified that across the platelet population, young platelets, identified by thiazole orange staining, had the highest levels of MHC-I/HLA-1. In our study using defined temporal labeling, we confirmed this observation and found novel differential expression patterns according to platelet age for several other receptors associated with collagen responsiveness. We identified higher surface expression levels of GPVI and PECAM-1 in younger compared with older platelets, whereas, conversely, the levels of CD9 increased as platelets age in the circulation. The origin of this increase in surface CD9 is unclear, but *de novo* production or accumulation from exosomes³⁰ are 2 possibilities. Despite its functional role remaining unclear,³¹ CD9 expression is generally high on platelets³² and it has been shown to localize with GPVI³³ and ADAM10³⁴ within tetraspanin-enriched microdomains. Both GPVI and PECAM-1 undergo proteolytic cleavage mediated by ADAM-10 and MMP2, respectively, under physiological and pathological conditions,³⁵⁻³⁷ and therefore, constitutive or triggered

activity may contribute to their decline in surface expression as platelets age. With relevance to the decline in platelet responsiveness associated with aging, a decrease in GPVI expression would correlate with the impaired CRP-XL-induced activation observed in our functional studies.

In addition to changes in receptor expression, we sought to examine the mRNA content of platelets. Young platelets are well accepted as having the highest levels of mRNA and that this mRNA content decreases with time.³⁸ Indeed, the results of our measurement, from the selection of mRNA, would support this view. Platelets have been reported as containing ~10 000 different mRNA transcripts³⁹ and that this mRNA has a half-life of ~6 hours.^{38,40} We wished to examine whether this loss of mRNA was universal using our temporal labeling approach. Unbiased principal component analysis confirmed that young platelets had higher overall levels of mRNA but also revealed a differential reduction in specific platelet mRNAs. Further analysis revealed the rate of loss for mRNAs to vary greatly, with those encoding granule proteins experiencing the least reduction. This variation in rates of loss strongly suggests that previous reports of 6 hours as the decay half-life of platelet mRNA^{38,40} is a general number, whereas the decay of particular mRNAs is under more refined control. The molecular nature and purpose of this apparent conservation will be important to elucidate, but research by Mills et al⁴⁰ has indicated that ribosomes protect mRNA through binding, which may enhance specific translation. Following our observation that mRNAs that encode granule proteins are particularly conserved, it is possible to hypothesize that this will support their specific protein synthesis. Having been reported for over 50 years,⁴¹ *de novo* protein synthesis in platelets, particularly in response to activation,⁴²⁻⁴⁴ is not a new concept. However, owing to the technically challenging nature of these studies, much interrogation of this process has occurred in an *ex vivo* setting and questions remain over the quantity of proteins produced⁴⁵ and their physiological and/or pathophysiological significance. Our labeling approach will help to unravel whether such *de novo* synthesis is part of a maturation step of platelets *in vivo* or following reports of selective protein packaging and release of granular contents and extracellular vesicles,⁴⁶⁻⁴⁸ whether selective retention of granular mRNA for protein synthesis prolongs their period of reactivity.

In summary, we present a novel approach for discovering age-related alterations in platelet form and function. Our approach has discovered changes in platelet GPVI, PECAM-1, and CD9 expression as platelets age in the circulation and provides insights into differential regulation and retention of platelet mRNAs. These changes may well underpin alterations in platelet functionality during the aging process. Therefore, this study reports novel insights into fundamental assumptions of platelet size and reactivity as they age in the circulation and provides an approach that will act as an important tool for future research.

Acknowledgments

This work was supported by the British Heart Foundation (RG/19/8/34500) (T.D.W. and P.C.A.), (FS/18/60/34181) (C.G.), and (FS/16/31699) (N.S.K.) and the Wellcome Trust (101604/Z/13/Z).

Authorship

Contribution: P.C.A. designed the research, performed the assays and collected data, analyzed and interpreted data,

performed statistical analysis, and wrote the manuscript; H.E.A. performed assays, analyzed and interpreted the data, performed statistical analysis, and revised the manuscript; C.G. and A.J. performed analysis and collected and analyzed data; M.C. assisted with the assays and collecting data; and N.S.K., J.A.M., M.M., and T.D.W. analyzed and interpreted the data and revised the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

ORCID profiles: P.C.A., 0000-0003-0904-677X; H.E.A., 0000-0002-2104-4972; N.S.K., 0000-0003-3509-8549; C.G., 0000-0003-0675-8632; M.C., 0000-0003-3164-512X; J.A.M., 0000-0001-9879-546X; M.M., 0000-0002-0597-829X; T.D.W., 0000-0003-3988-4408.

Correspondence: Paul C. Armstrong, Centre for Immunobiology, Blizard Institute, Faculty of Medicine and Dentistry, Queen Mary University of London, 4 Newark St, London E1 2AT, United Kingdom; email: p.c.armstrong@qmul.ac.uk.

References

1. Cecchetti L, Tolley ND, Michetti N, Bury L, Weyrich AS, Gresle P. Megakaryocytes differentially sort mRNAs for matrix metalloproteinases and their inhibitors into platelets: a mechanism for regulating synthetic events. *Blood*. 2011;118(7):1903-1911.
2. Rowley JW, Schwartz H, Weyrich AS. Platelet mRNA: the meaning behind the message. *Curr Opin Hematol*. 2012;19(5):385-391.
3. Kienast J, Schmitz G. Flow cytometric analysis of thiazole orange uptake by platelets: a diagnostic aid in the evaluation of thrombocytopenic disorders. *Blood*. 1990;75(1):116-121.
4. Hille L, Cederqvist M, Hromek J, Stratz C, Trenk D, Nuhrenberg TG. Evaluation of an alternative staining method using SYTO 13 to determine reticulated platelets. *Thromb Haemost*. 2019;119(5):779-785.
5. Bongiovanni D, Santamaria G, Klug M, et al. Transcriptome analysis of reticulated platelets reveals a prothrombotic profile. *Thromb Haemost*. 2019;119(11):1795-1806.
6. Allan HE, Hayman MA, Marcone S, et al. Proteome and functional decline as platelets age in the circulation. *J Thromb Haemost*. 2021;19(12):3095-3112.
7. Meintker L, Haimerl M, Ringwald J, Krause SW. Measurement of immature platelets with Abbott CD-Sapphire and Sysmex XE-5000 in haematology and oncology patients. *Clin Chem Lab Med*. 2013;51(11):2125-2131.
8. Dale G, Friese P, Hynes L, Burstein S. Demonstration that thiazole-orange-positive platelets in the dog are less than 24 hours old. *Blood*. 1995;85(7):1822-1825.
9. Himmelfarb J, Holbrook D, McMonagle E, Ault K. Increased reticulated platelets in dialysis patients. *Kidney Int*. 1997;51(3):834-839.
10. Lee EY, Kim SJ, Song YJ, Choi SJ, Song J. Immature platelet fraction in diabetes mellitus and metabolic syndrome. *Thromb Res*. 2013;132(6):692-695.
11. Ibrahim H, Schutt RC, Hannawi B, DeLao T, Barker CM, Kleiman NS. Association of immature platelets with adverse cardiovascular outcomes. *J Am Coll Cardiol*. 2014;64(20):2122-2129.
12. Cesari F, Marcucci R, Caporale R, et al. Relationship between high platelet turnover and platelet function in high-risk patients with coronary artery disease on dual antiplatelet therapy. *Thromb Haemost*. 2008;99(5):930-935.
13. Bernlochner I, Goedel A, Plischke C, et al. Impact of immature platelets on platelet response to ticagrelor and prasugrel in patients with acute coronary syndrome. *Eur Heart J*. 2015;36(45):3202-3210.
14. Armstrong PC, Hoefer T, Knowles RB, et al. Newly formed reticulated platelets undermine pharmacokinetically short-lived antiplatelet therapies. *Arterioscler Thromb Vasc Biol*. 2017;37(5):949-956.
15. Koike Y, Miyazaki K, Higashihara M, et al. Clinical significance of detection of immature platelets: comparison between percentage of reticulated platelets as detected by flow cytometry and immature platelet fraction as detected by automated measurement. *Eur J Haematol*. 2010;84(2):183-184.
16. Robinson MS, Mackie IJ, Khair K, et al. Flow cytometric analysis of reticulated platelets: evidence for a large proportion of non-specific labelling of dense granules by fluorescent dyes. *Br J Haematol*. 1998;100(2):351-357.
17. Nieswandt B, Bergmeier W, Rackebrandt K, Gessner JE, Zirngibl H. Identification of critical antigen-specific mechanisms in the development of immune thrombocytopenic purpura in mice. *Blood*. 2000;96(7):2520-2527.
18. Gupta S, Cherpokova D, Spindler M, Morowski M, Bender M, Nieswandt B. GPVI signaling is compromised in newly formed platelets after acute thrombocytopenia in mice. *Blood*. 2018;131(10):1106-1110.
19. Armstrong PC, Kirkby NS, Chan MV, et al. Novel whole blood assay for phenotyping platelet reactivity in mice identifies ICAM-1 as a mediator of platelet-monocyte interaction. *Blood*. 2015;126(10):e11-e18.
20. Armstrong PC, Kirkby NS, Zain ZN, Emerson M, Mitchell JA, Warner TD. Thrombosis is reduced by inhibition of COX-1, but unaffected by inhibition of COX-2, in an acute model of platelet activation in the mouse. *PLoS One*. 2011;6(5):e20062.
21. McBane RD 2nd, Gonzalez C, Hodge DO, Wysokinski WE. Propensity for young reticulated platelet recruitment into arterial thrombi. *J Thromb Thrombolysis*. 2014;37(2):148-154.
22. Karpatkin S. Heterogeneity of human platelets. II. Functional evidence suggestive of young and old platelets. *J Clin Invest*. 1969;48(6):1083-1087.
23. McDonald TP, Odell TT Jr, Gosslee DG. Platelet size in relation to platelet age. *Proc Soc Exp Biol Med*. 1964;115(3):684-689.

24. Thompson CB, Jakubowski JA, Quinn PG, Deykin D, Valeri CR. Platelet size as a determinant of platelet function. *J Lab Clin Med*. 1983;101(2):205-213.
25. Thompson CB, Jakubowski JA, Quinn PG, Deykin D, Valeri CR. Platelet size and age determine platelet function independently. *Blood*. 1984;63(6):1372-1375.
26. Thompson CB, Love DG, Quinn PG, Valeri CR. Platelet size does not correlate with platelet age. *Blood*. 1983;62(2):487-494.
27. Reddy EC, Wang H, Bang KWA, Packham MA, Rand ML. Young steady-state rabbit platelets do not have an enhanced capacity to expose procoagulant phosphatidylserine. *Platelets*. 2018;29(1):27-33.
28. Corash L, Shafer B. Use of asplenic rabbits to demonstrate that platelet age and density are related. *Blood*. 1982;60(1):166-171.
29. Angenieux C, Dupuis A, Gachet C, de la Salle H, Maitre B. Cell surface expression of HLA I molecules as a marker of young platelets. *J Thromb Haemost*. 2019;17(9):1511-1521.
30. Mathieu M, Nevo N, Jouve M, et al. Specificities of exosome versus small ectosome secretion revealed by live intracellular tracking of CD63 and CD9. *Nat Commun*. 2021;12(1):4389.
31. Matthews AL, Noy PJ, Reyat JS, Tomlinson MG. Regulation of A disintegrin and metalloproteinase (ADAM) family sheddases ADAM10 and ADAM17: the emerging role of tetraspanins and rhomboids. *Platelets*. 2017;28(4):333-341.
32. Higashihara M, Takahata K, Yatomi Y, Nakahara K, Kurokawa K. Purification and partial characterization of CD9 antigen of human platelets. *FEBS Lett*. 1990;264(2):270-274.
33. Prottly MB, Watkins NA, Colombo D, et al. Identification of Tspan9 as a novel platelet tetraspanin and the collagen receptor GPVI as a component of tetraspanin microdomains. *Biochem J*. 2009;417(1):391-400.
34. Arduise C, Abache T, Li L, et al. Tetraspanins regulate ADAM10-mediated cleavage of TNF-alpha and epidermal growth factor. *J Immunol*. 2008;181(10):7002-7013.
35. Chatterjee M, Gawaz M. Clinical significance of receptor shedding-platelet GPVI as an emerging diagnostic and therapeutic tool. *Platelets*. 2017;28(4):362-371.
36. Vulliamy P, Montague SJ, Gillespie S, et al. Loss of GPVI and GPIIb/alpha contributes to trauma-induced platelet dysfunction in severely injured patients. *Blood Adv*. 2020;4(12):2623-2630.
37. Montague SJ, Andrews RK, Gardiner EE. Mechanisms of receptor shedding in platelets. *Blood*. 2018;132(24):2535-2545.
38. Angenieux C, Maitre B, Eckly A, Lanza F, Gachet C, de la Salle H. Time-dependent decay of mRNA and ribosomal RNA during platelet aging and its correlation with translation activity. *PLoS One*. 2016;11(1):e0148064.
39. Eicher JD, Wakabayashi Y, Vitseva O, et al. Characterization of the platelet transcriptome by RNA sequencing in patients with acute myocardial infarction. *Platelets*. 2016;27(3):230-239.
40. Mills EW, Green R, Ingolia NT. Slowed decay of mRNAs enhances platelet specific translation. *Blood*. 2017;129(17):e38-e48.
41. Warshaw AL, Laster L, Shulman NR. The stimulation by thrombin of glucose oxidation in human platelets. *J Clin Invest*. 1966;45(12):1923-1934.
42. Schwartz H, Tolley ND, Foulks JM, et al. Signal-dependent splicing of tissue factor pre-mRNA modulates the thrombogenicity of human platelets. *J Exp Med*. 2006;203(11):2433-2440.
43. Weyrich AS, Dixon DA, Pabla R, et al. Signal-dependent translation of a regulatory protein, Bcl-3, in activated human platelets. *Proc Natl Acad Sci U S A*. 1998;95(10):5556-5561.
44. Lindemann S, Tolley ND, Dixon DA, et al. Activated platelets mediate inflammatory signaling by regulated interleukin 1beta synthesis. *J Cell Biol*. 2001;154(3):485-490.
45. Joshi A, Schmidt LE, Burnap SA, et al. Neutrophil-derived protein S100A8/A9 alters the platelet proteome in acute myocardial infarction and is associated with changes in platelet reactivity. *Arterioscler Thromb Vasc Biol*. 2021;42(1):49-62, ATVBHA121317113.
46. Kamykowski J, Carlton P, Sehgal S, Storrie B. Quantitative immunofluorescence mapping reveals little functional coclustering of proteins within platelet alpha-granules. *Blood*. 2011;118(5):1370-1373.
47. Le Blanc J, Fleury S, Boukhatem I, Belanger JC, Welman M, Lordkipanidze M. Platelets selectively regulate the release of BDNF, but not that of its precursor protein, proBDNF. *Front Immunol*. 2020;11:575607. <https://doi.org/10.3389/fimmu.2020.575607>
48. Aatonen MT, Ohman T, Nyman TA, Laitinen S, Gronholm M, Siljander PR. Isolation and characterization of platelet-derived extracellular vesicles. *J Extracell Vesicles*. 2014;3(1). <https://doi.org/10.3402/jev.v3.24692>