

Please cite the Published Version

Lou, Yingqiang, Zhao, Qingshan, Hu, Yunbiao, Chen, Lijun, Liu, Pengfei, Fang, Yun, Lloyd, Huw and Sun, Yuehua (2023) Personality-dependent nest site selection and nest success during incubation in wild chestnut thrushes. *iScience*, 26 (8). p. 107419. ISSN 2589-0042

DOI: <https://doi.org/10.1016/j.isci.2023.107419>

Publisher: Elsevier

Version: Published Version

Downloaded from: <https://e-space.mmu.ac.uk/632493/>

Usage rights:  [Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0](https://creativecommons.org/licenses/by-nc-nd/4.0/)

Additional Information: This is an Open Access article which appeared in *iScience*, published by Elsevier

Data Access Statement: All data reported in this paper are available at Mendeley, and the DOI is listed in the key resources table.

- All original code in this paper are available at Mendeley, and the DOI is listed in the key resources table.
- Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

Article

Personality-dependent nest site selection and nest success during incubation in wild chestnut thrushes

Yingqiang Lou,
Qingshan Zhao,
Yunbiao Hu, ...,
Yun Fang, Huw
Lloyd, Yuehua Sun

sunyh@ioz.ac.cn

Highlights

We measured two
personality traits in a wild
population of female
chestnut thrushes

More active females chose
nest sites with great nest
lateral concealment

Females with higher
breathing rate laid small
clutch sizes

Nests of females with
lower breathing rate had
higher nest success during
incubation



Article

Personality-dependent nest site selection and nest success during incubation in wild chestnut thrushes

Yingqiang Lou,¹ Qingshan Zhao,² Yunbiao Hu,¹ Lijun Chen,¹ Pengfei Liu,³ Yun Fang,¹ Huw Lloyd,⁴ and Yuehua Sun^{1,5,*}

SUMMARY

In birds, little is known about how individuals choose nest sites based on their personality traits. Here, we investigate whether a female's personality (activity and breathing rate) can affect patterns of nest site selection at different spatial scales in a wild population of chestnut thrush (*Turdus rubrocanus*) and determine whether nest site characteristics and female personality traits affect clutch size and nest success during incubation. We found that neither activity nor breathing rate were associated with large-scale nesting habitat variables. At the fine-scale level, more active females chose nest sites with greater nest lateral concealment. Females with higher breathing rates laid smaller clutch sizes than individuals with lower breathing rates. Nests of females with lower breathing rate had higher nest success during incubation. This work highlights the relationships between personality and nest site selection in birds, and the important role of female personality traits in reproductive success.

INTRODUCTION

Animal personality is defined as individual behavioral differences that are consistent over time and/or across contexts.^{1–3} Having been studied across many animal taxa,⁴ animal personality is now known to play important roles in sexual selection,^{5–7} parental care,^{8,9} and other life history traits.^{10,11} For example, female zebra finches (*Taeniopygia guttata*) choose mates with similar personality traits,¹² and parents with similar personality traits have higher reproductive success.^{13–15} Recently, quantifying spatial distribution patterns of individuals within a population based on animal personality has attracted increasing interest.^{16–18} Personality could reflect individual ability to cope with environmental conditions, which may in turn lead to personality-dependent distribution,^{19,20} which forms the premise of the personality-matching habitat choice hypothesis.^{17,19,21}

Nest site selection at different spatial scales (i.e., large- and fine-scale) is an important process for successful breeding,²² and is affected by many factors such as food resources and predator abundance.^{23,24} Recent studies have found that individual differences in personality traits are known to be related to large-scale nest site selection.^{17,25,26} For example, bold female common eider (*Somateria mollissima*) select nest sites with greater concealment and which are located further from landscape features.²⁷ To our knowledge, less is known about how personality traits affect patterns of fine-scale nest site selection in birds. Therefore, to gain an understanding of personality-dependent nest site selection requires a comprehensive examination of animal personality differences at different spatial scales.

Fine-scale nest site selection also affects individual reproductive strategy.^{28–30} Nests with better concealment can have larger clutch sizes,^{23,31} which may covary with nest fate. Large clutch size could produce more nestlings which would require adults to make more foraging trips, and nesting in areas with greater concealment could reduce the risk of nest predation.³² Furthermore, clutch size could also be affected by personality traits.³³ For example, Araya-Ajoy et al.³⁴ found that more explorative female great tit (*Parus major*) laid larger clutch sizes.

Personality traits may be a significant factor that affects avian reproductive success.^{17,35,36} Nest defense is one kind of personality traits, and more intense nest defense is associated with lower risk of predation,^{37,38}

¹Key Laboratory of Animal Ecology and Conservation Biology, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China

²State Key Laboratory of Urban and Regional Ecology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085, China

³School of Life Sciences and Technology, Longdong University, Qingyang 745000, China

⁴Department of Natural Sciences, Manchester Metropolitan University, Manchester M15GD, UK

⁵Lead contact

*Correspondence: sunyh@ioz.ac.cn

<https://doi.org/10.1016/j.isci.2023.107419>



Table 1. Repeatability of activity and breathing rate in chestnut thrush

Trait	N _{ind}	Range	R	P
Activity	153	0–18.385	0.337	0.003
Breathing rate	165	55–180	0.477	<0.001

N_{ind} = total number of individuals, and R = repeatability (R), and range = range of traits.

Activity is represented as the square root of the number of movements in the simple cage test. Breathing rate is defined as the number of breast movements during 60 s while being handled. The LMMs of activity and breathing rate were used to calculate the repeatability and can be found in Table A1.

and different personality traits may cause different responses to nest predators³⁹; for example, more proactive individuals are shown to respond more boldly toward human intruders or natural predators^{37,40} during incubation and nestling periods. Furthermore, nest site characteristics are also associated with reproductive success, since individuals who choose to select nest sites with greater nest concealment could increase the difficulty for predators to find the nests.³⁰ A recent study found that nest success was positively associated with nest concealment across 21 species on the Tibetan Plateau.³⁰ To date, few studies have investigated how multi-scale nest site characteristics (fine- and large-scale) and personality traits affect nest success during incubation.⁴¹

In this study, we extend the previous research of Zhao et al.²⁵ of a wild breeding population of chestnut thrush to investigate whether nest site selection and nest success during incubation are related to females' personality traits. We use two measures of animal personality—breathing rate and activity—which are easily measurable and repeatable in our study population.^{15,25} Although breathing rate is considered by some as a physiological trait-reflecting stress response,^{42,43} many authors have argued a strong case for boldness to be measured (at least as a proxy) by breathing rate (lower breathing rate reflecting greater boldness), as the two appear to be very closely linked.^{44–49} In chestnut thrushes, only females participate in nest site selection, incubation, and in brooding nestlings,¹⁵ and both sexes participate in nest defense and provisioning to offspring. Furthermore, our previous study found that nest site characteristics, such as concealment of nests from below, were positively associated with nest daily survival rate.⁵⁰ Therefore, the wild chestnut thrush population is a suitable model species with which to explore the relationship between personality with nest site selection and nest success during incubation.

Our objectives of this study are 3-fold: (1) to examine whether large- and fine-scale nest site characteristics were associated with female personality traits. Based on the previous results of Seltmann et al.²⁷ and Holtmann et al.,¹⁹ we predicted that females with lower breathing rate and activity select nest sites nearer from areas of human disturbance (e.g., human residence, roads, and farmland) and in more concealed areas (higher concealment of lateral, above and below around the nests). (2) To examine whether the nest site characteristics, and female personality traits were associated with clutch size. We predicted that nests with more concealment and females with lower breathing rate would lay larger clutches. (3) To examine whether the nest site characteristics, and female personality traits are related to nest success during incubation. Based on the results of Liu et al.³⁰ and Tamin et al.,³⁶ we predicted that nests with more concealment and nests of females with lower breathing rate would have higher nest survival rates during incubation.

RESULTS

Personality traits of females, clutch size and nest site characteristics

In 2013–2016, we measured the activity test 211 times in 153 female chestnut thrushes and breathing rate test 230 times in 165 females. Both activity and breathing rate were repeatable with individuals across time (Table 1). In total, we also collected the nest site characteristics of 125 female chestnut thrushes that were measured for activity and breathing rate from 2013 to 2016.

The results of pSEM showed that two hypothesized unidirectional relationships were statistically significant (Figure 1). Breathing rate was negatively associated with clutch size (Figure 1): females with higher breathing rate laid smaller clutch sizes. Neither activity nor PC1 scores of large- and fine-scale nest site characteristics were associated with clutch size (Figure 1). Activity was positively associated with PC1 score of fine-scale nest site characteristics (Figure 1): more active individuals chose nests with greater lateral concealment. We did not find any relationships between breathing rate and any nest site characteristics (Figure 1).




Figure 1. Effects of personality traits and nest site characteristics on clutch size in chestnut thrushes from piecewise structural equation model with individual identity as a random effect

Each box represents measured variables. Arrows represent relationships among variables. The dashed lines represent non-significant paths and solid lines represent significant paths. The gray lines represent negative relationships and black lines represent positive relationship. Significant paths are shown in bold. The piecewise structural equation models (pSEM) were conducted by piecewiseSEM package in R.⁶⁰

Personality traits of females and nest success during incubation

From 2013 to 2016, nest fate during incubation was recorded in 129 nests (24 nests failed and 105 nests successfully hatched). Female breathing rate was the only significant predictor of nesting success during incubation. Nests of females with lower breathing rate had higher nest success during incubation (Table 2).

DISCUSSION

Our results show that there is proof for correlation between female personality trait and nest-site characteristics: more active females nested in areas with higher PC1 score of fine-scale nest site characteristics. Moreover, we found that clutch size was negatively associated with female's breathing rate, but we did not find that activity or any nest site characteristics were associated with clutch size. We also found that nest success during incubation was associated with female personality traits in chestnut thrushes.

We found that activity of female chestnut thrushes was positively related to lateral nest concealment, which mainly represents PC1 score in fine-scale nest site characteristics. Increased activity may make individuals more conspicuous to visual predators,⁵¹ whereas greater lateral concealment of nests could reduce the ability of predators to locate nests. In our study area, numerous natural nest predators, such as Large Hawk Cuckoo (*Cuculus sparveroides*) and Elliot's Laughingthrush (*Garrulax elliotii*),⁵² pose a particular risk for the eggs and nesting. Therefore, we hypothesize that more active female chestnut thrushes may be nesting in areas with greater concealment to overcome the constraint of higher visibility due to increased activity (e.g., entering and leaving of nests).

We found that female chestnut thrushes with lower breathing rate laid larger clutch size, while the activity was not associated with clutch size, a finding similar to that of Zhao et al.⁵³ However, some studies have found that female personality traits are not related with either clutch or brood size.^{9,54,55} This variation in findings may be affected by the interaction of individual age and personality types. Dingemanse et al.³³ found that individuals with lower exploratory rates produced moderate-sized clutches throughout their lives, whereas faster-paced explorers produced moderate-sized clutch when young, but clutch size decreased with age. Currently, there is no suitable method to age chestnut thrushes in the field because we recaptured so few individuals across years, but further studies relating age and personality types are needed in the future.

We failed to find any relationships between nest site characteristics and clutch size in chestnut thrushes, in contrast to several other passerine studies.^{30,56} One potential explanation for this is that clutch size may be

Table 2. Predictors of nest success during incubation of chestnut thrushes

Predictors	Estimate	χ^2	P
Breathing rate	−0.051	4.246	0.039
Activity	0.107	0.638	0.424
Julian date	0.016	1.626	0.202
PC1 score of large-scale nest site characteristics	−0.344	2.831	0.092
PC1 score of fine-scale nest site characteristics	0.314	2.393	0.122

Significant predictors are highlighted in bold.

The nest success during incubation was defined as the nests that survived until at least one offspring hatched.

mediated by predation risk.^{57,58} Hu et al.⁵² found that there are many predators in our study site, and almost 50% of chestnut thrush nests fail due to nest predation.⁵⁰ Chestnut thrushes produce greater clutch size at nesting sites with lower predation risk but lay smaller clutches in nesting sites with higher predation risk. In this scenario, such high nest predation risk may dilute the positive effects of nest site characteristics, and this needs to be verified in future research.

The nests of female chestnut thrushes with lower breathing rates were more likely to survival during incubation. Lower breathing rate females may be more successful at defending their nests against predators, thus contributing to lower nest failure and higher nest success rates.^{40,59} Furthermore, boldness can covary with other personality traits, comprising a suite of behaviors or “behavioral syndromes”,² such as aggression and exploration, which may further promote individuals to participate in nest defense.³⁷ Further investigations are needed to identify the direct relationship between breathing rate and the intensity of nest defense in chestnut thrushes.

In conclusion, this study provides some of the first evidence to show the relationship between personality traits and fine-scale nest site selection. Breathing rate was associated with clutch size, thus providing some support for the pace-of-life syndrome hypothesis. Our study also identified a relationship between nest success during incubation and personality traits. Collectively, these findings highlight an important linkage between fine-scale nest site selection and nest success with certain personality traits in birds, and make a valuable contribution to addressing the knowledge gap in our understanding of the roles of personality traits in animal reproduction.

Limitations of the study

Our findings revealed that female personality traits were associated with clutch size, nest site selection, and nest success during incubation, so the major limitation of the present study is the lack of data concerning the roles of both parents’ personality traits during nestling period. Further studies are needed to explore how male and female personality traits affect the reproductive fitness.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- [KEY RESOURCES TABLE](#)
- [RESOURCE AVAILABILITY](#)
 - Lead contact
 - Materials availability
 - Data and code availability
- [EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS](#)
 - Study area and study species
 - Ethical statement
- [METHOD DETAILS](#)
 - Nest site characteristics
 - Measurement of personality traits
- [QUANTIFICATION AND STATISTICAL ANALYSIS](#)

SUPPLEMENTAL INFORMATION

Supplemental information can be found online at <https://doi.org/10.1016/j.isci.2023.107419>.

ACKNOWLEDGMENTS

This research was supported by the National Natural Science Foundation of China, China (Project 31472012 and 32070452).

AUTHOR CONTRIBUTIONS

Y.L. and Y.S. designed the experiment; Y.L., Q.Z., L.C., Y.H., and Y.F. performed the experiments in the wild, helped with data analysis and editing the manuscript; Y.L. and Y.S. wrote the first draft manuscript; H.L. helped with manuscript preparation and improved the English; all authors reviewed the work and approved the final version of the manuscript.

DECLARATION OF INTERESTS

The authors declare that they have no competing interests.

INCLUSION AND DIVERSITY

We support inclusive, diverse, and equitable conduct of research.

Received: April 28, 2023

Revised: June 8, 2023

Accepted: July 14, 2023

Published: July 19, 2023

REFERENCES

- Dall, S.R.X., Houston, A.I., and McNamara, J.M. (2004). The behavioural ecology of personality: consistent individual differences from an adaptive perspective. *Ecol. Lett.* 7, 734–739.
- Sih, A., Bell, A., and Johnson, J.C. (2004). Behavioral syndromes: an ecological and evolutionary overview. *Trends Ecol. Evol.* 19, 372–378.
- Møller, A.P. (2019). Head size and personality in great tits *Parus major*. *Anim. Biodivers. Conserv.* 42, 135–142.
- Carere, C., and Maestripietri, D. (2013). *Animal Personalities: Behavior, Physiology, and Evolution* (University of Chicago Press).
- Schuett, W., Tregenza, T., and Dall, S.R.X. (2010). Sexual selection and animal personality. *Biol. Rev. Camb. Phil. Soc.* 85, 217–246.
- Munson, A.A., Jones, C., Schraft, H., and Sih, A. (2020). You're just my type: Mate choice and behavioral types. *Trends Ecol. Evol.* 35, 823–833.
- Roth, A.M., Dingemanse, N.J., Nakagawa, S., McDonald, G.C., Løvlie, H., Robledo-Ruiz, D.A., and Pizzari, T. (2021). Sexual selection and personality: Individual and group-level effects on mating behaviour in red junglefowl. *J. Anim. Ecol.* 90, 1288–1306.
- Mutzel, A., Dingemanse, N.J., Araya-Ajoy, Y.G., and Kempenaers, B. (2013). Parental provisioning behaviour plays a key role in linking personality with reproductive success. *SAVE Proc. Royal Soc. B.* 280, 1–9.
- Thys, B., Eens, M., Pinxten, R., Iserbyt, A., and Naguib, M. (2021). Pathways linking female personality with reproductive success are trait- and year-specific. *Behav. Ecol.* 32, 114–123.
- Biro, P.A., and Stamps, J.A. (2008). Are animal personality traits linked to life-history productivity? *Trends Ecol. Evol.* 23, 361–368.
- Debecker, S., Sanmartín-Villar, I., de Guinea-Luengo, M., Cordero-Rivera, A., and Stoks, R. (2016). Integrating the pace-of-life syndrome across species, sexes and individuals: covariation of life history and personality under pesticide exposure. *J. Anim. Ecol.* 85, 726–738.
- Schuett, W., Godin, J.-G.J., and Dall, S.R.X. (2011). Do Female Zebra Finches, *Taeniopygia guttata*, Choose Their Mates Based on Their 'Personality'. *Ethology* 117, 908–917.
- Burtka, J.L., and Grindstaff, J.L. (2015). Similar nest defence strategies within pairs increase reproductive success in the eastern bluebird, *Sialia sialis*. *Anim. Behav.* 100, 174–182.
- Laubu, C., Dechaume-Moncharmont, F.X., Motreuil, S., and Schweitzer, C. (2016). Mismatched partners that achieve postpairing behavioral similarity improve their reproductive success. *Sci. Adv.* 2, e1501013.
- Lou, Y.Q., Chen, L.J., Hu, Y.B., Komdeur, J., Sun, Y.H., and Lyu, N. (2021). Animal personality can modulate sexual conflict over offspring provisioning. *Anim. Behav.* 177, 91–97.
- Spiegel, O., Leu, S.T., Bull, C.M., and Sih, A. (2017). What's your move? Movement as a link between personality and spatial dynamics in animal populations. *Ecol. Lett.* 20, 3–18.
- Rabdeau, J., Arroyo, B., Mougeot, F., Badenhauer, I., Bretagnolle, V., and Monceau, K. (2021). Do human infrastructures shape nest distribution in the landscape depending on individual personality in a farmland bird of prey? *J. Anim. Ecol.* 90, 2848–2858.
- Stuber, E.F., Carlson, B.S., Jesmer, B.R., and Pinter-Wollman, N. (2022). Spatial personalities: a meta-analysis of consistent individual differences in spatial behavior. *Behav. Ecol.* 33, 477–486.
- Holtmann, B., Santos, E.S.A., Lara, C.E., and Nakagawa, S. (2017). Personality-matching habitat choice, rather than behavioural plasticity, is a likely driver of a phenotype–environment covariance. *Proc. Biol. Sci.* 284, 20170943.
- Serrano-Davies, E., Araya-Ajoy, Y.G., Dingemanse, N.J., Sanz, J.J., and Tregenza, T. (2017). Personality-related differences in response to habitat in Mediterranean blue tits. *Ethology* 123, 861–869.
- Jacob, S., Bestion, E., Legrand, D., Clobert, J., and Cote, J. (2015). Habitat matching and spatial heterogeneity of phenotypes: implications for metapopulation and metacommunity functioning. *Evol. Ecol.* 29, 851–871.
- Citta, J.J., and Lindberg, M.S. (2007). Nest-site selection of passerines: effects of

geographic scale and public and personal information. *Ecology* 88, 2034–2046.

23. Latif, Q.S., Heath, S.K., and Rotenberry, J.T. (2012). How avian nest site selection responds to predation risk: testing an 'adaptive peak hypothesis'. *J. Anim. Ecol.* 81, 127–138.
24. Cunningham, J.A., Kesler, D.C., and Lanctot, R.B. (2016). Habitat and social factors influence nest-site selection in Arctic-breeding shorebirds. *The Auk* 133, 364–377.
25. Zhao, Q.S., Hu, Y.B., Liu, P.F., Chen, L.J., and Sun, Y.H. (2016). Nest site choice: a potential pathway linking personality and reproductive success. *Anim. Behav.* 118, 97–103.
26. DiNuzzo, E.R., and Griffen, B.D. (2020). The effects of animal personality on the ideal free distribution. *Proc. Biol. Sci.* 287, 20201095.
27. Selmann, M.W., Jaatinen, K., Steele, B.B., Öst, M., and Wright, J. (2014). Boldness and Stress Responsiveness as Drivers of Nest-Site Selection in a Ground-Nesting Bird. *Ethology* 120, 77–89.
28. Davis, S.K. (2005). Nest-site selection patterns and the influence of vegetation on nest survival of mixed-grass prairie passerines. *Condor* 107, 605–616.
29. Keyel, A.C., Strong, A.M., Perlut, N.G., and Reed, J.M. (2013). Evaluating the roles of visual openness and edge effects on nest-site selection and reproductive success in grassland birds. *The Auk* 130, 161–170.
30. Liu, J., Yan, H., Li, G., and Li, S. (2021). Nest concealment is associated with reproductive traits across sympatric bird species. *Ecol. Evol.* 11, 14079–14087.
31. Forstmeier, W., and Weiss, I. (2004). Adaptive plasticity in nest-site selection in response to changing predation risk. *Oikos* 104, 487–499.
32. Conway, C.J., and Martin, T.E. (2000). Evolution of passerine incubation behavior: influence of food, temperature, and nest predation. *Evolution* 54, 670–685.
33. Dingemanse, N.J., Moiron, M., Araya-Ajoy, Y.G., Mouchet, A., and Abbey-Lee, R.N. (2020). Individual variation in age-dependent reproduction: Fast explorers live fast but senesce young? *J. Anim. Ecol.* 89, 601–613.
34. Araya-Ajoy, Y.G., Kuhn, S., Mathot, K.J., Mouchet, A., Mutzel, A., Nicolaus, M., Wijmenga, J.J., Kempenaers, B., and Dingemanse, N.J. (2016). Sources of (co) variation in alternative siring routes available to male great tits (*Parus major*). *Evolution* 70, 2308–2321.
35. Santicchia, F., Gagnaison, C., Bisi, F., Martinoli, A., Matthysen, E., Bertolino, S., and Wauters, L.A. (2018). Habitat-dependent effects of personality on survival and reproduction in red squirrels. *Behav. Ecol. Sociobiol.* 72, 134.
36. Tamin, T., Morinay, J., Germain, M., Récapet, C., and Doligez, B. (2023). Behavioural syndrome between boldness and aggressiveness and link with reproductive success in a wild bird population. *Anim. Behav.* 197, 27–41.
37. Hollander, F.A., Van Overveld, T., Tokka, I., and Matthysen, E. (2008). Personality and Nest Defence in the Great Tit (*Parus major*). *Ethology* 114, 405–412.
38. Betini, G.S., and Norris, D.R. (2012). The relationship between personality and plasticity in tree swallow aggression and the consequences for reproductive success. *Anim. Behav.* 83, 137–143.
39. Grunst, A.S., Grunst, M.L., Pinxten, R., and Eens, M. (2019). Personality and plasticity in neophobia levels vary with anthropogenic disturbance but not toxic metal exposure in urban great tits: urban disturbance, metal pollution and neophobia. *Sci. Total Environ.* 656, 997–1009.
40. Vrublevska, J., Krama, T., Rantala, M.J., Mierauskas, P., Freeberg, T.M., and Krams, I.A. (2014). Personality and density affect nest defence and nest survival in the great tit. *Acta Ethol.* 18, 111–120.
41. Suckow, N.M., Pollock, H.S., Hauber, M.E., Kastner, M., Savidge, J., Baker, K., Rogers, H.S., and Goymann, W. (2022). Nest defense, personality, and fitness of a locally endangered island passerine. *Ethology* 128, 499–507.
42. David, M., Auclair, Y., Dechaume-Moncharmont, F.X., and Cézilly, F. (2012). Handling stress does not reflect personality in female zebra finches (*Taeniopygia guttata*). *J. Comp. Psychol.* 126, 10–14.
43. Dubuc-Messier, G., Réale, D., Perret, P., and Charmanier, A. (2017). Environmental heterogeneity and population differences in blue tits personality traits. *Behav. Ecol.* 28, 448–459.
44. Boon, A.K., Réale, D., and Boutin, S. (2007). The interaction between personality, offspring fitness and food abundance in North American red squirrels. *Ecol. Lett.* 10, 1094–1104.
45. Brommer, J.E., and Klun, E. (2012). Exploring the genetics of nestling personality traits in a wild passerine bird: testing the phenotypic gambit. *Ecol. Evol.* 2, 3032–3044.
46. Hall, M.L., van Asten, T., Katsis, A.C., Dingemanse, N.J., Magrath, M.J.L., and Mulder, R.A. (2015). Animal personality and pace-of-life syndromes: do fast-exploring fairy-wrens die young? *Front. Ecol. Evol.* 3.
47. Class, B., and Brommer, J.E. (2016). Senescence of personality in a wild bird. *Behav. Ecol. Sociobiol.* 70, 733–744.
48. Haigh, A., O'Riordan, R., and Butler, F. (2017). Variations in aggression and activity levels amongst squirrels inhabiting low and high density areas. *Ecol. Res.* 32, 931–941.
49. Piquet, J.C., López-Darias, M., van der Marel, A., Nogales, M., and Waterman, J. (2018). Unraveling behavioral and pace-of-life syndromes in a reduced parasite and predation pressure context: personality and survival of the Barbary ground squirrel. *Behav. Ecol. Sociobiol.* 72, 147.
50. Zhao, Q., and Sun, Y. (2018). Nest-Site Characteristics and Nesting Success of the Chestnut Thrush. *Ornithol. Sci.* 17, 3–9.
51. Houston, A.I. (2010). Evolutionary models of metabolism, behaviour and personality. *Philos. Trans. R. Soc. B* 365, 3969–3975.
52. Hu, Y.B., Zhao, Q.S., Lou, Y.Q., Chen, L.J., González, M.A., and Sun, Y.H. (2017). Parental attendance of Chestnut Thrush reduces nest predation during the incubation period: compensation for low nest concealment? *J. Ornithol.* 158, 1111–1117.
53. Zhao, Q.S., Hu, Y.B., Liu, P.F., Chen, L.J., and Sun, Y.H. (2016). Nest site choice: a potential pathway linking personality and reproductive success. *Anim. Behav.* 118, 97–103.
54. Patrick, S.C., Chapman, J.R., Dugdale, H.L., Quinn, J.L., and Sheldon, B.C. (2012). Promiscuity, paternity and personality in the great tit. *Proc. Biol. Sci.* 279, 1724–1730.
55. Nicolaus, M., Pialt, R., Ubels, R., Tinbergen, J.M., and Dingemanse, N.J. (2016). The correlation between coloration and exploration behaviour varies across hierarchical levels in a wild passerine bird. *J. Evol. Biol.* 29, 1780–1792.
56. Borgmann, K.L., and Conway, C.J. (2015). The nest-concealment hypothesis: new insights from a comparative analysis. *Wilson J. Ornithol.* 127, 646–660.
57. Eggers, S., Griesser, M., Nystrand, M., and Ekman, J. (2006). Predation risk induces changes in nest-site selection and clutch size in the Siberian jay. *Proc. Biol. Sci.* 273, 701–706.
58. Dillon, K.G., and Conway, C.J. (2018). Nest predation risk explains variation in avian clutch size. *Behav. Ecol.* 29, 301–311.
59. Trnka, A., and Grim, T. (2013). To compensate or not to compensate: testing the negotiation model in the context of nest defense. *Behav. Ecol.* 24, 223–228.
60. Lefcheck, J., Byrnes, J., and Grace, J. (2016). Package "piecewise SEM". R Package Version. 1.
61. Sun, Y.H., Swenson, J.E., Fang, Y., Klaus, S., and Scherzinger, W. (2003). Population ecology of the Chinese grouse, *Bonasa sewerzowi*, in a fragmented landscape. *Biol. Conserv.* 110, 177–184.
62. Colombelli-Négrel, D., and Kleindorfer, S. (2009). Nest height, nest concealment, and predator type predict nest predation in superb fairy-wrens (*Malurus cyaneus*). *Ecol. Res.* 24, 921–928.
63. Guan, H., Wen, Y., Wang, P., Lv, L., Xu, J., and Li, J. (2018). Seasonal increase of nest height of the Silver-throated Tit (*Aegithalos glaucogularis*): can it reduce predation risk? *Avian Res.* 9, 42.
64. Klun, E., Siitari, H., and Brommer, J.E. (2014). Testing for between individual correlations of

- personality and physiological traits in a wild bird. *Behav. Ecol. Sociobiol.* **68**, 205–213.
65. Blumstein, D.T., Daniel, J.C., and Evans, C.S. (2006). JWatcherTM 1.0 an Introductory User's Guide. 2-71.
66. Carere, C., and van Oers, K. (2004). Shy and bold great tits (*Parus major*): body temperature and breath rate in response to handling stress. *Physiol. Behav.* **82**, 905–912.
67. Gelman, A., Su, Y.-S., Yajima, M., Hill, J., Pittau, M., and Kerman, J. (2021). arm: data analysis using regression and multilevel/hierarchical models. R package version 1.11–2. 2020.
68. Anderson, D.R., and Burnham, K.P. (2002). Avoiding pitfalls when using information-theoretic methods. *J. Wildl. Manag.* **66**, 912–918.
69. Johnston, R., Jones, K., and Manley, D. (2018). Confounding and collinearity in regression analysis: a cautionary tale and an alternative procedure, illustrated by studies of British voting behaviour. *Qual. Quantity* **52**, 1957–1976.
70. Bates, D., Maechler, M., Bolker, B., and Walker, S. (2014). Introduction to Statistical Analysis Using R. R Package. version: 1-23.
71. R, C., Team. (2015). R: A Language and Environment for Statistical Computing.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental models: Organisms/strains		
Chestnut thrushes (<i>Turdus rubrocanus</i>)	Wild-caught in Lianhuashan Nature Reserve, Gansu Province, in China	N/A
Software and algorithms		
R software (v. 3.2.2)	R Core Team	https://www.rproject.org/
Deposited data		
Nest site selection data	This paper	Mendeley Data, V1, https://doi.org/10.17632/wtyvxf7dh.1

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources should be directed to and will be fulfilled by the lead contact, Yuehua Sun (sunyh@ioz.ac.cn).

Materials availability

DOIs are listed in the [key resources table](#).

Data and code availability

- All data reported in this paper are available at Mendeley, and the DOI is listed in the [key resources table](#).
- All original code in this paper are available at Mendeley, and the DOI is listed in the [key resources table](#).
- Any additional information required to reanalyze the data reported in this paper is available from the [lead contact](#) upon request.

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

Study area and study species

This study was conducted during 2013-2016 on a wild breeding chestnut thrush population located at the north edge of the Lianhuashan Nature Reserve, Gansu Province, in central China (34°40'N, 103°30'E). More detailed information about the reserve and local climate can be found in Sun et al.⁶¹ In our study area, only female chestnut thrushes select nest sites, in which they build open-cup nests in trees and shrubs beside the roads and farmlands^{50,52} and also near human residence during early May to late June. The nest height ranges from 0.90 m to 4.50 m.⁵⁰ We searched for nests of chestnut thrushes from late April to early July in each year of the study. The clutch size ranges from 1 to 5 (typically 3- 4 eggs), and incubation period and nestling period were 13-15 days, respectively. The chestnut thrushes re-nest when their nests are depredated, but only few individuals produce more than one brood in a year.

Ethical statement

All birds were captured under a bird ringing license issued from the China Bird Banding Center. All methods were carried out in the study were in accordance with the guidelines and regulations of the Animal Care and Use Committee of the Institute of Zoology, the Chinese Academy of Sciences. All experimental protocols were approved by Institute of Zoology, the Chinese Academy of Sciences, Permission number 2013/ 108. To reduce human disturbance during the incubation period, only one person checked each nest at a distance of over four m from the nest. We only checked nests during good weather conditions, when the ambient temperature was relatively high. Mist netting was done only under suitable weather conditions (no precipitation and low wind speed). Birds were trapped and released into the wild within

70 minutes near the trapping locations. No trapping-related mortality or nest abandon were recorded for females.

METHOD DETAILS

Nest site characteristics

Generally, the nests of chestnut thrushes were positioned 1-3 m above the ground,⁵² and we collected seven nest site characteristics when the nest failed or successfully fledged.⁵² During previous field observations, we found that chestnut thrushes nested closer to human settlements than other sympatric bird species, and are able to use some anthropogenic food resources. We also found that chestnut thrushes often foraged for food alongside the roads following rain. Furthermore, the nest height and nest concealment were also associated with reproductive success.^{30,62,63} Therefore, in our study, two scales of nest site characteristics were measured: large-scale which included distance of nest to human residences (m), distance of nest to farmland (m), distance of nest to the nearest road (m); and fine-scale measurements which included nest height (to the nearest 0.1 m), nest lateral concealment, and concealment above and below the nest. To minimize variation estimated by different observers, the same person (YB Hu) measured all nest site characteristics. All measurements followed Hu et al.⁵²: nest height was defined as the distance from the nest to the ground, to the nearest 0.1 m; nest lateral concealment was measured as the estimated vegetation cover from the most visible direction at a distance of 3 m after the observer checked from all directions. Concealment above and below the nest were recorded by estimating the percentage of vertical vegetation cover at 1 m above and 1 m below each nest, respectively. In addition, we recorded the location of all nests using a Global Positioning System (GPS) receiver, and then used the ArcGIS 10.2 (ESRI, Inc., Redlands, CA, USA) to measure the distances to human residences, roads and farmland habitat (to the nearest 1 m).

Measurement of personality traits

We captured birds by using mist nets which we positioned two meters away from the nest. In order to calculate the repeatability of personality traits, we tried to catch individuals twice during three breeding stages, including before clutch initiation, incubation and nestling period. The interval between two measurements was at least 7 days. Two personality traits, activity and breathing rate, were measured in 2013-2016. Activity was measured between 0900 and 1700 by using a simple cage test based on Klun et al.⁶⁴ and adapted by Zhao et al.²⁵ and Lou et al.¹⁵ We recorded the activity behaviors (such as walk, hop and fly) for 5 min after a 10 min acclimatization in the cage (size: L 60 × W 36 × H 60 cm) using JVC (GZ-E265AC). JWatcher software was used to analyse all recorded activity data.⁶⁵ We calculated weighted activity scores using the following formula, to account for differences in energy expenditure: activity score = walks × 1 + hops × 2 + flights × 3.^{15,25} Breathing rate was measured by using the handling stress test, and we recorded the total number of breast movements within 60 seconds.^{15,25} In general, we assumed that birds with lower breathing rates are bolder than those with higher rates.^{25,66} The body mass (to the nearest 0.1 g) was measured before release.

QUANTIFICATION AND STATISTICAL ANALYSIS

We constructed linear mixed models (LMMs) with individual identity as a random variable to calculate the repeatability of breathing rate and activity.^{15,53} Before the analysis, activity was square root transformed to meet Gaussian distribution of residuals. To control for confounding factors, seven predictor variables were included in each model: date (Julian date), year, sex, test sequence of activity and breathing rate (i.e., 1: first, 2: second or 3: third test), time of day (where 1200 = 0, 1300 = 1, 1100 = -1, etc.), context (i.e., 1: before breeding, 2: incubation or 3: nestling period) and body mass. All continuous variables were mean centered and standardized. The results were shown in the Table S1. Likelihood ratio test (LRT) between the models with and without the random effect (individual identity) was used to test the statistical significance of the repeatability. We used the LMMs of breathing rate and activity to run 1000 simulations using the *arm* package,⁶⁷ and the average best linear unbiased predictors (BLUPs) for the intercepts of each individual were used as personality profiles (breathing rate and activity) in the following data analysis, since a recent study found that the estimates of average BLUPs was less precise but unbiased.³³

To minimize the number of nest site characteristics, the principal component analysis (PCA) was performed on the large- and fine-scale nest site characteristics, respectively. We kept the first components extracted with eigenvalues > 1 (see table below), which could reflect important information in nest site characteristics. We used the factor scores from PCA (hereafter PC1 score) for individuals in the further analysis. In

large-scale nest site characteristics, higher PC1 score corresponded to longer distance among nests, human settlement and farmland. In the fine-scale nest site characteristics, higher PC1 score corresponded to greater lateral concealment.

Principal component analysis (PCA) of large- and fine-scale nest site characteristics in chestnut thrushes

Rotated factor loading

Large-scale nest site characteristics	PC1	PC2	PC3	PC4
Distance to human settlement	0.656	−0.147	−0.740	
Distance to farmland (m)	0.602	−0.488	0.631	
Distance to road (m)	0.454	0.860	0.232	
Eigenvalues	1.088	0.980	0.926	
% of variance	0.394	0.714	1	
Fine-scale nest site characteristics				
Nest height	0.289	0.925	−0.190	0.154
Nest lateral concealment	0.601	−0.003	0.296	−0.742
Concealment above the nest	0.572	−0.192	0.463	0.649
Concealment below the nest	0.477	−0.327	−0.814	0.063
Eigenvalues	1.356	0.972	0.851	0.702
% of variance	0.460	0.696	0.877	1

In order to investigate the relationships among female's personality traits (breathing rate and activity), large- and fine-scale nest site characteristics and clutch size, a piecewise structural equation model (pSEM) was used. The pSEM has the ability to propose causal relationships between predictor and response variables.⁶⁰ Individual identity was incorporated in all models as a random factor.

The pSEM was conducted by piecewiseSEM package in R,⁶⁰ which represents a list of generalized linear mixed models (GLMMs), and each pSEM was represented by a list of models which considered different variables as response variables. For the response variables, we built models which included potential paths between key variables, and the models were selected by Akaike information criterion (AIC). The best-fit model has the lowest AIC value⁶⁸ (Table S2). Fisher's C was used to assess the goodness of fit of the pSEM⁶⁰ and $P > 0.05$ indicated no missing paths among variables and that these models represented the data appropriately.

A GLMM with binomial distribution was used to investigate whether nest success during the incubation period (success was defined as at least one egg successfully hatched; failure was defined as the nest was depredated) was affected by PC1 scores of large- and fine-scale nest-site characteristics, Julian date, females' activity and breathing rate. Collinearity of fixed effects were examined to ensure the Variable Inflation Factor (VIF) with values < 2.5 .⁶⁹ We included female ID as random effect because of some repeated measurements of females among years. We also added year as a random effect to reduce the effects of environmental variations among years. All LMMs and GLMMs were fitted with *lmer* and *glmer* functions from the *lme4* package.⁷⁰ All statistics were performed with R software (v. 3.2.2).⁷¹