

Please cite the Published Version

Emamvirdi, Mahsa, Hosseinzadeh, Mahdi, Letafatkar, Amir, Thomas, Abbey C, Dos'Santos, Thomas, Smania, Nicola and Rossettini, Giacomo (2023) Comparing kinematic asymmetry and lateral step-down test scores in healthy, chronic ankle instability, and patellofemoral pain syndrome female basketball players: a cross-sectional study. Scientific Reports, 13 (1). p. 12412. ISSN 2045-2322

DOI: https://doi.org/10.1038/s41598-023-39625-1

Publisher: Nature Portfolio

Version: Published Version

Downloaded from: https://e-space.mmu.ac.uk/632554/

Usage rights: (cc) BY

Creative Commons: Attribution 4.0

Additional Information: This is an Open Access article which appeared in Scientific Reports.

Data Access Statement: The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines)

scientific reports

Check for updates

OPEN Comparing kinematic asymmetry and lateral step-down test scores in healthy, chronic ankle instability, and patellofemoral pain syndrome female basketball players: a cross-sectional study

Mahsa Emamvirdi^{®1}, Mahdi Hosseinzadeh^{®2⊠}, Amir Letafatkar^{®3}, Abbey C. Thomas^{®4}, Thomas Dos'Santos^{®5,6}, Nicola Smania^{®7,8} & Giacomo Rossettini^{®9}

We aimed to understand whether ankle dorsiflexion range of motion (ROM) and dynamic knee valgus (DKV) kinematic inter-limb asymmetries would be associated with the Lateral Step-Down Test (LSD) in basketball players with chronic ankle instability (CAI), patellofemoral pain (PFP) and healthy controls (HC). An observational cross-sectional study with a between-subject design was employed. Female basketball athletes with CAI (n = 20), PFP (n = 20) and HC (n = 20) were recruited. Ankle dorsiflexion-ROM, DKV angle during a single-limb squat, and LSD quality were measured bilaterally. The Asymmetry index (ASI) was calculated to identify between-limb percentage imbalances. The correlation matrix between the tasks was calculated. Ankle dorsiflexion-ROM was less in the CAI and PFP than in the HC group regardless of limb (p < 0.001). DKV angle was greater in the CAI and PFP than in the HC group (p = 0.001). LSDs were similar between the PFP and CAI groups (p = 0.698) but worse than the HC group (p = 0.001). The ASI showed asymmetry across all tasks (p < 0.001), with the greatest asymmetry for the DKV angle. The correlation matrix between tasks on both limbs was significant (p < 0.05). Our findings suggest significant asymmetries in ankle dorsiflexion-ROM and frontal plane knee control are present in female basketball athletes with CAI and PFP, and thus, highlights need to evaluate and reduce limb asymmetries in these populations.

Many health benefits accrue from consistent physical activity across the lifespan. Participation in physical activities, however, carries an inherent risk for acute and chronic musculoskeletal injury¹. Musculoskeletal injuries, particularly of the lower extremity, cause short-term disability, interfere with participation in physical activity, and are associated with joint disorders in later life². Thus, the general goal of preventing acute and chronic musculoskeletal injuries and reducing the associated burden among high-risk individuals is of considerable interest. Many studies report a higher prevalence of musculoskeletal pain among females in the general population³. When scholastic and colleagiate athletes are considered, females continue to experience greater rates of injury, particularly to the knee and ankle experiencing higher prevalences of Patellofemoral pain (PFP) and chronic ankle instability (CAI) in weight-bearing sports (e.g. basketball) compared to their male counterparts⁴. PFP is

¹Department of Biomechanics and Sport Injuries, Kharazmi University, Tehran, Iran. ²Department of Sport Injuries and Corrective Exercises, Sport Sciences Research Institute, No. 3, 5th Alley, Miremad Street, Motahhari Street, P O Box: 1587958711, Tehran, Iran. ³Sports Injury and Corrective Exercises, Faculty of Physical Education and Sport Sciences, Kharazmi University, Tehran, Iran. ⁴Department of Applied Physiology, Health, and Clinical Sciences, University of North Carolina at Charlotte, Charlotte, NC, USA. ⁵Department of Sport and Exercise Sciences, Musculoskeletal Science and Sports Medicine Research Centre, Manchester Metropolitan University, Manchester, UK. ⁶Manchester Institute of Sport, Manchester Metropolitan University, Manchester, OK. ⁷Department of Neurosciences, Biomedicine and Movement Sciences, University of Verona, Verona, Italy. ⁸Neuromotor and Cognitive Rehabilitation Research Centre (CRRNC), University of Verona, Verona, Italy. ⁹School of Physiotherapy, University of Verona, Verona, Italy. ^{Email:} Metti@ssrc.ac.ir a multifactorial clinical condition resulting from abnormal patellofemoral joint loading, increasing joint stress and producing retropatellar pain⁵. Conversely, CAI represents both mechanical and functional instability of the ankle joint resulting in repetitive bouts of instability following an initial lateral ankle sprain injury⁶. PFP and CAI represent two of the most prevalent musculoskeletal injuries among physically active individuals⁴ and, while these conditions affect different joints, a commonality among them is that patients with these musculoskeletal disorders experience aberrant movement patterns that may contribute to/arise from the disorder's presence⁷.

Previous studies have reported that dynamic knee valgus (DKV), an aberrant movement pattern during weight-bearing activities, is a significant risk factor in patients with PFP⁷. Though few researchers have investigated DKV angle in persons with CAI, those who have studied it found DKV angle to be greater in this population compared to their health counterparts^{8,9}. Accordingly, an increased DKV angle is not only associated with painful knee conditions but can also be related to dysfunctions in other lower limb joints (e.g. the ankle)¹⁰. Reduced ankle dorsiflexion range of motion (ROM) has been reported as a risk factor for some conditions, such as patellar tendinopathy¹¹, Achilles tendinopathy^{12,13}, chronic ankle instability¹⁴, metatarsal stress fractures¹¹, and anterior knee pain¹⁵.

In a closed kinetic chain, the ankle acts as a firm base of support such that its movement restriction and/ or instability can affect the function of proximal joints¹⁴. Therefore, restriction of ankle dorsiflexion ROM represents a possible risk factor for excessive DKV angle^{16,17} during squat and jump landing tasks¹⁸ and is linked to injurious landing mechanics¹⁹. With restricted ankle dorsiflexion ROM, individuals may try to compensate with movement in the frontal or transverse plane throughout the overall kinetic chain^{16,17}, thus, creating an increased DKV angle^{16,17,20}. This compensation may present as foot pronation²⁰, internal tibial rotation²⁰, hip internal rotation and adduction²¹, pelvic drop¹⁵, or gastrocnemius and soleus tightness²². Patients with PFP²² and CAI¹⁴ commonly display a limited ankle dorsiflexion ROM with adverse functional consequences for the knee joint biomechanics during landing⁹, ascending/descending stairs, squatting, jumping and running²³. Restricted dorsiflexion will also increase impact forces / mechanical loads and redistribute them proximally to the knee and hip in weight-bearing activities such as the Lateral Step-Down test (LSD)¹⁴. The LSD is a simple, clinicianfriendly tool designed to assess lower extremity movement quality during a functional activity²⁴. The test allows clinicians to identify faulty movement patterns and evaluate the trunk, hip and knee behaviour during the task²⁴. Since its creation, the LSD has been used to assess patients' quality of movement with CAI and PFP^{14,15}. There are several advantages of using LSD to assess movement quality, mainly because it is a quick and easy test to perform in a clinical environment²⁴. Moreover, the performance of the LSD score is influenced by a reduced ankle dorsiflexion ROM in individuals with PFP and CAI^{14,15}. Therefore, ankle dorsiflexion ROM represents a key assessment when patients demonstrate a lower movement quality during an LSD^{14,15}.

Quantifying neuromuscular control between legs (e.g. reduced function or performance in one limb compared to the other) is critical to identify individuals potentially at risk of injury, establishing when an athlete can return to play following injury, and optimizing strength and conditioning training²⁵. Neuromuscular asymmetry of the lower limbs is associated with potential injury and can be used to predict future injury or reinjury²⁶. Inter-limb asymmetries (ILA) may potentially place both legs at an increased risk of injury in sports; the strong leg may sustain excessive stress due to high dependence and loading (i.e. overuse), whereas the weak leg may be compromised to a potentially lower load tolerance capacity²⁷. In addition, ILA have been associated with an increased risk of sports injury because the asymmetries may result in unequal force attenuation or a loss of frontal plane stability, which are important to sustain the impacting forces²⁶. For example, research has examined lower ILA, mainly in healthy basketball players, during functional movements (e.g. cutting, pivoting, running)²⁸, leaving uncertainty about their role in athletes with CAI and PFP. Individuals with PFP may have gluteus medius muscle activation asymmetry, which may be associated with pain severity²⁹. Also, Nakagawa et al.³⁰ showed that male military recruits with greater asymmetry on the Y-Balance Test (YBT) posterolateral direction and Frontal Plane Knee Projection Angle (FPKPA) during single-leg squat were at a greater risk of developing PFP³⁰. However, Plastaras et al.³¹ showed early stages of unilateral PFP in female runners was not associated with hip abduction strength asymmetry. Regarding patients with CAI, the asymmetry between right and left anterior reach distances (>4 cm) and limitations in posterolateral reach distances (<80% normalized reach distance) represent risk factors for lateral ankle sprains and may contribute to CAI³². Tajdini et al. showed that patients with CAI walk with greater ILA in vertical ground reaction force and muscle activity across the gait cycle compared to the non-CAI group³³.

Although limited dorsiflexion ROM and its effect on excessive DKV and lower movement quality have been reported in PFP and CAI^{10,14}, to our best knowledge, there is a lack of research concomitantly investigating all three tasks (DKV angle, ankle dorsiflexion ROM and movement quality) in both lower limbs (injured and contralateral) among basketball players presenting PFP and CAI. From a clinical perspective, a better understanding of these phenomena would inform researchers and clinicians (e.g. physical therapists and physicians) to decide on the assessment, treatment evolution, and improving movement control of these patients considering both limbs and functional relationship of the lower kinetic chain³⁴.

Therefore, this study aims to understand whether dorsiflexion ROM and DKV kinematic ILA would be associated with the LSD scores in basketball players with CAI or PFP compared to healthy controls, and whether differences exist in these measures between populations. We hypothesized that: (1) there would be significant differences in DKV angle, ankle dorsiflexion ROM, and LSD scores in CAI and PFP compared to healthy controls; (2) there would be no significant difference between CAI and PFP in DKV angle, ankle dorsiflexion ROM, and LSD scores; (3) ankle dorsiflexion ROM and DKV asymmetry would be associated with LSD scores asymmetry in PFP and CAI; and (4) there would be an inverse association between ankle dorsiflexion ROM and LSD scores and a direct relation between DKV angle and LSD scores.

Materials and methods

Study design. This observational cross-sectional study was conducted and reported following the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guidelines. The study conformed to the Declaration of Helsinki. All participants provided written informed consent. The ethics board of the Sport Science Research Institute (IR.SSRC.REC.1400.120) approved this study.

Participants. Participants in this study were recreational, female, basketball athletes recruited from sports clubs in Iran. Participants were between 20 and 30 years of age (CAI: 22.70 ± 1.94 years, PFP: 23.05 ± 2.08 years, Healthy: 22.75 ± 2.31 years). The population of female basketball players were chosen due to the high prevalence of CAI and PFP^{4,35}. A physiotherapist with 13 years of experience in diagnosing and treating patients with musculoskeletal disorders assessed all the athletes, classifying them eligible when presenting with unilateral CAI, unilateral PFP or healthy conditions based on the inclusion and exclusion criteria reported in Table 1. Further, the physiotherapist confirmed the absence of PFP in the CAI group and the absence of CAI in the PFP group. Patients of the CAI and PFP groups were matched in age with healthy individuals. Moreover, during the research period, the participants were not receiving medical or rehabilitative care for CAI and PFP.

Sample size. G-Power software (G*Power©, University of Dusseldorf, Germany) was used to estimate the sample size according to a recent similar study that assessed DKV angle and LSD. The analysis revealed that to perform repeated measures, between factors Multivariate Analysis of Variance (MANOVA) with an effect size of 0.30, power of 0.30, and alpha of < 0.05, we would need at least 19 participants. To account for potential dropout, we determined a minimum sample size of 60 participants was required to be assigned into three groups: 20 participants with CAI, 20 with PFP, and 20 healthy females.

Setting and procedure. In the pre-screening phase, the research group distributed recruitment flyers by hand (brochures) to basketball clubs' meetings in the metropolitan area of Tehran. Before participating in the study, all athletes were briefed about the objectives and read and signed the informed consent form. Then, the included participants attended a baseline assessment as follows:

- (1) The CAI group completed a series of patient-reported outcomes surveys to confirm a history of significant ankle sprains/giving way/recurrent ankle sprains, including the Iranian version of the Cumberland Ankle Instability Tool Questionnaire (CAIT)³⁶ and the Persian version of Foot and Ankle Ability Measure (FAAM) for activities of daily living (ADL) and SPORTS subscales³⁷.
- (2) The PFP group were assessed by the Numerical Pain Rating Scale (NPRS)³⁸ and the Clark test³⁹.
- (3) The Healthy group completed the CAIT³⁶, FAAM-ADL and FAAM-SPORTS³⁷, NPRS³⁸ and Clark test³⁹.

Next, participants reported their age and involved limb, had their height (cm) and body mass (kg) measured using a tape measure and a digital scale, respectively, and their body mass index (BMI) calculated.

Moreover, the ankle dorsiflexion ROM and DKV angle were measured, and the LSD was performed for both lower limbs. The first limb measured (right or left) was randomized, but ankle dorsiflexion ROM measures were always obtained before the DKV angle and LSD assessments. All the assessments were conducted with the participants barefoot at Kharazmi University in January 2020. An examiner with 13 years of experience that received

	Inclusion criteria	Exclusion criteria
CAI group	 CAI was defined as incurring at least one ankle sprain and at least one subsequent episode of giving way occurring at least 12 months before the study We used the Iranian version of CAIT, FAAM- ADL and the FAAM-Sport Reporting a score: <24 on the (CAIT)³⁶, <90% on the FAAM- ADL <80% on the FAAM-Sport^{36,37} 	- A recent history (<6 months) of lower extremity injury/surgery (includ- ing lateral ankle sprain), diagnosis of ankle osteoarthritis, history of ankle surgery involving intra-articular fixation, or current pregnancy ¹⁴
PFP group	-Participants were included in the PFP who showed anterior knee pain for at least three months during the performance of at least two of the following tasks: ascending and descending stairs, squatting, running, jumping or remaining seated for a long time, besides showing a minimum score of three points in the NPRS ³⁸ -Participants also must have presented with an insidious onset of symptoms unrelated to trauma and a positive Clark test ³⁹	-Tenderness to palpation of the patellar tendon, the inferior pole of the patella, or tibial tubercle as the primary complaint. Tenderness to palpation of the patellar tendon, the inferior pole of the patella, or tibial tubercle as the primary complaint ¹⁰ -Other diagnoses of the knee including: patellar tendinitis, iliotibial band syndrome, Osgood- Schlatter's disease, Sinding-Larsen's Johansson's disease, fracture, or ligamentous injuries. Prior knee surgery, history of patellar subluxation or dislocation ¹⁰
Control healthy group	-Participants had to be free of lower extremity symptoms -Without any history of ankle sprains -No pain in the NPRS -Negative Clark test	 History of pathology involving the comparable knee, ankle or other joints of the lower extremity Reported any history of lower extremity surgery, neuropathies, diabetes, balance disorder, Raynaud's diseases, cold-induced circulatory problems, and other conditions known to affect balance

Table 1. Inclusion and exclusion criteria. *CAI* chronic ankle instability, *PFP* patellofemoral syndrome, *CAIT* FAAM-*ADL* foot and ankle ability measure-activities of daily, *CAIT* cumberland ankle instability tool, *FAAM-ADL* foot and ankle ability measure-activities of daily, *FAAM-sport* foot and ankle measure-sport, *NPRS* numeric pain rating scale.

1 0

5 h of training in ankle dorsiflexion ROM, DKV angle, and LSD assessments performed all the measurements. The examiner was blinded to the group allocation to avoid bias during the evaluation.

Instruments and measurements. Ankle dorsiflexion ROM. A digital inclinometer was used to record weight-bearing ankle dorsiflexion ROM by measuring the tibia angle to the ground during a two-point staggered upright position⁴⁰. Ankle dorsiflexion ROM in the weight-bearing lunge (WBLT) was measured with an Acumar^{**} digital inclinometer (model ACU360 Lafayette Instrument Company, Lafayette, IN, USA Single Digital Inclinometer). The participant put the tested foot along a 50-cm-long line drawn on the ground, and a continuous 60-cm-long line was drawn on a wall. A test taker placed an inclinometer on the anterior aspect of the tibia, 15 cm distal to the tibial tuberosity, so that the line bisected the heel, and the second toe was on the line. The subject lunged forward and directed their patella as close as the line drawn on the wall while the heel remained in contact with the floor. This puts the ankle joint in maximal dorsiflexion. Once maximal dorsiflexion was reached, the ankle dorsiflexion ROM was recorded (Fig. 1a)⁴⁰. Three assessments were recorded, and the average of the three trials was used for further analysis⁴¹. This method results in higher reliability coefficients (ICC = 0.96 to 0.99), representing a valuable assessment strategy⁴¹. In this study, the tests were conducted for the intraclass reliability of the examiner, resulting in a correlation coefficient (within-session reliability) of 0.95 (0.92–0.97).

Dynamic knee valgus (DKV) angle measurement. We measured the DKV angle during a single-leg squat. The single-leg squat was chosen because it allows for easy visualization of poor neuromuscular control (e.g. increased DKV angle) in patients with PFP and $CAI^{7,8}$. Participants were asked to stand on the test limb, facing the video camera and then to squat down at an angle of at least 45° knee flexion but not greater than 60°, for 5 s. The knee



Figure 1. (a) Weight-bearing lunge test ankle dorsiflexion ROM. (b) Dynamic Knee Valgus (DKV) angle measurement. (c) Lateral step-down test (LSD).

flexion angle was checked during practice trials (maximum of three) using a standard goniometer observed by the same examiner throughout all trials⁴². Trials were only accepted if the participant squatted within the desired range of knee flexion and maintained their balance while keeping their hands on their iliac crests throughout the trial⁴². Two-dimensional videos of the single-leg squat were captured using a Canon Vixia HF R42 digital camera (Canon USA), sampling at 60 frames per second. The camera was placed at the height of the subject's knee, 3 m anterior to the participant's squatting area, and aligned perpendicular to the frontal plane. DKV angle was calculated as the angle formed between the thigh and shank segments (Fig. 1b). Specifically, a line drawn from the anterior superior iliac spine to the midpoint of the knee, bisecting the thigh, defined the thigh segment. A line drawn from the knee's midpoint to the ankle's midpoint defined the shank segment. The DKV angle was calculated as 180° minus the angle between the thigh and shank segments⁴². DKV angles were obtained at peak knee flexion for each trial, determined visually by the investigator. The same individual who experienced palpation placed the markers on all participants. The average DKV angle value for both measures from three trials was analyzed⁴² and processed using Kinovea Software (v0.8.15; Kinovea Open Source Project, www.kinov ea.org). Within-day ICCs showed good reliability and ranged from 0.59 to 0.88, and between-days ICCs were good to excellent, ranging from 0.72 to 0.91⁴². In this study, the tests were conducted for the intraclass reliability of the examiner, resulting in a correlation coefficient (within-session reliability) of approximately DKV 0.92 (0.90 - 0.95).

Lateral step-down (LSD). The LSD measured the quality of functional movements²⁴. Participants stood in a unilateral stance atop a 20-cm step with the medial border of their foot near the stair edge and their hands on the hips. The limb not being tested was held off the step over the floor. In the next stage, they were asked to slowly descend the stair by flexing the test limb knee so that the contralateral heel touched the ground and returned to the initial position. This maneuver was repeated for 5 repetitions. Participants received no feedback on performing the task and no information on performance errors. A video camera (Canon Vixia HF R42 digital camera [Canon USA]; 60 fps sampling rate) aligned with the frontal plane was used to record the assessments for subsequent scoring (Fig. 1c). There were no limits on the number of times a video was watched or constraints on the playback speed when scoring the LSD. All participants began with a score of 0, and 1 point was added for each error committed¹⁵. Errors included: picking up hands from the hips, rotating or lifting the pelvis, flexing the trunk, moving into DKV (tibia protrusion deviating outwards from the second finger), and inability to sustain unilateral balance (e.g. bearing weight given the opposite limb)¹⁵. If the DKV angle exceeded the foot midline, an extra point was given for a total of 6 points. Higher scores (more errors) indicated poorer movement quality (0-1: "good", 2-3: "moderate", and 4-6: "poor")¹⁵. Pilot data from our lab suggest inter-rater agreement on the LSD to be substantial (kappa = 0.74), which aligns with previously established values in the literature (k = 0.67- $(0.81)^{43}$.

Data analysis. The Asymmetry index (ASI) was calculated to identify functional imbalances between limbs using the following formula:⁴⁴ ASI = (contralateral (or left) leg- involved (or right) leg/contralateral leg) \times 100.

To confirm that inter-limb differences are meaningful, we calculated the coefficient of variation $(CV)^{45}$ as $(SD/mean) \times 100$ for each participant and then averaged across all participants.

We considered the right side as the involved leg for the healthy group when calculating the ASI.

Statistical analysis. The independent variables in this study were group (PFP, CAI, or healthy) and limb (involved, contralateral). Dependent variables were LSD, DKV angle, and ankle dorsiflexion ROM. The SPSS software (ver. 23 for Windows; SPSS Inc., Chicago, IL, USA) was used to analyze the data collected in this study. Descriptive statistics were adopted to summarize the data with measures of central tendency. The Shapiro-Wilk and also Kolmogorov-Smirnov tests assessed the normal distribution of the tasks. A Two-way mixed model ANOVA was used to compare the interaction of group on limb on each dependent variable. The Scheffe test was used for multiple comparisons for the post hoc test. Paired t tests were adopted to further elucidate differences between the involved and contralateral legs in three tasks (ankle dorsiflexion ROM, DKV angle, and LSD). For non-parametric data (LSD), the Kruskal Wallis and Mann-Whitney U tests were used to determine the difference between groups. To determine the association between tasks on the involved and contralateral limbs, the relationship between ankle dorsiflexion ROM measures, DKV angle, and the LSD score was determined using a Spearman rank correlation coefficient. Correlation values were interpreted as follows: <0.25 little or no relationship, 0.25–0.50 fair relationship, 0.50–0.75 moderate to good relationship, >0.75 good to excellent relationship⁴⁶. The alpha value was set at p < 0.05 for all analyses. Finally, effect sizes (ES) (bias-corrected Hedges' g) and 95% confidence interval (CI) were also computed to estimate the precision and magnitude of group differences, given the multiple t tests ran. Effect sizes were interpreted as trivial (≤ 0.19), small (0.20–0.59), moderate (0.60–1.19), large (1.20–1.99), very large (2.00–3.99), and extremely large (\geq 4.00) for paired and independent comparisons⁴⁷.

Ethics declarations. The study was approved by Research Ethics Committees of Iran institute of sport science (Approval IR.SSRC.REC.1400.120). This study was performed in accordance with the standards of ethics outlined in the Declaration of Helsinki.

Results

Participants. One hundred and twenty-nine basketball players were screened, 76 participants were assessed for eligibility based on the inclusion/exclusion criteria. Next, we selected 24 participants from the PFP group, 26 from the CAI group, and 26 from the healthy group. Finally, considering the dropout and based on the exclusion

criteria, we included 60 participants (20 participants in each group). The flowchart of the participants' selection is reported in Fig. 2.

Descriptive data. There were no significant differences between groups in age, height, mass, and BMI (p > 0.05; Table 2). The CAI group reported lower CAIT, FAAM-ADL, and FAAM-Sport scores (p < 0.001) compared to the healthy group (Table 2). The PFP group reported higher NPRS (p < 0.001) than the healthy group.

Outcome data. The results demonstrated differences between the three groups for ankle dorsiflexion ROM, DKV angle and LSD in both the involved and contralateral lower limbs (p < 0.001; Tables 3 and 5). In addition, Two -way mixed model ANOVA revealed significant differences in ankle dorsiflexion involved*LSD contralateral in PFP, LSD contralateral* ankle dorsiflexion ROM contralateral in group CAI, and LSD involved* DKV angle involved in CAI group. Regarding the other factors in the three groups, there was no significant difference (Table 4).



Figure 2. Flowchart for enrollment of the participants in the current study.

						One-way	Multiple di	fference	comparison			
Groups	CAI (n=20)	PFP (n=20)	Healthy (n=20)	Kolmogorov–Smirnov test	Shapiro–Wilk's test	Anova (p-value)	CAI vs PFP	ES	CAI vs Healthy	ES	PFP vs healthy	ES
Age (y)	22.70 ± 1.94	23.05 ± 2.08	22.75 ± 2.31	0.109	0.056	0.853	0.932	0.174	0.357	0.023	0.445	0.136
Mass (kg)	56.65 ± 3.49	56.25 ± 2.80	57.60±3.13	0.097	0.056	0.388	0.128	0.126	0.517	0.286	0.318	0.454
Height (cm)	166.65±4.45	166.90±3.98	168.85±3.89	0.200	0.196	0.190	0.513	0.059	0.403	0.526	0.842	0.724
BMI (Kg/ m ²)	20.40 ± 1.22	20.19 ± 0.68	20.19 ± 0.64	0.200	0.560	0.679	0.60	0.212	0.041	0.215	0.817	0.000
Involved limb (right/ left)	14/6	15/5	-	-	-	-	-	-	-	-	-	-
CAIT	17.30 ± 2.38	-	30.00 ± 0.00	-	-	0.000	-	-	0.000	-	-	-
FAAM- ADL	76.36±2.89	-	100.00 ± 0.00	-	-	0.000	-	-	0.000	-	-	-
FAAM- Sport	69.01±4.96	-	100.00 ± 0.00	-	-	0.000	-	-	0.000	-	-	-
NPRS	-	5.95 ± 1.05	0.00 ± 0.00	-	-	0.000	-	-	-	-	0.000	-

Table 2. Participant characteristics at the baseline (mean \pm SD). *CAI* chronic ankle instability, *PFP* patellofemoral syndrome, *n* number, *y* year, *kg* kilogram, *cm* centimeter, *BMI* body mass index, *CAIT* cumberland ankle instability Tool, *FAAM*-ADL foot and ankle ability measure-activities of daily, *FAAM* sport foot and ankle measure-sport, *NPRS* numeric pain rating scale, *ES* effect size.

.....

Ankle dorsiflexion ROM. Ankle dorsiflexion ROM was significantly greater in healthy controls compared to CAI and PFP groups with very large and moderate effect sizes, respectively (Fig. 1a, Tables 3 and 4). The contralateral ankle dorsiflexion ROM was significantly greater in the CAI compared to the PFP group with a moderate effect size and significantly greater in healthy controls compared to the PFP group with a large effect size (Fig. 3a, Tables 3 and 4).

DKV. Contralateral limb DKV angle was significantly greater in the CAI and PFP groups compared to healthy controls with very large effect sizes. (Fig. 3b, Tables 3 and 4).

LSD. A significant main effect of DKV involved was observed for LSD involved in individuals with CAI (Table 4). Involved limb DKV angle was significantly greater in the CAI and PFP groups than in healthy controls with large effect sizes (Tables 3 and 4). A significant main effect of ankle dorsiflexion ROM was observed for LSD contralateral in individuals with PFP. A significant main effect of ankle dorsiflexion ROM contralateral was also observed for LSD contralateral in individuals with CAI (Table 4). LSD scores revealed no significant difference in movement quality based on the test in the involved limb in the CAI group compared to the PFP group with a trivial effect size. However, three scores differed between groups with a very large effect size (Tables 3 and 4). Contralateral limb LSD scores were 1.15 points greater in the CAI compared to the healthy groups with a very large effect size. Similarly, contralateral LSD scores were 1.85 points greater in the PFP group, compared to the healthy group with a very large effect size (Fig. 3c, Tables 3 and 4).

Asymmetry index. Testing of homogeneity of variance revealed that data were not normally distributed. There was a significant difference in asymmetry index across all tasks (ankle dorsiflexion ROM, DKV angle) in each group separately (Table 5). The greatest asymmetry was presented in the DKV angle for the CAI and PFP groups (-23%, -18%, respectively). It is worth noting that PFP and healthy groups in ankle dorsiflexion ROM had a CV value greater than the asymmetry score; thus, as a group, these athletes did not have a meaningful between-limb imbalance in ankle dorsiflexion ROM.

Correlation. The relation between all tasks on both the involved and contralateral sides was significant (Table 6). Furthermore, an inverse moderate to good linear relation existed between ankle dorsiflexion ROM with DKV angle and LSD scores. Alternatively stated, as dorsiflexion ROM increased, the DKV angle and LSD score decreased (better movement quality). Finally, a good to excellent direct linear relation was also observed between the DKV angle and the LSD scores. That is, a greater ankle dorsiflexion ROM, DKV angle, and LSD, on the involved limb was associated with greater values on the contralateral.

Discussion

This study aimed to quantify differences in ankle dorsiflexion ROM, DKV angle, and LSD across groups as well as determine the association between limb and ankle dorsiflexion ROM, DKV angles, and the LSD ILAs scores in basketball players with CAI, PFP and healthy controls. We observed that female basketball players with PFP and CAI have impaired ankle dorsiflexion ROM, greater DKV angle, poorer scores in the LSD, and greater ILAs than healthy controls (Tables 3, 5 and Fig. 3a–c). Differences in ankle dorsiflexion ROM, DKV angle, and LSD

								Multiple co	mparisons	differences (p	(
Variable	Kolmogorov-Smirnov test	Shapiro-Wilk's test	Group	Kolmogorov-Smirnov test	Shapiro-Wilk's test	mean±SD	ANOVA	CAI vs PFP	ES	CAI vs healthy	ES	PFP vs healthy	ES
Ankle			CAI	0.200	0.858	37.99 ± 3.36							
dorsitlex- ion ROM	0.200	0.167	PFP	0.200	0.427	40.97 ± 6.22	F = 17.84 p-value = 0.001	-2.98 (0.153)	-0.59	-8.86 (0.001)	-2.29	-5.88 (0.001)	-1.10
involved			Healthy	0.200	0.927	46.90 ± 4.33	4					Ì	
Ankle			CAI	0.116	0.054	44.44 ± 3.62							
dorsiflex- ion ROM	0.200	0.306	PFP	0.200	0.466	40.04 ± 5.29	F = 12.90 p-value = 0.001	(0.011)	0.97	-2.67 (0.174)	-0.67	- 7.07 (0.001)	-1.47
contralateral			Healthy	0.200	0.313	47.11 ± 4.25	4						
			CAI	0.200	0.887	16.36 ± 2.62							
DKV involved	0.063	0.035	PFP	0.200	0.295	16.86 ± 3.26	F = 82.03 p-value = 0.001	(0.583)	-0.16	7.70 (0.001)	3.43	8.20 (0.001)	3.12
			Healthy	0.200	0.358	8.66±1.78		(2000)		(1000)		(
			CAI	0.200	0.305	13.24 ± 1.85							
DKV con- tralateral	0.200	0.236	PFP	0.73	0.104	14.26 ± 3.04	F = 53.90 p-value = 0.001	-1 (0.647)	-0.40	5.16 (0.001)	2.81	6.16 (0.001)	2.46
			Healthy	0.197	0.175	8.09 ± 1.81	L L					Ì	
			CAI	0.000	0.006	3.45 ± 0.94							
LSD involved	0.000	0.001	PFP	0.000	0.002	3.40 ± 1.09	F = 70.28 p-value = 0.001	0.05 (0.698)	0.04	2.90 (0.001)	3.83	2.85 (0.001)	3.34
			Healthy	0.000	0.000	0.55 ± 0.51	4					Ì	
			CAI	0.000	0.000	1.40 ± 0.59							
LSD con- tralateral	0.000	0.000	PFP	0.000	0.002	2.10 ± 1.16	F = 27.36 p-value = 0.001	-0.7 (0.049)	-0.76	(0.001)	2.20	1.85 (0.001)	2.10
			Healthy	0.000	0.000	0.25 ± 0.44	4						
Table 3. R motion, <i>DI</i>	tesult of one-way ANOV «V dynamic knee valgus	/A on data in healt \$, LSD lateral step-	hy particif. down test,	vants and those presentiv ES effect size, SD stands	ng with CAI and P 11d deviation, <i>RON</i>	FP. CAI chro 1 range of me	nic ankle instab otion.	ility, <i>PFP</i> p	atellofem	oral syndrc	ome, ROM	[range of	

	LSD inv	olved		LSD con	itralatera	1	LSD in contral	volved*L ateral	.SD
Dependent variables	CAI	PFP	Healthy	CAI	PFP	Healthy	CAI	PFP	Healthy
Ankle dorsiflexion ROM involved	0.253	0.146	0.570	0.306	0.042*	0.124	0.732	0.837	-
Ankle dorsiflexion ROM contralateral	0.264	0.730	0.739	0.016*	0.404	0.269	0.841	0.823	-
DKV involved	0.016*	0.669	0.093	0.305	0.120	0.684	0.888	0.848	-
DKV contralateral	0.051	0.669	0.144	0.751	0.120	0.614	0.934	0.753	-

Table 4. Results of the two-way mixed model ANOVA. Data are presented as P-values. The mean difference is significant at the 0.05 level, *CAI* chronic ankle instability, *PFP* patellofemoral syndrome, *ROM* range of motion, *DKV* dynamic knee valgus, *LSD* lateral step-down test, *ROM* range of motion. * indicates significant difference at the p < 0.05 level.



Figure 3. (a) Comparison of the ankle dorsiflexion ROM in groups with CAI, PFP and the healthy group. *ADF ROM* ankle dorsiflexion range of motion, *CAI* chronic ankle instability, *PFP* patellofemoral pain. (b) Comparison the DKV angle in the groups with the CAI, PFP, and the healthy group. *DKV* dynamic knee valgus angle, *CAI* chronic ankle instability, *PFP* patellofemoral syndrome. (c) Comparison of the LSD scores in the groups with the CAI, PFP and the healthy group. *LSD* lateral step-down test, *CAI* chronic ankle instability, *PFP* patellofemoral syndrome.

scores in individuals with CAI and PFP were observed compared to the healthy controls, supporting our first hypothesis and substantiating the findings of previous studies^{14,15,30}.

The ankle dorsiflexion ROM in the involved limb was 9° less in the CAI group and 6° less in the PFP group compared to the healthy control group (Table 3, Fig. 3a), while differences in the uninvolved limb were 4° greater in the CAI compared to the PFP group and 7° less in the PFP compared to healthy controls. These ROM findings agree with Grindstaff et al.¹⁴, who reported that patients with CAI had reduced ankle dorsiflexion ROM

	Group					Multiple comparison (post-hoc test) on asymmetry between limbs						
ASI (%)	CAI	PFP	Healthy	Total	P-value	(I) Group	(J) Group	I-J mean ± SD ASI (%)	I-J p-value			
Ankle dorsiflexion	14 39 + 5 57*†	-246 ± 9.09	-0.43 + 2.27	4 12 + 9 0 66		CAL	PFP	16.86±2.38	< 0.001*			
ROM mean ± SD	14.57 ± 5.57	2.40 ± 9.09	0.45 ± 2.27	4.12 ± 2.0.00		0/11	Healthy	13.96±1.34	< 0.001*			
EC	1.04	0.16	0.03	2.16	0.001 E = 41.02	DED	CAI	-16.86 ± 2.38	< 0.001*			
Lo	1.74	0.10	0.05	5.10	0.001, r = 41.02	FFF	Healthy	-2.89 ± 2.09	0.440			
CV04	11.00 + 4.56	5 92 + 2 90	1 29 + 0 76 6	1.29 ± 0.76	6 10 + 5 06		Healthy	CAI	-13.96±1.384	< 0.001*		
C V 70	11.09±4.50	3.82 ± 2.89	1.38±0.76	0.10 ± 3.00		Treatiny	PFP	2.89 ± 2.09	0.440			
DKV angle (°)	22 26 + 9 42	$18.72 \pm 4.05 \pm$	7 86 + 10 05	16.65 ± 10.20		CAL	PFP	-4.63 ± 2.18	0.119			
mean ± SD	-23.30±8.45	-18.72±4.931	-7.80±10.05	-10.03 ± 10.30		C/H	Healthy	-15.49 ± 2.93	< 0.001*			
EC	1.27	0.82	0.21	2.22	0.001 E 10.27	10.27	CAI	4.63±2.18	0.119			
LO	1.57	0.82	0.51	2.52	0.001, F = 19.27	PFP	Healthy	-10.85 ± 2.50	< 0.001*			
CV04	14 62 + 4 80	12.04 + 2.05	7.02 + 4.44	11 22 + 5 17		Healthy	CAI	15.49±2.93	< 0.001*			
C V 70	14. 02 ± 4.80	12.04 ± 2.95	7.02 ± 4.44	11.22 ± 5.17		ricality	PFP	10.85 ± 2.50	< 0.001*			

Table 5. Asymmetry between limbs. The mean diffrence is siggnificant at the 0.05 level, *ASI* % asymmetry index between legs, *ROM* range of motion, *DKV*: dynamic knee valgus, *CV* coefficient of variation, ES effect size, *CAI* chronic ankle instability, *PFP* patellofemoral syndrome, *ROM* range of motion. *Indicates significant post-hoc comparison between CAI and PFP groups (p < 0.001). [†]Indicates significant difference on post-hoc testing compared to healthy group (p < 0.001).

Tasks		Ankle dorsiflexion ROM involved	DKV involved	LSD involved	Ankle dorsiflexion ROM contralateral	DKV contralateral	LSD contralateral
Ankle dorsiflexion ROM			-0.667*	-0.594*	0.737*	-0.674*	-0.550*
involved			< 0.001	< 0.001	< 0.001	< 0.001	< 0.001
DKV involved	ρ	-0.667*		0.796*	-0.530*	0.974*	0.707*
DK v liivoived	Р	< 0.001		< 0.001	< 0.001	< 0.001	< 0.001
LSD involved	ρ	-0.594*	0.796*		-0.494*	0.785*	0.776*
	Р	< 0.001	< 0.001		< 0.001	< 0.001	< 0.001
Ankle dorsiflexion ROM	ρ	0.737*	-0.530*	-0.494*		-0.590*	-0.614*
contralateral	Р	< 0.001	< 0.001	< 0.001		< 0.001	< 0.001
DKV controlotorol	ρ	-0.674*	0.974*	0.785*	-0.590*		0.719*
DK v contrataterat	Р	< 0.001	< 0.001	< 0.001	< 0.001		< 0.001
LSD controlotorol	ρ	-0.550*	0.707*	0.776*	-0.614*	0.737*	
	Р	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	

testing compared to neuriny group (p (0.001)

Table 6. Spearman's rho correlations between the different tasks. *DKV* dynamic knee valgus test, *LSD* lateral step-down test. *Correlation is significant at the 0.01 level (2-tailed).

compared to healthy subjects¹⁴. Similarly, in the study by Rabin et al.¹⁵ movement pattern was assessed visually during an LSD and rated as "good" or "moderate in PFP patients¹⁵. They demonstrated that ankle dorsiflexion ROM was more limited among participants with a moderate versus a good quality of movement¹⁵. Importantly, our ankle dorsiflexion ROM values exceeded the previously established minimum detectable difference of 4° and 6°, suggesting these differences may impact patient function^{14,41}.

The DKV angle in the single-leg squat test revealed that the involved limb angle was nearly 8° greater in the CAI group and approximately 9° greater in the PFP group compared to healthy controls (Table 3, Fig. 3b). The DKV angle for the contralateral limb was almost 5° greater in the CAI and 6° greater in the PFP versus the healthy control group (Table 3, Fig. 3b). The minimal detectable change (MDC) of the DKV angle ranged from approximately 3.5–10° for the single-leg squat⁴⁸. Accordingly, the mean difference observed in patients with CAI and PFP of over 8° between groups is clinically relevant and in agreement with the results of previous investigators who demonstrated greater DKV angle in females with than without PFP⁴⁹. Although many studies emphasize that DKV angle is a significant risk factor in patients with PFP⁷, to our best knowledge, only one study⁸ has confirmed that patients with CAI demonstrated increased DKV angle compared with coper and control subjects⁸. The paucity of studies investigating DKV angle in CAI offers future research opportunities. Additionally, these results collectively suggest the need to evaluate and improve ankle dorsiflexion ROM as well as DKV angle in patients with both PFP and CAI.

In terms of LSD results, our study revealed that CAI and PFP performed, on average, moderate quality movement (3.45), while healthy groups performed on average good quality movement (0.55) (Table 3, Fig. 3c). On the contralateral side, we observed moderate quality movement for CAI and PFP groups (1.40, 2.10 respectively) and good quality movement for the healthy group (0.25). There is a paucity of studies directly comparing the scores of movement quality between asymptomatic and musculoskeletal pain individuals. Ferreira et al. showed that women with PFP show muscle coordination and motor control alterations, which correlate with kinematic alterations during LSD⁵⁰. There is, however, limited research investigating LSD in women with CAI. In one study that was conducted, the authors reported individuals with poor movement quality had significantly less ankle dorsiflexion ROM compared to the good movement quality group¹⁴.

Our second hypothesis was only partially supported as contralateral ankle dorsiflexion ROM was 4° greater in the CAI (44°) compared to the PFP group (40°) (Table 3, Fig. 3a). In rehabilitation, inter-clinician changes of 4.6° and intra-clinician changes of 4.7° are needed for a change in dorsiflexion ROM to be considered outside of the measurement error of the WBLT⁵¹. Our data did not exceed the MDC; accordingly, the current study's findings should be treated with caution⁵¹. Moreover, in our study, the difference in ankle dorsiflexion ROM did not affect the significance of the difference between the DKV angle and LSD scores in both groups observed on the contralateral side, contrasting other previous evidence^{14,40}. For example, Grindstaff et al.¹⁴ showed that patients with CAI presenting a poor movement quality had, on average, 6° less ankle dorsiflexion ROM than participants with good movement quality and almost 3° less than the participants with moderate movement quality in involved and contralateral side. Also, Rabin et al. showed that decreased ankle dorsiflexion ROM impacted the movement quality of the LSD in healthy females⁴⁰. A possible explanation of the study discrepancy could be related to the inter-individual variability of neuromuscular control observed among patients with musculoskeletal pain⁵². Furthermore, the lack of studies directly comparing patients with PFP and CAI makes comparing our results and other evidence difficult. Thus, future studies should investigate the combined role of kinematic, muscle activation, strength, and movement quality deficits in individuals with CAI and PFP to determine whether dorsiflexion ROM and kinematic asymmetry exist in these populations impacting functional performance.

Our asymmetry analyses revealed a significant association between ankle dorsiflexion ROM and DKV with the LSD scores in female basketball athletes with PFP and CAI in support of our third hypothesis (Tables 3, 5). Our research agrees with Herrington, who demonstrated that healthy national league basketball players showed significant asymmetry in DKV angle during bilateral drop jump landings⁵³. Moreover, a study examining the performance of patients with PFP during the single-leg squat and Y-balance test showed results consistent with our findings³⁰. They showed that male military recruits with greater asymmetry on the Y-Balance test posterolateral direction, and frontal plane knee projection angle were at a greater risk of developing PFP³⁰. This asymmetry may suggest a neuromuscular imbalance between limbs. Side-to-side imbalances in neuromuscular strength, flexibility, and coordination represent significant predictors of increased injury risk⁵⁴. Asymmetry between two limbs is considered a risk factor in athletes²⁸. In the present study, comparing athletes with PFP and CAI, it seems that athletes with PFP experience more asymmetry and motor deficits on the contralateral side. Therefore, including the entire motor chain on both lower limbs is significant in evaluating and designing prevention and treatment programs. In addition, further prospective studies are needed to explore whether there is a cause-effect relationship between inter-limb asymmetry and injury development.

In addition, we aimed to determine which task has the greatest sensitivity to identify asymmetries based on ASI calculation. ASI varied among ankle dorsiflexion ROM, DKV angle, and LSD tasks and indicated that the percentage of asymmetry could differ depending on the task performed. Accordingly, this finding indicates that obtaining measurements on various independent tasks is important to assess ASI. The largest ASI detected in our study was found in DKV for CAI and PFP. This ASI is higher than the 10–15% threshold describing the potential risk of injury among participants as reported in the literature²⁸. Therefore, the DKV is likely a critical task for detecting ILA between legs. In addition, prospective research is needed to determine if there is a direct correlation between DKV angle asymmetry and injury rate over time. Moreover, it is challenging to establish an ASI threshold for typical values on the other tasks examined in this research because of the scarcity of research on this topic.

The fourth hypothesis's results demonstrated a moderate to good relationship between ankle dorsiflexion ROM and LSD scores and a good to strong relationship between DKV angle and LSD scores. These findings agree with similar studies investigating ankle dorsiflexion ROM and LSD scores in CAI and PFP^{14,15,21,40}. During the LSD, maximal ankle dorsiflexion ROM is required to allow the heel to remain in contact with the ground¹⁴. A limitation in ankle dorsiflexion ROM or the proximal lower limb joints may require compensation from other joints¹⁶. Decreased ankle dorsiflexion ROM will limit the forward progression of the tibia over the talus, resulting in compensatory subtalar pronation, which, in turn, may displace the knee medially⁵⁵. Previous studies suggest that limited ankle dorsiflexion ROM may be associated with a greater medial knee displacement during various functional activities¹⁶⁻¹⁸; such compensation may increase DKV angle. Less research has been conducted to investigate the influence of limited ankle dorsiflexion ROM on a frontal and transverse plane lower limb alignment. Bell et al. found that ankle ROM and strength rather than hip ROM and strength were associated with DKV angle during a controlled squat⁵⁶. In the above study, the DKV angle was corrected by adding a heel lift, confirming the role of limited ankle dorsiflexion ROM in dynamic knee valgus during a loading task⁵⁶. Bell-Jenje et al.²¹ demonstrated that participants with restricted dorsiflexion ROM had greater dynamic knee valgus during the LSD^{21} . Also, they showed that the association between an increase in hip adduction ROM and decreased dorsiflexion ROM during a functional loading task such as the LSD test emphasizes the importance of the kinetic chain in lower limb alignment²¹. In the closed kinetic chain, the relative angle, magnitude, and direction of one motion-dependent body segment affect another segment, leading to differences in how these segments interact^{14,21}. This finding is significant in a clinical context where limited dorsiflexion ROM is associated with CAI¹⁴, PFP¹⁵, and dorsiflexion ROM may, therefore, play a key role in predicting injuries²¹, but further longitudinal research is needed. Additionally, these results collectively suggest the need to evaluate and improve ankle dorsiflexion ROM to improve movement quality in patients with both PFP and CAI. There are few studies investigating this relationship in CAI. So, future research is needed to investigate a relationship between CAI and other lower limb injuries during functional tasks (e.g. jump landing/cutting test, single leg countermovement jumps); possible use of posture software for the movement analysis is further suggested. Also, a future prospective study is needed to examine whether increased DKV angle with reduced dorsiflexion ROM may contribute to a greater risk of knee injury in patients with CAI.

Limitations. This study presents some limitations that should be acknowledged. Firstly, we cannot obtain causative conclusions from our study by adopting a cross-sectional design⁵⁷. Accordingly, prospective studies should consider the cause-effect relationship between inter-limb asymmetry and injury development. Secondly, we have recruited only a sample of female non-professional basketball players with CAI and PFP. Thus, our findings could not be generalized to male participants of different ages, professional athletes, players of other contact sports or in the presence of different clinical conditions (e.g. ACL reconstruction). Thirdly, the small effect size and power used in sample size estimation is another limitation that should be considered in future research. Fourthly, this study did not examine hormone profiles and status or inquire about contraception usage, which can have important implications for physiological function in female participants⁵⁸. Lastly, we assessed the quality of movement only using the LSD and adopted a 2D analysis of limb motion. Therefore, future studies should consider investigating a motor task similar to basketball-specific movement techniques (e.g. jump landing/cutting test, single-leg countermovement jumps) and also integrating a 3D analysis (e.g. hip, knee and ankle joints simultaneously) combined with a real-time electromyography data acquisition to gain further insight into the importance of ankle dorsiflexion ROM and DKV angle during sport-specific tasks.

Conclusions

Female basketball players with PFP and CAI have impaired ankle dorsiflexion ROM, greater DKV angles, poorer scores in the LSD, and greater ILAs compared to healthy controls. Ankle dorsiflexion ROM and DKV were associated with LSD. These findings inform clinicians and researchers of the need to evaluate ROM and function throughout the kinetic chain and in both lower limbs in injured athletes tailoring specific rehabilitation programs.

Data availability

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Received: 6 February 2023; Accepted: 27 July 2023 Published online: 31 July 2023

References

- 1. Finch, C., Owen, N. & Price, R. Current injury or disability as a barrier to being more physically active. *Med. Sci. Sports Exerc.* 33, 778–782 (2001).
- Waterman, B. R., Owens, B. D., Davey, S., Zacchilli, M. A. & Belmont, P. J. Jr. The epidemiology of ankle sprains in the United States. *JBJS* 92, 2279–2284 (2010).
- Wijnhoven, H. A., de Vet, H. C. & Picavet, H. S. J. Explaining sex differences in chronic musculoskeletal pain in a general population. *Pain* 124, 158–166 (2006).
- Deitch, J. R., Starkey, C., Walters, S. L. & Moseley, J. B. Injury risk in professional basketball players: A comparison of Women's National Basketball Association and National Basketball Association athletes. Am. J. Sports Med. 34, 1077–1083 (2006).
- Powers, C. M., Witvrouw, E., Davis, I. S. & Crossley, K. M. Evidence-based framework for a pathomechanical model of patellofemoral pain: 2017 patellofemoral pain consensus statement from the 4th International Patellofemoral Pain Research Retreat, Manchester, UK: Part 3. Br. J. Sports Med. 51, 1713–1723 (2017).
- 6. Delahunt, E. *et al.* Inclusion criteria when investigating insufficiencies in chronic ankle instability. *Med. Sci. Sports Exerc.* **42**, 2106–2121 (2010).
- Emamvirdi, M., Letafatkar, A. & Khaleghi Tazji, M. The effect of valgus control instruction exercises on pain, strength, and functionality in active females with patellofemoral pain syndrome. Sports Health 11, 223–237 (2019).
- 8. Han, S. *et al.* Prelanding movement strategies among chronic ankle instability, coper, and control subjects. *Sports Biomech.* **21**, 391–407 (2022).
- 9. Terada, M., Pietrosimone, B. & Gribble, P. A. Individuals with chronic ankle instability exhibit altered landing knee kinematics: Potential link with the mechanism of loading for the anterior cruciate ligament. *Clin. Biomech.* **29**, 1125–1130 (2014).
- 10. Coelho, B. A. L. *et al.* Immediate effect of a single ankle mobilization session in range of motion, pain and dynamic knee valgus of women with patellofemoral pain and restricted dorsiflexion range of motion: A randomized controlled trial. *Phys. Ther. Sport* **47**, e3–e4 (2021).
- 11. Lima, Y. L., de Paula Lima, P. O., Bezerra, M. A., de Oliveira, R. R. & Almeida, G. P. L. WITHDRAWN: The association of ankle dorsiflexion range of motion and dynamic knee valgus: A systematic review with meta-analysis. *Phys. Ther. Sport* **45**, e4 (2020).
- 12. Arora, N. K., Sharma, S., Sharma, S. & Arora, I. K. Physical modalities with eccentric exercise are no better than eccentric exercise alone in the treatment of chronic *Achilles Tendinopathy*: A systematic review and meta-analysis. *Foot* **53**, 101927 (2022).
- 13. Rabin, A., Kozol, Z. & Finestone, A. S. Limited ankle dorsiflexion increases the risk for mid-portion *Achilles tendinopathy* in infantry recruits: A prospective cohort study. *J. Foot Ankle Res.* 7, 1–7 (2014).
- Grindstaff, T. L., Dolan, N. & Morton, S. K. Ankle dorsiflexion range of motion influences lateral step down test scores in individuals with chronic ankle instability. *Phys. Ther. Sport* 23, 75–81 (2017).
- 15. Rabin, A. et al. Factors associated with visually assessed quality of movement during a lateral step-down test among individuals with patellofemoral pain. J. Orthop. Sports Phys. Ther. 44, 937–946 (2014).
- Macrum, E., Bell, D. R., Boling, M., Lewek, M. & Padua, D. Effect of limiting ankle-dorsiflexion range of motion on lower extremity kinematics and muscle-activation patterns during a squat. J. Sport Rehabil. 21, 144–150 (2012).
- 17. Sigward, S. M., Ota, S. & Powers, C. M. Predictors of frontal plane knee excursion during a drop land in young female soccer players. J. Orthop. Sports Phys. Ther. 38, 661–667 (2008).
- Lima, Y. L. et al. The association of ankle dorsiflexion and dynamic knee valgus: A systematic review and meta-analysis. Phys. Ther. Sport 29, 61–69 (2018).
- Mason-Mackay, A., Whatman, C. & Reid, D. The effect of reduced ankle dorsiflexion on lower extremity mechanics during landing: A systematic review. J. Sci. Med. Sport 20, 451–458 (2017).

- Dill, K. E., Begalle, R. L., Frank, B. S., Zinder, S. M. & Padua, D. A. Altered knee and ankle kinematics during squatting in those with limited weight-bearing–lunge ankle-dorsiflexion range of motion. J. Athl. Train. 49, 723–732 (2014).
- 21. Bell-Jenje, T. *et al.* The association between loss of ankle dorsiflexion range of movement, and hip adduction and internal rotation during a step down test. *Man. Ther.* **21**, 256–261 (2016).
- 22. Piva, S. R., Goodnite, E. A. & Childs, J. D. Strength around the hip and flexibility of soft tissues in individuals with and without patellofemoral pain syndrome. J. Orthop. Sports Phys. Ther. 35, 793–801 (2005).
- 23. Doherty, C. et al. Single-leg drop landing movement strategies in participants with chronic ankle instability compared with lateral ankle sprain 'copers'. Knee Surg. Sports Traumatol. Arthrosc. 24, 1049–1059 (2016).
- E Silva, R. L., Pinheiro, Y. T., de Almeida Lins, C. A., de Oliveira, R. R. & Silva, R. S. Assessment of quality of movement during a lateral step-down test: Narrative review. J. Bodyw. Mov. Ther. 23, 835–843 (2019).
- 25. Hewit, J., Cronin, J. & Hume, P. Multidirectional leg asymmetry assessment in sport. Strength Cond. J. 34, 82-86 (2012).
- 26. Paterno, M. V. *et al.* Biomechanical measures during landing and postural stability predict second anterior cruciate ligament injury after anterior cruciate ligament reconstruction and return to sport. *Am. J. Sports Med.* **38**, 1968–1978 (2010).
- Ford, K. R., Myer, G. D. & Hewett, T. E. Valgus knee motion during landing in high school female and male basketball players. Med. Sci. Sports Exerc. 35, 1745–1750 (2003).
- Fort-Vanmeerhaeghe, A., Montalvo, A. M., Sitja-Rabert, M., Kiefer, A. W. & Myer, G. D. Neuromuscular asymmetries in the lower limbs of elite female youth basketball players and the application of the skillful limb model of comparison. *Phys. Ther. Sport* 16, 317–323 (2015).
- Payne, K., Payne, J. & Larkin, T. A. Patellofemoral pain syndrome and pain severity is associated with asymmetry of gluteus medius muscle activation measured via ultrasound. Am. J. Phys. Med. Rehabil. 99, 595–601 (2020).
- Nakagawa, T. H., Dos Santos, A. F., Lessi, G. C., Petersen, R. S. & Silva, R. S. Y-balance test asymmetry and frontal plane knee projection angle during single-leg squat as predictors of patellofemoral pain in male military recruits. *Phys. Ther. Sport* 44, 121–127 (2020).
- 31. Plastaras, C. et al. Is hip abduction strength asymmetry present in female runners in the early stages of patellofemoral pain syndrome?. Am. J. Sports Med. 44, 105–112 (2016).
- 32. Plisky, P. J., Raul, M. J., Kaminski, T. W. & Underwood, F. B. Star excursion balance test as a predictor of lower extremity injury in high school basketball players. J. Orthop. Sports Phys. Ther. 36, 911–919 (2006).
- Tajdini, H., Mantashloo, Z., Thomas, A. C., Letafatkar, A. & Rossettini, G. Inter-limb asymmetry of kinetic and electromyographic during walking in patients with chronic ankle instability. Sci. Rep. 12, 1–10 (2022).
- 34. Piccoli, A. *et al.* Effect of attentional focus instructions on motor learning and performance of patients with central nervous system and musculoskeletal disorders: A systematic review. *J. Funct. Morphol. Kinesiol.* **3**, 40 (2018).
- 35. Attenborough, A. S. et al. Chronic ankle instability in sporting populations. Sports Med. 44, 1545–1556 (2014).
- Mirshahi, M., Halabchi, F., Golbakhsh, M. & Saadat, S. Reliability and recalibration of the Persian version of Cumberland Ankle Instability Tool cut-off score in athletes with functional ankle instability. Adv. J. Emerg. Med. 3, e26 (2019).
- Mazaheri, M. et al. Reliability and validity of the Persian version of foot and ankle ability measure (FAAM) to measure functional limitations in patients with foot and ankle disorders. Osteoarthr. Cart. 18, 755–759 (2010).
- Da Cunha, R. A. et al. Translation, cross-cultural adaptation, and clinimetric testing of instruments used to assess patients with patellofemoral pain syndrome in the Brazilian population. J. Orthop. Sports Phys. Ther. 43, 332–339 (2013).
- Doberstein, S. T., Romeyn, R. L. & Reineke, D. M. The diagnostic value of the Clarke sign in assessing chondromalacia patella. J. Athl. Train. 43, 190–196 (2008).
- Rabin, A. & Kozol, Z. Measures of range of motion and strength among healthy women with differing quality of lower extremity movement during the lateral step-down test. J. Orthop. Sports Phys. Ther. 40, 792–800 (2010).
- 41. Konor, M. M., Morton, S., Eckerson, J. M. & Grindstaff, T. L. Reliability of three measures of ankle dorsiflexion range of motion. Int. J. Sports Phys. Ther. 7, 279 (2012).
- Munro, A., Herrington, L. & Carolan, M. Reliability of 2-dimensional video assessment of frontal-plane dynamic knee valgus during common athletic screening tasks. J. Sport Rehabil. 21, 7–11 (2012).
- Piva, S. R. et al. Reliability of measures of impairments associated with patellofemoral pain syndrome. BMC Musculoskelet. Disord. 7, 1–13 (2006).
- Carpes, F. P., Mota, C. B. & Faria, I. E. On the bilateral asymmetry during running and cycling–A review considering leg preference. *Phys. Ther. Sport* 11, 136–142 (2010).
- Exell, T., Irwin, G., Gittoes, M. & Kerwin, D. Strength and performance asymmetry during maximal velocity sprint running. Scand. J. Med. Sci. Sports 27, 1273–1282 (2017).
- 46. Portney, L. G. & Watkins, M. P. Foundations of Clinical Research: Applications to Practice Vol. 892 (Pearson/Prentice Hall Upper Saddle River, 2009).
- 47. Hopkins, W. G. A scale of magnitudes for effect statistics. N. View Stat. 502, 411 (2002).
- Werner, D. M., Di Stasi, S., Lewis, C. L. & Barrios, J. A. Test-retest reliability and minimum detectable change for various frontal plane projection angles during dynamic tasks. *Phys. Ther. Sport* 40, 169–176 (2019).
- Willson, J. D. & Davis, I. S. Utility of the frontal plane projection angle in females with patellofemoral pain. J. Orthop. Sports Phys. Ther. 38, 606–615 (2008).
- 50. Ferreira, C. L. *et al.* Women with patellofemoral pain show altered motor coordination during lateral step down. *J. Biomech.* **110**, 109981 (2020).
- Powden, C. J., Hoch, J. M. & Hoch, M. C. Reliability and minimal detectable change of the weight-bearing lunge test: A systematic review. *Man. Ther.* 20, 524–532 (2015).
- 52. Pincus, T., Smeets, R. J., Simmonds, M. J. & Sullivan, M. J. The fear avoidance model disentangled: Improving the clinical utility of the fear avoidance model. *Clin. J. Pain* 26, 739–746 (2010).
- Herrington, L. Knee valgus angle during landing tasks in female volleyball and basketball players. J. Strength Cond. Res. 25, 262–266 (2011).
- Knapik, J. J., Bauman, C. L., Jones, B. H., Harris, J. M. & Vaughan, L. Preseason strength and flexibility imbalances associated with athletic injuries in female collegiate athletes. Am. J. Sports Med. 19, 76–81 (1991).
- Kaufman, K. R., Brodine, S. K., Shaffer, R. A., Johnson, C. W. & Cullison, T. R. The effect of foot structure and range of motion on musculoskeletal overuse injuries. *Am. J. Sports Med.* 27, 585–593 (1999).
- Bell, D. R., Padua, D. A. & Clark, M. A. Muscle strength and flexibility characteristics of people displaying excessive medial knee displacement. Arch. Phys. Med. Rehabil. 89, 1323–1328 (2008).
- Von Elm, E. et al. The strengthening the reporting of observational studies in epidemiology (STROBE) statement: Guidelines for reporting observational studies. Ann. Intern. Med. 147, 573–577 (2007).
- Elliott-Sale, K. J. et al. Methodological considerations for studies in sport and exercise science with women as participants: A working guide for standards of practice for research on women. Sports Med. 51, 843–861 (2021).

Acknowledgements

The authors would like to thank all the athletes for participation in this study.

Author contributions

M.E., A.L., M.H., G.R., N.S., TH.A. and TH.D. conceptualized the research; M.E., G.R., N.S., TH.D. and A.L. designed the methodology; M.E. conducted the research; M.E. and M.H. performed the statistical analysis; M.E. wrote the main manuscript; M.H., A.L., G.R., N.S., TH.A., TH.D. reviewed and edited the draft manuscript; All authors have read and agreed to the published version of the manuscript.

Funding

This research received no external funding.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to M.H.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

© The Author(s) 2023