

Please cite the Published Version

Alruhaimi, RS, Mostafa-Hedeab, G, Abduh, MS, Bin-Ammar, A, Hassanein, EHM, Kamel, EM and Mahmoud, AM (2023) A flavonoid-rich fraction of *Euphorbia peplus* attenuates hyperglycemia, insulin resistance, and oxidative stress in a type 2 diabetes rat model. *Frontiers in Pharmacology*, 14. 1204641 ISSN 1663-9812

DOI: <https://doi.org/10.3389/fphar.2023.1204641>

Publisher: Frontiers Media S.A.

Version: Published Version

Downloaded from: <https://e-space.mmu.ac.uk/633076/>

Usage rights:  [Creative Commons: Attribution 4.0](https://creativecommons.org/licenses/by/4.0/)

Additional Information: This is an Open Access article published in *Frontiers in Pharmacology*, by Frontiers Media S.A.

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)



OPEN ACCESS

EDITED BY

Chika Ifeanyi Chukwuma,
Central University of Technology, South
Africa

REVIEWED BY

Jude Akinyelu,
Federal University Oye-Ekiti, Nigeria
Londiwe Mbatha,
Durban University of Technology, South
Africa

*CORRESPONDENCE

Ayman M. Mahmoud,
✉ ayman.mahmoud@science.bsu.edu.eg,
✉ a.mahmoud@mmu.ac.uk

RECEIVED 12 April 2023

ACCEPTED 01 June 2023

PUBLISHED 16 June 2023

CITATION

Alruhaimi RS, Mostafa-Hedeab G,
Abduh MS, Bin-Amman A,
Hassanein EHM, Kamel EM and
Mahmoud AM (2023), A flavonoid-rich
fraction of *Euphorbia peplus* attenuates
hyperglycemia, insulin resistance, and
oxidative stress in a type 2 diabetes
rat model.

Front. Pharmacol. 14:1204641.

doi: 10.3389/fphar.2023.1204641

COPYRIGHT

© 2023 Alruhaimi, Mostafa-Hedeab,
Abduh, Bin-Amman, Hassanein, Kamel
and Mahmoud. This is an open-access
article distributed under the terms of the
[Creative Commons Attribution License
\(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or
reproduction in other forums is
permitted, provided the original author(s)
and the copyright owner(s) are credited
and that the original publication in this
journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

A flavonoid-rich fraction of *Euphorbia peplus* attenuates hyperglycemia, insulin resistance, and oxidative stress in a type 2 diabetes rat model

Reem S. Alruhaimi¹, Gomaa Mostafa-Hedeab^{2,3},
Maisa Siddiq Abduh^{4,5}, Albandari Bin-Amman⁶,
Emad H. M. Hassanein⁷, Emadeldin M. Kamel⁸ and
Ayman M. Mahmoud^{9,10*}

¹Department of Biology, College of Science, Princess Nourah Bint Abdulrahman University, Riyadh, Saudi Arabia, ²Pharmacology Department, Medical College, Jouf University, Sakaka, Saudi Arabia, ³Pharmacology Department, Faculty of Medicine, Beni-Suef University, Beni-Suef, Egypt, ⁴Immune Responses in Different Diseases Research Group, Department of Medical Laboratory Sciences, Faculty of Applied Medical Sciences, King Abdulaziz University, Jeddah, Saudi Arabia, ⁵Center of Excellence in Genomic Medicine Research, King Abdulaziz University, Jeddah, Saudi Arabia, ⁶Department of Clinical Nutrition, College of Applied Medical Sciences, University of Hail, Hail, Saudi Arabia, ⁷Department of Pharmacology and Toxicology, Faculty of Pharmacy, Al-Azhar University, Assiut, Egypt, ⁸Chemistry Department, Faculty of Science, Beni-Suef University, Beni-Suef, Egypt, ⁹Department of Life Sciences, Faculty of Science and Engineering, Manchester Metropolitan University, Manchester, United Kingdom, ¹⁰Physiology Division, Zoology Department, Faculty of Science, Beni-Suef University, Beni-Suef, Egypt

Background: Type 2 diabetes (T2D) is a metabolic disorder characterized by insulin resistance (IR) and hyperglycemia. Plants are valuable sources of therapeutic agents for the management of T2D. *Euphorbia peplus* has been widely used as a traditional medicine for the treatment of various diseases, but its beneficial role in T2D has not been fully explored.

Methods: The anti-diabetic efficacy of *E. peplus* extract (EPE) was studied using rats with T2D induced by high-fat diet (HFD) and streptozotocin (STZ). The diabetic rats received 100, 200, and 400 mg/kg EPE for 4 weeks.

Results: Phytochemical fractionation of the aerial parts of *E. peplus* led to the isolation of seven known flavonoids. Rats with T2D exhibited IR, impaired glucose tolerance, decreased liver hexokinase and glycogen, and upregulated glycogen phosphorylase, glucose-6-phosphatase (G-6-Pase), and fructose-1,6-bisphosphatase (F-1,6-BPase). Treatment with 100, 200, and 400 mg/kg EPE for 4 weeks ameliorated hyperglycemia, IR, liver glycogen, and the activities of carbohydrate-metabolizing enzymes. EPE attenuated dyslipidemia, serum transaminases, tumor necrosis factor (TNF)- α , interleukin (IL)-1 β and liver lipid accumulation, nuclear factor (NF)- κ B p65, and lipid peroxidation, nitric oxide and enhanced antioxidants. All EPE doses upregulated serum adiponectin and liver peroxisome proliferator-activated receptor γ (PPAR γ) in HFD/STZ-induced rats. The isolated flavonoids showed *in silico* binding affinity toward hexokinase, NF- κ B, and PPAR γ .

Conclusion: *E. peplus* is rich in flavonoids, and its extract ameliorated IR, hyperglycemia, dyslipidemia, inflammation and redox imbalance, and upregulated adiponectin and PPAR γ in rats with T2D.

KEYWORDS

Euphorbia, diabetes, insulin resistance, oxidative stress, inflammation

1 Introduction

Diabetes mellitus (DM) is a common metabolic disorder associated with several complications, including nephropathy, neuropathy, and cardiomyopathy. This disorder includes type 1 (T1DM) and type 2 (T2DM) forms of the disease where T1DM is characterized by insulin insufficiency, whereas insulin resistance (IR) is the characteristic feature of T2DM. Both insulin deficiency and IR lead to the accumulation of glucose in the blood (hyperglycemia) (American Diabetes Association, 2021). DM is a fast-increasing disease worldwide, and the number of patients with diabetes is expected to reach 700 million by 2045 (Saeedi et al., 2019). T2DM is the most common form of the disease characterized by hyperglycemia and IR (Kahn et al., 2014). IR increases the risk of hypertension, dyslipidemia, and atherosclerosis (Guzik and Cosentino, 2018). Oxidative stress (OS) and inflammation mediated by excess reactive oxygen species (ROS) and inflammatory mediators produced under hyperglycemic conditions are implicated in the pathophysiology of DM and its complications (Mahmoud et al., 2012). Excess ROS can damage cellular macromolecules and work in concert with inflammatory mediators to provoke cell death. ROS and inflammatory mediators impair insulin signaling by provoking β -cell death, alter peripheral glucose uptake, and increase gluconeogenesis (Jheng et al., 2012). Therefore, mitigation of OS and inflammation could be beneficial to prevent IR and hyperglycemia in T2DM.

Plants of the genus *Euphorbia* include numerous known species with chemical diversity and multiple biological and commercial uses (Shi et al., 2008). The latex of plants of the family *Euphorbiaceae* was acknowledged for its various phytoconstituents that possess both commercial and pharmacological importance such as triterpene alcohols (Giner and Schroeder, 2015). Because of its toxic nature and unpleasantness, the latex protects the plants against the attack of animals (Al-Sultan and Hussein, 2006). Steroids, flavonoids, sesquiterpenoids, glycerols, and cerebrosides are among the phytoconstituents reported in plants of the genus *Euphorbia* (Shi et al., 2008; Kamel et al., 2022). With this rich content, *Euphorbia* plants found their way to be employed in folkloric medicine to treat migraine, intestinal parasites, gonorrhoea, and skin disorders (Singla and Kamla, 1990), and studies have reported their wound-healing potential (Pattanaik et al., 2014; Ahmed et al., 2016). Recent work from our laboratory revealed the inhibitory activity of *E. peplus* on xanthine oxidase (XO) and hyperuricemia in rats (Kamel et al., 2022). Other studies showed the possible beneficial effects of *E. royleana* stem extract (Zafar et al., 2021) and *E. hirta* flower extract (Kumar et al., 2010) in rats with streptozotocin (STZ)- and alloxan-induced diabetes, respectively. These studies revealed the ability of *E. royleana* and *E. hirta* to ameliorate hyperglycemia and oxidative damage. Another recent study highlighted the anti-hyperglycemic effect of *E. helioscopia* methanolic extract in sucrose-supplemented rats (Mustafa et al., 2022). Owing to the promising therapeutic value of plants of this genus, this study explored the phytochemical constituents and the effect of *E. peplus* extract (EPE) on hyperglycemia, IR, OS, and inflammation in rats with T2D induced by high-fat diet (HFD) and STZ.

2 Materials and methods

2.1 Phytochemical investigation

2.1.1 General

Proton nuclear magnetic resonance (^1H NMR) and ^{13}C NMR (500 MHz and 125 MHz, respectively) spectra were recorded on the Bruker AV-500 spectrometer using TMS as an internal standard. The optical rotation of isolated flavonoids was obtained using a Rudolph Autopol III polarimeter. Ultraviolet (UV) spectral data were measured using the Shimadzu UV-vis 160i spectrophotometer, and the HREIMS and EIMS spectral data were recorded using the Finnigan MAT TSQ 700 mass spectrometer. Infrared spectral data were obtained through KBr pellets on the Shimadzu FTIR-8400 instrument.

2.1.2 Plant collection, extraction, and isolation

The plant was collected from Beni Suef Governorate in March 2021 and identified by a taxonomist and a voucher specimen (EP-038021-2) was stored. The aerial parts (2.75 kg) were extracted four times using 70% acetone followed by the removal of the solvent under reduced pressure, resulting in 904 g of extract. Thereafter, the extract was dissolved in water and successively partitioned using chloroform, ethyl acetate (EA), and *n*-butanol (3L x 2, each). The EA fraction (69.7 g) was subjected to chromatographic fractionation over a silica gel column (120 × 4 cm, 1.1 kg) and eluted with dichloromethane (DCM)/acetone mixture of increasing polarity. To track the movement of the bands along the column and to regulate the collection of fractions, a UV lamp was employed. A total of 22 fractions were collected and combined into seven main subfractions (F1–F7) according to their similar thin-layer chromatography (TLC) profiles. Subfraction F3 was chromatographed over silica gel using chloroform–EA of gradient elution to afford nine subfractions (F3.1–F3.9). Subfractions F3.3–F3.7 were combined and applied to the Sephadex LH-20 column eluted with methanol (MeOH):water (50:50→100:0) to give seven TLC-monitored subfractions (E1–E7). Subfractions (E2–E5) were combined and purified over a Sephadex LH-20 column eluted with MeOH to yield the purified compounds 2 (22 mg), 3 (17 mg), and 4 (14 mg). Subfraction F4 was fractionated over a polyamide 6S column eluted with the MeOH–water solvent mixture of increasing polarity to afford eleven subfractions (F4.1–F4.11). Subfraction F4.6 was purified over the Sephadex LH-20 column eluted with MeOH to yield purified compound 1 (23 mg). Subfractions F4.8–F4.10 were combined and re-chromatographed using the Sephadex LH-20 column eluted with MeOH to give compound 5 (19 mg). Subfraction F5 was partitioned by means of the Sephadex LH-20 column using MeOH–water (2:8, 3:7→10:0) to afford six subfractions (F5.1–F5.6). Compound 6 (23 mg) was obtained from the chromatographic fractionation of F5.3–F5.5 over two consecutive Sephadex LH-20 columns using 30% MeOH as an eluent. Subfraction F7 was subjected to silica gel column chromatography eluted with the solvent system chloroform–MeOH–water (lower layer, 28:9:6 and 6:3:1) to yield

five subfractions (F7.1–F7.5). Compound 7 (24 mg) was obtained from the recombination and purification of subfractions F7.1–F7.3 over Sephadex LH-20 and eluted with MeOH (Supplementary Figure S1).

2.2 *In vitro* radical-scavenging activity

The RSA activity of EPE was measured using 2,2-diphenyl-1-picrylhydrazyl (DPPH) and 2,2-azinobis (3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) assays following the methods of Brand-Williams et al. (1995) and Re et al. (1999), respectively, using ascorbic acid as a standard.

2.3 Experimental animals and treatments

Male Wistar rats weighing 170–190 g were included in this investigation. The rats were maintained under standard conditions of temperature ($23^{\circ}\text{C} \pm 1^{\circ}\text{C}$) and humidity (50%–60%) on a 12-h light/dark cycle with free access to food and water. The animal study protocol was approved by the Research Ethics Committee of Al-Azhar University (ZA-AS/PH/18/C/2023). The rats received a normal diet and a single intraperitoneal (i.p.) injection of freshly prepared citrate buffer (pH 4.5) to serve as a control. Other rats were fed a HFD (58% fat, 17% carbohydrate, and 25% protein) for 28 days and received a single i.p. dose of STZ (35 mg/kg; Sigma, United States) dissolved in freshly prepared citrate buffer (pH 4.5) to induce T2D (Germoush et al., 2019). After 7 days, T2D was confirmed by measuring blood glucose (BG) for 2 h after supplementing the overnight fasted rats with 3 g/kg glucose orally. The rats exhibited BG higher than 250 mg/kg were included in the investigation.

To investigate the antidiabetic effects of the EA fraction of *E. pepplus* extract (EPE; dissolved in 0.5% carboxymethyl cellulose (CMC) as a vehicle), 30 diabetic and 12 normal rats were allocated into seven groups ($n = 6$) as follows:

- Group I (control): received 0.5% CMC.
- Group II (EPE): received 400 mg/kg EPE.
- Group III (diabetic): received 0.5% CMC.
- Group IV (diabetic + 100 mg/kg EPE): received 100 mg/kg EPE.
- Group V (diabetic + 200 mg/kg EPE): received 200 mg/kg EPE.
- Group VI (diabetic + 400 mg/kg EPE): received 400 mg/kg EPE.
- Group VII (diabetic + PIO): received 10 mg/kg of the antidiabetic pioglitazone (PIO) (Abd El-Twab et al., 2016).

EPE, 0.5% CMC, and PIO were supplemented orally for 4 weeks. A day before the end of the experiment, the rats were fasted overnight and then supplemented with 3 g/kg glucose solution, and the blood was collected from the tail vein over 2 h for the determination of BG using a Spinreact (Spain) kit (Trinder, 1969). At the end of treatments, the animals were euthanized under ketamine anesthesia (100 mg/kg i.p.), and blood and liver samples were collected. Serum was separated following centrifugation of blood, and samples from the liver were homogenized in Tris-HCl buffer (pH = 7.4). Other samples

were fixed in 10% neutral buffered formalin (NBF) or stored at -80°C .

2.4 Biochemical assays

Serum insulin, transaminases (ALT and AST), adiponectin and cytokines (TNF- α and IL-1 β) were determined using kits from RayBiotech (United States), Spinreact (Spain), and R&D Systems (United States), respectively. NF- κ B p65 in liver homogenate was determined using the kit from R&D Systems (United States). All assays were performed according to the manufacturers' instructions.

The homeostasis model assessment of IR (HOMA-IR) was calculated as previously described by Haffner (2000) using the following equation:

$$\text{HOMA-IR} = \frac{\text{Fasting insulin } \left(\frac{\mu\text{U}}{\text{ml}}\right) \times \text{Fasting glucose } \left(\frac{\text{mmol}}{\text{L}}\right)}{22.5}$$

Liver glycogen content was determined as previously described (Seifter and Dayton, 1950). Liver homogenate was centrifuged, and the clear supernatant was used to assess the activities of hexokinase (Brandstrup et al., 1957), G-6-Pase (Koide and Oda, 1959), F-1,6-BPase (Freedland and Harper, 1959), and glycogen phosphorylase (Stalmans and Hers, 1975). Malondialdehyde (MDA) (Ohkawa et al., 1979), nitric oxide (NO) (Green et al., 1982), reduced glutathione (GSH) (Beutler et al., 1963), and the activities of superoxide dismutase (SOD) (Marklund and Marklund, 1974), catalase (CAT) (Aebi, 1984), and glutathione peroxidase (GPx) (Flohé and Günzler, 1984) were determined in the supernatant of the homogenized liver. Liver triglycerides (TGs) and cholesterol were assayed using Spinreact (Spain) kits after extracting the lipids using chloroform/MeOH mixture (2:1, v/v) as described by Folch et al. (1957). Serum TG, total cholesterol (TC), and high-density lipoprotein (HDL)-C were assayed using kits from Spinreact (Spain). Low-density lipoprotein (LDL)-C, very low-density lipoprotein (vLDL)-C, and atherogenic index of plasma (AIP) were calculated according to the following equations:

$$\text{vLDL.C} = \text{TG}/5,$$

$$\text{LDL.C} = \text{TC} - (\text{HDL.C} + \text{vLDL.C}),$$

$$\text{AIP} = \text{Log} \left(\frac{\text{TG}}{\text{HDL}} \right).$$

2.5 Histopathological study

Samples from the liver fixed in 10% NBF for 24 h were dehydrated, cleared, and embedded in paraffin wax. Sections of 5 μm thickness were cut for routine staining with hematoxylin and eosin (H&E) (Bancroft and Gamble, 2008) and examined under a light microscope.

2.6 Quantitative real-time polymerase chain reaction

To determine the changes in PPAR γ mRNA, RNA was isolated from the frozen liver using TRIzol and quantified, and samples with




FIGURE 1
(A) Chemical structure of the isolated compounds. (B, C) DPPH and ABTS radical-scavenging activities of EPE. Data are mean \pm SD (N = 3).

OD_{260/280} \geq 1.8 were reverse-transcribed into cDNA using a cDNA synthesis kit (Thermo Scientific, United States). cDNA amplification was carried out using SYBR Green Master Mix (Thermo Scientific, United States), and the primers used in the qRT-PCR experiment are listed in [Supplementary Table S1](#). The Ct values were analyzed by using the $2^{-\Delta\Delta Ct}$ method (Livak and Schmittgen, 2001).

2.7 *In silico* molecular docking

The binding of *E. peplus* compounds with hexokinase II (PDB ID: 2NZT), NF- κ B-DNA complex (PDB ID 1LE9), and PPAR γ (PDB ID: 2PRG) was investigated as previously reported ([Supplementary Material](#)) (Sami et al., 2022; Abduh et al., 2023).

2.8 Statistical analysis

The obtained results are presented as mean \pm standard deviation (SD), and all statistical comparisons were made using one-way ANOVA followed by *post hoc* Tukey's test on GraphPad Prism 8 software. *p*-values < 0.05 were considered statistically significant.

3 Results

3.1 Phytochemical investigation and *in vitro* RSA

The analysis of the EA fraction of *E. peplus* led to the isolation of seven known flavonoids. Structures of isolated




FIGURE 2 EPE ameliorated glucose intolerance (A and B), serum insulin (C), and HOMA-IR (D) in diabetic rats. Data are mean ± SD (n = 6). *p < 0.05 and ***p < 0.001 vs. control. #p < 0.05, ##p < 0.01, and ###p < 0.001 vs. diabetic.

compounds (1-7) were elucidated based on spectroscopic data (Supplementary Figures S2–15) and by comparison with those previously reported. The isolated flavonoids (Figure 1A) were identified as isoquercetin (1) (Han et al., 2004), myricitrin (2) (Fu et al., 2013), astragaloside (3) (Wei et al., 2011), quercitrin (4) (Tatsis et al., 2007), quercetin (5) (Mabry et al., 1970), kaempferol (6) (Mabry et al., 1970; Elsayed et al., 2020), and methyl gallate (7) (Ma et al., 2005). The *in vitro* RSA showed a concentration-dependent antioxidant activity of EPE against DPPH (Figure 1C) and ABTS (Figure 1D) radicals with IC₅₀ values of 51.30 and 27.78 µg/ml, respectively.

3.2 EPE ameliorates glucose intolerance and IR in diabetic rats

OGTT was performed, and insulin was measured to determine the anti-hyperglycemic effect of EPE. The HFD/STZ-induced diabetic rats exhibited significant elevation in BG (Figure 2A, B). Treatment with EPE and PIO effectively ameliorated BG levels in diabetic rats (p < 0.001). Insulin was declined in diabetic rats (p < 0.001; Figure 2C), and the value of HOMA-IR was elevated (Figure 2D). All doses of EPE effectively alleviated insulin and HOMA-IR (p < 0.001). EPE didn't alter glucose and insulin in normal animals.

3.3 EPE modulates carbohydrate-metabolizing enzymes in diabetic rats

The activity of hexokinase (Figure 3A) was decreased, and G-6-Pase (Figure 3B), F-1,6-BPase (Figure 3C), and glycogen phosphorylase (Figure 3D) were activated in the diabetic rat liver (p < 0.001). Liver glycogen was decreased in diabetic rats as compared to the non-diabetic animals (p < 0.001; Figure 3E). EPE remarkably increased hexokinase and glycogen and suppressed other enzymes in diabetic rats.

MD simulations showed the binding affinity of EPE flavonoids with hexokinase as shown in Table 1 and Figure 4 and Supplementary Figure S16. Compounds 3, 4, 5, and 6 exhibited the lowest binding energy (-7.4, -7.3, -8.1, and -7.8 kcal/mol, respectively) and formed multiple polar bonding and hydrophobic interactions with different amino acid residues (Table 1).

3.4 EPE ameliorates dyslipidemia and liver lipid accumulation in diabetic rats

TG, TC, LDL-C, and vLDL-C were increased in the serum of diabetic rats (p < 0.001) as shown in Figures 5A–D. HDL-C was decreased (p < 0.01; Figure 5E), and AIP was elevated (p < 0.001;




FIGURE 3
EPE increased hexokinase (A), decreased G-6-Pase (B), F-1,6-BPase (C), and glycogen phosphorylase (D) and enhanced glycogen (E) in the liver of diabetic rats. Data are mean ± SD (n = 6). *p < 0.05, **p < 0.01, and ***p < 0.001 vs. control. ##p < 0.01 and ###p < 0.001 vs. diabetic.

TABLE 1 Binding affinities, interacting polar residues, and hydrophobic interactions of the compounds isolated from *E. peplus* with hexokinase.

Compound	Binding energy (kcal/mol)	Polar bond	Hydrophobic interaction
1	-6.9	Gly535, Thr536, Gly747, Glu783, Thr784, and Thr863	Thr680, Met748, and Gly780
2	-7.0	Leu617 and Gln739	Ala505, Ser506, Ala507, Pro508, Lys510, Pro605, Lys618, Glu708, Ala711, Asp714, Asn715, and Lys738
3	-7.4	Asp532, Asp657, Glu864, and Ser897	Arg539, Ile677, Asp861, Thr863, Lys866, Asp895, and Gly896
4	-7.3	Glu877, Cys886, Val888, and Ser886	Lys873, His876, Lys880, and Asp887
5	-8.1	Asp657, Thr680, Thr863, and Ser897	Asp532, Thr536, Arg539, Ile677, Gly679, Asp861, Asp895, and Gly896
6	-7.8	Asp657, Thr680, Thr863, and Ser897	Asp532, Thr536, Arg539, Ile677, Gly679, Asp861, Asp895, and Gly896
7	-6.1	Asp532, Thr661, and Ser897	Gly535, Thr536, Ile677, Gly679, Thr680, Asp861, Gly862, and Thr863




Figure 5F) in diabetic animals. All doses of EPE decreased serum lipids and AIP and increased HDL-C in diabetic rats. Dyslipidemia was associated with increased liver TG

(Figure 6A) and cholesterol (Figure 6B). Likewise, the stained sections of the liver of diabetic rats revealed the deposition of lipids (Figure 6C) along with increased circulating transaminases




FIGURE 5

EPE decreased serum TG (A), TC (B), LDL-C (C), vLDL-C (D), and AIP (F) and increased HDL-C (E) in diabetic rats. Data are mean \pm SD ($n = 6$). * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$ vs. control. ## $p < 0.01$ and ### $p < 0.001$ vs. diabetic.

(Figures 6D,E; $p < 0.001$). All doses of EPE decreased liver lipids and serum transaminases in diabetic rats.

3.5 EPE mitigates oxidative stress in diabetic rats

MDA and NO were elevated in HFD/STZ-induced rats ($p < 0.001$) as compared to the control rats (Figures 7A,B). In contrast, GSH (Figure 7C), SOD (Figure 7D), CAT (Figure 7E), and GPx (Figure 7F) were decreased in diabetic animals. EPE decreased MDA and NO and increased antioxidants effectively in diabetic rats while showing no effect on normal animals.

3.6 EPE attenuates inflammation in diabetic rats

Liver NF- κ B p65 and serum TNF- α and IL-1 β were upregulated in diabetic rats as depicted in Figures 8A–C. Treatment with EPE noticeably decreased the assayed inflammatory markers in rats with diabetes. The binding affinity of the isolated flavonoids toward NF- κ B was investigated with MD (Table 2; Figure 9 and Supplementary Figure S17). All compounds showed binding affinity marked by the polar bonding and hydrophobic interactions, and compounds 2, 3, 4, and 5 showed the lowest binding energy (–9.5, –10.6, –9.8, and –9.6 kcal/mol, respectively).




FIGURE 6 EPE decreased liver TG (A) and cholesterol (B), prevented lipid deposition (C), and ameliorated serum ALT (D) and AST (E) in diabetic rats. Data are mean ± SD (n = 6). *p < 0.05, **p < 0.01, and ***p < 0.001 vs. control. ###p < 0.001 vs. diabetic.

3.7 EPE upregulates adiponectin and PPARγ in diabetic rats

Circulating adiponectin was declined in rats with diabetes, and all EPE doses effectively restored its levels (Figure 10A). The effect of EPE on PPARγ and the binding affinity of the isolated flavonoids were determined using qRT-PCR and MD, respectively. As shown in Figure 10B, diabetic rats exhibited significant downregulation of liver PPARγ, an effect that was reversed following treatment with all doses of EPE and the PPARγ agonist PIO. MD revealed the affinity of *E. peplus* flavonoids toward PPARγ, and compounds 3, 4, and 5 exhibited the lowest binding energy (-8.7, -8.0, and -8.0 kcal/mol, respectively) (Table 3; Figure 10C and Supplementary Figure S18I).

4 Discussion

Plants of the genus *Euphorbia* showed a very promising anti-diabetic effect in STZ-, alloxan-, and sucrose-induced DM in rats (Kumar et al., 2010; Zafar et al., 2021; Mustafa et al., 2022), and the LD₅₀ of most *Euphorbia* species was estimated to exceed 5,000 mg/kg (Abd-Elhakim et al., 2019). Herein, we explored the ameliorative effect of the flavonoid-rich fraction of *E. peplus* on hyperglycemia, IR, OS, and inflammation in HFD/STZ-induced T2D rats. The *in vitro* assays showed that EPE scavenged DPPH radicals in a concentration-dependent manner. Previous studies showed the DPPH radical-scavenging efficacy of plants of the genus *Euphorbia* such as *E. royleana* (Zafar et al., 2021). The DPPH assay data were supported by the ability of EPE to scavenge ABTS radicals, demonstrating its powerful RSA.




FIGURE 7
 EPE decreased liver MDA (A) and NO (B) and increased GSH (C), SOD (D), CAT (E), and GPx (F) in diabetic rats. Data are mean ± SD (n = 6). *p < 0.05, **p < 0.01, and ***p < 0.001 vs. control. #p < 0.05, ##p < 0.01, and ###p < 0.001 vs. diabetic.

ABTS assay is more reliable and accurate for the evaluation of RSA of phytoconstituents than DPPH (Floegel et al., 2011). The RSA of EPE could be directly related to the rich content of flavonoids that possess potent scavenging properties against free radicals (Kamel et al., 2016; Elsayed et al., 2020).

The effect of EPE on glucose intolerance and IR was investigated *in vivo* in rats with HFD/STZ-induced diabetes. HFD and STZ were employed to induce T2D as this model showed similarities to the disease in humans. Feeding a HFD results in IR, and STZ decreases insulin release by damaging β-cells (Breyer et al., 2005; Lee et al., 2011), leading to hyperglycemia. Together with IR, hyperglycemia is a characteristic feature of T2D and should be managed to prevent complications in different

organs (Jellinger, 2007). Here, HFD/STZ-challenged animals showed hyperglycemia marked by glucose intolerance and IR. The developed T2D was consistent with our previous investigations, showing IR and hyperglycemia in HFD/STZ-induced rats (Mahmoud et al., 2012; Germoush et al., 2019; Elsayed et al., 2020; Abduh et al., 2023). The chronic hyperglycemia in this model was supported by the values of HbA1c%, a reliable marker for both diagnosis and prognosis of DM (American Diabetes Association, 2014) reported in our recent work (Abduh et al., 2023). Hyperglycemia was associated with hypoinsulinemia, and the development of IR as the value of HOMA-IR was revealed. Similar to these findings, elevated glucose, HbA1c%, and HOMA-IR along with decreased insulin




FIGURE 8 EPE decreased liver NF-κB p65 (A) and serum TNF-α (B) and IL-1β (C) in diabetic rats. Data are mean ± SD (n = 6). *p < 0.05, **p < 0.01, and ***p < 0.001 vs. control. ###p < 0.001 vs. diabetic.

TABLE 2 Binding affinities, interacting polar residues, and hydrophobic interactions of the compounds isolated from *E. pepus* with the NF-κB-DNA complex.

Compound	Binding energy (kcal/mol)	Polar bond	Hydrophobic interaction
1	-8.9	Gln274 and two DNA units	Five DNA units
2	-9.5	Four DNA units	Gln274 and four DNA units
3	-10.6	Asp217, Lys218, and four DNA units	Asn186, Arg187, Arg305, and one DNA unit
4	-9.8	Glu222 and four DNA units	Three DNA units
5	-9.6	Lys241, Ser246, Arg246, Asn247, and three DNA units	Asp271, Lys272, and one DNA unit
6	-9.1	Lys241, Asp271, Arg246, Asn247, and three DNA units	Ser246, Lys272, and one DNA unit
7	-6.4	Lys241 and three DNA units	Arg246, Lys272, and two DNA units

were reported in HFD/STZ-challenged rats (Abduh et al., 2023). The declined insulin is due to damage caused to the pancreatic islets induced by STZ-mediated ROS generation and DNA damage (Lenzen, 2008). Although the early phase of damage is associated with increased insulin release as a compensatory mechanism, prolonged hyperglycemia and ROS release deteriorate the

pancreatic islets and promote more β-cell damage and ultimately reduced insulin release (Ntimbane et al., 2016). The effects of excessive ROS include enhanced lipid peroxidation (LPO), massively increased cytosolic Ca²⁺, and diminished pancreatic antioxidants, effects that enhance the destruction of β-cells (Nahdi et al., 2017).




FIGURE 9

Molecular docking showing the binding modes of compounds 2, 3, 4, and 5 with the NF- κ B-DNA complex.

Treatment with EPE effectively ameliorated glucose intolerance and HOMA-IR, denoting its anti-hyperglycemic and insulin-sensitizing effects. These effects added support to

the previously reported anti-hyperglycemic activity of plants of the same genus. For instance, *E. royleana* stem extract decreased fasting BG (FBG) and ameliorated glucose intolerance in diabetic




FIGURE 10
 EPE increased serum adiponectin (A) and upregulated liver PPAR γ mRNA (B) in diabetic rats. Data are mean \pm SD (n = 6). *p < 0.05 and ***p < 0.001 vs. control. ##p < 0.01 and ###p < 0.001 vs. diabetic. (C) Molecular docking showing the binding modes of compounds 2, 3, and 4 with PPAR γ .

rats (Zafar et al., 2021), and *E. helioscopia* alleviated BG and insulin in sucrose-fed rats (Mustafa et al., 2022). The ameliorative effect of EPE on hyperglycemia is a direct result of increased

insulin secretion. Impaired insulin release and IR increase hepatic glucose output due to suppressed glycolysis and glycogenesis. Impaired insulin release and IR can also impair

TABLE 3 Binding affinities, interacting polar residues, and hydrophobic interactions of the compounds isolated from *E. peplus* with PPAR γ .

Compound	Binding energy (kcal/mol)	Polar bond	Hydrophobic interaction
1	-7.9	Gln271, Glu272, and Ser342	Ile262, Leu270, Gly284, Phe287, Arg288, Ile341, and Met348
2	-8.7	Ile281	Leu270, Gln271, Arg280, Gln283, Gly284, Cys285, Phe287, Arg288, Ile341, Ser342, and Met348
3	-8.0	Ser342 and Glu343	Ile262, Leu270, Gly284, Phe287, Arg288, Leu330, Leu333, Leu340, and Ile341
4	-8.0		Ile262, Thr268, Leu270, Gln271, Arg280, Gly284, Cys285, Phe287, Arg288, Ile341, and Ser342
5	-7.5	Glu259, Gln271, Arg280, and Glu291	Leu270, Gly284, Phe287, and Arg288
6	-7.4		Leu270, Gln271, Arg280, Gln283, Gly284, Phe287, Arg288, and Glu291
7	-5.8	Ser289, His323, and Met364	Cys285, Tyr327, Phe363, His449, Leu469, and Tyr473

peripheral glucose uptake and hepatic gluconeogenesis, resulting in hyperglycemia (Nordlie et al., 1999). By alleviating insulin release and IR, EPE effectively ameliorated hyperglycemia possibly by modulating enzymes involved in glycogenesis and gluconeogenesis. This notion was supported by the findings of this study where EPE increased hexokinase and suppressed F-1,6-BPase, G-6-Pase, and glycogen phosphorylase, resulting in increased liver glycogen content. Hexokinase is involved in glucose oxidation and suppressed by IR and insulin deficiency. Suppressed hexokinase activity decreases glycolysis, and hence glucose accumulates in the blood (Gupta et al., 1999). Along with hexokinase suppression, insulin insufficiency activates G-6-Pase, F-1,6-BPase, and glycogen phosphorylase, resulting in enhanced gluconeogenesis and glycogenolysis (Roden and Bernroider, 2003). The improved insulin sensitivity and levels of EPE decreased glycogenolysis and gluconeogenesis and enhanced liver glycogen by modulating the activity of the involved enzymes. In addition to the determined enzymes, insulin activates glycogen synthase and suppresses glycogen phosphorylase (Postic et al., 2004), and this explains the alleviated glycogen levels following treatment with EPE. Owing to its role in glucose oxidation, the ameliorated FBG following EPE supplementation is a result of enhanced hexokinase activity. To further explore the effect of EPE on hexokinase activity, we carried out MD simulations of the binding affinity of the contained flavonoids toward the enzyme. All flavonoids revealed binding affinity marked by polar bonding toward important residues in the active site and dense hydrophobic interactions. Recent findings showed improvements in glycemic status and insulin sensitivity by plant extracts that modulate the carbohydrate-metabolizing enzymes (Germoush et al., 2019; Elsayed et al., 2020). In this context, Mustafa et al. (2022) related the anti-hyperglycemic effect of *E. helioscopia* in sucrose-fed rats to its ability to modulate the activities of pyruvate kinase, glucokinase, and phosphofructokinase.

In addition to hyperglycemia, dyslipidemia is found in T2D and can increase atherogenicity and the risk of cardiovascular disease (Reaven, 2005). Elevated serum lipids and decreased

HDL-C in this study represent an atherogenic profile as previously described (Germoush et al., 2019). AIP, a marker of lipoprotein particle size that possesses a predictive value beyond that of the assayed lipids (Dobiášová, 2006), was increased in diabetic rats. The observed dyslipidemia is a direct result of IR and the enhanced lipolysis and decreased lipogenesis (Carpentier, 2021). Increased lipolysis provokes liver lipid accumulation which is also promoted by increased synthesis of free fatty acids (FFAs) that provoke lipogenesis within hepatocytes (Mohamed et al., 2016). Lipid accumulation in hepatocytes causes cell injury, thereby aggravating IR, hyperglycemia, and dyslipidemia (Levinthal and Tavill, 1999). Herein, lipids were increased in the liver, and circulating transaminases were elevated in diabetic rats as previously reported (Elsayed et al., 2020; Abduh et al., 2023). EPE effectively ameliorated serum and liver lipids, effects that were directly related to the enhanced insulin release and sensitivity.

Owing to the involvement of OS and inflammation in provoking IR and the complications of DM (Mahmoud et al., 2012; Mahmoud, 2017), we explored the ability of EPE to suppress these pathological processes. Diabetic rats showed OS and inflammatory reactions marked by elevated MDA, NO, NF- κ B, TNF- α , and IL-1 β and declined antioxidants. OS, defined by excess ROS and decreased antioxidants, is a key mechanism in IR and can damage cells and alter multiple signaling pathways. Hyperglycemia can increase the production of ROS and lead to OS by activating NADPH oxidases and promoting mitochondrial dysfunction (Jimenez et al., 2018). Excess ROS can activate pathways related to increased pro-inflammatory cytokines, and both can impair insulin signaling, leading to IR and glucose accumulation in the blood (Rösen et al., 2001). The altered insulin levels shift the signaling where PI3K phosphorylates Rac, resulting in increased NADPH oxidase 4-mediated ROS generation (Campa et al., 2015). Excess ROS activates casein kinase-2 followed by retromer that alters glucose transporter-4 membrane translocation and impair glucose uptake (Ma et al., 2014). ROS can also increase mitochondrial fission that stimulates stress responses and

impairs insulin signaling and has been linked to IR as well as apoptosis (Jheng et al., 2012). Pro-inflammatory cytokines trigger IR by altering insulin signaling and many kinases. The elevated IL-1 β and TNF- α reported in this study can impair insulin-stimulated uptake of glucose, stimulate lipolysis and gluconeogenesis, and inhibit tyrosine phosphorylation of insulin receptor substrate-1 and protein kinase B activation (Green et al., 1994; Del Aguila et al., 1999; Jager et al., 2007). Therefore, attenuation of OS and pro-inflammatory cytokines can attenuate IR and increase insulin signaling, activity, and stimulated glucose uptake.

EPE enhanced antioxidants and prevented OS and inflammation in diabetic rats in this investigation. In addition to its *in vitro* RSA, EPE prevented LPO, enhanced antioxidants, and suppressed NF- κ B and cytokines in diabetic rats. The suppression of inflammation following EPE supplementation was supported by *in silico* investigations that showed the ability of flavonoids to bind strongly with NF- κ B through multiple polar bonding and hydrophobic interactions. The attenuation of these pathological processes contributed to the anti-hyperglycemic and insulin-sensitizing effects of EPE. Numerous studies showed the beneficial effects of antioxidants and plant extracts that are rich in antioxidant phytochemicals against hyperglycemia and IR (Mahmoud, 2013; Mahmoud et al., 2017; Germoush et al., 2019). The antioxidant and anti-inflammatory role of EPE is related to its content of flavonoids which possess potent RSA and showed benefits against DM (Mahmoud, 2013; Mahmoud et al., 2017; Germoush et al., 2019; Abukhalil et al., 2021). In diabetic patients, the supplementation of flavonoids improved glycemic and lipidemic statuses and antioxidants and decreased inflammatory markers (Li et al., 2015). In obese patients, the consumption of flavonoids positively affected the metabolic status by lowering systemic oxidation and enhancing insulin sensitivity (Suliburska et al., 2012).

The beneficial effects of EPE could also be linked to the upregulation of adiponectin and PPAR γ . EPE increased serum adiponectin that participated, at least in part, in the amelioration of hyperglycemia. Adiponectin exerts insulin-sensitizing effects and possesses anti-inflammatory activity, and experimental evidence revealed that it ameliorated hyperglycemia in HFD-fed rodents (Fruebis et al., 2001; Yamauchi et al., 2001). Despite its ameliorated hyperglycemia in T1D and T2D in rodents, high adiponectin doses didn't affect BG in normal animals. These findings suggested that the downregulation of glycogenolysis and gluconeogenesis mediated its anti-hyperglycemic effects. Accordingly, adiponectin decreased glucose production in rat hepatocytes and G-6-Pase mRNA abundance in mice (Berg et al., 2001; Combs et al., 2001). It can also upregulate liver CD36, PPAR α , and UCP-2, effects that were related to the increase in insulin sensitivity (Yamauchi et al., 2001). EPE upregulated liver PPAR γ , and its flavonoids were shown to dock into the PPAR γ active site through polar bonding and hydrophobic interactions. The activation of PPAR γ is a key mechanism for ameliorating hyperglycemia, and IR and PPAR γ agonists, such as PIO, increase insulin sensitivity and ameliorate hyperglycemia,

dyslipidemia, OS, and inflammation (Tontonoz and Spiegelman, 2008). PPAR γ suppresses OS and inflammation by enhancing antioxidant enzymes (Okuno et al., 2010), inhibiting the activation of NF- κ B both directly and indirectly (Kersten et al., 2000; Remels et al., 2009), and preventing ROS generation from NADPH oxidases (Hwang et al., 2005). However, the lack of PPAR γ protein expression data could be considered a limitation to this study.

5 Conclusion

This investigation introduces new information that *E. peplus* is rich in flavonoids and possesses potent radical-scavenging and anti-diabetic efficacies. EPE ameliorated hyperglycemia, IR, OS, dyslipidemia, and inflammation in rats with T2D. In addition, EPE modulated carbohydrate-metabolizing enzymes and enhanced antioxidants, adiponectin, and PPAR γ . *In silico* findings revealed the binding affinity of *E. peplus* constituents toward hexokinase, NF- κ B, and PPAR γ . Therefore, *E. peplus* could be a promising candidate for the development of a potent anti-hyperglycemic and insulin-sensitizing agent. However, further investigations to determine other molecular mechanism(s) of action are needed.

Data availability statement

The original contributions presented in the study are included in the article/Supplementary Material; further inquiries can be directed to the corresponding author.

Ethics statement

The animal study was reviewed and approved by the Research Ethics Committee of Al-Azhar University.

Author contributions

Conceptualization, AM, EK, and RA; methodology, AM, RA, MA, GM-H, AB-A, and EK; software, AM and EK; validation, AM; formal analysis, RA, MA, AM, and EK; investigation, RA, AM, GM-H, MA, AB-A, and EK; resources, RA, GM-H, MA, and EH; data curation, AM, EK, and RA; writing—original draft and preparation, AM, GM-H, and EK; writing—review and editing, AM; visualization, AM; supervision, AM; project administration, RA and AM; funding acquisition, RA. All authors contributed to the article and approved the submitted version.

Funding

This research was funded by the Nourah Bint Abdulrahman University Researchers Supporting Project (grant number PNURSP

2023R381), Princess Nourah Bint Abdulrahman University, Riyadh, Saudi Arabia.

Acknowledgments

Princess Nourah bint Abdulrahman University Researchers Supporting Project number (PNURSP2023R381), Princess Nourah bint Abdulrahman University, Riyadh, Saudi Arabia.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

- Abd El-Twab, S. M., Mohamed, H. M., and Mahmoud, A. M. (2016). Taurine and pioglitazone attenuate diabetes-induced testicular damage by abrogation of oxidative stress and up-regulation of the pituitary-gonadal axis. *Can. J. Physiol. Pharmacol.* 94, 651–661. doi:10.1139/cjpp-2015-0503
- Abd-Elhakim, Y. M., Abdo Nassan, M., Salem, G. A., Sasi, A., Aldhahrani, A., Ben Issa, K., et al. (2019). Investigation of the *in-vivo* cytotoxicity and the *in silico*-prediction of MDM2-p53 inhibitor potential of *Euphorbia peplus* methanolic extract in rats. *Toxins (Basel)* 11, 642. doi:10.3390/toxins11110642
- Abduh, M. S., Alzogaibi, M. A., Alzogaibi, A. M., Bin-Ammar, A., Alotaibi, M. F., Kamel, E. M., et al. (2023). Arbutin ameliorates hyperglycemia, dyslipidemia and oxidative stress and modulates adipocytokines and PPAR γ in high-fat diet/streptozotocin-induced diabetic rats. *Life Sci.* 321, 121612. doi:10.1016/j.lfs.2023.121612
- Abukhalil, M. H., Althunibat, O. Y., Aladaileh, S. H., Al-Amarat, W., Obeidat, H. M., Al-Khawalde, A. a. A., et al. (2021). Galangin attenuates diabetic cardiomyopathy through modulating oxidative stress, inflammation and apoptosis in rats. *Biomed. Pharmacother.* 138, 111410. doi:10.1016/j.biopha.2021.111410
- Aebi, H. (1984). “[13] catalase *in vitro*,” in *Methods in enzymology* (Cambridge: Academic Press), 121–126.
- Ahmed, S., Yousaf, M., Mothana, R. A., and Al-Rehaily, A. J. (2016). Studies on wound healing activity of some EUPHORBIA species on experimental rats. *Afr. J. Tradit. Complement. Altern. Med.* 13, 145–152. doi:10.21010/ajtcam.v13i5.19
- Al-Sultan, S., and Hussein, Y. A. (2006). Acute toxicity of *Euphorbia helioscopia* in rats. *Pak. J. Nutr.* 5, 135–140. doi:10.3923/pjn.2006.135.140
- American Diabetes Association (2021). 2. Classification and diagnosis of diabetes: Standards of medical care in diabetes-2021. *Diabetes Care* 44, S15–S33. doi:10.2337/dc21-S002
- American Diabetes Association (2014). Standards of medical care in diabetes-2014. *Diabetes Care* 37, S14–S80. doi:10.2337/dc14-S014
- Bancroft, J. D., and Gamble, M. (2008). *Theory and practice of histological techniques*. Netherlands: Elsevier health sciences.
- Berg, A. H., Combs, T. P., Du, X., Brownlee, M., and Scherer, P. E. (2001). The adipocyte-secreted protein Acrp30 enhances hepatic insulin action. *Nat. Med.* 7, 947–953. doi:10.1038/90992
- Beutler, E., Duron, O., and Kelly, B. M. (1963). Improved method for the determination of blood glutathione. *J. laboratory Clin. Med.* 61, 882–888.
- Brand-Williams, W., Cuvelier, M. E., and Berset, C. (1995). Use of a free radical method to evaluate antioxidant activity. *LWT - Food Sci. Technol.* 28, 25–30. doi:10.1016/s0023-6438(95)80008-5
- Brandstrup, N., Kirk, J. E., and Bruni, C. (1957). The hexokinase and phosphoglucosomerase activities of aortic and pulmonary artery tissue in individuals of various ages. *J. Gerontol.* 12, 166–171. doi:10.1093/geronj/12.2.166
- Breyer, M. D., Böttinger, E., Brosius, F. C., Iii, Coffman, T. M., Harris, R. C., Heilig, C. W., et al. (2005). Mouse models of diabetic nephropathy. *J. Am. Soc. Nephrol.* 16, 27–45. doi:10.1681/ASN.2004080648
- Campa, C. C., Ciralo, E., Ghigo, A., Germena, G., and Hirsch, E. (2015). Crossroads of PI3K and rac pathways. *Small GTPases* 6, 71–80. doi:10.4161/21541248.2014.989789

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors, and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fphar.2023.1204641/full#supplementary-material>

Carpentier, A. C. (2021). 100th anniversary of the discovery of insulin perspective: Insulin and adipose tissue fatty acid metabolism. *Am. J. Physiology-Endocrinology Metabolism* 320, E653–E670. doi:10.1152/ajpendo.00620.2020

Combs, T. P., Berg, A. H., Obici, S., Scherer, P. E., and Rossetti, L. (2001). Endogenous glucose production is inhibited by the adipose-derived protein Acrp30. *J. Clin. Invest.* 108, 1875–1881. doi:10.1172/JCI14120

Del Aguila, L. F., Claffey, K. P., and Kirwan, J. P. (1999). TNF-alpha impairs insulin signaling and insulin stimulation of glucose uptake in C2C12 muscle cells. *Am. J. Physiology - Endocrinol. Metabolism* 276, E849–E855. doi:10.1152/ajpendo.1999.276.5.E849

Dobiášová, M. (2006). AIP-atherogenic index of plasma as a significant predictor of cardiovascular risk: From research to practice. *Vnitr Lek.* 52, 64–71.

Elsayed, R. H., Kamel, E. M., Mahmoud, A. M., El-Bassuony, A. A., Bin-Jumah, M., Lamsabhi, A. M., et al. (2020). Rumex dentatus L. phenolics ameliorate hyperglycemia by modulating hepatic key enzymes of carbohydrate metabolism, oxidative stress and PPAR γ in diabetic rats. *Food Chem. Toxicol.* 138, 111202. doi:10.1016/j.fct.2020.111202

Floegel, A., Kim, D.-O., Chung, S.-J., Koo, S. I., and Chun, O. K. (2011). Comparison of ABTS/DPPH assays to measure antioxidant capacity in popular antioxidant-rich US foods. *J. food Compos. analysis* 24, 1043–1048. doi:10.1016/j.jfca.2011.01.008

Flohé, L., and Günzler, W. A. (1984). Assays of glutathione peroxidase. *Methods Enzymol.* 105, 114–121. doi:10.1016/s0076-6879(84)05015-1

Folch, J., Lees, M., and Sloane Stanley, G. H. (1957). A simple method for the isolation and purification of total lipides from animal tissues. *J. Biol. Chem.* 226, 497–509. doi:10.1016/s0021-9258(18)64849-5

Freedland, R. A., and Harper, A. E. (1959). Metabolic adaptations in higher animals. *J. Biol. Chem.* 234, 1350–1354. doi:10.1016/s0021-9258(18)70010-0

Fruebis, J., Tsao, T. S., Javarschi, S., Ebbets-Reed, D., Erickson, M. R., Yen, F. T., et al. (2001). Proteolytic cleavage product of 30-kDa adipocyte complement-related protein increases fatty acid oxidation in muscle and causes weight loss in mice. *Proc. Natl. Acad. Sci. U. S. A.* 98, 2005–2010. doi:10.1073/pnas.041591798

Fu, Y., Li, Z., Si, J., Chang, Q., Li, Z., and Pan, R. (2013). Separation and purification of myricitrin from bayberry tree bark by high-speed counter-current chromatography. *J. Liq. Chromatogr. Relat. Technol.* 36, 1503–1512. doi:10.1080/10826076.2012.692147

Germoush, M. O., Elgebaly, H. A., Hassan, S., Kamel, E. M., Bin-Jumah, M., and Mahmoud, A. M. (2019). Consumption of terpenoids-rich padina pavonia extract attenuates hyperglycemia, insulin resistance and oxidative stress, and upregulates PPAR γ in a rat model of type 2 diabetes. *Antioxidants (Basel)* 9, 22. doi:10.3390/antiox9010022

Giner, J. L., and Schroeder, T. N. (2015). Polygonifoliol, a new tirucallane triterpene from the latex of the seaside sandmat *Euphorbia polygonifolia*. *Chem. Biodivers.* 12, 1126–1129. doi:10.1002/cbdv.201400426

Green, A., Dobias, S. B., Walters, D. J. A., and Brasier, A. R. (1994). Tumor necrosis factor increases the rate of lipolysis in primary cultures of adipocytes without altering levels of hormone-sensitive lipase. *Endocrinology* 134, 2581–2588. doi:10.1210/endo.134.6.8194485

Green, L. C., Wagner, D. A., Glogowski, J., Skipper, P. L., Wishnok, J. S., and Tannenbaum, S. R. (1982). Analysis of nitrate, nitrite, and [15N]nitrate in biological fluids. *Anal. Biochem.* 126, 131–138. doi:10.1016/0003-2697(82)90118-x

- Gupta, D., Raju, J., Prakash, J., and Baquer, N. Z. (1999). Change in the lipid profile, lipogenic and related enzymes in the livers of experimental diabetic rats: Effect of insulin and vanadate. *Diabetes Res. Clin. Pract.* 46, 1–7. doi:10.1016/s0168-8227(99)00067-4
- Guzik, T. J., and Cosentino, F. (2018). Epigenetics and immunometabolism in diabetes and aging. *Antioxid. Redox Signal* 29, 257–274. doi:10.1089/ars.2017.7299
- Haffner, S. M. (2000). Coronary heart disease in patients with diabetes. *N. Engl. J. Med.* 342, 1040–1042. doi:10.1056/NEJM200004063421408
- Han, J.-T., Bang, M.-H., Chun, O.-K., Kim, D.-O., Lee, C.-Y., and Baek, N.-I. (2004). Flavonol glycosides from the aerial parts of *Aceriphyllum rossii* and their antioxidant activities. *Archives Pharmacol. Res.* 27, 390–395. doi:10.1007/BF02980079
- Hwang, J., Kleinhenz, D. J., Lassegue, B., Griendling, K. K., Dikalov, S., and Hart, C. M. (2005). Peroxisome proliferator-activated receptor-gamma ligands regulate endothelial membrane superoxide production. *Am. J. Physiol. Cell Physiol.* 288, C899–C905. doi:10.1152/ajpcell.00474.2004
- Jager, J., Grémeaux, T., Cormont, M., Le Marchand-Brustel, Y., and Tanti, J. F. (2007). Interleukin-1beta-induced insulin resistance in adipocytes through down-regulation of insulin receptor substrate-1 expression. *Endocrinology* 148, 241–251. doi:10.1210/en.2006-0692
- Jellinger, P. S. (2007). Metabolic consequences of hyperglycemia and insulin resistance. *Clin. Cornerstone* 8, S30–S42. doi:10.1016/s1098-3597(07)80019-6
- Jheng, H. F., Tsai, P. J., Guo, S. M., Kuo, L. H., Chang, C. S., Su, I. J., et al. (2012). Mitochondrial fission contributes to mitochondrial dysfunction and insulin resistance in skeletal muscle. *Mol. Cell Biol.* 32, 309–319. doi:10.1128/MCB.05603-11
- Jimenez, R., Toral, M., Gómez-Guzmán, M., Romero, M., Sanchez, M., Mahmoud, A. M., et al. (2018). The role of Nrf2 signaling in ppar β/δ -mediated vascular protection against hyperglycemia-induced oxidative stress. *Oxid. Med. Cell Longev.* 2018, 5852706. doi:10.1155/2018/5852706
- Kahn, S. E., Cooper, M. E., and Del Prato, S. (2014). Pathophysiology and treatment of type 2 diabetes: Perspectives on the past, present, and future. *Lancet* 383, 1068–1083. doi:10.1016/S0140-6736(13)62154-6
- Kamel, E. M., Ahmed, N. A., El-Bassuony, A. A., Hussein, O. E., Alrashdi, B., Ahmed, S. A., et al. (2022). Xanthine oxidase inhibitory activity of *Euphorbia peplus* L. Phenolics. *Comb. Chem. High. Throughput Screen* 25, 1336–1344. doi:10.2174/1386207324666210609104456
- Kamel, E. M., Mahmoud, A. M., Ahmed, S. A., and Lamsabhi, A. M. (2016). A phytochemical and computational study on flavonoids isolated from *Trifolium resupinatum* L. and their novel hepatoprotective activity. *Food Funct.* 7, 2094–2106. doi:10.1039/c6fo00194g
- Kersten, S., Desvergne, B., and Wahli, W. (2000). Roles of PPARs in health and disease. *Nature* 405, 421–424. doi:10.1038/35013000
- Koide, H., and Oda, T. (1959). Pathological occurrence of glucose-6-phosphatase in serum in liver diseases. *Clin. Chim. Acta* 4, 554–561. doi:10.1016/0009-8981(59)90165-2
- Kumar, S., Malhotra, R., and Kumar, D. (2010). Antidiabetic and free radicals scavenging potential of *Euphorbia hirta* flower extract. *Indian J. Pharm. Sci.* 72, 533–537. doi:10.4103/0250-474X.73921
- Lee, Y. S., Li, P., Huh, J. Y., Hwang, I. J., Lu, M., Kim, J. I., et al. (2011). Inflammation is necessary for long-term but not short-term high-fat diet-induced insulin resistance. *Diabetes* 60, 2474–2483. doi:10.2337/db11-0194
- Lenzen, S. (2008). The mechanisms of alloxan- and streptozotocin-induced diabetes. *Diabetologia* 51, 216–226. doi:10.1007/s00125-007-0886-7
- Levinthal, G. N., and Tavill, A. S. (1999). Liver disease and diabetes mellitus. *Clin. DIABETES* 17, 73–81.
- Li, D., Zhang, Y., Liu, Y., Sun, R., and Xia, M. (2015). Purified anthocyanin supplementation reduces dyslipidemia, enhances antioxidant capacity, and prevents insulin resistance in diabetic patients. *J. Nutr.* 145, 742–748. doi:10.3945/jn.114.205674
- Livak, K. J., and Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 25, 402–408. doi:10.1006/meth.2001.1262
- Ma, J., Nakagawa, Y., Kojima, I., and Shibata, H. (2014). Prolonged insulin stimulation down-regulates GLUT4 through oxidative stress-mediated retromer inhibition by a protein kinase CK2-dependent mechanism in 3T3-L1 adipocytes. *J. Biol. Chem.* 289, 133–142. doi:10.1074/jbc.M113.533240
- Ma, X., Wu, L., Ito, Y., and Tian, W. (2005). Application of preparative high-speed counter-current chromatography for separation of methyl gallate from *Acer truncatum* Bunge. *J. Chromatogr. A* 1076, 212–215. doi:10.1016/j.chroma.2005.04.077
- Mabry, T. J., Markham, K., Thomas, M., Mabry, T. J., Markham, K., and Thomas, M. (1970). "The NMR spectra of flavonoids," in *The systematic identification of flavonoids* (Berlin, Heidelberg: Springer), 274–343.
- Mahmoud, A. M., Abd El-Twab, S. M., and Abdel-Reheim, E. S. (2017). Consumption of polyphenol-rich morus alba leaves extract attenuates early diabetic retinopathy: The underlying mechanism. *Eur. J. Nutr.* 56, 1671–1684. doi:10.1007/s00394-016-1214-0
- Mahmoud, A. M., Ashour, M. B., Abdel-Moneim, A., and Ahmed, O. M. (2012). Hesperidin and naringin attenuate hyperglycemia-mediated oxidative stress and proinflammatory cytokine production in high fat fed/streptozotocin-induced type 2 diabetic rats. *J. Diabetes its Complicat.* 26, 483–490. doi:10.1016/j.jdiacomp.2012.06.001
- Mahmoud, A. M. (2017). "Exercise ameliorates metabolic disturbances and oxidative stress in diabetic cardiomyopathy: Possible underlying mechanisms," in *Exercise for cardiovascular disease prevention and treatment: From molecular to clinical, Part 1*. Editor J. Xiao (Singapore: Springer Singapore), 207–230.
- Mahmoud, A. M. (2013). Hematological alterations in diabetic rats - role of adipocytokines and effect of citrus flavonoids. *Excli J.* 12, 647–657.
- Marklund, S., and Marklund, G. (1974). Involvement of the superoxide anion radical in the autoxidation of pyrogallol and a convenient assay for superoxide dismutase. *FEBS Eur. J. Biochem.* 47, 469–474. doi:10.1111/j.1432-1033.1974.tb03714.x
- Mohamed, J., Nazratun Nafizah, A. H., Zariyantey, A. H., and Budin, S. B. (2016). Mechanisms of Diabetes-Induced Liver Damage: The role of oxidative stress and inflammation. *Sultan Qaboos Univ. Med. J.* 16, e132–e141. doi:10.18295/squmj.2016.16.02.002
- Mustafa, I., Anwar, H., Irfan, S., Muzaffar, H., and Ijaz, M. U. (2022). Attenuation of carbohydrate metabolism and lipid profile by methanolic extract of *Euphorbia helioscopia* and improvement of beta cell function in a type 2 diabetic rat model. *BMC Complementary Med. Ther.* 22, 23. doi:10.1186/s12906-022-03507-2
- Nahdi, A. M. T. A., John, A., and Raza, H. (2017). Elucidation of molecular mechanisms of streptozotocin-induced oxidative stress, apoptosis, and mitochondrial dysfunction in rin-5F pancreatic β -cells. *Oxidative Med. Cell Longev.* 2017, 7054272. doi:10.1155/2017/7054272
- Nordlie, R. C., Foster, J. D., and Lange, A. J. (1999). Regulation of glucose production by the liver. *Annu. Rev. Nutr.* 19, 379–406. doi:10.1146/annurev.nutr.19.1.379
- Ntimbane, T., Mailhot, G., Spahis, S., Rabasa-Lhoret, R., Kleme, M. L., Mellou, D., et al. (2016). CFTR silencing in pancreatic β -cells reveals a functional impact on glucose-stimulated insulin secretion and oxidative stress response. *Am. J. Physiol. Endocrinol. Metab.* 310, E200–E212. doi:10.1152/ajpendo.00333.2015
- Ohkawa, H., Ohishi, N., and Yagi, K. (1979). Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal. Biochem.* 95, 351–358. doi:10.1016/0003-2697(79)90738-3
- Okuno, Y., Matsuda, M., Miyata, Y., Fukuhara, A., Komuro, R., Shimabukuro, M., et al. (2010). Human catalase gene is regulated by peroxisome proliferator activated receptor-gamma through a response element distinct from that of mouse. *Endocr. J.* 57, 303–309. doi:10.1507/endocrj.k09e-113
- Pattanaik, S., Si, S. C., Pal, A., Panda, J., and Nayak, S. S. (2014). Wound healing activity of methanolic extract of the leaves of *Crataeva magna* and *Euphorbia nerifolia* in rats. *J. Appl. Pharm. Sci.* 4, 046–049. doi:10.7324/japs.2014.40310
- Postic, C., Dentin, R., and Girard, J. (2004). Role of the liver in the control of carbohydrate and lipid homeostasis. *Diabetes Metabolism* 30, 398–408. doi:10.1016/s1262-3636(07)70133-7
- Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M., and Rice-Evans, C. (1999). Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic. Biol. Med.* 26, 1231–1237. doi:10.1016/s0891-5849(98)00315-3
- Reaven, G. M. (2005). Compensatory hyperinsulinemia and the development of an atherogenic lipoprotein profile: The price paid to maintain glucose homeostasis in insulin-resistant individuals. *Endocrinol. Metab. Clin. North Am.* 34, 49–62. doi:10.1016/j.ecl.2004.12.001
- Remels, A. H., Langen, R. C., Gosker, H. R., Russell, A. P., Spaapen, F., Voncken, J. W., et al. (2009). PPARgamma inhibits NF-kappaB-dependent transcriptional activation in skeletal muscle. *Am. J. Physiol. Endocrinol. Metab.* 297, E174–E183. doi:10.1152/ajpendo.90632.2008
- Roden, M., and Bernroider, E. (2003). Hepatic glucose metabolism in humans-its role in health and disease. *Best. Pract. Res. Clin. Endocrinol. Metab.* 17, 365–383. doi:10.1016/s1521-690x(03)00031-9
- Rösen, P., Nawroth, P. P., King, G., Möller, W., Tritschler, H. J., and Packer, L. (2001). The role of oxidative stress in the onset and progression of diabetes and its complications: A summary of a congress series sponsored by UNESCO-MCBN, the American diabetes association and the German diabetes society. *Diabetes/Metabolism Res. Rev.* 17, 189–212. doi:10.1002/dmrr.196
- Saeedi, P., Petersohn, I., Salpea, P., Malanda, B., Karuranga, S., Unwin, N., et al. (2019). Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: Results from the international diabetes federation diabetes atlas, 9th edition. *Diabetes Res. Clin. Pract.* 157, 107843. doi:10.1016/j.diabres.2019.107843
- Sami, D. H., Soliman, A. S., Khawailed, A. A., Hassanein, E. H., Kamel, E. M., and Mahmoud, A. M. J. L. S. (2022). 7-hydroxycoumarin modulates Nrf2/HO-1 and microRNA-34a/SIRT1 signaling and prevents cisplatin-induced oxidative stress, inflammation, and kidney injury in rats. *Life Sci.* 310, 121104. doi:10.1016/j.lfs.2022.121104
- Seifter, S., Dayton, S., et al. (1950). The estimation of glycogen with the anthrone reagent. *Arch. Biochem.* 25, 191–200.

- Shi, Q. W., Su, X. H., and Kiyota, H. (2008). Chemical and pharmacological research of the plants in genus *Euphorbia*. *Chem. Rev.* 108, 4295–4327. doi:10.1021/cr078350s
- Singla, A., and Kamla, P. (1990). Phytoconstituents of *Euphorbia* species. *Fitoterapia* 41, 483–516.
- Stalmans, W., and Hers, H. G. (1975). The stimulation of liver phosphorylase b by AMP, fluoride and sulfate. A technical note on the specific determination of the a and b forms of liver glycogen phosphorylase. *Eur. J. Biochem.* 54, 341–350. doi:10.1111/j.1432-1033.1975.tb04144.x
- Suliburska, J., Bogdanski, P., Szulinska, M., Stepien, M., Pupek-Musialik, D., and Jablecka, A. (2012). Effects of green tea supplementation on elements, total antioxidants, lipids, and glucose values in the serum of obese patients. *Biol. Trace Elem. Res.* 149, 315–322. doi:10.1007/s12011-012-9448-z
- Tatsis, E. C., Boeren, S., Exarchou, V., Troganis, A. N., Vervoort, J., and Gerothanassis, I. P. (2007). Identification of the major constituents of *Hypericum perforatum* by LC/SPE/NMR and/or LC/MS. *Phytochemistry* 68, 383–393. doi:10.1016/j.phytochem.2006.11.026
- Tontonoz, P., and Spiegelman, B. M. (2008). Fat and beyond: The diverse biology of PPARgamma. *Annu. Rev. Biochem.* 77, 289–312. doi:10.1146/annurev.biochem.77.061307.091829
- Trinder, P. (1969). Determination of glucose in blood using glucose oxidase with an alternative oxygen acceptor. *Ann. Clin. Biochem. An Int. J. Biochem. laboratory Med. Ann. Clin. Biochem. An Int. J. Biochem. laboratory Med.* 6, 24–27. doi:10.1177/000456326900600108
- Wei, Y., Xie, Q., Fisher, D., and Sutherland, I. A. (2011). Separation of patuletin-3-O-glucoside, astragaloside, quercetin, kaempferol and isorhamnetin from *Flaveria bidentis* (L.) Kuntze by elution-pump-out high-performance counter-current chromatography. *J. Chromatogr. A* 1218, 6206–6211. doi:10.1016/j.chroma.2011.01.058
- Yamauchi, T., Kamon, J., Waki, H., Terauchi, Y., Kubota, N., Hara, K., et al. (2001). The fat-derived hormone adiponectin reverses insulin resistance associated with both lipotrophy and obesity. *Nat. Med.* 7, 941–946. doi:10.1038/90984
- Zafar, M., Sharif, A., Khan, D., Akhtar, B., Muhammad, F., Akhtar, M. F., et al. (2021). Preventive effect of *Euphorbia royleana* Boiss on diabetes induced by streptozotocin via modulating oxidative stress and deoxyribonucleic acid damage. *Toxin Rev.* 40, 777–790. doi:10.1080/15569543.2020.1780262