


**Please cite the Published Version**

Morgan, P , Smeuninx, B and Breen, L (2020) Exploring the Impact of Obesity on Skeletal Muscle Function in Older Age. *Frontiers in Nutrition*, 7. 569904 ISSN 2296-861X

**DOI:** <https://doi.org/10.3389/fnut.2020.569904>

**Publisher:** Frontiers Media S.A.

**Version:** Published Version

**Downloaded from:** <https://e-space.mmu.ac.uk/633293/>

**Usage rights:**  [Creative Commons: Attribution 4.0](https://creativecommons.org/licenses/by/4.0/)

**Additional Information:** This is an open access article published in *Frontiers in Nutrition*, by Frontiers Media S.A.

**Enquiries:**

If you have questions about this document, contact [openresearch@mmu.ac.uk](mailto:openresearch@mmu.ac.uk). Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)



# Exploring the Impact of Obesity on Skeletal Muscle Function in Older Age

Paul T. Morgan<sup>1</sup>, Benoit Smeuninx<sup>1,2</sup> and Leigh Breen<sup>1\*</sup>

<sup>1</sup> School of Sport, Exercise and Rehabilitation Sciences, University of Birmingham, Birmingham, United Kingdom, <sup>2</sup> Cellular & Molecular Metabolism Laboratory, Monash Institute of Pharmacological Sciences, Monash University, Parkville, VIC, Australia

## OPEN ACCESS

### Edited by:

Benjamin Toby Wall,  
University of Exeter, United Kingdom

### Reviewed by:

Thomas Elbenhardt Jensen,  
University of Copenhagen, Denmark  
Marlou L. Dirks,  
University of Exeter, United Kingdom

### \*Correspondence:

Leigh Breen  
L.breen@bham.ac.uk

### Specialty section:

This article was submitted to  
Nutrition and Metabolism,  
a section of the journal  
Frontiers in Nutrition

**Received:** 05 June 2020

**Accepted:** 06 November 2020

**Published:** 01 December 2020

### Citation:

Morgan PT, Smeuninx B and Breen L  
(2020) Exploring the Impact of Obesity  
on Skeletal Muscle Function in Older  
Age. *Front. Nutr.* 7:569904.  
doi: 10.3389/fnut.2020.569904

Sarcopenia is of important clinical relevance for loss of independence in older adults. The prevalence of obesity in combination with sarcopenia (“sarcopenic-obesity”) is increasing at a rapid rate. However, whilst the development of sarcopenia is understood to be multi-factorial and harmful to health, the role of obesity from a protective and damaging perspective on skeletal muscle in aging, is poorly understood. Specifically, the presence of obesity in older age may be accompanied by a greater volume of skeletal muscle mass in weight-bearing muscles compared with lean older individuals, despite impaired physical function and resistance to anabolic stimuli. Collectively, these findings support a potential *paradox* in which obesity may protect skeletal muscle mass in older age. One explanation for these paradoxical findings may be that the anabolic response to weight-bearing activity could be greater in obese vs. lean older individuals due to a larger mechanical stimulus, compensating for the heightened muscle anabolic resistance. However, it is likely that there is a complex interplay between muscle, adipose, and external influences in the aging process that are ultimately harmful to health in the long-term. This narrative briefly explores *some* of the potential mechanisms regulating changes in skeletal muscle mass and function in aging combined with obesity and the interplay with sarcopenia, with a particular focus on muscle morphology and the regulation of muscle proteostasis. In addition, whilst highly complex, we attempt to provide an updated summary for the role of obesity from a protective and damaging perspective on muscle mass and function in older age. We conclude with a brief discussion on treatment of sarcopenia and obesity and a summary of future directions for this research field.

**Keywords:** sarcopenic-obesity, muscle function, metabolic syndrome, sarcopenia, obesity paradox, anabolic resistance, intramuscular lipids

## INTRODUCTION

The degenerative, generalized, and precipitous loss of skeletal muscle mass, quality and strength associated with aging is termed “sarcopenia” and is characterized by muscle loss of ~0.5–1.0% per year (1). The consequences of muscle mass and strength loss with aging are well-documented and encompass a number of physiological and non-physiological outcomes (1–7). In addition, sarcopenia is associated with decreased mobility and impaired whole-body metabolic health, as

well as impaired function during locomotion, reduced resting energy expenditure, a reduction in non-structured free-living physical activity and increased fat mass (8, 9), factors that have also been found to be present with obesity. Evidence now exists to implicate the progressive deterioration of muscle quality, defined as muscle strength relative to a given quantity of muscle mass, to functional impairment with aging (2, 10, 11), including reductions in fiber size, number, and contractile function (12, 13), the degree of lipid infiltration (14, 15), and impaired neurological modulation of contraction (16). However, whilst worldwide rates of obesity have reached an all-time high (World Health Organization, WHO) and is associated with a plethora of comorbidities (17), research on how obesity might affect the progression of sarcopenia is in its relative infancy. Therefore, the purpose of this review is to explicitly focus on skeletal muscle deterioration in the presence of excess adiposity in older age and will explore some of the potential underlying mechanisms regulating changes in skeletal muscle function, with a particular focus on *in-vivo* models. The reader should be directed to recent reviews elsewhere for in-depth critical discussions on the potential multi-faceted causes of sarcopenia, obesity, and the confluence of these two conditions (7, 18–21). Whilst not a direct focus of the present review, the effects of obesity on molecular signaling associated with skeletal muscle contraction, bone health and the contribution of specific genetic and gut microbiome fingerprints on obesity as well to muscle mass/loss and its possible pleiotropic effects on adipose tissue are also reviewed elsewhere (22–25).

## Causes of Sarcopenia and Obesity

It is now widely acknowledged that the development of sarcopenia and loss of muscle strength is caused by the complex interplay of multiple factors beyond muscle atrophy (10, 26, 27) including; declines in neural function, hormonal changes, chronic low-grade inflammation, mitochondrial dysfunction, impaired/reduced satellite cell function, and lifestyle factors [e.g., malnutrition and physical inactivity; (2, 7, 21, 28–31)]. Similar to sarcopenia, the presence of obesity is also typically accelerated by chronic inactivity. Obesity is primarily caused by a progressive decline in total energy expenditure (i.e., decreased physical activity and reduced basal metabolic rate) in the presence of excessive caloric intake (20, 32). As opposed to the traditional model of sarcopenia, the high energy intakes observed with obesity may offer some protection against sarcopenic development by ensuring sufficient protein intake and supporting elevated muscle protein synthesis (20, 32–37). Growing evidence supports an underlying genetic component to the development of obesity (38–40). Whilst the major factors involved in obesity seem to be linked with dietary and physical activity habits, these factors have also been associated with the genome that may, independently, influence energy expenditure, fuel metabolism, muscle fiber function, gut microbiome, and hormone/appetite regulation (i.e., ghrelin and leptin) and/or food preferences (38–40).

## Health Consequences of Obesity

As aforementioned, alongside an aging population, levels of global obesity are on a progressive rise, with >30% of men and >40% of women currently classified as being overweight, and >10% as obese (41). An increase in whole-body adiposity is typically accompanied by a concomitant increase in ectopic fat deposition within skeletal muscle, termed myosteatosis (42, 43), both of which are associated with a plethora of comorbidities (17), including the progression of sarcopenia (44–47) and physical disability (48, 49). Independent of the effects of obesity on skeletal muscle health, adiposity is also a strong risk factor for poor overall health, reduced functional capacity and quality of life in older age (50–52). Furthermore, obesity has well-known metabolic effects that can lead to significant health complications such as metabolic syndrome (defined by the existence of a cluster of conditions including hypertension, high blood glucose levels, high serum triglyceride levels), Type II diabetes, and an increased risk of cancer (53, 54). In addition, obesity is a strong risk factor for atherosclerosis and other cardiovascular complications (55), as well as for many other chronic and acute diseases involving end-stage organ failure and infection (54–57), leading to further acute complications and prolonged periods of hospitalization. Unsurprisingly, the prevalence of many of the aforementioned medical complications are strongly associated with obesity in older age (58). Obesity is known to accelerate the risk of developing cardiovascular disease by increasing circulating lipids and glucose as well as impairing lipid and glucose metabolism and increasing systolic and diastolic blood pressure (59–61). In addition, obesity is strongly associated with elevated chronic systemic inflammation and progressive declines in physical activity and cardiorespiratory fitness, which all contribute to a worse prognosis of ill health and accelerated sarcopenia risk (59–62). Obesity is associated with a number of endocrinological impairments (63). For example, in obese adipose tissue adipocytes undergo hypertrophy, hyperplasia and activation resulting in accumulation of pro-inflammatory macrophages and other immune cells as well as dysregulated production of various adipokines that together further exacerbate inflammation (23). The excessive production and impaired capacity to store lipids in old obese individuals is also known to increase reactive oxygen species and impair metabolic function that is ultimately capable of inducing metabolic dysfunction (notably, impaired mitochondrial and skeletal muscle function via elevated inflammation) (23). Whilst adiponectin, an adipokine produced primarily by adipose tissue, is known to hold a number of important metabolic functions (e.g., insulin-sensitizer and anti-inflammatory), these responses become dysregulated with obesity, making skeletal muscle particularly susceptible to metabolic impairments and increased risk of morbidity (64, 65). It has been suggested that a vicious circle of maintaining a high volume of adipose tissue and skeletal muscle inflammation, triggers the development and acceleration of sarcopenic-obesity with a complex interplay between a number of mechanisms that are beyond the scope of this review (i.e. adipocyte-like phenotype of muscle progenitor cells, insulin resistance, impaired neuromodulation of contraction, leptin

resistance, impaired endocrine function, reduced release of adiponectin, increased glycation products and oxidative stress etc.) (23, 63, 66). However, less is known on how obesity, *per se*, might affect the progression of sarcopenia in older age and/or in pre-sarcopenic individuals. Undoubtedly though, myosteatosis has the potential to exert detrimental local effects on muscle contractility given its close proximity (67) and this is discussed below.

## Diagnosis and Prevalence of Sarcopenic-Obesity

The combined presence of sarcopenia and obesity (i.e., sarcopenic-obesity) is rapidly increasing in older adults. Changes to body composition, including reduced muscle mass concomitant with an increase in fat mass and simultaneous reduction in resting metabolic rate, are evident with aging (68). Whole-body muscle mass and strength start to decline progressively as early as 30 years of age, with a more accelerated loss after the age of ~60–70 years (69–71). The prevalence of metabolic syndrome also increases with age, with a rise after the 3rd decade of life, reaching a peak between 50 and 80 years (72, 73). Importantly, as skeletal muscle plays a critical role in glycaemic control and metabolic homeostasis (74, 75), the combination of a loss of lean body mass and a concomitant increase in visceral adiposity will likely exacerbate the risk of developing metabolic disease, particularly in the older adult, accelerating skeletal muscle deterioration, reduced physical performance, an increased risk for disability, hospitalizations, morbidity, and early mortality (76–78). The development of sarcopenic-obesity is thought to be brought about by a combination of factors that contribute to the development of both conditions (21) and is associated with significantly greater body fat, oxidative stress and inflammation and intracellular lipotoxicity, glucose and endocrine dysregulation, as well as impaired muscle strength and lower lean body mass (63, 66, 79–81). Further, when obesity and impaired muscle function co-exist, they act synergistically in a “vicious cycle” on the risk of developing multiple health-related outcomes (19, 21, 42, 82–84).


Multiple definitions exist for sarcopenic-obesity (21, 63), however, the use of gross metrics such as body mass and BMI, often provide an incomplete understanding of the person's actual body composition. This paradigm is depicted in **Figure 1** which illustrates a cross-section of the thigh taken by magnetic resonance imagery (MRI) from a young healthy individual (**Figure 1A**) and pre-sarcopenic older individual (**Figure 1B**) with similar thigh circumferences (~60–65 cm) and similar BMI (~26 kg·m<sup>2</sup>). Despite a similar initial appearance, the differences in the composition of the muscle between individuals are clear. In the young-healthy individual, the portion of muscle mass is relatively large with minimal adiposity and lipid infiltration. In contrast, the pre-sarcopenic older tissue is significantly smaller and contains a much larger layer of subcutaneous fat and lipid infiltration within and between muscles of the thigh. In this example, the pre-sarcopenic individual presents with a relatively “normal” body mass and BMI but stores higher and relatively lower amounts of adipose and lean tissue,

respectively. Therefore, the pre-sarcopenic individual presents a significantly elevated risk of developing metabolic complications. **Figures 1C–E** demonstrate a significant greater lipid droplet number and area in type II fibers of obese older compared with healthy young and healthy old individuals, which could be indicative of poorer muscle quality and impaired oxidative function in older obese individuals. The precise identification of sarcopenia and obesity is of vital importance for accurate diagnosis and treatment of sarcopenic-obesity (84). Importantly, as obese older adults typically experience a reduction in relative (i.e., %) muscle mass as opposed to a loss in absolute (i.e., kg) muscle mass (44), sarcopenia in obese individuals is often characterized based upon changes to muscle quality and/or relative rather than absolute levels of skeletal muscle mass (86–88). A number of tools are, therefore, used in combination to assess body composition (i.e., MRI, CT, DEXA, BIA), muscle strength (i.e., dynamometry, 1- and 12-repetition maximum testing, grip strength), and physical performance [i.e., gait speed, stair climb, chair rise time, standard physical performance battery tests (SPPB)]. The following section will discuss the potential implications of obesity for aging muscle, with a specific focus on muscle morphology and the existence of an “obesity-paradox.”

## IMPLICATIONS OF OBESITY FOR AGING MUSCLE

### Obesity and Muscle Morphology

It is well-recognized that aging is typically characterized by a slowing of muscle phenotype toward type I and a concomitant reduction in type II muscle fiber area and number [i.e., (89)]. However, in contrast to these typical symptoms of aging, obesity seems to be associated with a more predominant faster phenotype (particularly type IIx) and smaller percentage of type I muscle fibers (85, 90, 91), with a negative relationship between the degree of adiposity and the relative percentage of type I muscle fibers, comparable to that observed with reduced physical activity (92–95). In further support of this phenomenon, following a weight loss intervention, a positive relationship between the percentage of excess weight loss and the percentage of type I fibers has been found in morbidly obese individuals (91). Whilst there is evidence to suggest that a relative reduction in type I fibers is a result of excess adiposity, it is also possible that this may reflect an intrinsic defect predisposing individuals toward obesity (91). This is pertinent to note as, aside from any impact on functional outcomes, a relative reduction in type I muscle fibers is also related to impaired metabolic health, increased LDL content, decreased insulin sensitivity and decreased arterial elasticity (90). In contrast to type I muscle fibers, type II muscle fibers possess impaired lipid disposal capabilities, thus contributing to a reduction in whole-body lipid oxidation and an increased storage of lipids (96–98). Type II muscle fibers are more closely associated with higher oxidative stress which may, in turn, negatively impact the integrity of mitochondria and lead to an apoptotic cascade that may ultimately result in cell death (99). Others have also reported that skeletal muscle from obese individuals has markedly lower mitochondrial



**FIGURE 1** | Cross-sectional MRI of the midpoint of quadriceps muscle from a young healthy individual (**A**) and pre-sarcopenic older individual (**B**) with a similar thigh circumference (~60–65 cm) and BMI (~26 kg·m<sup>2</sup>). The young-healthy individual exhibits a larger quantity of skeletal muscle mass (**A**) whereas high infiltration by adipose tissue is observed in the pre-sarcopenic older individual (**B**). (**C–E**) Provides an illustration of representative fiber cross-sectional images stained for Myosin Heavy Chain I (red) and lipid droplets (bodipy, green) in young lean (YL), old lean (OL), and older obese (OO), respectively. Notably, type I fiber lipid droplet number is significantly greater in young lean ( $n = 616$ ) compared with older lean ( $n = 412$ ) and older obese ( $n = 533$ ) and greater in older obese compared with older lean (**C–E**, respectively) indicative of an ‘athlete paradox’, whereby young, healthy individuals exhibit higher levels of intramuscular lipids but with the superior capacity for oxidation. In addition, type II fiber lipid droplet number and area is significantly greater in the young lean ( $n = 377$ ) and old obese ( $n = 415$ ) compared with old lean ( $n = 242$ ) individuals, despite poor whole-body metabolic health in the former (i.e., old obese), indicative of poorer muscle quality and impaired oxidative function. Further, in this study, type I and II muscle fiber cross-sectional area was greater in young lean ( $4,031 \pm 1,978 \mu\text{m}^2$ ,  $4,009 \pm 1,733 \mu\text{m}^2$ , respectively) compared with older obese ( $3,421 \pm 1,528 \mu\text{m}^2$ ,  $3,390 \pm 1,199 \mu\text{m}^2$ , respectively) and older lean ( $3,009 \pm 1,251 \mu\text{m}^2$ ,  $2,170 \pm 1,243 \mu\text{m}^2$ , respectively) groups and significantly greater in old obese compared with older lean [from Smeuninx et al. (85)]. White bars for (**C–E**) represent 50  $\mu\text{m}$ .

content and a concomitant lower oxidative capacity, which is also linked with insulin resistance (43, 96, 100–103). A recent review has found adiposity to have profound negative impacts on cellularity, secretory profiles, and inflammatory status, which drive lipotoxicity of skeletal muscle and negatively impact muscle fiber contractility (78). In contrast, an increase in the absolute number and size of type II muscle fibers is associated with indices of muscle strength and power, and, thus, may contribute to a reduced risk of falls and fracture which can have important implications for muscle loss in aging (104). However, both aging and obesity are independently associated with muscle atrophy and reduced myogenesis, as well as impaired excitation-contraction coupling (105, 106). Obese individuals are also known to possess impaired functional capacity (48, 107) and fat mass is negatively associated with jump height in middle-aged and older adults (108). An important consideration in these individuals is the significantly higher intramuscular lipid content in type II fiber (compared with older lean controls) combined with impaired muscle fiber contraction, lower relative muscle strength and impaired oxidative function, indicative of poorer muscle quality (85, 109–112). At least in old sarcopenic-obese rodents, ectopic fat deposition seems to also contribute to anabolic resistance (113). We have also previously shown

type I fiber intramuscular lipid content to be equivalent between young lean and old obese, but not old lean, individuals (85) (**Figure 1**). However, type I fiber intramuscular lipid content was negatively associated with insulin sensitivity for old lean and old obese, but not young lean individuals (85), reinforcing the notion that the capacity for intramuscular lipid oxidation is likely greater in young lean and, in particular, significantly impaired in older obese individuals (14, 85). It is also pertinent to note that such observations are also likely to be related to aerobic fitness/physical activity levels rather than obesity, *per se*. Indeed, both impaired oxidative capacity and increased muscle IIX fiber number are both typical outcomes of physical inactivity (114). Nevertheless, as reduced physical activity is typically an inherent characteristic of obesity, it is often difficult to distinguish between the two. It is plausible that a physically active older obese individual might possess greater intramuscular lipid oxidation capabilities compared with an inactivity older obese individual (85).

Obesity is also characterized by the induction of skeletal muscle remodeling as well as mitochondria-mediated apoptosis (115). On the contrary, exercise has been reported as a positive regulator of skeletal muscle remodeling and apoptosis, however, as aforementioned, obese individuals typically present

with low levels of physical activity (115). Interestingly, it has been suggested that the impaired function of mitochondria with obesity may be triggered by altering the expression of mitochondrial proteins regulating key metabolic processes in skeletal muscle likely due, in-part, to the subcellular localization of lipid droplets and a reduction in the amount of skeletal muscle perilipin 2 protein per intramyocellular lipid, particularly in type II muscle fibers (116, 117). A recent review has also demonstrated that such effects of obesity on muscle morphology and contractile performance may be muscle specific, such that obesity may increase the absolute force producing capacity of the postural and antigravity muscles (22, 118). Further, studies examining the contractile performance of isolated muscle suggest that obesity may accelerate the age-related decline of respiratory, but not locomotory weight-bearing, muscles (119). However, whilst a high-fat diet (and an associated increase in body and fat mass in rodents) may contribute to increased muscle mass and strength, the increase in absolute strength is smaller than the magnitude of weight gain, meaning that *in-vivo* locomotor function is likely to be impaired in old obese adults (119). In further support of this notion, a recent study demonstrated an additive effect of obesity, induced by a high-fat diet on the reduction in contractile function with aged rodents, demonstrating that the increase in intramyocellular lipid levels were associated with the degree of impaired muscle contractile force (120, 121).

## A Paradox of Obesity in Older Age?

Whilst many studies have shown that muscle strength and, to some extent, muscle mass are strong predictors of survival (2, 71, 122–124), the relationship between obesity and mortality remains equivocal (125–127). Studies have shown that obese individuals seemingly maintain a higher absolute quantity of muscle mass compared with their lean counterparts (85), supporting the existence of an aging “obesity paradox” (128). The existence of an obesity paradox has been observed in a range of populations including, for example, patients undergoing lung cancer surgery (129), with overall morbidity and in-hospital mortality significantly decreased in obese patients (129). Whilst a number of studies have shown a BMI in the “normal” range (18.5–25 kg·m<sup>2</sup>) to be associated with the lowest risk of mortality (130, 131), other large cohort studies have demonstrated a “survival benefit” for overweight (25–30 kg·m<sup>2</sup>) and even obese ( $\geq 30$  kg·m<sup>2</sup>) individuals, suggesting a protective effect of obesity against mortality in older age (125, 132–134). By contrast, a recent study found that whilst waist circumference was associated with greater muscle size, this was also associated with impaired muscle function (assessed via leg extension strength, chair stands and stair climb time, gait speed, and SPPB scores) as well as with measures of muscle quality (135). It is widely accepted that, with increasing body mass, absolute skeletal muscle mass increases, and this might explain these consequential findings in obese individuals (87). In support of this notion, reduced survival for individuals with a ‘normal’ (18.5–25 kg·m<sup>2</sup>) or ‘low’ ( $\leq 18.5$  kg·m<sup>2</sup>) BMI, compared with overweight ( $\geq 25$  kg·m<sup>2</sup>), might be explained by loss of muscle mass in the former (136). A recent study found that skeletal muscle mass mediates associations

of BMI with adiposity and mortality, such that irrespective of adiposity and BMI, muscle mass was inversely associated with the risk of mortality (137). Excluding participants with low muscle mass across all groups also alleviated the risk associated with a low BMI, but magnified the risk associated with a high BMI, such that survival was greatest with a normal BMI (137). These data are in agreement with a recent cohort study sampling >38,000 men with >12,000 deaths, suggesting that the obesity paradox is likely explained by low levels of lean mass, rather than low fat mass, in the lower range of BMI, such that a “J” shaped association was consistently observed between BMI and all-cause mortality (138). In contrast, a “U” shaped association was found between predicted lean body mass and all-cause mortality (138). Thus, more recent observations are not consistent with a survival advantage related to overweight or obesity, but rather to elevated levels (or better preservation) of muscle mass in aging and “at risk” groups. Individuals with a BMI of  $\geq 30$  kg·m<sup>2</sup> have, indeed, been shown to be at a significantly higher risk of functional impairment (139).

The principle factors driving individuals toward either the sarcopenic-obese or healthy-obese phenotype are currently unknown but might include differences in physical activity and caloric intake as these are known regulators and associating factors to many of the potential causes of sarcopenia and obesity (i.e. reduced lean mass, elevated oxidative stress and inflammation, impaired muscle regeneration, anabolic and insulin resistance, nutrient ‘overload’ and metabolic dysfunction etc.) (66, 80). Furthermore, the inability to delineate sarcopenic-obesity based on BMI alone, adds to the misclassification of certain individuals (140). Indeed, whilst obese individuals are typically less physically active, observations of higher levels of muscle mass in obese groups are likely a response to an elevated overload stimulus due to the requirement to move an increased body mass. It is reasonable to speculate that the role of other factors, such as genetics, may also contribute to the discrepancies observed between these two aging groups. As aforementioned, whilst Smeuninx et al., amongst others, have shown that obese individuals seemingly maintain a higher absolute quantity of muscle mass, greater insulin resistance is also observed in these individuals compared with lean age-matched controls, suggesting that the quality, and not necessarily quantity, of skeletal muscle may play a more relevant role in whole-body metabolic health, particularly in aging (85). Observations to date question the role of obesity to skeletal muscle anabolism, and, interestingly, whether a “threshold” exists after which obesity becomes deleterious. Based on our observations, and those of others, we postulate the existence of an aging-obesity “muscle quality threshold” beyond which, obesity-induced muscle anabolic resistance results in a precipitous, rapid decline in muscle mass and function (Figure 2), centered on muscle quality rather than absolute levels of skeletal muscle mass. Research suggests that obesity may have differential effects on muscle metabolism and preservation of muscle mass depending on age (88) and it is likely that being young, physically active and not suffering from the longer-term impacts of obesity, may also mediate some level of protection against any obesity-related anabolic resistance (141). Thus, obesity may

not be problematic for muscle mass and function unless present under situations of aging, chronic inactivity, and/or other known risk factors for anabolic resistance (142–147). Undoubtedly, more studies are required to examine the metabolic, morphological, and functional characteristics related to obesity, particularly in pre-sarcopenic individuals and/or in the earlier stages of the aging process to better understand the principle factors driving individuals toward either the sarcopenic-obese or ‘healthy’-obese phenotype.

## DYSREGULATION OF SKELETAL MUSCLE IN OBESITY AND AGING

### Regulation of Muscle Proteostasis

The regulation of skeletal muscle mass is a complex process that involves the precise coordination of several metabolic and intracellular signaling pathways, ultimately affecting the dynamic balance between muscle protein synthesis (MPS) and muscle protein breakdown (MPB). In order to increase skeletal muscle mass, MPS must exceed MPB, which can be achieved by performing regular resistance exercise training in combination with adequate protein nutrition (148–151). By contrast, dysregulation of MPB in combination with impaired and/or normal MPS is observed in numerous conditions such as aging and prolonged immobilization (152, 153). In a young, healthy individual, skeletal muscle turnover is ~1–2% per day (154). Indeed, following the intake of as little as ~10 g of dietary protein, a transient robust increase in MPS is observed (155–158) and with the addition of exercise, the duration and the magnitude of this anabolic response can be potentiated (158–160). Whilst the addition of regular physical activity elevates MPB, when physical activity is combined with sufficient protein intake in healthy individuals, the outcome will be a favorable elevation in MPS and net protein accretion (155). Indeed, physical activity is an important locus of control in the regulation of skeletal muscle mass in both young and older individuals, with increased physical activity levels stimulating MPS (161) and decreased physical activity levels blunting MPS (142).


### Old Age and Muscle Proteostasis

Whilst young individuals demonstrate a pronounced response to anabolic stimuli, a blunted response has been observed in older adults (157, 161–163). These studies amongst others, indicate the importance of increasing daily, per meal and quality of protein doses as well as the intensity and volume of exercise to restore muscle anabolic sensitivity in the older adult (164–167). Recent studies have also demonstrated anabolic resistance (defined as a diminished MPS response to food intake and exercise) in older adults following a chronic exercise training program (168), which suggests that by extension, aspects of chronological aging are inevitable, and that regular physical activity becomes a less effective stimulus for muscle remodeling and maintenance. It has been suggested that the post-prandial inhibition of MPB may also be impaired in older adults in combination with a blunted post-prandial MPS stimulation (169). This age-related anabolic resistance is likely an important driving factor in age-related muscle loss and appears to be exacerbated by aspects of

biological aging such as intermittent periods of musculoskeletal disuse and reduced physical activity (87, 104, 142, 144, 145, 157, 165, 170, 171). Aging is also associated with impaired molecular regulation of skeletal muscle signaling pathways. Growing evidence suggests that mTOR complex 1 [mTORC1] (mammalian target of rapamycin) signaling, which is thought to have a regulating impact on protein synthesis, influences the aging process as acute inhibition of this pathway (by rapamycin) seems to extend the lifespan in model organisms and provides protection against a number of age-related pathologies (172–177). In addition to impaired mTORC1 signaling, impaired satellite cell proliferation is associated with aging (99, 178). This has important implications in the aging adult as satellite cells are thought to play a critical role in muscle fiber growth, muscle tissue turnover, and regeneration and are activated and proliferate in response to anabolic stimuli (179). Changes in the secretion of sex-hormones (i.e., growth hormone, testosterone, estrogen) are also associated with chronological aging (180). There has also been a recent interest in the role of a number of other factors involved in an individuals’ dietary status, beyond protein intake, that have been implicated in the age-associated deterioration of skeletal muscle mass. For example, whilst the role of vitamin D status in sarcopenia is not well-understood, it has been suggested that older adults are at higher risk of lower levels of vitamin D as a result of decreased cutaneous synthesis and dietary intake of vitamin D (181, 182) and epidemiological evidence indicates an association between low levels of vitamin D and diseases associated with aging (182). Specifically, vitamin D induces myogenic differentiation in skeletal muscle derived stem cells (183). Similarly, there has also been a recent rise in the interest of the role of vitamin D receptors in the nervous, cardiovascular, and endocrine systems that may interact to accelerate sarcopenia progression (184, 185).

### Obesity and Muscle Proteostasis

As aforementioned, infiltration of fat is evident within muscle (termed “myosteatorsis”) and this typically increases with age. Similar to chronological aging, it is generally agreed that rates of basal MPS are similar between obese and lean individuals (186). However, muscle anabolic resistance is exacerbated with obesity and thus the combination of obesity with older age leads to a near non-existent elevation of the MPS response to amino acid provision beyond basal levels, which may be associated with a plethora of factors including; reduced physical activity and a low relative proportion of lean mass, intracellular lipotoxicity and elevated inflammation, impaired muscle regeneration, insulin resistance, impaired endocrine function, impaired anabolic signaling etc. (85, 187, 188). Whilst little is known of the impact of obesity on MPB, there is evidence to suggest that MPB may be elevated due to a higher inflammatory burden associated with excessive adiposity and overfeeding (188–199). It has been suggested that the muscle anabolic resistance observed in obese individuals may be underpinned by impaired/inhibited muscle anabolic signaling phosphorylation (200, 201). Taken together, the available data suggest the possibility of an obesity-induced inability to regulate muscle protein turnover in response to nutrition, which may lead to impaired skeletal muscle




remodeling. In contrast to suggestions of elevated MPB, the impairment of MPS following protein nutrition in obese may be offset by inhibition of MPB, implicating a possible protective mechanism to explain the apparent preservation of muscle mass in obese vs. lean older adults (188). It is also worthy of note that, as obese individuals are subjected to greater loading forces and increased muscle contractile work during activities of daily living (due to the requirement of moving more inert mass), this might elicit elevations in MPS and offer a degree of “protection” against the loss age-related muscle mass loss [(85, 187, 188, 202, 203); **Figure 3**]. Indeed, there is currently not a clear consensus for the impacts of obesity on MPS. There is no obvious explanation for the discrepancies observed between studies on the impacts of obesity on post-prandial MPS (85, 186–188, 204, 205). However, differences in habitual levels of physical activity as well as the immediate effects of exercise on indices of skeletal muscle anabolism, might explain in part, some of these contrasting observations between lean and obese individuals. Indeed, we showed that the MPS response to protein ingestion correlated with physical activity (as assessed via daily step count,  $r = 0.57$ ) (85). Further, some of these discrepant findings might be explained by the assessment of MPS under hyperinsulinemic-hyperaminoacidemic conditions, which are not representative of a normal physiological response (186, 187). Anabolic resistance to protein provision has also been associated with obesity due to overfeeding and/or lipid oversupply (187, 188, 204), and therefore, the predominant cause(s) of obesity should also be considered. Nonetheless, irrespective of these discrepant findings, the magnitude of any potential “training”

effect of physical activity, as discussed above, is typically lower than the increase in body weight observed with excess adiposity and is likely insufficient in offsetting sarcopenia (119, 206).

Whilst high-fat diets, lipid administration, obesity, and ectopic fat deposition have all been shown to induce whole-body and muscle anabolic resistance (113, 187, 204, 207, 208), as aforementioned, data also implicates obesity in impairing MPS through compromised muscle quality and/or elevated levels of adipose-derived inflammatory cytokines (139, 209). Indeed, a number of lipid species and metabolites have been implicated in impairing muscle function as well as developing anabolic resistance (210–216). However, despite previously observing a negative association between the net post-prandial MPS response following ingestion of a moderate protein dose (15g milk protein) and leg fat mass, at least in our hands, we were unable to demonstrate any association with muscle fiber-specific intramuscular lipid content (85). This was despite a two-fold higher type II fiber intramuscular lipid content in older obese compared with older lean and young lean individuals. This, however, is in contrast to previous reports of associations between intramuscular lipids and muscle anabolic resistance in old rodents (113). Type I fiber intramuscular lipid content was, though, negatively associated with insulin sensitivity for old lean and old obese, but not young lean. This reinforces the notion that the capacity for intramuscular lipid oxidation is likely greater in young lean individuals and that this may contribute to metabolic resistance in the former (14). Activity-matched older lean and older obese comparisons are required to confirm such speculation (14). The subcellular characteristics, location of





intramuscular and/or specific class of lipid intermediates (i.e., ceramides, diacylglycerol, etc.) may associate more closely with age-related muscle anabolic resistance (217). The specific role of perilipins, a lipid droplet-associated protein, requires further research (213, 215), particularly as there is lack of research investigating anabolic resistance in response to load carriage work in obese older vs. lean age-matched individuals. Taken together, these data may support an obesity-induced blunting of post-prandial MPS due to intramuscular lipid accumulation. We speculate that any potential protection of adiposity on skeletal muscle mass is likely solely driven by physical activity and that decreases in muscle mass and function in individuals diagnosed with sarcopenia and obesity combined are likely to be accelerating at a rate whereby levels of adiposity fail to provide any protection, but rather compound disease progression. Importantly, and as aforementioned, the causes of obesity and muscle loss are likely interrelated and interact. For example, research suggests that increasing storage of fat may also influence

sedentary behavior, producing a vicious cycle of skeletal muscle deterioration and impaired whole-body metabolic health (218). Therefore, appropriate treatment for obesity (and sarcopenic-obesity) is imperative in improving muscle function and quality of life in older age.

## TREATMENT OF SARCOPENIA AND OBESITY

Treating obesity appropriately in older age is of upmost importance in order to prevent any further loss of skeletal muscle associated with aging (219–221). Treatment of obesity in older age (and sarcopenic-obesity) requires a combination of personalized nutrition and physical activity approaches as, under certain conditions, inappropriate changes to these paradigms may actually accelerate the progression of sarcopenia given the susceptibility to energy restriction and, thus, a potential catabolic state (222). As the pathogenesis of these three conditions (sarcopenia, obesity, and sarcopenic-obesity) is multifaceted, understanding the optimal treatment is highly complex. Optimal treatment of sarcopenic-obesity likely incorporates a combination of resistance and aerobic training in the presence of a small dietary calorie deficit with sufficient protein intake, as severe weight reduction strategies may also compromise the ability to preserve muscle function and mass which may subsequently lead to frailty, disability, and increased morbidity and mortality (220, 223). It is also noteworthy that bone density may also be impaired with weight-loss programs which may, in-turn, contribute to fracture risk, and subsequent further risk of hospitalization, periods of prolonged inactivity and further muscle loss and functional impairment (87, 104, 222). Nevertheless, weight loss should be considered safe in older adults, if managed appropriately (222, 223). As such, treatment of sarcopenia alone may be easier than treatment of (pre)-sarcopenic obese individuals. Accumulating evidence also indicates that diet-induced weight loss can improve physical function among obese older adults, independent of any changes in physical activity (224–226). Incorporation of resistance exercise, in particular, into a training program is extremely important in reversing the symptoms of obesity in older age. Studies suggest that when a diet and exercise intervention is administered to obese individuals, subsequent improvements in muscle strength and muscle quality, are observed (227, 228). More recent attempts to assess the effectiveness of chronic training paradigms in obese older individuals have investigated the potential of a combined aerobic and resistance training approach and found positive outcomes when combined to either alone (229). Resistance exercise is known to have a number of effects that have the potential to prevent and/or overcome anabolic resistance including: enhanced post-prandial muscle protein synthesis, sensitized muscle insulin action, increased number and size of muscle fibers (particularly fast twitch) and reduced inflammation [i.e., IL-6 and TNF- $\alpha$ ; (160, 230, 231)]. Aerobic exercise has also been shown to improve oxidative capacity, reduce intramyocellular lipid accumulation, enhance lipolysis and improve glucose utilization and insulin sensitivity

(232), all of which have the potential to improve symptoms of sarcopenia and/or obesity in older age. Whilst we recognize that research suggests that weight loss in obese individuals is also typically followed by a further period of weight gain (233–236), it is important to note that the metabolic health benefits of weight loss and increased physical activity are likely sustained for a prolonged period (234, 235). Indeed, even when weight has been regained, long lasting benefits of weight loss are observed on a number of important metrics including insulin sensitivity (237). Accordingly, a multi-faceted approach to the prevention, management and treatment of sarcopenic-obesity remains the most promising.

Taken together, as undoubtedly older individuals with higher amounts of muscle mass are less likely to suffer from the inevitable impacts of age-related muscle loss, it would be prudent to increase basal levels of muscle mass, preferably, via a well-programmed resistance exercise training programme possibly combined with small increases in fat mass and body weight. It is, however, important to recognize that excess weight gain via increases in fat mass, whilst may be associated with higher volumes of muscle, are likely to impair muscle function and accelerate the age-related impairment in muscle function by significantly impairing muscle quality. In contrast, given the potential for muscle mass in obese older adults to be somewhat protected from sarcopenia, it *may* be prudent to suggest individuals carry a small amount of extra weight in older age, at a consequence of a small sacrifice in muscle quality, in support of a “muscle quality threshold” (Figure 2). Whilst speculative at this stage, this may help achieve a delicate balance of increasing lean and fat mass without the deliberating effects of excess adiposity. However, whilst the optimal strategy in pre-sarcopenic individuals is unclear, older individuals who are under weight and seemingly at an elevated risk, may benefit from small increases in fat mass. Undoubtedly though, promotion of a healthy lifestyle for chronic prevention (or attenuation), rather than treatment, *per se*, of sarcopenia, obesity, and sarcopenic-obesity should be considered the *best* approach by promoting adaptations of higher basal levels of muscle mass. Therefore, a combination of progression physical activity and gradual well-planned caloric restriction are likely key to achieving improved health outcomes in overweight individuals (84, 219, 238, 239). For more comprehensive reviews on the treatment of sarcopenia, obesity and sarcopenic-obesity in older age, the reader is directed to the following [i.e., (216, 232, 233)].

## CONCLUSIONS

### Future Directions

Throughout this review, we have provided some suggestions on how/where this research field could benefit with further research. From a broad perspective, further work is required to delineate the precise mechanisms through which obesity exacerbates the age-related loss of muscle mass and function to better understand the cause(s) and treatments of sarcopenic-obesity. For example, there is a need to understand the specific role of myosteatosis, lipid species and metabolites, in age-related muscle anabolic resistance, particularly in obese individuals. Indeed,

further exploration of the role of intramuscular lipids in obesity-induced muscle anabolic resistance and sarcopenia is pivotal, particularly with regard to the co-localization of lipid droplets and mitochondria. In addition, it is plausible to suggest that the MPS response to weight-bearing activity could be greater in obese vs. lean older individuals, due to the larger mechanical stimulus. This could provide some explanation for the compensation for the blunted MPS response to amino acids, despite an apparent better preservation of muscle mass in obese older individuals. Future studies should also attempt longitudinal observations of obesity-induced changes to skeletal muscle as well as the effects of lifelong high calorie dietary patterns and sedentarism, and the reversal of obesity on skeletal muscle to truly understand the complex interactions between aging, obesity, skeletal muscle mass regulation and metabolic health parameters. Finally, as reduced physical activity is typically an inherent characteristic of obesity, further research is required to delineate the mechanisms associated with obesity and physical activity, independently, to the age-associated deterioration of skeletal muscle, as it is often difficult to distinguish between the two.

## Conclusion

The prevalence of sarcopenia, obesity, and sarcopenic-obesity are increasing globally, posing a considerable challenge to public health, healthcare resources, and the global economy. Sarcopenic-obesity is associated with accelerated functional decline, chronic disease risk, and increased risk of mortality. In contrast, increased levels of muscle mass, irrespective of adiposity, are associated with reduced risk of mortality. Despite evidence reporting greater amounts of muscle mass, but not muscle quality, in obese vs. normal weight older individuals in support of an “age-associated obesity paradox,” the current available data are inconclusive. Instead, obesity may impair muscle anabolic processes, potentially through compromised muscle quality and/or elevated levels of adipose-derived inflammatory cytokines. When combined with the age-related loss of muscle mass, this may further exacerbate the loss of muscle function in aging as well as the development of metabolic disease and disability. Older individuals are also likely disproportionately impacted by obesity at a functional level compared with younger obese individuals. The loss of muscle quality likely represents an important contributor to the impairment in physical function in older age. Indeed, intramyocellular lipid accumulation, a common characteristic of obesity, may interfere with contractile force production. Currently, however, the mechanisms through which obesity in older age may protect muscle quantity, yet seemingly impair muscle quality and function are unclear and warrant further investigation, as the complex interplay between obesity and sarcopenia makes it difficult to determine precise cause and effect.

## AUTHOR CONTRIBUTIONS

PM completed the literature searches, review, and drafted the manuscript. PM, BS, and LB reviewed, edited, and approved the manuscript for its intellectual content. All authors contributed to the article and approved the submitted version.

## FUNDING

This work was not sponsored by any funding body external to University of Birmingham. Dunhill Medical Trust support the salary of PM (RPGF1802\1).

## REFERENCES

- Cruz-Jentoft AJ, Bahat G, Bauer J, Boirie Y, Bruyère O, Cederholm T, et al. Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing*. (2019) 48:16–31. doi: 10.1093/ageing/afy169
- Newman AB, Kupelian V, Visser M, Simonsick EM, Goodpaster BH, Kritchevsky SB, et al. Strength, but not muscle mass, is associated with mortality in the health, aging and body composition study cohort. *J Gerontol A Biol Sci Med Sci*. (2006) 61:72–7. doi: 10.1093/gerona/61.1.72
- Metter EJ, Conwit R, Tobin J, Fozard JL. Age-associated loss of power and strength in the upper extremities in women and men. *J Gerontol Ser A Biol Sci Med Sci*. (1997) 52A:B267–76. doi: 10.1093/gerona/52A.5.B267
- Landi F, Cruz-Jentoft AJ, Liperoti R, Russo A, Giovannini S, Tosato M, et al. Sarcopenia and mortality risk in frail older persons aged 80 years and older: results from the SIRENTE study. *Age Ageing*. (2013) 42:203–9. doi: 10.1093/ageing/afs194
- Rosenberg IH. Sarcopenia: origins and clinical relevance. *J Nutr*. (1997) 127:990S–1S. doi: 10.1093/jn/127.5.990S
- Rosenberg IH. Sarcopenia: origins and clinical relevance. *Clin Geriatr Med*. (2011) 127(5 Suppl.):990S–1S. doi: 10.1016/j.cger.2011.03.003
- Walston JD. Sarcopenia in older adults. *Curr Opin Rheumatol*. (2012) 24:623–7. doi: 10.1097/BOR.0b013e328358d59b
- Hunter GR, Gower BA, Kane BL. Age related shift in visceral fat. *Int J Body Compos Res*. (2010) 8:103–8.
- Hunter GR, Singh H, Carter SJ, Bryan DR, Fisher G. Sarcopenia and its implications for metabolic health. *J Obes*. (2019) 2019:1–10. doi: 10.1155/2019/8031705
- Lauretani F, Russo CR, Bandinelli S, Bartali B, Cavazzini C, Di Iorio A, et al. Age-associated changes in skeletal muscles and their effect on mobility: an operational diagnosis of sarcopenia. *J Appl Physiol*. (2003) 95:1851–60. doi: 10.1152/jappphysiol.00246.2003
- Visser M, Newman AB, Nevitt MC, Kritchevsky SB, Stamm EB, Goodpaster BH, et al. Reexamining the sarcopenia hypothesis. Muscle mass versus muscle strength, health, aging, and body composition study research group. *Ann N Y Acad Sci*. (2000) 904:456–61. doi: 10.1111/j.1749-6632.2000.tb06500.x
- Larsson L, Degens H, Li M, Salvati L, Lee Y il, Thompson W, et al. Sarcopenia: aging-related loss of muscle mass and function. *Physiol Rev*. (2019) 99:427–511. doi: 10.1152/physrev.00061.2017
- Lexell J, Taylor CC, Sjöström M. What is the cause of the ageing atrophy? *J Neurol Sci*. (1988) 84:275–94. doi: 10.1016/0022-510X(88)90132-3
- Goodpaster BH, He J, Watkins S, Kelley DE. Skeletal muscle lipid content and insulin resistance: evidence for a paradox in endurance-trained athletes. *J Clin Endocrinol Metab*. (2001) 86:5755–61. doi: 10.1210/jcem.86.12.8075
- Sipilä S, Suominen H. Knee extension strength and walking speed in relation to quadriceps muscle composition and training in elderly women. *Clin Physiol*. (1994) 14:433–42. doi: 10.1111/j.1475-097X.1994.tb00402.x
- Delbono O. Neural control of aging skeletal muscle. *Aging Cell*. (2003) 2:21–9. doi: 10.1046/j.1474-9728.2003.00011.x
- Swinburn BA, Sacks G, Hall KD, McPherson K, Finegood DT, Moodie ML, et al. The global obesity pandemic: shaped by global drivers and local environments. *Lancet*. (2011) 378:804–14. doi: 10.1016/S0140-6736(11)60813-1
- Narici MV, Maffulli N. Sarcopenia: characteristics, mechanisms and functional significance. *Br Med Bull*. (2010) 95:139–59. doi: 10.1093/bmb/ldq008
- Roubenoff R. Sarcopenic obesity: the confluence of two epidemics. *Obes Res*. (2004) 12:887–88. doi: 10.1038/oby.2004.107
- Wright SM, Aronne LJ. Causes of obesity. *Abdom Radiol*. (2012) 37:730–2. doi: 10.1007/s00261-012-9862-x
- Stenholm S, Harris TB, Rantanen T, Visser M, Kritchevsky SB, Ferrucci L. Sarcopenic obesity: definition, cause and consequences. *Curr Opin Clin Nutr Metab Care*. (2008) 11:693–700. doi: 10.1097/MCO.0b013e328312c37d
- Tallis J, James RS, Seebacher F. The effects of obesity on skeletal muscle contractile function. *J Exp Biol*. (2018) 221:jeb163840. doi: 10.1242/jeb.163840
- Kalinkovich A, Livshits G. Sarcopenic obesity or obese sarcopenia: a cross talk between age-associated adipose tissue and skeletal muscle inflammation as a main mechanism of the pathogenesis. *Ageing Res Rev*. (2017) 35:200–21. doi: 10.1016/j.arr.2016.09.008
- Jiang BC, Villareal DT. Weight loss-induced reduction of bone mineral density in older adults with obesity. *J Nutr Gerontol Geriatr*. (2019) 38:100–14. doi: 10.1080/21551197.2018.1564721
- Lustgarten MS. The role of the gut microbiome on skeletal muscle mass and physical function: 2019 update. *Front Physiol*. (2019) 10:1435. doi: 10.3389/fphys.2019.01435
- Baumgartner RN, Koehler KM, Gallagher D, Romero L, Heymsfield SB, Ross RR, et al. Epidemiology of Sarcopenia among the elderly in New Mexico. *Am J Epidemiol*. (1998) 147:755–63. doi: 10.1093/oxfordjournals.aje.a009520
- Campbell MJ, McComas AJ, Petito F. Physiological changes in ageing muscles. *J Neurol Neurosurg Psychiatry*. (1973) 36:174–82. doi: 10.1136/jnnp.36.2.174
- Brown JC, Harhay MO, Harhay MN. Sarcopenia and mortality among a population-based sample of community-dwelling older adults. *J Cachexia Sarcopenia Muscle*. (2016) 7:290–8. doi: 10.1002/jcsm.12073
- Dufour AB, Hannan MT, Murabito JM, Kiel DP, McLean RR. Sarcopenia definitions considering body size and fat mass are associated with mobility limitations: the framingham study. *J Gerontol Ser A Biol Sci Med Sci*. (2013) 68:168–74. doi: 10.1093/gerona/gls109
- Marsh AP, Rejeski WJ, Espeland MA, Miller ME, Church TS, Fielding RA, et al. Muscle strength and BMI as predictors of major mobility disability in the lifestyle interventions and independence for elders pilot (LIFE-P). *J Gerontol Ser A Biol Sci Med Sci*. (2011) 66A:1376–83. doi: 10.1093/gerona/glr158
- Ziaaldini MM, Marzetti E, Picca A, Murlasits Z. Biochemical pathways of sarcopenia and their modulation by physical exercise: a narrative review. *Front Med*. (2017) 4:167. doi: 10.3389/fmed.2017.00167
- Martinez JA. Body-weight regulation: causes of obesity. *Proc Nutr Soc*. (2000) 59:337–45. doi: 10.1017/S0029665100000380
- Calvani R, Miccheli A, Landi F, Bossola M, Cesari M, Leeuwenburgh C, et al. Current nutritional recommendations and novel dietary strategies to manage sarcopenia. *J Frailty Aging*. (2013) 2:38–53.
- Granic A, Mendonça N, Sayer AA, Hill TR, Davies K, Siervo M, et al. Effects of dietary patterns and low protein intake on sarcopenia risk in the very old: the newcastle 85+ study. *Clin Nutr*. (2020) 39:166–73. doi: 10.1016/j.clnu.2019.01.009
- Okamura T, Miki A, Hashimoto Y, Kaji A, Sakai R, Osaka T, et al. Shortage of energy intake rather than protein intake is associated with sarcopenia in elderly patients with type 2 diabetes: a cross-sectional study of the KAMOGAWA-DM cohort. *J Diabetes*. (2019) 11:477–83. doi: 10.1111/1753-0407.12874
- Robinson S, Reginster J, Rizzoli R, Shaw S, Kanis J, Bautmans I, et al. Does nutrition play a role in the prevention and management of sarcopenia? *Clin Nutr*. (2018) 37:1121–32. doi: 10.1016/j.clnu.2017.08.016
- Robinson S, Cooper C, Aihie Sayer A. Nutrition and sarcopenia: a review of the evidence and implications for preventive strategies. *J Aging Res*. (2012) 2012:1–6. doi: 10.1155/2012/510801
- Albuquerque D, Nóbrega C, Manco L, Padez C. The contribution of genetics and environment to obesity. *Br Med Bull*. (2017) 123:159–73. doi: 10.1093/bmb/ldx022

## ACKNOWLEDGMENTS

The authors would like to acknowledge the contributions of BS in the development of **Figure 2**. The authors would also like to thank Dunhill medical trust who support the salary of PM.

39. Cheung WW, Mao P. Recent advances in obesity: genetics and beyond. *ISRN Endocrinol.* (2012) 2012:536905. doi: 10.5402/2012/536905
40. Herrera BM, Lindgren CM. The genetics of obesity. *Curr Diab Rep.* (2010) 10:498–505. doi: 10.1007/s11892-010-0153-z
41. WHO. *Obesity WHO/Europe* (2016). Available online at: <http://www.euro.who.int/en/healthtopics/noncommunicable-diseases/obesity>
42. Baumgartner RN. Body composition in healthy aging. *Ann N Y Acad Sci.* (2000) 904:437–48. doi: 10.1111/j.1749-6632.2000.tb06498.x
43. Goodpaster BH, Theriault R, Watkins SC, Kelley DE. Intramuscular lipid content is increased in obesity and decreased by weight loss. *Metabolism.* (2000) 49:467–72. doi: 10.1016/S0026-0495(00)80010-4
44. Anton SD, Karabetian C, Naugle K, Buford TW. Obesity and diabetes as accelerators of functional decline: can lifestyle interventions maintain functional status in high risk older adults? *Exp Gerontol.* (2013) 48:888–97. doi: 10.1016/j.exger.2013.06.007
45. Jarosz PA, Bellar A. Sarcopenic obesity: an emerging cause of frailty in older adults. *Geriatr Nurs (Minneapolis).* (2009) 30:64–70. doi: 10.1016/j.gerinurse.2008.02.010
46. Park SW, Goodpaster BH, Strotmeyer ES, Kuller LH, Broudeau R, Kammerer C, et al. Accelerated loss of skeletal muscle strength in older adults with type 2 diabetes: the health, aging, and body composition study. *Diabetes Care.* (2007) 30:1507–12. doi: 10.2337/dc06-2537
47. Villareal DT, Banks M, Siener C, Sinacore DR, Klein S. Physical frailty and body composition in obese elderly men and women. *Obes Res.* (2004) 12:913–20. doi: 10.1038/oby.2004.111
48. Lang IA, Llewellyn DJ, Alexander K, Melzer D. Obesity, physical function, and mortality in older adults. *J Am Geriatr Soc.* (2008) 56:1474–8. doi: 10.1111/j.1532-5415.2008.01813.x
49. Figaro MK, Kritchevsky SB, Resnick HE, Shorr RI, Butler J, Shintani A, et al. Diabetes, inflammation, and functional decline in older adults: findings from the health, aging and body composition (ABC) study. *Diabetes Care.* (2006) 29:2039–25. doi: 10.2337/dc06-0245
50. Alley DE, Chang VW. The changing relationship of obesity and disability, 1988–2004. *JAMA.* (2007) 298:2020. doi: 10.1001/jama.298.17.2020
51. Houston DK, Ding J, Nicklas BJ, Harris TB, Lee JS, Nevitt MC, et al. Overweight and obesity over the adult life course and incident mobility limitation in older adults: the health, aging and body composition study. *Am J Epidemiol.* (2009) 169:927–36. doi: 10.1093/aje/kwp007
52. Stenholm S, Rantanen T, Alanen E, Reunanen A, Sainio P, Koskinen S. Obesity history as a predictor of walking limitation at old age. *Obesity.* (2007) 15:929–38. doi: 10.1038/oby.2007.583
53. Muoio DM, Newgard CB. Obesity-related derangements in metabolic regulation. *Annu Rev Biochem.* (2006) 75:367–401. doi: 10.1146/annurev.biochem.75.103004.142512
54. Alberti KGMM, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JI, Donato KA, et al. Harmonizing the metabolic syndrome. *Circulation.* (2009) 120:1640–5. doi: 10.1161/CIRCULATIONAHA.109.192644
55. Frühbeck G, Toplak H, Woodward E, Yumuk W, Maislos M, Oppert J-M. Obesity: the gateway to ill health - an EASO position statement on a rising public health, clinical and scientific challenge in Europe. *Obes Facts.* (2013) 6:117–20. doi: 10.1159/000350627
56. Bauer UE, Briss PA, Goodman RA, Bowman BA. Prevention of chronic disease in the 21st century: elimination of the leading preventable causes of premature death and disability in the USA. *Lancet.* (2014) 384:45–52. doi: 10.1016/S0140-6736(14)60648-6
57. Bischoff SC, Boirie Y, Cederholm T, Chourdakis M, Cuerda C, Delzenne NM, et al. Towards a multidisciplinary approach to understand and manage obesity and related diseases. *Clin Nutr.* (2017) 36:917–38. doi: 10.1016/j.clnu.2016.11.007
58. Daviglus ML, Stamler J, Pirzada A, Yan LL, Garside DB, Liu K, et al. Favorable cardiovascular risk profile in young women and long-term risk of cardiovascular and all-cause mortality. *JAMA.* (2004) 292:1588–92. doi: 10.1001/jama.292.13.1588
59. Lavie CJ, De Schutter A, Parto P, Jahangir E, Kokkinos P, Ortega FB, et al. Obesity and prevalence of cardiovascular diseases and prognosis—the obesity paradox updated. *Prog Cardiovasc Dis.* (2016) 58:537–47. doi: 10.1016/j.pcad.2016.01.008
60. Lavie CJ, Oktay AA, Milani R V. The obesity paradox and obesity severity in elderly STEMI patients. *Eur Hear J Qual Care Clin Outcomes.* (2017) 3:166–7. doi: 10.1093/ehjqcco/qcx018
61. Oktay AA, Lavie CJ, Kokkinos PF, Parto P, Pandey A, Ventura HO. The interaction of cardiorespiratory fitness with obesity and the obesity paradox in cardiovascular disease. *Prog Cardiovasc Dis.* (2017) 60:30–44. doi: 10.1016/j.pcad.2017.05.005
62. Ortega FB, Lavie CJ, Blair SN. Obesity and cardiovascular disease. *Circ Res.* (2016) 118:1752–70. doi: 10.1161/CIRCRESAHA.115.306883
63. Roh E, Choi KM. Health consequences of sarcopenic obesity: a narrative review. *Front Endocrinol.* (2020) 11:332. doi: 10.3389/fendo.2020.00332
64. Martinez-Huenschullán SF, Tam CS, Ban LA, Ehrenfeld-Slater P, McLennan SV, Twigg SM. Skeletal muscle adiponectin induction in obesity and exercise. *Metabolism.* (2020) 102:154008. doi: 10.1016/j.metabol.2019.154008
65. Meex RCR, Blaak EE, Loon LJC. Lipotoxicity plays a key role in the development of both insulin resistance and muscle atrophy in patients with type 2 diabetes. *Obes Rev.* (2019) 20:1205–17. doi: 10.1111/obr.12862
66. Kob R, Bollheimer LC, Bertsch T, Fellner C, Djukic M, Sieber CC, et al. Sarcopenic obesity: molecular clues to a better understanding of its pathogenesis? *Biogerontology.* (2015) 16:15–29. doi: 10.1007/s10522-014-9539-7
67. Levine ME, Crimmins EM. The impact of insulin resistance and inflammation on the association between sarcopenic obesity and physical functioning. *Obesity.* (2012) 20:2101–6. doi: 10.1038/oby.2012.20
68. Baumgartner RN, Stauber PM, McHugh D, Koehler KM, Garry PJ. Cross-sectional age differences in body composition in persons 60 + years of age. *J Gerontol Ser A Biol Sci Med Sci.* (1995) 50A:M307–16. doi: 10.1093/gerona/50A.6.M307
69. Frontera WR, Hughes VA, Fielding RA, Fiatarone MA, Evans WJ, Roubenoff R. Aging of skeletal muscle: A 12-yr longitudinal study. *J Appl Physiol.* (2000) 88:1321–6. doi: 10.1152/jappl.2000.88.4.1321
70. Rantanen T, Masaki K, Foley D, Izmirlian G, White L, Guralnik JM. Grip strength changes over 27 yr in Japanese-American men. *J Appl Physiol.* (1998) 85:2047–53. doi: 10.1152/jappl.1998.85.6.2047
71. Rantanen T, Harris T, Leveille SG, Visser M, Foley D, Masaki K, et al. Muscle strength and body mass index as long-term predictors of mortality in initially healthy men. *J Gerontol Ser A Biol Sci Med Sci.* (2000) 55:168–73. doi: 10.1093/gerona/55.3.M168
72. Goodpaster BH, Krishnaswami S, Harris TB, Katsiaras A, Kritchevsky SB, Simonsick EM, et al. Obesity, regional body fat distribution, and the metabolic syndrome in older men and women. *Arch Intern Med.* (2005) 165:777. doi: 10.1001/archinte.165.7.777
73. Park YW, Zhu S, Palaniappan L, Heshka S, Carnethon MR, Heymsfield SB. The metabolic syndrome: prevalence and associated risk factor findings in the US population from the third national health and nutrition examination survey, 1988–1994. *Arch Intern Med.* (2003) 163:427–36. doi: 10.1001/archinte.163.4.427
74. DeFronzo RA, Jacot E, Jequier E, Maeder E, Wahren J, Felber JP. The effect of insulin on the disposal of intravenous glucose: results from indirect calorimetry and hepatic and femoral venous catheterization. *Diabetes.* (1981) 30:1000–7. doi: 10.2337/diab.30.12.1000
75. Zurlo F, Larson K, Bogardus C, Ravussin E. Skeletal muscle metabolism is a major determinant of resting energy expenditure. *J Clin Invest.* (1990) 86:1423–7. doi: 10.1172/JCI114857
76. Koopman JJE, van Bodegom D, van Heemst D, Westendorp RGJ. Handgrip strength, ageing and mortality in rural Africa. *Age Ageing.* (2015) 44:465–70. doi: 10.1093/ageing/afu165
77. Zhang X, Wang C, Dou Q, Zhang W, Yang Y, Xie X. Sarcopenia as a predictor of all-cause mortality among older nursing home residents: a systematic review and meta-analysis. *BMJ Open.* (2018) 8:e021252. doi: 10.1136/bmjopen-2017-021252
78. Carter CS, Justice JN, Thompson L. Lipotoxicity, aging, and muscle contractility: does fiber type matter? *GeroScience.* (2019) 41:297–308. doi: 10.1007/s11357-019-00077-z
79. da Cunha Nascimento D, da Cunha Oliveira S, Leite Vieira DC, Schwerz Funghetto S, Silva AO, Valduga R, et al. The impact of sarcopenic obesity on inflammation, lean body mass, and muscle strength in elderly women. *Int J Gen Med.* (2018) 11:443–9. doi: 10.2147/IJGM.S187285
80. Collins KH, Herzog W, MacDonald GZ, Reimer RA, Rios JL, Smith IC, et al. Obesity, metabolic syndrome, and musculoskeletal disease: common inflammatory pathways suggest a central role for loss of muscle integrity. *Front Physiol.* (2018) 9:112. doi: 10.3389/fphys.2018.00112

81. Jura M, Kozak LP. Obesity and related consequences to ageing. *Age*. (2016) 38:23. doi: 10.1007/s11357-016-9884-3
82. Aubertin-Leheudre M, Lord C, Goulet ÉDB, Khalil A, Dionne IJ. Effect of sarcopenia on cardiovascular disease risk factors in obese postmenopausal women. *Obesity*. (2006) 14:2277–83. doi: 10.1038/oby.2006.267
83. Dominguez LJ, Barbagallo M. The cardiometabolic syndrome and sarcopenic obesity in older persons. *J Cardiometab Syndr*. (2007) 2:183–9. doi: 10.1111/j.1559-4564.2007.06673.x
84. Batsis JA, Villareal DT. Sarcopenic obesity in older adults: aetiology, epidemiology and treatment strategies. *Nat Rev Endocrinol*. (2018) 14:513–37. doi: 10.1038/s41574-018-0062-9
85. Smeuninx B, Mckendry J, Wilson D, Martin U, Breen L. Age-related anabolic resistance of myofibrillar protein synthesis is exacerbated in obese inactive individuals. *J Clin Endocrinol Metab*. (2017) 102:3535–45. doi: 10.1210/jc.2017-00869
86. Koster A, Ding J, Stenholm S, Caserotti P, Houston DK, Nicklas BJ, et al. Does the amount of fat mass predict age-related loss of lean mass, muscle strength, and muscle quality in older adults? *J Gerontol Ser A Biol Sci Med Sci*. (2011) 66 A:888–95. doi: 10.1093/gerona/66a/10/888
87. Janssen I, Heymsfield SB, Wang Z, Ross R. Skeletal muscle mass and distribution in 468 men and women aged 18–88 yr. *J Appl Physiol*. (2000) 89:81–88. doi: 10.1152/jappl.2000.89.1.81
88. Baumgartner RN, Wayne SJ, Waters DL, Janssen I, Gallagher D, Morley JE. Sarcopenic obesity predicts instrumental activities of daily living disability in the elderly. *Obes Res*. (2004) 12:1995–2004. doi: 10.1038/oby.2004.250
89. Rubbieri G, Mossello E, Di Bari M. Techniques for the diagnosis of sarcopenia. *Clin Cases Miner Bone Metab*. (2014) 11:181–4. doi: 10.11138/ccmbm/2014.11.3.181
90. Fisher G, Windham ST, Griffin P, Warren JL, Gower BA, Hunter GR. Associations of human skeletal muscle fiber type and insulin sensitivity, blood lipids, and vascular hemodynamics in a cohort of premenopausal women. *Eur J Appl Physiol*. (2017) 117:1413–22. doi: 10.1007/s00421-017-3634-9
91. Tanner CJ, Barakat HA, Dohm GL, Pories WJ, MacDonald KG, Cunningham PRG, et al. Muscle fiber type is associated with obesity and weight loss. *Am J Physiol Metab*. (2002) 282:E1191–6. doi: 10.1152/ajpendo.00416.2001
92. Kriketos AD, Pan DA, Lillioja S, Cooney GJ, Baur LA, Milner MR, et al. Interrelationships between muscle morphology, insulin action, and adiposity. *Am J Physiol Integr Comp Physiol*. (1996) 270:R1332–9. doi: 10.1152/ajpregu.1996.270.6.R1332
93. Krotkiewski M, Björntorp P. Muscle tissue in obesity with different distribution of adipose tissue. Effects of physical training. *Int J Obes*. (1986) 10:331–41.
94. Lillioja S, Young AA, Culter CL, Ivy JL, Abbott WG, Zawadzki JK, et al. Skeletal muscle capillary density and fiber type are possible determinants of *in vivo* insulin resistance in man. *J Clin Invest*. (1987) 80:415–24. doi: 10.1172/JCI113088
95. Marin P, Andersson B, Krotkiewski M, Björntorp P. Muscle fiber composition and capillary density in women and men with NIDDM. *Diabetes Care*. (1994) 17:382–6. doi: 10.2337/diacare.17.5.382
96. He J, Watkins S, Kelley DE. Skeletal muscle lipid content and oxidative enzyme activity in relation to muscle fiber type in type 2 diabetes and obesity. *Diabetes*. (2001) 50:817–23. doi: 10.2337/diabetes.50.4.817
97. Kempen KP, Saris WH, Kuipers H, Glatz JF, Van Der Vusse GJ. Skeletal muscle metabolic characteristics before and after energy restriction in human obesity: fibre type, enzymatic beta-oxidative capacity and fatty acid-binding protein content. *Eur J Clin Invest*. (1998) 28:1030–7. doi: 10.1046/j.1365-2362.1998.00383.x
98. Wade AJ, Marbut MM, Round JM. Muscle fibre type and aetiology of obesity. *Lancet*. (1990) 335:805–8. doi: 10.1016/0140-6736(90)90933-V
99. Alway SE, Myers MJ, Mohamed JS. Regulation of satellite cell function in sarcopenia. *Front Aging Neurosci*. (2014) 6:246. doi: 10.3389/fnagi.2014.00246
100. Consitt LA, Bell JA, Houmard JA. Intramuscular lipid metabolism, insulin action, and obesity. *IUBMB Life*. (2009) 61:47–55. doi: 10.1002/iub.142
101. Kelley DE, Goodpaster B, Wing RR, Simoneau JA. Skeletal muscle fatty acid metabolism in association with insulin resistance, obesity, and weight loss. *Am J Physiol Metab*. (1999) 277:E1130–41. doi: 10.1152/ajpendo.1999.277.6.E1130
102. Simoneau JA, Veerkamp JH, Turcotte LP, Kelley DE. Markers of capacity to utilize fatty acids in human skeletal muscle: relation to insulin resistance and obesity and effects of weight loss. *FASEB J*. (1999) 13:2051–60. doi: 10.1096/fasebj.13.14.2051
103. Simoneau JA, Bouchard C. Skeletal muscle metabolism and body fat content in men and women. *Obes Res*. (1995) 3:23–29. doi: 10.1002/j.1550-8528.1995.tb00117.x
104. English KL, Paddon-Jones D. Protecting muscle mass and function in older adults during bed rest. *Curr Opin Clin Nutr Metab Care*. (2010) 13:34–39. doi: 10.1097/MCO.0b013e328333aa66
105. Chuang SY, Hsu YY, Chen RCY, Liu WL, Pan WH. Abdominal obesity and low skeletal muscle mass jointly predict total mortality and cardiovascular mortality in an elderly asian population. *J Gerontol Ser A Biol Sci Med Sci*. (2016) 71:1049–55. doi: 10.1093/gerona/66a/10/1049
106. Mathus-Vliegen EMH. Obesity and the elderly. *J Clin Gastroenterol*. (2012) 46:533–44. doi: 10.1097/MCG.0b013e32825692ce
107. Pataky Z, Armand S, Müller-Pinget S, Golay A, Allet L. Effects of obesity on functional capacity. *Obesity*. (2014) 22:56–62. doi: 10.1002/oby.20514
108. Moore BA, Bembem DA, Lein DH, Bembem MG, Singh H. Fat mass is negatively associated with muscle strength and jump test performance. *J Frailty Aging*. (2020) 9:214–18. doi: 10.14283/jfa.2020.11
109. Choi SJ, Files DC, Zhang T, Wang Z-M, Messi ML, Gregory H, et al. Intramyocellular lipid and impaired myofiber contraction in normal weight and obese older adults. *J Gerontol Ser A Biol Sci Med Sci*. (2016) 71:557–64. doi: 10.1093/gerona/66a/10/557
110. Trommelen J, Holwerda AM, Kouw IWK, Langer H, Halson SL, Rollo I, et al. Resistance exercise augments postprandial overnight muscle protein synthesis rates. *Med Sci Sport Exerc*. (2016) 48:2517–25. doi: 10.1249/MSS.0000000000001045
111. Moro C, Bajpeyi S, Smith SR. Determinants of intramyocellular triglyceride turnover: implications for insulin sensitivity. *Am J Physiol Metab*. (2008) 294:E203–13. doi: 10.1152/ajpendo.00624.2007
112. van Loon LJC, Goodpaster BH. Increased intramuscular lipid storage in the insulin-resistant and endurance-trained state. *Pflügers Arch*. (2006) 451:606–16. doi: 10.1007/s00424-005-1509-0
113. Tardif N, Salles J, Guillet C, Tordjman J, Reggio S, Landrier JF, et al. Muscle ectopic fat deposition contributes to anabolic resistance in obese sarcopenic old rats through eIF2 $\alpha$  activation. *Aging Cell*. (2014) 13:1001–11. doi: 10.1111/acel.12263
114. Bogdanis GC. Effects of physical activity and inactivity on muscle fatigue. *Front Physiol*. (2012) 3:142. doi: 10.3389/fphys.2012.00142
115. Heo JW, Yoo SZ, No MH, Park DH, Kang JH, Kim TW, et al. Exercise training attenuates obesity-induced skeletal muscle remodeling and mitochondria-mediated apoptosis in the skeletal muscle. *Int J Environ Res Public Health*. (2018) 15:2301. doi: 10.3390/ijerph15102301
116. Kras KA, Langlais PR, Hoffman N, Roust LR, Benjamin TR, De Filippis EA, et al. Obesity modifies the stoichiometry of mitochondrial proteins in a way that is distinct to the subcellular localization of the mitochondria in skeletal muscle. *Metabolism*. (2018) 89:18–26. doi: 10.1016/j.metabol.2018.09.006
117. Kristensen MD, Petersen SM, Møller KE, Lund MT, Hansen M, Hansen CN, et al. Obesity leads to impairments in the morphology and organization of human skeletal muscle lipid droplets and mitochondrial networks, which are resolved with gastric bypass surgery-induced improvements in insulin sensitivity. *Acta Physiol*. (2018) 224:e13100. doi: 10.1111/apha.13100
118. Tomlinson DJ, Erskine RM, Morse CI, Winwood K, Onambélé-Pearson G. The impact of obesity on skeletal muscle strength and structure through adolescence to old age. *Biogerontology*. (2016) 17:467–83. doi: 10.1007/s10522-015-9626-4
119. Hill C, James R, Cox V, Tallis J. Does dietary-induced obesity in old age impair the contractile performance of isolated mouse soleus, extensor digitorum longus and diaphragm skeletal muscles? *Nutrients*. (2019) 11:505. doi: 10.3390/nu11030505
120. Eshima H, Tamura Y, Kakehi S, Kurebayashi N, Murayama T, Nakamura K, et al. Long-term, but not short-term high-fat diet induces fiber composition changes and impaired contractile force in mouse fast-twitch skeletal muscle. *Physiol Rep*. (2017) 5:e13250. doi: 10.14814/phy2.13250

121. Eshima H, Tamura Y, Kakehi S, Kakigi R, Hashimoto R, Funai K, et al. A chronic high-fat diet exacerbates contractile dysfunction with impaired intracellular Ca<sup>2+</sup> release capacity in the skeletal muscle of aged mice. *J Appl Physiol.* (2020) 128:1153–62. doi: 10.1152/jappphysiol.00530.2019
122. Gale CR, Martyn CN, Cooper C, Sayer AA. Grip strength, body composition, and mortality. *Int J Epidemiol.* (2007) 36:228–35. doi: 10.1093/ije/dyl224
123. Rantanen T, Volpato S, Ferrucci L, Heikkinen E, Fried LP, Guralnik JM. Handgrip strength and cause-specific and total mortality in older disabled women: exploring the mechanism. *J Am Geriatr Soc.* (2003) 51:636–41. doi: 10.1034/j.1600-0579.2003.00207.x
124. Rantanen T. Muscle strength, disability and mortality. *Scand J Med Sci Sport.* (2003) 13:3–8. doi: 10.1034/j.1600-0838.2003.00298.x
125. Flegal KM, Kit BK, Orpana H, Graubard BI. Association of all-cause mortality with overweight and obesity using standard body mass index categories: a systematic review and meta-analysis. *JAMA.* (2013) 309:71–82. doi: 10.1001/jama.2012.113905
126. Flegal KM, Graubard BI, Williamson DF, Gail MH. Excess deaths associated with underweight, overweight, and obesity. *JAMA.* (2005) 293:1861–7. doi: 10.1001/jama.293.15.1861
127. Zamboni M, Mazzali G, Zoico E, Harris TB, Meigs JB, Di Francesco V, et al. Health consequences of obesity in the elderly: a review of four unresolved questions. *Int J Obes.* (2005) 29:1011–29. doi: 10.1038/sj.ijo.0803005
128. Gruberg L, Weissman NJ, Waksman R, Fuchs S, Deible R, Pinnow EE, et al. The impact of obesity on the short-term and long-term outcomes after percutaneous coronary intervention: the obesity paradox? *J Am Coll Cardiol.* (2002) 39:578–84. doi: 10.1016/S0735-1097(01)01802-2
129. Li S, Wang Z, Huang J, Fan J, Du H, Liu L, et al. Systematic review of prognostic roles of body mass index for patients undergoing lung cancer surgery: does the ‘obesity paradox’ really exist? *Eur J Cardio Thoracic Surg.* (2016) 51:ezw386. doi: 10.1093/ejcts/ezw386
130. Aune D, Sen A, Prasad M, Norat T, Janszky I, Tonstad S, et al. BMI and all cause mortality: systematic review and non-linear dose-response meta-analysis of 230 cohort studies with 3.74 million deaths among 30.3 million participants. *BMJ.* (2016) 353:i2156. doi: 10.1136/bmj.i2156
131. Calle EE, Thun MJ, Petrelli JM, Rodriguez C, Heath CW. Body-mass index and mortality in a prospective cohort of U.S. Adults. *N Engl J Med.* (1999) 341:1097–105. doi: 10.1056/NEJM199910073411501
132. Bowman K, Delgado J, Henley WE, Masoli JA, Kos K, Brayne C, et al. Obesity in older people with and without conditions associated with weight loss: follow-up of 955,000 primary care patients. *J Gerontol Ser A Biol Sci Med Sci.* (2017) 72:203–9. doi: 10.1093/gerona/glw147
133. Carnethon MR, De Chavez PJD, Biggs ML, Lewis CE, Pankow JS, Bertoni AG, et al. Association of weight status with mortality in adults with incident diabetes. *JAMA.* (2012) 308:581–90. doi: 10.1001/jama.2012.9282
134. Costanzo P, Cleland JGF, Pellicori P, Clark AL, Hepburn D, Kilpatrick ES, et al. The obesity paradox in type 2 diabetes mellitus: relationship of body mass index to prognosis. *Ann Intern Med.* (2015) 162:610–8. doi: 10.7326/M14-1551
135. Mesinovic J, McMillan L, Shore-Lorenti C, De Courten B, Ebeling P, Scott D. Metabolic syndrome and its associations with components of sarcopenia in overweight and obese older adults. *J Clin Med.* (2019) 8:145. doi: 10.3390/jcm8020145
136. Prado CM, Gonzalez MC, Heymsfield SB. Body composition phenotypes and obesity paradox. *Curr Opin Clin Nutr Metab Care.* (2015) 18:535–51. doi: 10.1097/MCO.0000000000000216
137. Abramowitz MK, Hall CB, Amodu A, Sharma D, Androga L, Hawkins M. Muscle mass, BMI, and mortality among adults in the United States: a population-based cohort study. *PLoS ONE.* (2018) 13:e0194697. doi: 10.1371/journal.pone.0194697
138. Lee DH, Keum N, Hu FB, Orav EJ, Rimm EB, Willett WC, et al. Predicted lean body mass, fat mass, and all cause and cause specific mortality in men: prospective US cohort study. *BMJ.* (2018) 362:k2575. doi: 10.1136/bmj.k2575
139. Schaap LA, Pluijm SMF, Deeg DJH, Visser M. Inflammatory markers and loss of muscle mass (sarcopenia) and strength. *Am J Med.* (2006) 119:526.e9–e17. doi: 10.1016/j.amjmed.2005.10.049
140. Cespedes Feliciano EM, Kroenke CH, Caan BJ. The obesity paradox in cancer: how important is muscle? *Annu Rev Nutr.* (2018) 38:357–79. doi: 10.1146/annurev-nutr-082117-051723
141. Hulston CJ, Woods RM, Dewhurst-Trigg R, Parry SA, Gagnon S, Baker L, et al. Resistance exercise stimulates mixed muscle protein synthesis in lean and obese young adults. *Physiol Rep.* (2018) 6:1–11. doi: 10.14814/phy2.13799
142. Breen L, Stokes KA, Churchward-Venne TA, Moore DR, Baker SK, Smith K, et al. Two weeks of reduced activity decreases leg lean mass and induces “anabolic resistance” of myofibrillar protein synthesis in healthy elderly. *J Clin Endocrinol Metab.* (2013) 98:2604–12. doi: 10.1210/jc.2013-1502
143. Breen L, Phillips SM. Interactions between exercise and nutrition to prevent muscle waste during ageing. *Br J Clin Pharmacol.* (2013) 75:708–15. doi: 10.1111/j.1365-2125.2012.04456.x
144. Drummond MJ, Dickinson JM, Fry CS, Walker DK, Gundermann DM, Reidy PT, et al. Bed rest impairs skeletal muscle amino acid transporter expression, mTORC1 signaling, and protein synthesis in response to essential amino acids in older adults. *Am J Physiol Metab.* (2012) 302:E1113–22. doi: 10.1152/ajpendo.00603.2011
145. Glover EI, Phillips SM, Oates BR, Tang JE, Tarnopolsky MA, Selby A, et al. Immobilization induces anabolic resistance in human myofibrillar protein synthesis with low and high dose amino acid infusion. *J Physiol.* (2008) 586:6049–61. doi: 10.1113/jphysiol.2008.160333
146. Volpi E, Nazemi R, Fujita S. Muscle tissue changes with aging. *Curr Opin Clin Nutr Metab Care.* (2004) 7:405–10. doi: 10.1097/01.mco.0000134362.76653.b2
147. Wall BT, Dirks ML, Snijders T, Senden JMG, Dolmans J, van Loon LJC. Substantial skeletal muscle loss occurs during only 5 days of disuse. *Acta Physiol.* (2014) 210:600–11. doi: 10.1111/apha.12190
148. Biolo G, Maggi SP, Williams BD, Tipton KD, Wolfe RR. Increased rates of muscle protein turnover and amino acid transport after resistance exercise in humans. *Am J Physiol Metab.* (1995) 268:E514–20. doi: 10.1152/ajpendo.1995.268.3.E514
149. Wolfe RR. The underappreciated role of muscle in health and disease. *Am J Clin Nutr.* (2006) 84:475–82. doi: 10.1093/ajcn/84.3.475
150. Zhang X, Chinkes DL, Wolfe RR. Measurement of muscle protein fractional synthesis and breakdown rates from a pulse tracer injection. *Am J Physiol Metab.* (2002) 283:E753–64. doi: 10.1152/ajpendo.00053.2002
151. Atherton PJ, Smith K. Muscle protein synthesis in response to nutrition and exercise. *J Physiol.* (2012) 590:1049–57. doi: 10.1113/jphysiol.2011.225003
152. Wilkinson DJ, Piasecki M, Atherton PJ. The age-related loss of skeletal muscle mass and function: measurement and physiology of muscle fibre atrophy and muscle fibre loss in humans. *Ageing Res Rev.* (2018) 47:123–32. doi: 10.1016/j.arr.2018.07.005
153. Wilkinson DJ, Bukhari SSI, Phillips BE, Limb MC, Cegielski J, Brook MS, et al. Effects of leucine-enriched essential amino acid and whey protein bolus dosing upon skeletal muscle protein synthesis at rest and after exercise in older women. *Clin Nutr.* (2018) 37:2011–21. doi: 10.1016/j.clnu.2017.09.008
154. Phillips SM, Tang JE, Moore DR. The role of milk- and soy-based protein in support of muscle protein synthesis and muscle protein accretion in young and elderly persons. *J Am Coll Nutr.* (2009) 28:343–54. doi: 10.1080/07315724.2009.10718096
155. Atherton PJ, Etheridge T, Watt PW, Wilkinson D, Selby A, Rankin D, et al. Muscle full effect after oral protein: time-dependent concordance and discordance between human muscle protein synthesis and mTORC1 signaling. *Am J Clin Nutr.* (2010) 92:1080–8. doi: 10.3945/ajcn.2010.29819
156. Breen L, Phillips SM. Skeletal muscle protein metabolism in the elderly: Interventions to counteract the “anabolic resistance” of ageing. *Nutr Metab.* (2011) 8:68. doi: 10.1186/1743-7075-8-68
157. Cuthbertson D, Smith K, Babraj J, Leese G, Waddell T, Atherton P, et al. Anabolic signaling deficits underlie amino acid resistance of wasting, aging muscle. *FASEB J.* (2005) 19:422–4. doi: 10.1096/fj.04-2640fje
158. Cuthbertson DJ, Babraj J, Smith K, Wilkes E, Fedele MJ, Esser K, et al. Anabolic signaling and protein synthesis in human skeletal muscle after dynamic shortening or lengthening exercise. *Am J Physiol Metab.* (2006) 290:E731–8. doi: 10.1152/ajpendo.00415.2005
159. Miller BF, Olesen JL, Hansen M, Døssing S, Cramer RM, Welling RJ, et al. Coordinated collagen and muscle protein synthesis in human patella tendon and quadriceps muscle after exercise. *J Physiol.* (2005) 567:1021–33. doi: 10.1113/jphysiol.2005.093690
160. Phillips SM, Tipton KD, Aarsland A, Wolf SE, Wolfe RR. Mixed muscle protein synthesis and breakdown after resistance exercise in humans. *Am J Physiol.* (1997) 273:E99–107. doi: 10.1152/ajpendo.1997.273.1.E99

161. Kumar V, Atherton PJ, Selby A, Rankin D, Williams J, Smith K, et al. Muscle protein synthetic responses to exercise: effects of age, volume, and intensity. *J Gerontol Ser A Biol Sci Med Sci.* (2012) 67:1170–7. doi: 10.1093/gerona/gls141
162. Breen L, Phillips SM. Nutrient interaction for optimal protein anabolism in resistance exercise. *Curr Opin Clin Nutr Metab Care.* (2012) 15:226–32. doi: 10.1097/MCO.0b013e3283516850
163. Kumar V, Selby A, Rankin D, Patel R, Atherton P, Hildebrandt W, et al. Age-related differences in the dose-response relationship of muscle protein synthesis to resistance exercise in young and old men. *J Physiol.* (2009) 587:211–17. doi: 10.1113/jphysiol.2008.164483
164. Morton RW, McGlory C, Phillips SM. Nutritional interventions to augment resistance training-induced skeletal muscle hypertrophy. *Front Physiol.* (2015) 6:245. doi: 10.3389/fphys.2015.00245
165. Moore DR. Keeping Older Muscle “Young” through Dietary. *Proc IUNS 20th Int Congr Nutr.* (2014) 5:599–607S. doi: 10.3945/an.113.005405
166. Morton RW, Murphy KT, McKellar SR, Schoenfeld BJ, Henselmans M, Helms E, et al. A systematic review, meta-analysis and meta-regression of the effect of protein supplementation on resistance training-induced gains in muscle mass and strength in healthy adults. *Br J Sports Med.* (2018) 52:376–84. doi: 10.1136/bjsports-2017-097608
167. Moore DR, Churchward-Venne TA, Witard O, Breen L, Burd NA, Tipton KD, et al. Protein ingestion to stimulate myofibrillar protein synthesis requires greater relative protein intakes in healthy older versus younger men. *J Gerontol Ser A Biol Sci Med Sci.* (2015) 70:57–62. doi: 10.1093/gerona/glu103
168. Brook MS, Wilkinson DJ, Phillips BE, Perez-Schindler J, Philp A, Smith K, et al. Skeletal muscle homeostasis and plasticity in youth and ageing: impact of nutrition and exercise. *Acta Physiol.* (2016) 216:15–41. doi: 10.1111/apha.12532
169. Wilkes EA, Selby AL, Atherton PJ, Patel R, Rankin D, Smith K, et al. Blunting of insulin inhibition of proteolysis in legs of older subjects may contribute to age-related sarcopenia. *Am J Clin Nutr.* (2009) 90:1343–50. doi: 10.3945/ajcn.2009.27543
170. Katsanos CS, Kobayashi H, Sheffield-Moore M, Aarsland A, Wolfe RR. Aging is associated with diminished accretion of muscle proteins after the ingestion of a small bolus of essential amino acids. *Am J Clin Nutr.* (2005) 82:1065–73. doi: 10.1093/ajcn/82.5.1065
171. Wall BT, Gorissen SH, Pennings B, Koopman R, Groen BBL, Verdijk LB, et al. Aging is accompanied by a blunted muscle protein synthetic response to protein ingestion. *PLoS ONE.* (2015) 10:e0140903. doi: 10.1371/journal.pone.0140903
172. Song Z, Moore DR, Hodson N, Ward C, Dent JR, O’Leary MF, et al. Resistance exercise initiates mechanistic target of rapamycin (mTOR) translocation and protein complex co-localisation in human skeletal muscle. *Sci Rep.* (2017) 7:5028. doi: 10.1038/s41598-017-05483-x
173. Johnson SC, Rabinovitch PS, Kaeblerlein M. mTOR is a key modulator of ageing and age-related disease. *Nature.* (2013) 493:338–45. doi: 10.1038/nature11861
174. Hands SL, Proud CG, Wytenbach A. mTOR’s role in ageing: protein synthesis or autophagy? *Ageing.* (2009) 1:586–97. doi: 10.18632/ageing.100070
175. Laplante M, Sabatini DM. mTOR signaling in growth control and disease. *Cell.* (2012) 149:274–93. doi: 10.1016/j.cell.2012.03.017
176. Hodson N, McGlory C, Oikawa SY, Jeromson S, Song Z, Rüegg MA, et al. Differential localization and anabolic responsiveness of mTOR complexes in human skeletal muscle in response to feeding and exercise. *Am J Physiol Physiol.* (2017) 313:C604–11. doi: 10.1152/ajpcell.00176.2017
177. Hodson N, Philp A. The importance of mTOR trafficking for human skeletal muscle translational control. *Exerc Sport Sci Rev.* (2019) 47:46–53. doi: 10.1249/JES.0000000000000173
178. Brack AS, Muñoz-Cánoves P. The ins and outs of muscle stem cell aging. *Skelet Muscle.* (2015) 6:1. doi: 10.1186/s13395-016-0072-z
179. Pallafacchina G, François S, Regnault B, Czarny B, Dive V, Cumano A, et al. An adult tissue-specific stem cell in its niche: a gene profiling analysis of *in vivo* quiescent and activated muscle satellite cells. *Stem Cell Res.* (2010) 4:77–91. doi: 10.1016/j.scr.2009.10.003
180. Decaroli MC, Rochira V. Aging and sex hormones in males. *Virulence.* (2017) 8:545–70. doi: 10.1080/21505594.2016.1259053
181. Boucher BJ. Is vitamin D status relevant to metabolic syndrome? *Dermatoendocrinol.* (2012) 4:212–24. doi: 10.4161/derm.20012
182. Meehan M, Penckofer S. The role of vitamin D in the aging adult. *J Aging Gerontol.* (2014) 2:60–71. doi: 10.12974/2309-6128.2014.02.02.1
183. Braga M, Simmons Z, Norris KC, Ferrini MG, Artaza JN. Vitamin D induces myogenic differentiation in skeletal muscle derived stem cells. *Endocr Connect.* (2017) 6:139–50. doi: 10.1530/EC-17-0008
184. Scimeca M, Centofanti F, Celi M, Gasbarra E, Novelli G, Botta A, et al. Vitamin D receptor in muscle atrophy of elderly patients: a key element of osteoporosis-sarcopenia connection. *Ageing Dis.* (2018) 9:952. doi: 10.14336/AD.2018.0215
185. Roth SM, Zmuda JM, Cauley JA, Shea PR, Ferrell RE. Vitamin D receptor genotype is associated with fat-free mass and sarcopenia in elderly men. *J Gerontol Ser A Biol Sci Med Sci.* (2004) 59:10–B15. doi: 10.1093/gerona/59.1.B10
186. Koww IWK, Van Dijk JW, Horstman AMH, Kramer IF, Goessens JPB, Van Dielen FMH, et al. Basal and postprandial myofibrillar protein synthesis rates do not differ between lean and obese middle-aged men. *J Nutr.* (2019) 149:1533–42. doi: 10.1093/jn/nxz104
187. Beals JW, Sukiennik RA, Nallabelli J, Emmons RS, van Vliet S, Young JR, et al. Anabolic sensitivity of postprandial muscle protein synthesis to the ingestion of a protein-dense food is reduced in overweight and obese young adults. *Am J Clin Nutr.* (2016) 104:1014–22. doi: 10.3945/ajcn.116.130385
188. Murton AJ, Marimuthu K, Mallinson JE, Selby AL, Smith K, Rennie MJ, et al. Obesity appears to be associated with altered muscle protein synthetic and breakdown responses to increased nutrient delivery in older men, but not reduced muscle mass or contractile function. *Diabetes.* (2015) 64:3160–71. doi: 10.2337/db15-0021
189. Castro AM, Macedo-de la Concha LE, Pantoja-Meléndez CA. Low-grade inflammation and its relation to obesity and chronic degenerative diseases. *Rev Médica del Hosp Gen México.* (2017) 80:101–5. doi: 10.1016/j.hgmx.2016.06.011
190. Monteiro R, Azevedo I. Chronic inflammation in obesity and the metabolic syndrome. *Mediators Inflamm.* (2010) 2010:1–10. doi: 10.1155/2010/289645
191. Rodríguez-Hernández H, Simental-Mendía LE, Rodríguez-Ramírez G, Reyes-Romero MA. Obesity and inflammation: epidemiology, risk factors, and markers of inflammation. *Int J Endocrinol.* (2013) 2013:1–11. doi: 10.1155/2013/678159
192. Dalle S, Rossmeislova L, Kopko K. The role of inflammation in age-related sarcopenia. *Front Physiol.* (2017) 8:1045. doi: 10.3389/fphys.2017.01045
193. Ellulu MS, Patimah I, Khaza’ai H, Rahmat A, Abed Y. Obesity and inflammation: the linking mechanism and the complications. *Arch Med Sci.* (2017) 4:851–63. doi: 10.5114/aoms.2016.58928
194. Erskine RM, Tomlinson DJ, Morse CI, Winwood K, Hampson P, Lord JM, et al. The inflammatory and combined effects of obesity- and ageing-induced systemic inflammation on human skeletal muscle properties. *Int J Obes.* (2017) 41:102–11. doi: 10.1038/ijo.2016.151
195. Haran PH, Rivas DA, Fielding RA. Role and potential mechanisms of anabolic resistance in sarcopenia. *J Cachexia Sarcopenia Muscle.* (2012) 3:157–62. doi: 10.1007/s13539-012-0068-4
196. Katsanos CS, Mandarino LJ. Protein metabolism in human obesity: a shift in focus from whole-body to skeletal muscle. *Obesity.* (2011) 19:469–75. doi: 10.1038/oby.2010.290
197. Londhe P, Guttridge DC. Inflammation induced loss of skeletal muscle. *Bone.* (2015) 80:131–42. doi: 10.1016/j.bone.2015.03.015
198. Lumeng CN, Saltiel AR. Inflammatory links between obesity and metabolic disease. *J Clin Invest.* (2011) 121:2111–17. doi: 10.1172/JCI57132
199. Guillet C, Masgrau A, Walrand S, Boirie Y. Impaired protein metabolism: interlinks between obesity, insulin resistance and inflammation. *Obes Rev.* (2012) 13:51–57. doi: 10.1111/j.1467-789X.2012.01037.x
200. Beals JW, Skinner SK, McKenna CF, Poozhikunnel EG, Farooqi SA, van Vliet S, et al. Altered anabolic signalling and reduced stimulation of myofibrillar protein synthesis after feeding and resistance exercise in people with obesity. *J Physiol.* (2018) 596:5119–33. doi: 10.1113/JP276210
201. Beals JW, Burd NA, Moore DR, van Vliet S. Obesity alters the muscle protein synthetic response to nutrition and exercise. *Front Nutr.* (2019) 6:87. doi: 10.3389/fnut.2019.00087
202. James WP, Davies HL, Bales J, Dauncey MJ. Elevated metabolic rates in obesity. *Lancet.* (1978) 1:1122–5. doi: 10.1016/S0140-6736(78)90300-8
203. Garcia-Vicencio S, Coudeyre E, Kluka V, Cardenoux C, Jegu AG, Fourot AV, et al. The bigger, the stronger? Insights from muscle architecture and

- nervous characteristics in obese adolescent girls. *Int J Obes.* (2016) 40:245–51. doi: 10.1038/ijo.2015.158
204. Stephens FB, Chee C, Wall BT, Murton AJ, Shannon CE, van Loon LJC, et al. Lipid-induced insulin resistance is associated with an impaired skeletal muscle protein synthetic response to amino acid ingestion in healthy young men. *Diabetes.* (2015) 64:1615–20. doi: 10.2337/db14-0961
205. Guillet C, Delcourt I, Rance M, Giraudet C, Walrand S, Bedu M, et al. Changes in basal and insulin and amino acid response of whole body and skeletal muscle proteins in obese men. *J Clin Endocrinol Metab.* (2009) 94:3044–50. doi: 10.1210/jc.2008-2216
206. Hill C, Tallis J. Is obesity a risk factor for skeletal muscle ageing? *Aging.* (2019) 11:2183–4. doi: 10.18632/aging.101941
207. Anderson SR, Gilge DA, Steiber AL, Previs SF. Diet-induced obesity alters protein synthesis: tissue-specific effects in fasted versus fed mice. *Metabolism.* (2008) 57:347–54. doi: 10.1016/j.metabol.2007.10.009
208. Chevalier S, Marliss EB, Morais JA, Lamarche M, Gougeon R. Whole-body protein anabolic response is resistant to the action of insulin in obese women. *Am J Clin Nutr.* (2005) 82:355–65. doi: 10.1093/ajcn/82.2.355
209. Rieu I, Magne H, Savary-Auzeloux I, Averous J, Bos C, Peyron MA, et al. Reduction of low grade inflammation restores blunting of postprandial muscle anabolism and limits sarcopenia in old rats. *J Physiol.* (2009) 587:5483–92. doi: 10.1113/jphysiol.2009.178319
210. Akhmedov D, Berdeaux R. The effects of obesity on skeletal muscle regeneration. *Front Physiol.* (2013) 4:371. doi: 10.3389/fphys.2013.00371
211. Corcoran MP, Lamon-Fava S, Fielding RA. Skeletal muscle lipid deposition and insulin resistance: Effect of dietary fatty acids and exercise. *Am J Clin Nutr.* (2007) 85:662–77. doi: 10.1093/ajcn/85.3.662
212. Dubé JJ, Amati F, Stefanovic-Racic M, Toledo FGS, Sauers SE, Goodpaster BH. Exercise-induced alterations in intramyocellular lipids and insulin resistance: the athlete's paradox revisited. *Am J Physiol Endocrinol Metab.* (2008) 294:882–8. doi: 10.1152/ajpendo.00769.2007
213. Morales PE, Bucarey JL, Espinosa A. Muscle lipid metabolism: role of lipid droplets and perilipins. *J Diabetes Res.* (2017) 2017:1789395. doi: 10.1155/2017/1789395
214. Nielsen J, Christensen AE, Nellemann B, Christensen B. Lipid droplet size and location in human skeletal muscle fibers are associated with insulin sensitivity. *Am J Physiol Metab.* (2017) 313:E721–30. doi: 10.1152/ajpendo.00062.2017
215. Shepherd SO, Cocks M, Meikle PJ, Mellett NA, Ranasinghe AM, Barker TA, et al. Lipid droplet remodelling and reduced muscle ceramides following sprint interval and moderate-intensity continuous exercise training in obese males. *Int J Obes.* (2017) 41:1745–54. doi: 10.1038/ijo.2017.170
216. Strissel KJ, Stancheva Z, Miyoshi H, Perfield JW, DeFuria J, Jick Z, et al. Adipocyte death, adipose tissue remodeling, and obesity complications. *Diabetes.* (2007) 56:2910–18. doi: 10.2337/db07-0767
217. Coen PM, Goodpaster BH. Role of intramyocellular lipids in human health. *Trends Endocrinol Metab.* (2012) 23:391–8. doi: 10.1016/j.tem.2012.05.009
218. Pietiläinen KH, Kaprio J, Borg P, Plasqui G, Yki-Järvinen H, Kujala UM, et al. Physical inactivity and obesity: a vicious circle. *Obesity.* (2008) 16:409–14. doi: 10.1038/oby.2007.72
219. Batsis JA, Gill LE, Masutani RK, Adachi-Mejia AM, Blunt HB, Bagley PJ, et al. Weight loss interventions in older adults with obesity: a systematic review of randomized controlled trials since 2005. *J Am Geriatr Soc.* (2017) 65:257–68. doi: 10.1111/jgs.14514
220. Bouchonville MF, Villareal DT. Sarcopenic obesity - how do we treat it? *Curr Opin Endocrinol Diabetes Obes.* (2013) 20:412–19. doi: 10.1097/01.med.0000433071.11466.7f
221. Gill LE, Bartels SJ, Batsis JA. Weight management in older adults. *Curr Obes Rep.* (2015) 4:379–88. doi: 10.1007/s13679-015-0161-z
222. Scott D, Seibel MJ, Cumming R, Naganathan V, Blyth F, Le Couteur DG, et al. Associations of body composition trajectories with bone mineral density, muscle function, falls, and fractures in older men: the concord health and ageing in men project. *J Gerontol Ser A.* (2020) 75:939–45. doi: 10.1093/gerona/ggz184
223. Weinheimer EM, Sands LP, Campbell WW. A systematic review of the separate and combined effects of energy restriction and exercise on fat-free mass in middle-aged and older adults: implications for sarcopenic obesity. *Nutr Rev.* (2010) 68:375–88. doi: 10.1111/j.1753-4887.2010.00298.x
224. Jensen GL, Roy MA, Buchanan AE, Berg MB. Weight loss intervention for obese older women: improvements in performance and function. *Obes Res.* (2004) 12:1814–20. doi: 10.1038/oby.2004.225
225. Jensen GL, Hsiao PY. Systematic review finds modest weight loss at 1 year but a lack of high-quality evidence to support the efficacy of programmes encouraging weight loss in older people. *Evid Based Med.* (2010) 15:41–42. doi: 10.1136/ebm1035
226. Miller GD, Nicklas BJ, Davis C, Loeser RF, Lenchik L, Messier SP. Intensive weight loss program improves physical function in older obese adults with knee osteoarthritis. *Obesity.* (2006) 14:1219–30. doi: 10.1038/oby.2006.139
227. Frimel TN, Sinacore DR, Villareal DT. Exercise attenuates the weight-loss-induced reduction in muscle mass in frail obese older adults. *Med Sci Sports Exerc.* (2008) 40:1213–19. doi: 10.1249/MSS.0b013e31816a85ce
228. Wang X, Miller GD, Messier SP, Nicklas BJ. Knee strength maintained despite loss of lean body mass during weight loss in older obese adults with knee osteoarthritis. *J Gerontol Ser A Biol Sci Med Sci.* (2007) 62:866–71. doi: 10.1093/gerona/62.8.866
229. Davidson LE, Hudson R, Kilpatrick K, Kuk JL, McMillan K, Janiszewski PM, et al. Effects of exercise modality on insulin resistance and functional limitation in older adults. *Arch Intern Med.* (2009) 169:122–31. doi: 10.1001/archinternmed.2008.558
230. Brook MS, Wilkinson DJ, Smith K, Atherton PJ. The metabolic and temporal basis of muscle hypertrophy in response to resistance exercise. *Eur J Sport Sci.* (2016) 16:633–44. doi: 10.1080/17461391.2015.1073362
231. McLeod M, Breen L, Hamilton DL, Philp A. Live strong and prosper: the importance of skeletal muscle strength for healthy ageing. *Biogerontology.* (2016) 17:497–510. doi: 10.1007/s10522-015-9631-7
232. Willis LH, Slentz CA, Bateman LA, Shields AT, Piner LW, Bales CW, et al. Effects of aerobic and/or resistance training on body mass and fat mass in overweight or obese adults. *J Appl Physiol.* (2012) 113:1831–7. doi: 10.1152/jappphysiol.01370.2011
233. Funk MD, Lee M, Vidoni ML, Reininger BM. Weight loss and weight gain among participants in a community-based weight loss challenge. *BMC Obes.* (2019) 6:2. doi: 10.1186/s40608-018-0225-1
234. Marchesini G, Montesi L, El Ghoch M, Brodosi L, Calugi S, Dalle Grave R. Long-term weight loss maintenance for obesity: a multidisciplinary approach. *Diabetes Metab Syndr Obes Targets Ther.* (2016) 9:37–46. doi: 10.2147/DMSO.S89836
235. Hall KD, Kahan S. Maintenance of lost weight and long-term management of obesity. *Med Clin North Am.* (2018) 102:183–97. doi: 10.1016/j.mcna.2017.08.012
236. Hussain SS, Bloom SR. The regulation of food intake by the gut-brain axis: implications for obesity. *Int J Obes.* (2013) 37:625–33. doi: 10.1038/ijo.2012.93
237. Haufe S, Haas V, Utz W, Birkenfeld AL, Jeran S, Bohnke J, et al. Long-lasting improvements in liver fat and metabolism despite body weight regain after dietary weight loss. *Diabetes Care.* (2013) 36:3786–92. doi: 10.2337/dc13-0102
238. Villareal DT, Apovian CM, Kushner RF, Klein S, American Society for Nutrition, NAASO TOS. Obesity in older adults: technical review and position statement of the American society for nutrition and NAASO, the obesity society. *Am J Clin Nutr.* (2005) 82:923–34. doi: 10.1093/ajcn/82.5.923
239. Villareal DT, Chode S, Parimi N, Sinacore DR, Hilton T, Armamento-Villareal R, et al. Weight loss, exercise, or both and physical function in obese older adults. *N Engl J Med.* (2011) 364:1218–29. doi: 10.1056/NEJMoa1008234

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Morgan, Smeuninx and Breen. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.