


Please cite the Published Version

Gupta, Garima, Maiya, G. Arun, Bhat, Shayamsunder N., Hande, Manjunatha, Jude, Edward and Reeves, Neil D  (2022) Effect of balance strategies on fall risk in type 2 diabetes mellitus with peripheral neuropathy: a systematic review and meta-analysis. *Critical Reviews in Physical and Rehabilitation Medicine*, 34 (4). pp. 1-22. ISSN 0896-2960

DOI: <https://doi.org/10.1615/CritRevPhysRehabilMed.2022046155>

Publisher: Begell House

Version: Accepted Version

Downloaded from: <https://e-space.mmu.ac.uk/633503/>

Usage rights:  In Copyright

Additional Information: This is an accepted manuscript of an article which appeared in final form in *Critical Reviews in Physical and Rehabilitation Medicine*, published by Begell House

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

52

53 **Introduction**

54 The global health care burden of Type 2 Diabetes Mellitus (T2DM) as estimated by the
55 International Diabetes Federation (IDF) is expected to exponentially increase to 10.4% by
56 2040. This estimation puts a spotlight on the risk for the development of secondary
57 complications related to Type 2 Diabetes mellitus[1] [2]. Among the various secondary
58 complication, Diabetic peripheral neuropathy (DPN) accounts for 3/4th of the total number
59 [3] which is experienced as sensory-motor deficits. The sensory deficits implicate as loss of
60 sensations of pain, pinprick, temperature, proprioception, vibration along with “pins and
61 needles”, “burning” and “electric shocks”. The motor deficits present as an alteration in
62 muscle structure, functional strength, and joint stiffness. An escalation of these severe
63 sensory-motor deficits of diabetic neuropathy results in impaired gait, balance, and postural
64 sway, which leads to an inability to perform various activities of daily living, thereby
65 increasing the risk of falls. [4–6]

66

67 Recent evidence suggests a 17-fold rise in the risk of falls among the DPN individuals as
68 compared to healthy elders. [7] Similar findings on fall risk due to balance deficits show a
69 36% rise among Type 2 Diabetes Mellitus without neuropathy and 53% with neuropathy.[8]

70

71 The consequences of fall not only include physical effects but also functional, social, and
72 cognitive effects forming a vicious cycle. This leads to mild to severe fear for activity
73 participation, resulting in deconditioning, social isolation, contributing to fall of risk, and
74 reduced Quality of Life. [9–11]

75

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

76 The IDF interventional guidelines recommend the use of both pharmacological and
77 nonpharmacological measures for fall prevention with exercises as a primary choice. [12]
78 A study by Mendes R et al 2015 on exercise prescriptions for T2DM suggests the
79 incorporation of an individualized aerobic, resistance, and flexibility program. [13] However
80 the study of Kim et al. 2015 emphasis the role of educational interventions with a
81 multifactorial approach for fall prevention. [14]

82

83 There exists an underlying difference between diabetes with neuropathy and without
84 neuropathy owing to the additional sensory-motor deficits. Thus exercise guidelines targeting
85 only diabetic population or older adults cannot be replicated for those with Type 2 diabetes
86 with neuropathy. Thus the current review aims to determine the effect of multifactorial
87 balance rehabilitation strategies on Quality of Life, fall risk, and balance on Type 2 Diabetes
88 Mellitus with neuropathy.

89

90

91 **METHODS**

92

93 Registration

94 PROSPERO Reference ID: CRD42020161868

95

96 Search strategy

97 Review was conducted following PRISMA guidelines. Two independent reviewers searched
98 the following six databases: Pub Med, Scopus, Web of Science, Cumulative Index to Nursing
99 and Allied Health Literature (CINAHL), COCHRANE central, Embase. The databases were
100 searched from the beginning years of the database up to 25th November 2019. We updated
101 the search on 15th March 2020. Principal keywords for search stratagem were diabetic
102 peripheral neuropathy, balance rehabilitation strategies, balance, fall risk, and Quality of Life.
103 MeSH terms were used to search these keywords. Boolean operators "OR" and "AND" were
104 used to create the combined search strategy. (Supplementary file 1)

105 Study selection:

106 Search from all the database was imported to online Rayyan software. Two reviewers (G.G
107 and S.B) independently resolved the duplicates and screened the titles and abstracts. The
108 selected articles were downloaded, read, and evaluated by the reviewers separately. In case of
109 conflict third independent reviewer (A.G) was contacted to resolve the disagreement. The
110 reference list of the included articles was screened to identify any other relevant study.

111 Eligibility criteria:

112 Only randomized controlled trials (RCTs) with balance rehabilitation strategies as an
113 intervention on diabetic neuropathy population were included. For the present review,
114 operational definition of balance rehabilitation strategies covers all the physiotherapeutic

115 exercises (proprioceptive exercises, aerobic exercises, strength exercises, visual training, task
116 training, gait training, weight shifting, or transfer exercises) aiming to improve balance or
117 fall risk or Quality of Life in diabetic neuropathy. RCTs with interventions out of the scope
118 of the operational definition of balance rehabilitation strategies were excluded. All types of
119 study settings were included. The review compares the effect of balance rehabilitation
120 strategy with standard diabetic care, diabetic self-care education, or no treatment. Outcome
121 measures related to balance, fall risk, and Quality of Life were considered for this review.

122 Method of Data collection

123 Data regarding study population, size, design, type of interventions, duration, outcome
124 measures, and study results were noted down and managed on an excel sheet.

125 (Supplementary file 2)

126 Estimation of risk of bias (ROB) in included studies

127 The modified Cochrane Collaboration ROB tool was used to detect the quality of evidence. It
128 was decided independently by the reviewer's judgment (G.G, S.B) and any agreement dispute
129 was resolute by the third reviewer (A.G). Each study was assessed on five criterions
130 (selection, performance, attrition, reporting, and other bias). Each criterion of ROB was
131 classified as having low, high, or unclear risk. Later the overall ROB for each study was
132 assessed and they were categorized into good, fair, and poor studies. If study met all ROB
133 criteria (all low ROB) then the study was categorized as a good quality study, if the study had
134 one high ROB or two unclear ROB and outcome of the study are unlikely to be biased it was
135 rated as a fair quality study, if the outcome of the study were likely to be biased then the
136 study was categorized as poor quality. Also if more than two criteria were high ROB or
137 unclear then the study was rated poor-quality. [15]

138 GRADE evaluation

139 GRADE system was used to examine the quality of evidence. It also helped to summarize the
140 recommendations. [16] GRADE evidence profile (www.gradeworkinggroup.org) with
141 GRADEpro GDT online service was prepared to assess the quality of evidence. Any
142 disagreements between the reviewers (G.G and M.H) were resolved by a third reviewer (A.G)

143 Data synthesis

144 A meta-analysis of pooled data using a random-effect model was done by Cochran review
145 manager software version 5.3. More than 50% of the variance in I^2 was considered
146 heterogeneity. Standard deviations and mean differences values were pooled for synthesizing
147 meta-analysis results. We conducted the narrative analysis whenever data cannot be pooled
148 due to the varied use of outcome measures.

149 RESULTS

150 Study selection

151 A total of 2371 citations appeared in the search, after duplicate removal, title, and abstract
152 screening; full-text screening for 54 articles was done. Seven RCTs were included for final
153 narrative synthesis and meta-analysis. (Figure 1) Out of these seven RCT's:

- 154 • Only one study was eligible for the narrative synthesis of Quality of Life.
- 155 • Due to the varied use of outcome measures of all seven RCT's were assessed for the
156 narrative synthesis of balance and fall risk.
- 157 • Four RCTs observing the Berg Balance Scale, Functional Reach Test, Timed Up-Go
158 test, and One-Leg Stance as their balance and fall risk outcome measures were
159 included for meta-analysis.

160 Risk of Bias (ROB):

161 In the present review, two included studies were of good quality, four were fair quality and
162 one included study was poor quality. Participant blinding (performance bias) and allocation
163 concealment (selection bias) were the two primary biases seen in the included studies.
164 (Figure 2)

165 The GRADE quality of evidence for Quality of Life was moderate. Evidence for balance and
166 fall risk (BBS, FRT, and TUG) was very low and for the one-leg stance, it varied from
167 moderate to very low. (Table 2)

168

169 Study Characteristics

170 Participants:

171 Seven RCTs with a total of 418 participants aged 30 years and above were included for the
172 review. [17–23] Due to an inconsistent pattern of reporting, diabetic duration, and glycemic
173 parameters could not be summarized for the present review. (Table 3)

174 Intervention:

175 Multi-factorial Balance rehabilitation strategies: Multifactorial nature of balance
176 rehabilitation was not studied in any of the included RCT. Balance rehabilitation strategies in
177 two of the included studies comprised of balance exercises along with lower limb
178 strengthening exercises. Another included study combined the balance exercises with health
179 care education while the other study combined the balance exercises with gait training. Two
180 studies evaluated multi-sensory exercises and one RCT evaluated the effect of task-oriented
181 balance training. (Table 3)

182 Control group interventions included standard medical care or diabetic self-care education or
183 traditional balance exercises or no treatment. (Table 3)

184 Description of outcome measures

185 *Quality of Life*: Out of seven included studies only one study by Venkatraman et al 2019

186 measured Quality of life in their outcome measures. It was measured by the EQ-5D-5L index
187 score and SF36v2.

188 *Balance and fall risk*: All the seven included RCTs measured the balance and fall risk in their
189 outcome measures. Wide varieties of outcome measures were used in the included studies. List
190 of various balance and fall risk outcome measures included were Berg Balance Scale (BBS),
191 Functional Reach Test (FRT), Timed Up and Go Test (TUG), one leg stance (OLS)/unipedal
192 stance, Activity Specific Balance Confidence scale (ABC), Fall Efficacy Scale- International
193 (FES-I), Romberg's test, Performance-Oriented Mobility Assessment (POMA), backward
194 release test, postural assessment, proprioceptive, outdoor gait assessment via gyroscope,
195 dynamic balance test on a 5m beam, static balance test via Biodex Balance System and
196 tandem stance time. Thus there was only a small similarity of balance outcome measures in
197 the included studies.

198
199 Effects of balance rehabilitation strategies:

200 Narrative synthesis: Effect of the balance rehabilitation strategies on Quality of Life:

201 Only one RCT evaluated the effect of a balance exercise intervention on QoL. Eight weeks of
202 lower limb strengthening and balance training once weekly was given. The study utilized two
203 generic QoL tools; the EQ-5D-5L index as a primary outcome and SF-36v2 as its secondary
204 outcome measure. On comparing EQ-5D-5L outcome measure over 6 months it showed a
205 non-significant difference (mean difference-0.02 [95% CI 0.01, 0.06]; p= 0.175). Domain
206 wise analysis of SF36v2 showed that the intervention group showed improvement in body
207 pain (mean difference 5.14 [95% CI 2.05, 8.23]; p= 0.001) and in general health but

208 improvements in general health was not statistically significant (mean difference 2.36 [95%
209 CI -0.28, 4.99]; p= 0.080)

210 Meta-analysis: Effect of balance rehabilitation strategies on balance and fall risk measures:

211 The balance and fall risk was measured by all seven RCTs but as the outcome measures were
212 not consistently studied in most of the RCTs; they did not qualify for meta-analysis. Only
213 four (BBS, FRT, TUG, and OLS) balance and fall risk outcomes measures were synthesized
214 using meta-analysis.

215 A meta-analysis of Berg Balance Scale as an outcome measure:

216 A meta-analysis of three RCTs with a total of 135 participants showed balance rehabilitation
217 has no effect as compared to control or diabetic education (self-care or foot care) or standard
218 care. (MD 1.45, 95% CI -0.47, 3.38; p =0.14; I²= 59%) (Supplementary file 3)

219 A meta-analysis of Functional Reach Measure as an outcome measure:

220 A meta-analysis of four RCTs with total 233 participants on FRT as an outcome showed
221 balance rehabilitation was effective compared to control or diabetic education (self-care or
222 foot care) or standard care (MD 3.82, 95% CI 0.82, 3.83; P=0.01; I²= 72%) (Supplementary
223 file 3)

224 A meta-analysis of Timed Up and Go Test as an outcome measure:

225 A meta-analysis of five RCTs with a total of 326 participants on TUG as an outcome showed
226 balance rehabilitation was effective compared to control or diabetic education (self-care/ foot
227 care) or standard care (MD -1.41, 95% CI -2.14, -0.69; P=0.0001; I²= 50%) (Supplementary
228 file 3)

229 A meta-analysis of one leg balance Test as an outcome measure:

230 Meta-analysis of two RCTs with total of 75 participants on OLS/ unipedal stance under four
231 testing conditions (right and left eyes open and closed) showed balance rehabilitation was
232 effective compared to control or diabetic education (self care or foot care) or standard
233 care {Right EO (MD 7.86, 95% CI 1.97, 13.94, ; $p < 0.009$; $I^2 = 34\%$), Left EO (MD 6.14 , 95%
234 CI 2.64,9.64 ; $p < 0.0006$; $I^2 = 1\%$), right EC (MD 2.45, 95% CI 0.61, 4.28; $p < 0.009$; $I^2 =$
235 56%), Left EC (MD 1.80, 95% CI 0.86, 2.75, ; $p < 0.0002$; $I^2 = 0\%$). Though Robin L Kruse et
236 al 2010 observed a one-leg stance test in their study, it was not included for the present meta-
237 analysis as separate data for right and left side of the leg was not available. (Supplementary
238 file 3)

239 Narrative synthesis: Effect of balance rehabilitation strategies on balance and fall risk:

240 Out of seven included RCTs; five RCTs were included for narrative synthesis. In all the five
241 studies fall risk was the indirect interpretation of balance. One study reported that after 8
242 weeks of supervised intervention, postural assessment, and proprioception significantly
243 improved in intervention groups.[18] One study measured Balance confidence via Activities-
244 Specific Balance Confidence (ABC) scale, reported improvement after eight weeks as well as
245 at six months follow up.[21] With twelve months follow up of leg strengthening and balance
246 exercise intervention another study did not report any improvement in patient's balance
247 confidence with fall efficacy scale (FES). Kruse et al 2010 also measured the one-leg stance
248 time as a balance outcome measure and reported no significant difference between the groups
249 except under the eyes-closed condition. The author did not report the detailed procedure of a
250 one-leg stance (OLS) test mentioning on which leg the participants performed the test; hence
251 OLS data is not included for meta-analysis in the present review. [22] Malik et al 2016
252 observed a backward release test (reactive balance) and Romberg's test (static balance) as an
253 outcome measure of their study with 8 weeks of task-oriented training but possibly due to

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

254 selective reporting bias the results of these outcomes were not addressed in their results.[19]

255 Another study measured the effect of 12 weeks of intervention by the wide range of balance

256 and fall risk outcome measures (POMA, Outdoor gait assessment, dynamic balance test,

257 static balance test by Biodex USA and FES-I) reported that in comparison with the control

258 group intervention group increased their habitual speed of walking by 0.15 m/s (p<0.001).

259 Also, they reported significant improvement in dynamic balance (time to walk over beam),

260 POMA (balance and gait measures), postural sway on biodex, and balance confidence (FES-

261 I). [21]

262 **Discussion:**

263 The review focused on the effect of multifactorial balance rehabilitation strategies on Quality

264 of Life, fall risk, and balance in diabetic neuropathy. Rehabilitation guidelines are available

265 for Type 2 Diabetes Mellitus but when it progresses to diabetic neuropathy, the added deficits

266 and complications require detailed exercise recommendations, were not well explored. [24–

267 26] We observed that included studies used varied balance and fall risk outcome measures

268 hence all the outcome measures could not be pooled for meta-analysis. Meta-analyses were

269 performed for berg balance scale, timed up and go test, functional reach test, and one-leg

270 stand test. For the rest of the outcome measures on fall risk, balance, and Quality of Life, a

271 narrative synthesis was done.

272 Out of four fall risk and balance outcome measures (BBS, FRT, TUG, and OLS) meta-

273 analysis of three (FRT, TUG, and OLS) shows positive therapeutic effects of balance

274 rehabilitation on fall risk and balance. Though there are no previous meta-analysis

275 recommendations available, our results are in line with previous narrative synthesis

276 conducted on the heterogenic neuropathy population. [27,28]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

277 A possible explanation for BBS to show no treatment effect could be that BBS assessment
278 covers more aspects (control of centre of gravity, lower limb strength, gaze stabilization, use
279 of proprioceptive inputs, flexibility of upper, lower body and trunk) of balance than the other
280 three tests. As compared to BBS, TUG mainly focuses on functional mobility, FRT focuses
281 on limits of stability and one-leg stance tests the balancing ability in a reduced base of
282 support condition. Hence to get significant treatment effect on BBS assessment, balance
283 rehabilitation needs to covers mores aspects of balance mechanism. Thus the difference in the
284 results of the treatment effect could be due to different assessment nature of outcome
285 measures.

286 The study by Venkatraman et al. 2019, included for the narrative synthesis of Quality of Life,
287 reported no overall effect on Health-related Quality of Life (HRQoL) with 8 weeks of
288 balance exercises. Although subgroup analysis showed that improvement in functional
289 measures (timed up and go, five-time sit to stand and balance confidence) is associated with
290 improvement in EQ 5D 5L index scores. The author concluded that to achieve meaningful
291 changes in Quality of Life in diabetic neuropathy, exercises must be more vigorous and
292 training must be given for a longer period. Due to lack of literature on DPN population, we
293 cannot be conclusive about these findings but two large RCT's conducted on the diabetic
294 population without neuropathy also reported similar observations that exercises have
295 beneficial effects over QoL in diabetic population but the intervention must have high
296 volume and must be given for longer duration (9 to 12 months) for it to show significant
297 changes.[29,30] As diabetic neuropathy population has various added deficits it may require
298 even more time to achieve the statistically or clinically significant changes in QoL

299 Study limitations

300 The inclusion of only English language articles was one of the limitations of the review.

1
2
3 301 Clinical implications:

4
5 302 Based on the review we found there is a dearth of evidence on a multifactorial balance
6
7 303 rehabilitation program. There is a need for high-quality RCT on a multifactorial balance
8
9 304 rehabilitation program in diabetic neuropathy. This will benefit people living with diabetic
10
11 305 neuropathy for management of fall risk, balance and improve overall Quality of Life.

12
13
14 306 **CONCLUSION:**

15
16
17 307 The present systematic review suggests that strategies specific/ targeting to balance have a
18
19 308 positive effect on balance issues and fall risk in diabetic neuropathy. There is not sufficient
20
21 309 data available to conclude the effect of multifactorial balance rehabilitation strategies on
22
23
24 310 Quality of Life in diabetic neuropathy.

25
26
27
28 311

29
30
31 312 **DECLARATION**

32
33
34 313 **ETHICS STATEMENT:**

35
36
37
38 314 The present study was a meta-analysis, which did not involve human participants and/or
39
40 315 animals. Besides, no informed consent was needed for the meta-analysis.

41
42
43
44 316

45
46
47 317 **FUNDING:**

48
49
50 318 This systematic review did not receive financial support from any funding agencies.

51
52
53 319 **CONFLICT OF INTEREST**

54
55
56
57 320 The authors declare no conflicts of interest.
58
59
60
61
62
63
64
65

321 AUTHOR CONTRIBUTION

1
2
3
4 322 A.G designed the study and led the study design. G.G and S.B. identified and acquired
5
6 323 reports of trials and extracted data. G.G. and R.S. performed all data analyses, checked for
7
8 324 statistical inconsistency, and interpreted data. A.G., G.G., R.S., S.B., N.R., and M.H.
9
10
11 325 contributed to data interpretation. G.G., N. R., and A.G. drafted the report, and all other
12
13 326 authors (R.S., S.B., and M.H.) critically reviewed the report. A.G. and G.G. are the
14
15
16 327 guarantors of this work and, as such, had full access to all the data in the study and take
17
18 328 responsibility for the integrity of the data and the accuracy of the data analysis
19
20
21

22 329 **ACKNOWLEDGMENT:**

23
24
25 330 Authors would wish to acknowledge the Centre for Diabetic Foot Care and Research, MCHP,
26
27 331 MAHE, Manipal for providing all the technical and logistical support during the review
28
29
30 332 process.
31
32

33 333 **REFERENCES:**

- 34
35
36
37 334 [1] Zheng Y, Ley SH, Hu FB. Global aetiology and epidemiology of type 2 diabetes
38 335 mellitus and its complications. *Nat Rev Endocrinol* 2018;14:88–98.
39 336 <https://doi.org/10.1038/nrendo.2017.151>.
40
41 337 [2] International Diabetes Federation. *IDF Diabetes Atlas Eighth edition 2017*. 2017.
42 338 [https://doi.org/http://dx.doi.org/10.1016/S0140-6736\(16\)31679-8](https://doi.org/http://dx.doi.org/10.1016/S0140-6736(16)31679-8).
43
44 339 [3] Juster-Switlyk K, Smith AG. Updates in diabetic peripheral neuropathy.
45 340 *F1000Research* 2016;5:738. <https://doi.org/10.12688/f1000research.7898.1>.
46
47
48 341 [4] Mustapa A, Justine M, Mohd Mustafah N, Jamil N, Manaf H. Postural Control and
49 342 Gait Performance in the Diabetic Peripheral Neuropathy: A Systematic Review.
50 343 *Biomed Res Int* 2016;2016. <https://doi.org/10.1155/2016/9305025>.
51
52 344 [5] Almurdhhi MM, Brown SJ, Bowling FL, Boulton AJM, Jeziorska M, Malik RA, et al.
53 345 Altered walking strategy and increased unsteadiness in participants with impaired
54 346 glucose tolerance and Type 2 diabetes relates to small-fibre neuropathy but not vitamin
55 347 D deficiency. *Diabet Med* 2017;34:839–45. <https://doi.org/10.1111/dme.13316>.
56
57
58 348 [6] Priya TMV, Rajarajeswari A, Sivakumar R. Effectiveness of training on postural
59 349 stability in mild to moderate diabetic neuropathy patients. *Indian J Public Heal Res Dev*
60
61
62
63
64
65

- 350 2019;10:780–5. <https://doi.org/10.5958/0976-5506.2019.01985.5>.
- 1
2 351 [7] Vinik AI, Camacho P, Reddy S, Valencia WM, Trence D, Matsumoto AM, et al.
3 352 Aging, Diabetes, and Falls. *Endocr Pract* 2017;23:1120–42.
4 353 <https://doi.org/10.4158/EP171794.RA>.
- 6
7 354 [8] Timar B, Timar R, Gaiță L, Oancea C, Levai C, Lungeanu D. The Impact of Diabetic
8 355 Neuropathy on Balance and on the Risk of Falls in Patients with Type 2 Diabetes
9 356 Mellitus: A Cross-Sectional Study. *PLoS One* 2016;11:e0154654.
10 357 <https://doi.org/10.1371/journal.pone.0154654>.
- 12 358 [9] Tander B, Atmaca A, Ulus Y, Tura Ç, Akyol Y, Kuru Ö. Balance performance and
13 359 fear of falling in older patients with diabetics: a comparative study with non-diabetic
14 360 elderly. *Turk J Phys Med Rehab* 2016;62:314–22.
15 361 <https://doi.org/10.5606/tftrd.2016.77861>.
- 18 362 [10] Pin S, Spini D. Impact of falling on social participation and social support trajectories
19 363 in a middle-aged and elderly European sample. *SSM - Popul Heal* 2016;2:382–9.
20 364 <https://doi.org/10.1016/j.ssmph.2016.05.004>.
- 22 365 [11] Atler KE, Schmid AA, Klinedinst TC, Grimm LA, Marchant TP, Marchant DR, et al.
23 366 The Relationship between Quality of Life, Activity and Participation among People
24 367 with Type 2 Diabetes Mellitus. *Occup Ther Heal Care* 2018;32:341–62.
25 368 <https://doi.org/10.1080/07380577.2018.1522017>.
- 28 369 [12] International Diabetes Federation. IDF Clinical Practice Recommendations for
29 370 managing Type 2 Diabetes in Primary Care International Diabetes Federation - 2017.
30 371 2017.
- 32 372 [13] Mendes R, Sousa N, Almeida A, Subtil P, Guedes-Marques F, Reis VM, et al.
33 373 Exercise prescription for patients with type 2 diabetes - A synthesis of international
34 374 recommendations: Narrative review. *Br J Sports Med* 2016;50:1379–81.
35 375 <https://doi.org/10.1136/bjsports-2015-094895>.
- 38 376 [14] Kim EJ, Arai H, Chan P, Chen LK, Hill KD, Kong B, et al. Strategies on fall
39 377 prevention for older people living in the community: A report from a round-table
40 378 meeting in IAGG 2013. *J Clin Gerontol Geriatr* 2015;6:39–44.
41 379 <https://doi.org/10.1016/j.jcgg.2015.02.004>.
- 44 380 [15] Higgins JP, Savović J, Page MJ, Elbers RG, Sterne JA. Assessing risk of bias in a
45 381 randomized trial. 2019. <https://doi.org/10.1002/9781119536604.ch8>.
- 47 382 [16] Granholm A, Alhazzani W, Møller MH. Use of the GRADE approach in systematic
48 383 reviews and guidelines. *Br J Anaesth* 2019;123:554–9.
49 384 <https://doi.org/10.1016/j.bja.2019.08.015>.
- 51 385 [17] Majeed Kutty NA, Majida NAL. Effects of Multisensory Training on Balance and Gait
52 386 in Persons with Type 2 Diabetes: A Randomised Controlled Trial. *Disabil CBR Incl*
53 387 *Dev* 2013;24:79. <https://doi.org/10.5463/dcid.v24i2.206>.
- 56 388 [18] Ahmad I, Noohu MM, Verma S, Singla D, Hussain ME. Effect of sensorimotor
57 389 training on balance measures and proprioception among middle and older age adults
58 390 with diabetic peripheral neuropathy. *Gait Posture* 2019;74:114–20.

- 391 <https://doi.org/10.1016/j.gaitpost.2019.08.018>.
- 1
2 392 [19] Ghazal J, Malik AN, Amjad I. Task oriented training improves the balance outcome
3 393 and reducing fall risk in diabetic population. *Pakistan J Med Sci* 2016;32:983–7.
4 394 <https://doi.org/10.12669/pjms.324.10092>.
- 5
6 395 [20] Song CH, Petrofsky JS, Lee SW, Lee KJ, Yim JE. Effects of an Exercise Program on
7 396 Balance and Trunk Proprioception in Older Adults with Diabetic Neuropathies.
8 397 *Diabetes Technol Ther* 2011;13:803–11. <https://doi.org/10.1089/dia.2011.0036>.
- 9
10 398 [21] Allet L, Armand S, Bie RA De, Golay A. The gait and balance of patients with
11 399 diabetes can be improved : a randomised controlled trial 2010:458–66.
12 400 <https://doi.org/10.1007/s00125-009-1592-4>.
- 13
14 401 [22] Kruse RL, Lemaster JW, Madsen RW. Fall and Balance Outcomes After an
15 402 Intervention to Promote Leg Strength, Balance, and Walking in People With Diabetic
16 403 Peripheral Neuropathy: “Feet First” Randomized Controlled Trial. *Phys Ther*
17 404 2010;90:1568–79.
- 18
19 405 [23] Venkataraman K, Tai BC, Khoo EYH, Tavintharan S, Chandran K, Hwang SW, et al.
20 406 Short-term strength and balance training does not improve quality of life but improves
21 407 functional status in individuals with diabetic peripheral neuropathy: a randomised
22 408 controlled trial. *Diabetologia* 2019;62:2200–10. <https://doi.org/10.1007/s00125-019-04979-7>.
- 23
24 409
25
26
27 410 [24] Riandini T, Wee HL, Khoo EYH, Tai BC, Wang W, Koh GCH, et al. Functional status
28 411 mediates the association between peripheral neuropathy and health-related quality of
29 412 life in individuals with diabetes. *Acta Diabetol* 2018;55:155–64.
30 413 <https://doi.org/10.1007/s00592-017-1077-8>.
- 31
32
33 414 [25] Chau RMW, Ng TKW, Kwan RLC, Choi C, Cheing GLY, Chau RMW, et al. Risk of
34 415 fall for people with diabetes. *Disabil Rehabil* 2013;35:1975–80.
35 416 <https://doi.org/10.3109/09638288.2013.770079>.
- 36
37 417 [26] Chapman A, Meyer C, Renehan E, Hill KD, Browning CJ. Exercise interventions for
38 418 the improvement of falls-related outcomes among older adults with diabetes mellitus:
39 419 A systematic review and meta-analyses. *J Diabetes Complications* 2017;31:631–45.
40 420 <https://doi.org/10.1016/j.jdiacomp.2016.09.015>.
- 41
42
43 421 [27] Majeedkuty NA, Jabbar MA, Sreenivasulu S. Physical therapy for diabetic peripheral
44 422 neuropathy: A narrative review. *Disabil CBR Incl Dev* 2019;30:112–25.
45 423 <https://doi.org/10.5463/dcid.v30i1.760>.
- 46
47
48 424 [28] Streckmann F, Zopf EM, Lehmann HC, May K, Rizza J, Zimmer P, et al. Exercise
49 425 intervention studies in patients with peripheral neuropathy: a systematic review. *Sports*
50 426 *Med* 2014;44:1289–304. <https://doi.org/10.1007/s40279-014-0207-5>.
- 51
52 427 [29] Nicolucci A, Balducci S, Cardelli P, Cavallo S, Fallucca S, Bazuro A, et al.
53 428 Relationship of exercise volume to improvements of quality of life with supervised
54 429 exercise training in patients with type 2 diabetes in a randomised controlled trial: The
55 430 Italian Diabetes and Exercise Study (IDES). *Diabetologia* 2012;55:579–88.
56 431 <https://doi.org/10.1007/s00125-011-2425-9>.
- 57
58
59
60
61
62
63
64
65

432 [30] Myers VH, McVay MA, Brashear MM, Johannsen NM, Swift DL, Kramer K, et al.
1 433 Exercise training and quality of life in individuals with type 2 diabetes. *Diabetes Care*
2 434 2013;36:1884–90. <https://doi.org/10.2337/dc12-1153>.
3

4 435
5

6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

436

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

FIGURE LEGENDS

Fig 1: Flow diagram for the article selection and screening process according to PRISMA guidelines

Fig 2: Summary of Risk of bias based on authors' judgments about each ROB domain.

442 **TABLE LEGENDS**

443 Table 1: Inclusion and Exclusion criteria

444 Table 2: Grade evidence Profile and Summary of findings table

445 Table 3: Description of included studies

446

447

448 Table 1: Inclusion and Exclusion criteria

Criteria	Inclusion	Exclusion
Study design	RCTs	Non-RCT, case series, case reports, pre-post study design, conference presentations, review articles, and cross-sectional studies
Population	Type 2 Diabetes Mellitus with diabetic neuropathy	Type-I diabetes, gestational diabetes, or where the type of diabetes was not specified. Cause of neuropathy was other than Type 2 diabetes,
Intervention	Multifactorial Balance Rehabilitation strategies	Exercises delivered through Expensive sophisticated instruments (e.g.: isokinetic exerciser), electrotherapy (monotherapy, light therapy, vibrating insole), or alternative interventions (yoga, tai chi, acupressure, dance, etc.)
Comparison	Standard Diabetic Care/Diabetic Self Care Education /No Treatment	Pharmacological interventions, electrotherapy interventions
outcomes	Balance and/or QoL related outcomes	Outcomes not related to Balance and/or QoL related outcomes

449

450

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 2: Grade evidence Profile and Summary of findings table

Question: Balance rehabilitation exercises compared to diabetic education/ self-care health education/ standard care or no treatment for diabetic neuropathy

Setting: out-patient/ rehab-clinic/ home exercises or combination of supervised exercises and home program

Number of studies	Study design	Risk of bias	Certainty Assessment				Other considerations	Number of patients		Effect		Certainty	Importance
			Inconsistency	Indirectness	Imprecision			Balance rehabilitation exercises	diabetic education/ self-care health education/ standard care or no treatment	Relative (95% CI)	Absolute (95% CI)		
Quality of Life (follow up: mean 8 weeks; assessed with: EQ-5D-5L index and SF-36v2)													
1	randomized trials	serious ^a	not serious	not serious	not serious	none	The overall study showed a non-significant difference between the groups, but the domain wise analysis of SF-36V2 showed improvement in the "body pain domain" with 8 weeks of once-weekly leg strengthening and balance training in DPN.				⊕⊕⊕○ MODERATE	CRITICAL	
Balance and Fall risk (follow up: mean 8-24 weeks; assessed with: Berg Balance Scale; Scale from- 0 to 56)													
3	randomised trials	very serious ^b	serious ^c	not serious	very serious ^d	none	68	67	-	MD 1.45 higher (0.47 lower to 3.38 higher)	⊕○○○ VERY LOW	IMPORTANT	
Balance and Fall Risk (follow up: mean 8 weeks; assessed with: Functional Reach Test)													
5	randomised trials	very serious ^e	serious ^c	not serious	serious ^f	none	114	119	-	MD 3.82 higher (0.82)	⊕○○○ VERY LOW	IMPORTANT	

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

higher to
6.83
higher)

Balance and Fall Risk (follow up: mean 6-24 weeks; assessed with: Timed Up & Go Test)

6	randomise d trials	seriou s ^g	serious ^c	not serious	serious ^h	none	163	163	-	MD 1.41 lower (2.14 lower to 0.69 lower)	⊕○○○ VERY LOW	IMPORTAN T
---	-----------------------	--------------------------	----------------------	-------------	----------------------	------	-----	-----	---	---	---------------------	---------------

Balance and Fall Risk (follow up: mean 8 weeks; assessed with: One Leg Stance (OLS) right eyes open)

3	randomise d trials	seriou s ⁱ	not serious ^j	not serious	serious ^h	none	39	36	-	MD 7.86 higher (1.97 higher to 13.74 higher)	⊕⊕○○ LOW	IMPORTAN T
---	-----------------------	--------------------------	--------------------------	-------------	----------------------	------	----	----	---	---	-------------	---------------

Balance and Fall Risk (follow up: mean 8 weeks; assessed with: One Leg Stance (OLS) left eyes open)

3	randomise d trials	seriou s ⁱ	not serious	not serious	serious ^h	none	39	36	-	MD 6.14 higher (2.64 higher to 9.64 higher)	⊕⊕○○ LOW	IMPORTAN T
---	-----------------------	--------------------------	-------------	-------------	----------------------	------	----	----	---	--	-------------	---------------

Balance and Fall Risk (follow up: mean 8 weeks; assessed with: One Leg Stance (OLS) right eyes closed)

3	randomise d trials	seriou s ⁱ	serious ^c	not serious	serious ^h	none	39	36	-	MD 2.45 higher (0.61 higher to 4.28 higher)	⊕○○○ VERY LOW	IMPORTAN T
---	-----------------------	--------------------------	----------------------	-------------	----------------------	------	----	----	---	--	---------------------	---------------

Balance and Fall risk (follow up: mean 8 weeks; assessed with: One Leg Stance (OLS) left eyes closed)

3	randomise d trials	seriou s ⁱ	not serious	not serious	not serious	none	39	36	-	MD 1.8 higher (0.86 higher to 2.75 higher)	⊕⊕⊕○ MODERAT E	IMPORTAN T
---	-----------------------	--------------------------	-------------	-------------	-------------	------	----	----	---	---	----------------------	---------------

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

CI: Confidence interval; **MD:** Mean difference

Explanations

a. ROB was fair

b. Out of 3 included studies ROB for Malik et al 2016 was poor, Song et al 2011 was fair, hence marked very serious

c. heterogeneity is between 50% to 75 % i.e. moderate heterogeneity

d. CI of Kruse et al 2010 was large and crosses the clinical decision threshold. Also, the overall pooled effect of also crosses the clinical decision threshold.

e. out of 4 included studies Malik et al 2016 was poor, Song et al 2011, Ahmed et al2019 and Venkatraman et al2019 were fair, hence marked very serious

f. Though the overall pooled effect does not have wide CI, 2 of the included individual studies had wide CI crossing the clinical decision threshold

g. all the 5 included studies were at fair ROB according to the author's judgment, hence marked serious

h. Though the overall pooled effect does not have wide CI, one of the included individual studies had wide CI crossing the clinical decision threshold

i. Both the included studies of Ahmed et al 2019 and Song et al 2011 were at fair ROB, hence marked serious

j. Heterogeneity is between 25% to50% i.e. Low heterogeneity

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 3: Description of included studies

Author	Journal & Year	Age (years)	Population	Study Design	Total Sample	No of Subjects Intervention Group	No of Subjects In Control	Intervention Group	Control Group	Duration	Outcome Measures Balance	Outcome Measures QOL
Ahmed et al	Gait & posture 2019	45-75	DPN	RCT	37	a) Less than 60 years: 8, b)more than 60 years: 12	a) Less than 60 years: 8, b) More than 60 years: 9	Sensory-motor training, Diabetic & Foot Care Education	Diabetic &Foot Care Education	8 weeks	FRT, TUG, OLS, Postural Assessment, Proprioception	Nil
Venkataraman et al	Diabetologia 2019	40-79	DPN	RCT	143	70	73	Balance retraining and strengthening interventions guided by a physiotherapist	Standard medical care	8 weeks	FRT, TUG, ABC	SF-36V2, EQ-5D-5L.
Malik et al	Pak J Med Sci 2016	30-70	DPN	RCT	18	8	10	Task-oriented training	Traditional balance training	8 weeks	FRT, BBS, Rhomberg's, Backward Release Test	Nil

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Song et al	Diabetes Technology & Therapeutics 2011	≥ 70	DPN	RCT	38	19	19	Balance Exercises	Health education	8 weeks	FRT, TUG, OLS, BBS	Nil
Kruse et al	Physical Therapy 2010	≥ 50	DPN	RCT	79	41	38	Part 1 leg strengthening and balance exercises, self-monitored walking program; part 2 telephone calls	8 visits by therapist for self-care in diabetes	12 months	BBS, OLS, TUG, FES-I	Nil
Majeed K et al	Disability, CBR & Inclusive Development 2013	55-75	DPN	RCT	32	16	16	multisensory training and diabetic education	diabetic education	6 weeks	TUG, 6MWT	Nil
Allet et al	Diabetologia 2010	> 60	DPN	RCT	71	35	36	Gait and balance exercises	No treatment	12 weeks	POMA, Out Door Gait Assessment, Dynamic Balance Test, Static Bal Test ByBiodex	Nil

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Fig 1: Flow diagram for the article selection and screening process according to PRISMA guidelines




Fig 2: Summary of Risk of bias based on authors' judgments about each ROB domain.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Ahmed et al 2019	+	?	●	+	+	+	+
Allet et al 2010	+	+	+	+	+	+	+
Kruse et al 2010	+	+	+	+	+	+	+
Majeed K et al	+	+	?	+	+	+	+
Malik et al 2016	+	?	?	?	?	●	?
Song et al 2011	+	?	?	+	+	+	+
Venketraman et al 2019	+	+	?	+	+	+	+