

Please cite the Published Version

Donelon, Thomas A, Edwards, Jamie, Brown, Mathew, Jones, Paul A, O'Driscoll, Jamie and Dos'Santos, Thomas (2024) Differences in biomechanical determinants of ACL injury risk in change of direction tasks between males and females: a systematic review and meta-analysis. *Sports Medicine - Open*, 10 (1). 29 ISSN 2199-1170

DOI: <https://doi.org/10.1186/s40798-024-00701-z>

Publisher: Springer

Version: Published Version

Downloaded from: <https://e-space.mmu.ac.uk/634265/>

Usage rights:  [Creative Commons: Attribution 4.0](https://creativecommons.org/licenses/by/4.0/)

Additional Information: This is an open access article which originally appeared in *Sports Medicine - Open*

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

SYSTEMATIC REVIEW

Open Access



Differences in Biomechanical Determinants of ACL Injury Risk in Change of Direction Tasks Between Males and Females: A Systematic Review and Meta-Analysis

Thomas A. Donelon^{1*}, Jamie Edwards¹, Mathew Brown¹, Paul A. Jones², Jamie O'Driscoll¹ and Thomas Dos'Santos³

Abstract

Background Change of direction (COD) movements are associated with non-contact anterior cruciate ligament (ACL) injuries in multidirectional sports. Females appear at increased risk compared to males, which could be attributable to whole body kinematic strategies and greater multiplanar knee joint loads (KJLs) during COD which can increase ACL loading.

Objective The aim of this systematic review and meta-analysis was to examine and quantitatively synthesise the evidence for differences between males and females regarding KJLs and their biomechanical determinants (whole body kinematic strategies determining KJLs) during COD tasks.

Methods Databases including SPORTDiscus, Web of Science, and PubMed were systematically searched (July 2021–June 2023) for studies that compared differences in knee joint loads and biomechanical determinants of KJLs during COD between males and females. Inclusion criteria were: (1) females and males with no prior history of ACL injury (18–40 years); (2) examined biomechanical determinants of KJLs and/or KJLs during COD tasks > 20°; (3) compared ≥ 1 outcome measure between males and females. Studies published between 2000 and 2023 examining a cutting task > 20° with a preceding approach run that compared KJLs or the whole body multiplanar kinematics associated with them, between sexes, using three-dimensional motion analysis.

Results This meta-analysis included 17 studies with a pooled sample size of 451 participants (227 males, 224 females). Meta-analysis revealed females displayed significantly less peak knee flexion during stance (SMD: 0.374, 95% CI 0.098–0.649, $p=0.008$, $I^2: 0\%$); greater knee abduction at initial contact (IC) (SMD: 0.687, 95% CI 0.299–1.076, $p=0.001$, $I^2: 55\%$); less hip internal rotation (SMD: 0.437, 95% CI 0.134–0.741, $p=0.005$, $I^2: 34\%$) and hip abduction at IC (SMD: -0.454, 95% CI 0.151–0.758, $p=0.003$, $I^2: 33\%$). No significant differences were observed between males and females for any internal or externally applied KJLs. All retrieved studies failed to control for strength, resistance training or skill history status.

Conclusion No differences were observed in KJLs between males and females despite females displaying greater knee abduction at IC and less peak knee flexion during the stance phase of CODs, which are visual characteristics of non-contact ACL injury. Further research is required to examine if this translates to a similar injury risk, considering

*Correspondence:

Thomas A. Donelon
td254@canterbury.ac.uk

Full list of author information is available at the end of the article



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

morphological differences in strain characteristics of the ACL between males and females. This observation may in part explain the disproportionate ACL injury incidence in female multidirectional athletes. Further higher quality controlled research is required whereby participants are matched by skill training history, resistance training history and strength status to ensure an appropriate comparison between males and females.

Key Points

- No differences between sexes were found for multiplanar KJLs despite previous research supporting this. Further research is required to identify if this is the case and if it affects injury risk between males and females.
- Females displayed greater knee abduction at initial contact, less peak knee flexion during stance, and less hip rotation and abduction at IC together with less peak hip flexion during stance during CODs.
- Previous research has failed to account for confounding factors such as skill training history, resistance training history and strength status together with gendered environmental factors that could contribute to the misinterpretation of COD differences between males and females.

Keywords Cutting, Pivoting, Sex-comparison, Gender-comparison, Anterior-cruciate-ligament

Background

Changes of direction (CODs) encompass an integral element of team sports and invasion games, which are associated with key decisive moments in sport, such as creating space, evading an opponent (i.e., tackle break success in rugby), and goal scoring [1]. Notational analysis has observed CODs to occur at frequencies of every four to six seconds across a number of sports such as netball [2], soccer [3] and hockey [4], with up to 700 CODs observed in soccer in a 90 min period [5–7]. Considering the rate and frequency of CODs in invasion games, it is of significant importance for coaching and sports medicine staff to condition athletes appropriately to meet these physical demands [8]. Although CODs are a key component of effective performance in multidirectional sports, COD actions are also a primary mechanism of non-contact ACL injury [9–11].

A substantial amount of anterior cruciate ligament (ACL) injuries are reported annually with an estimated global incidence of 68.6 per 100,000 person-years [12] alongside an observed increase in ACL reconstructions globally [12–15]. Despite recent advancements in sport technology, medicine and coaching practice, ACL injury rates are projected to rise [16]. Of clinical importance, 70% of ACL injuries are non-contact [17], with approximately 0.62 ACL injuries reported per 1000 player exposures [18]. It is of concern that a gynocentric pattern in the epidemiological literature is prevalent, with females being 1.7 times more likely to incur an ACL injury compared to male counterparts when matched for playing exposures [19], alongside a significantly greater risk of ACL re-injury [20]. ACL injuries are debilitating by nature, often requiring surgical intervention and

a lengthy return to play time of 6–24 months [21, 22]. Substantial social and health related implications have also been identified following ACL reconstruction. These include a financial burden to public services (e.g., £63 million GBP spent on reconstructions annually in the United Kingdom [23]) associated with a decline in mental health [24]. An increased susceptibility to osteoarthritis has also been observed [25], coupled with 50% failing to return to sport within a year post-surgery [26]. The above necessitates a greater understanding of injury mitigation strategies in order to attenuate the financial, social and health related issues associated with ACL injury.

ACL ruptures occur when a catastrophic load is applied to the ACL, whereby the strain exceeds the ligament's mechanical tolerance, resulting in tissue failure [27]. Previous cadaveric [28] and modelling [29–31] research have identified knee abduction, shear, and internal rotational loads (alternatively known as externally applied moments or torques) to substantially increase ACL strain independently, although observed strain is greatest when a combination of these loads is applied together [32]. Such have been termed multiplanar knee joint loads (KJLs) in the literature, and have been identified as a predictor of ACL injury [33] and also considered surrogate measures of non-contact ACL injury risk [34–36]. There is a propensity to generate large and potentially hazardous multiplanar KJLs during COD actions that are commonly performed in team sports, which are amplified with specific sub-optimal postures (e.g., knee abduction, extended knee posture, lateral trunk flexion), warranting their investigation in relation to non-contact ACL injury risk. A substantial amount of research has been undertaken in order to identify the biomechanical determinants of these KJLs in CODs to understand the kinetic, kinematics, and

technical parameters associated with increased KJLs and potential non-contact ACL injury risk [33, 37–44]. This has provided practitioners with guidelines for optimal technique and how to mitigate KJLs in COD tasks [45], with researchers demonstrating a reduction in KJLs through targeted COD technique modification training interventions [37, 46, 47].

Females have been identified as exhibiting these sub-optimal body postures and greater relative KJLs during CODs more than their male counterparts [48–52]. This observation may provide some explanation of the disparity in reported injury rates. A previous systematic review and meta-analysis [53] identified females as displaying an increased knee abduction angle across a range of weight-bearing tasks, including running, landing and cutting. Knee abduction angle itself would only represent one component of the resultant knee abduction moment, and does not take into account the other segmental and kinetic contributions to multiplanar KJLs [36]. To date, two systematic reviews have examined differences between males and females in biomechanical surrogates of non-contact ACL injuries in landing [54] and cutting [55]. Benjaminse et al. [55] identified greater knee abduction angles in two of their retrieved studies (one with a large effect size (ES) of 0.99), and knee abduction moments, whilst Beaulieu et al. [54] identified females exhibited greater knee abduction angles during landing. A limitation of both these systematic studies is that they did not meta-analyse the data to examine and establish the magnitude of differences between males and females. Meta-analysis would previously not have been possible in the aforementioned review [55], due to insufficient literature available (seven retrieved studies from the years 1947–2008). In recent years there has been a substantial research effort to further understand biomechanical differences between males and females during cutting [50, 51, 56–64], allowing for meta-analysis to be undertaken.

Therefore, the aim of this systematic review and meta-analysis is threefold: firstly, to synthesise the evidence regarding differences between males and females regarding KJLs and their biomechanical determinants during CODs measured through 3-dimensional (3D) motion (and ground reaction force [GRF]) analysis; secondly, to identify relevant effect modifiers in uninjured athletes and thirdly, to provide recommendations and directions for future research examining sex differences in COD tasks. It was hypothesised that females would exhibit less knee flexion and greater knee abduction angles, and greater knee abduction moments during COD tasks. The findings of this meta-analysis may assist in ACL injury mitigation strategies, injury screening protocols, and

physical preparation and management of female and male athletes.

Methods

A systematic review and meta-analysis were performed in accordance with the recently updated PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) guidelines [65]. The study was also registered with PROSPERO on 17th November 2021 (CRD42021266215) and adhered to the ethical recommendations for the publishing of systematic reviews in accordance with Wager and Wiffen [66]. Some minor amendments were made to the review completion date and inclusion criteria; namely, the population element of the Population, Intervention, Comparator, Outcomes and Study design (PICOS) framework was amended to include healthy and recreationally active participants and removed the performance level stipulation of playing twice a week in given sport. The rationale for amendment was due to the majority of retrieved papers failing to specify competition playing frequency, and a paucity of literature examining elite athletes [67]. The registration document was amended to reflect these changes appropriately (4th October 2022).

Study Inclusion and Exclusion Criteria

A PICOS framework was constructed to define the inclusion and exclusion criteria for this study and is presented in Table 1. This article is part of a wider project and search strategy identifying a number of neuromuscular and biomechanical differences between sexes in COD tasks. It was decided to solely focus on biomechanical injury risk surrogates obtained from 3D motion analysis, with the neuromuscular surrogates and data obtained from modelling and simulation to be disseminated in another project. The aim of this paper is to analyse the sex differences in biomechanical surrogates of non-contact ACL injury risk (biomechanical determinants of KJLs and KJLs themselves) obtained from 3D motion and GRF analysis.

In randomised controlled trials or studies examining the effect of an intervention such as bracing [59] or fatigue [58, 77], only baseline data from the control group were extracted for subsequent analysis. Exclusion criteria consisted of studies examining injured or ACL reconstructed populations, case studies, and poster presentations/ conference proceedings. Studies that did not meet the PICOS criteria were excluded from the review.

Search Strategy

A literature search was performed using PubMed, Web of Science, and SPORTDiscus databases from July 2021

Table 1 PICOS framework used to define inclusion and exclusion criteria of studies

Population	For the purpose of this review, male, female and sex refers to individuals assigned male or female at birth based on biological characteristics Healthy women – No restrictions were placed regarding reproductive status and hormonal contraception (HC) usage Healthy males (as a control for sex comparison of biomechanics) Age 18–40 years (children and pre/adolescent populations were excluded to control for the effect of puberty or changes during adolescence) Competing in or familiar with jump landing / cutting dominant, field / court based invasion games / sports Performance level: Tier 1 minimum – Tier 5 [67]—Healthy, recreationally active and elite athletes No history of ACL injury
Intervention/method	A specific intervention was not investigated but participants were required to meet the population criteria above Studies must have examined biomechanical surrogates of non-contact ACL injury risk during pre-planned or unplanned change of direction tasks, with 3D motion and/or GRF analysis (inclusive of pre and/or unplanned tasks) Studies which adhered to the following change of direction task criteria were included: A preceding approach run of a minimum of 3 steps containing a subsequent change in direction > 20° The decision was made to omit tasks that included a split/ false step or hop or that omitted an approach run in line with previous COD definitions [68, 69] Omission of an approach run would not truly replicate the loading parameters of non-contact ACL injury situations due to the absence of a deceleration (as deceleration has been identified as the component where most noncontact ACL ruptures occur) (Donelon et al., 2020 [45])
Comparator	To determine the effect of sex: A direct between group comparison of a biomechanical surrogate of non-contact ACL injury risk between females and males (acting as a control)
Outcome	Precise mean and SD provided for injury risk factors between males and females Biomechanics: Knee abduction, rotation, flexion moments / impulse (knee joint loads) [45, 70–74] Proximal anterior tibial shear [75] Technical, kinetic, or kinematic determinants of surrogates of injury risk (knee joint loads) [36, 70, 76] at initial contact (IC) (first instance of ground contact in COD) and peak value obtained during stance (across the full cutting cycle 0–100%) related to quadriceps, ligament, trunk, and leg dominance, such as: Vertical/posterior GRF/ impulse <i>Initial or peak</i> Lateral trunk flexion/rotation angle Hip internal rotation angle Knee valgus / internal rotation angle Knee flexion / hip flexion Foot progression angles Rearfoot/ forefoot strike Coordination changes Asymmetries
Study design	Peer review full article in English, examining humans from the year 2000 onwards due to a lack of 3D motion analysis research before this date Direct assessment of change of direction biomechanics (with surrogates of injury) between healthy males and females
Other data extraction	The following data were extracted and recorded in a spreadsheet: (1) Author names, publication year and country of origin (2) Sample size and participant characteristics including sport(s), playing level/status, training history, strength history/status/profile, reproductive status(females), hormonal contraception usage (3) Angle of COD task (4) Anticipatory nature of COD task (planned/unplanned) (5) If unanticipated, method and timing of stimulus presentation (6) Approach velocity prior to COD (7) How approach velocity was controlled for / calculated (8) Rest period between trials (if stated) (9) How risk factor (ACL surrogate) was assessed (methods) (10) Reliability and familiarisation stated for outcome measures / tasks (11) Outcome measures (mean, SD) (12) Any other empirical data available for a variable that could mitigate any sex differences in surrogate injury risk identified (e.g. strength or experience / playing time) (13) For female populations, information relating to: a. Reproductive status b. Menstrual cycle phase c. Hormonal contraception use

ACL—Anterior Cruciate Ligament, COD—Change of Direction, GRF—Ground Reaction Force, HC—Hormonal Contraception, MSK—Musculoskeletal, SD—Standard Deviation

to June 2023 with the final search date of 1st June 2023. A schematic of search methodology in accordance with established PRISMA guidelines [44] is presented in the results below. Search terms that were used are presented in Table 2:

Articles retrieved by this search were then title, abstract and then full-text screened against the PICOS framework to examine their suitability for inclusion by the lead researcher (TD) and another researcher (TDS). Should disagreement have arisen surrounding the inclusion of an article, a third researcher (PJ) was consulted and their decision deemed final. Bibliographies of prospectively eligible (full texts reviewed) were then hand searched in order to identify further eligible studies.

Methodological Quality and Publication Bias

An assessment of methodological quality was independently undertaken by two of the researchers (TD and TDS) as per previously established methods [70, 78, 79] using a COD specific scale constructed by Brown et al. [80]. This is deemed to be more suitable for assessing the methodological quality of COD studies due to the omission of criteria such as random allocation, assessor blinding and subject blinding that are present in more commonly used scales such as the Cochrane or Delphi, Physiotherapy, Evidence Database scales [79, 81]. Change of direction specific protocols were rigorously assessed by the tool due to specific criteria present, such as the allowance of practice trials, duration of rest between trials and velocity of COD tasks. Each component was individually scored from 0 to 2 (where 0 = clearly no, 1 = maybe or insufficient information; and 2 clearly yes). Any disagreement was resolved through consensus and discussion involving a third researcher (PJ). The methodological assessment tool is presented in Table 3.

Data Extraction

The following data were extracted by the lead author in Table 4 (TD): quantitative data pertaining to study methodology, participant characteristics (age, height, mass) and verification, biomechanical ACL injury risk surrogates during initial contact (IC) (defined as the first instance of foot-contact during the cut), range of motion

(ROM; defined as point of IC to maximum knee flexion) and peak stance (defined as the peak value obtained across 100% of the cutting cycle), reliability measures and measured outcomes and results (means and SDs of both male and female conditions). Once extracted, data were pooled together for COD angle and anticipation status of the COD task due to insufficient data for separate angle-dependent and anticipatory analysis. The authors acknowledge that anticipated and unanticipated COD conditions have been identified as producing significantly different outcome metrics, although sex differences in kinematics and KJLS have been observed under both anticipated [48, 52] and unanticipated [50, 56] conditions. This is attributable to a time constraint to orientate the body in preparation for the COD and therefore the biomechanical demands to complete the task remain the same [82]. This decision was made based on the above and considering the aim of this meta-analysis is to identify differences between males and females in COD tasks. Data were then systematically separated by variable timing (IC, ROM or during peak stance), moment convention (internal / external) and reference frame prior to analysis.

All variables that were extracted are present in Table 4.

Statistics

All statistical analysis was performed using Comprehensive Meta-Analysis (Comprehensive Meta-Analysis Version 3, Biostat, Englewood, NJ, USA). Separate pooled analyses for initial contact and peak variables obtained during stance were performed for each independent parameter to establish the standardised mean difference (SMD) between male and female groups. SMD was selected as the appropriate outcome measure due to inter-study variance in data reporting approaches. SMD effect thresholds were as follows: 0.2–0.5 small effect, 0.5–0.8 medium effect and above 0.8 as a large effect [83]. Where sufficient data were available, subgroup analyses were also performed on COD angle, anticipation, and sport. Further meta-regression moderator analyses were performed on methodological quality, age, mass, achieved velocity and playing experience. Statistical heterogeneity was

Table 2 Search strategy used for literature searching

Search strategy	
"Anterior cruciate ligament" NOT reconstruction OR "ACL" NOT reconstruction	AND
"Sex-differences", OR "gender-differences" OR "Sex-comparison" OR "gender-comparison" OR "sex" OR "gender"	AND
"biomechanics" OR "biomechanical-determinants" OR "technical-determinants" OR "kinetics" OR "kinematics" OR "Neuromuscular" OR "electromyography" OR "muscle activation"	AND
"Change of direction", or "Cutting manoeuvre", or "Run and cut", or "Run-and-cut", or "Sidestepping", or "Side-stepping" or "Shuttle-run"	AND

Table 3 Brown assessment of methodological quality

Question	Criteria	Beaulieu et al. [56]	Condello et al. [57]	Iguchi et al. [58]	Ihmels et al. [59]	Khalid et al. [60]	McLean et al. [49]	McLean et al. [48]	McLean et al. [88]	Nagano et al. [61]	O'Connor et al. [62]	Pollard et al. [89]	Pollard et al. [86]	Schreurs et al. [63]	Sigward and Powers 2006 [52]	Sigward et al. [51]	Sigward et al. [50]	Tanikawa et. [64]
1	Power analysis was performed and justification of study sample size	0	0	1	2	0	2	2	2	2	2	2	2	2	0	2	0	2
2	Athlete demographics were clearly defined: gender, age, body height, and body mass at time of test	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	1
3	Athlete characteristics were clearly defined: sport, experience or activity/level and level of play	2	2	1	1	2	0	2	2	1	2	2	2	1	2	2	2	1
4	Inclusion and exclusion criteria were clearly stated for athletes	0	0	0	2	0	2	2	2	2	2	2	2	2	2	2	2	2
5	Proper training and practice trials of the test were given to the athletes allowing for adequate familiarisation	2	2	2	0	2	0	0	0	2	2	2	2	2	2	0	0	1

Table 3 (continued)

Question	Criteria	Beaulieu et al. [56]	Condello et al. [57]	Iguchi et al. [58]	Ihmels et al. [59]	Khalid et al. [60]	McLean et al. [49]	McLean et al. [48]	McLean et al. [88]	Nagano et al. [61]	O'Connor et al. [62]	Pollard et al. [89]	Pollard et al. [86]	Schreurs et al. [63]	Sigward and Powers 2006 [52]	Sigward et al. [51]	Sigward et al. [50]	Tanikawa et. [64]
6	Methods were described in great detail to allow replication of the test. Testing devices, n of trials, n and duration of rest, speed, angle of COD	2	2	1	1	2	1	1	1	1	1	2	1	2	1	1	1	1
7	Test-retest reliability of measurement device reported	0	1	0	0	0	0	0	0	1	0	0	0	0	2	0	0	0
8	Outcome variables clearly defined	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	1
9	Statistical analyses were appropriate	2	2	1	1	1	2	2	2	2	1	2	2	1	2	1	1	1
	Total score (maximum 18)	12	13	10	11	12	11	12	13	15	15	16	15	14	15	11	10	10
	Score as %	67%	72%	56%	61%	67%	61%	67%	72%	83%	83%	89%	83%	78%	83%	61%	56%	56%

COD—Change of Direction, n—number

assessed alongside the pooled analysis and reported as the I^2 statistic. The application of fixed or random effects analysis was determined by the absence or presence of significant statistical heterogeneity. The threshold for heterogeneity significance was set at an I^2 statistic of $>40\%$ in which an Egger's regression test was systematically planned to assess the presence of funnel plot asymmetry to account for potential publication bias [84]. Pooled analysis results were considered significant if $p < 0.05$ and the Z-value > 2 .

Results

Search Results

Figure 1 presents a flow chart summarising the results of the systematic search process. Following duplicate removal and application of eligibility criteria, 14 studies [48–51, 57–62, 64, 85–88] were initially included for meta-analysis. Following reference list screening of eligible studies, a further three studies were deemed eligible for inclusion [52, 63, 89], resulting in 17 studies overall used for quantitative analysis. The authors had access to the full text for all included studies.

Characteristics of the Studies

The 17 included studies (Table 5) evaluated sex differences in surrogates of non-contact ACL injury risk. A total of 451 participants were included in the analysis (227 males, 224 females). The mean (SD) age, height and mass of males and females were 21.10 (1.74) years, 1.79 (0.05) m, 74.5 (6.84) kg and 20.82 (1.70) years, 1.67 (0.05) m and 62.83 (6.55) kg, respectively. In total, 254 participants played soccer (132 males and 122 females) and 109 were defined as physically active or recreationally active (54 males, 55 females). Forty participants played basketball (20 males 20 females) whilst 29 (13 males, 16 females) were of a non-defined team sporting background [63]. Finally, 16 participants were termed “proficient” [49] in cutting manoeuvres in addition to 3 female lacrosse players used in a study [61]. Mean sporting experience in years was stated in 11 studies [48, 51, 52, 56, 57, 60, 62, 86–89] and was 12.18 years (2.58) for males and 11.53 (2.91) for females. Only one study [56] controlled for menstrual cycle phase in female participants. No study reported resistance training history, hormonal contraceptive use or reproductive status. Thirteen studies [50–52, 56–58, 60–64, 86, 89] reported familiarisation prior to data collection consisting of numerous practice trials [50–52, 57, 58, 61, 63, 64, 86] or a specific session delivered separately [56, 62, 77, 89]. Only two studies [52, 57] reported reliability statistics for their outcome measures, namely intra-class correlation coefficients (ICCs)

[57], and coefficients of multiple correlations (CMCs) [52].

Nine studies utilised a 45° COD task [48, 51, 52, 56, 59, 62, 77, 86, 89], with two adopting a 60° sidestep [57, 58]. Two studies adopted multiple COD angles including 45° and 110° [50], 45° and 180° [88], and 45–180° in four 45° increments [63]. One study each used isolated 90° [64] and 180° [61] CODs. Ten of the included studies included an anticipated COD task [48, 49, 52, 57, 59, 61, 63, 64, 86, 88]; with one study examining both unanticipated and anticipated CODs [77]. Of the six studies utilising unanticipated conditions [50, 51, 58, 62, 85, 89], a task choice approach was utilised requiring a decision between a 45°, 110° or straight run [50, 51]; a 30° crossover cut and 60° sidestep [58]; a straight-line run, “hard stop” [62, 85] or stop jump [89].

Assessment of Methodological Quality

Assessment of methodological quality is presented in Table 3. The mean score for the Brown Methodological Quality of Assessment for the 17 included studies was 12.65 (70.26%) \pm 1.97 (10.94%). Scores ranged from 10 (55.5%) [50, 58, 64] to 16 (88.8%) [89]. Nine studies were below this mean score [48–51, 56, 58, 59, 64, 77] with eight studies presenting greater methodological quality greater than the mean [52, 57, 61–63, 86, 88, 89]. Between day reliability was reported in only one study [52]. All but one study [64] clearly defined athlete demographics and outcome variables. Six studies did not include information surrounding familiarisation of the athletes to the COD task to be performed [48–51, 59, 88]. Eight [50, 51, 58, 59, 62–64, 77] of the 17 included studies conducted inappropriate statistical processes when examining differences between males and females in COD biomechanics, through pooling data across tasks [50, 62–64], maturation [51], fatigue [58, 77] and bracing [59] protocols prior to performing statistical analysis.

Quantitative Synthesis: Kinematics

Hip

In the sagittal plane, meta-analysis indicated that females exhibited less peak hip flexion during stance than males (SMD: 0.504, 95% CI 0.134–0.741, $p=0.06$ I^2 : 22%; 7 measured groups with 155 participants (78 males, 77 females)) (Fig. 2). No statistical differences were found in hip flexion at initial contact (SMD: 0.487 95% CI –0.160 to 1.133, $p=0.140$, I^2 : 56%; 4 studies with 93 participants (47 males, 46 females)) (Fig. 2). In the frontal plane, it was revealed that females produced significantly less hip abduction at initial contact (SMD: –0.454, 95% CI 0.151–0.758, $p=0.003$, I^2 : 33%; 4 studies (two with multiple angle conditions) with 176 participants (93 males, 83 females)) (Fig. 2). No significant statistical differences in

Table 4 Variables extracted from retrieved studies

Outcome measures	Precise mean and SD provided for injury risk factors between males and females Biomechanics: Knee abduction, rotation, flexion moments / impulse (knee joint loads) [70–74] Proximal anterior tibial shear [75] Technical, kinetic, or kinematic determinants of surrogates of injury risk (knee joint loads) [36, 70, 76] at initial contact (IC) (first instance of ground contact in COD), range of motion (ROM; defined as point of IC to maximum knee flexion) and peak stance (peak value obtained during the full cutting cycle 0–100%) related to quadriceps, ligament, trunk, and leg dominance, such as: Vertical / posterior GRF/ impulse <i>Initial or peak</i> Lateral trunk flexion/rotation angle Hip internal rotation angle Knee valgus / internal rotation angle Knee flexion / hip flexion Foot progression angles Rearfoot/ forefoot strike Coordination changes Asymmetries
Other data extraction	The following data were extracted and recorded in a spreadsheet: (1) Author names, publication year and country of origin (2) Sample size and participant characteristics including sport(s), playing level/status, training history, strength history/status/profile, reproductive status(females), hormonal contraception usage (3) Angle of COD task (4) Anticipatory nature of COD task (planned/unplanned) (5) If unanticipated, method and timing of stimulus presentation (6) Approach velocity prior to COD (7) How approach velocity was controlled for / calculated (8) Rest period between trials (if stated) (9) How risk factor (ACL surrogate) was assessed (methods) (10) Reliability and familiarisation stated for outcome measures / tasks (11) Outcome measures (mean, SD) (12) Any other empirical data available for a variable that could mitigate for any sex differences in surrogate injury risk identified (e.g. strength or experience / playing time?) (13) For female populations, information relating to: a. Reproductive status b. Menstrual cycle phase c. Hormonal Contraception use

ACL—Anterior Cruciate Ligament, COD—Change of Direction, GRF—Ground Reaction Force, HC—Hormonal Contraception, MSK—Musculoskeletal, SD—Standard Deviation

peak hip abduction during stance were observed (SMD: 0.273, 95% CI 0.223–0.770, $p=0.281$, $I^2: 57%$; 7 measured groups with 156 participants (78 males, 78 females)) (Fig. 2). In the transverse plane, meta-analysis identified females produced significantly less internal rotation at initial contact (SMD: 0.437, 95% CI 0.134–0.741, $p=0.005$, $I^2: 34%$; 6 measured groups with 176 participants (93 males, 83 females)) while no statistical difference was found for peak hip internal rotation during stance (SMD: 0.093, 95% CI –0.436 to 0.623, $p=0.730$, $I^2: 62%$; 7 measured groups with 156 participants (78 males, 78 females)) (Fig. 2).

Knee

In the sagittal plane, meta-analysis indicated that females produced significantly less peak knee flexion during stance than males (SMD: 0.374, 95% CI 0.098–0.649), $p=0.008$, $I^2: 0%$; 9 measured groups with 209 participants (105 males, 104 females)), while no statistical differences were found for knee flexion at initial contact

(SMD: 0.304, 95% CI 0.098–0.649, $p=0.220$, $I^2: 64%$; 9 measured groups with 204 participants (102 males, 102 females)) (Fig. 2). No statistical difference was found for knee flexion range of motion (SMD: 0.184, 95% CI –0.222 to 0.591, $p=0.374$, $I^2: 43%$; 6 measured groups with 169 participants (78 males, 91 females)) (Fig. 2). In the frontal plane, females displayed significantly greater knee abduction angles at initial contact (SMD: 0.687, 95% CI 0.299–1.076, $p=0.001$, $I^2: 55%$; 9 measured groups with 259 participants (134 males, 125 females)), but no statistical difference was found for peak knee abduction angle during stance (SMD: 0.140, 95% CI 0.690–0.971, $p=0.740$, $I^2: 87%$; 9 measured groups with 206 participants (103 males, 103 females)) (Fig. 2). In the transverse plane, knee rotation at initial contact and peak rotation during stance both indicated no statistical differences between males and females (SMD: 0.125, 95% CI –0.179 to 0.428, $p=0.421$, $I^2: 0%$; 7 measured groups with 169 participants (84 males, 85 females); (SMD: 0.068, 95% CI –0.224 to 0.360, $p=0.646$, $I^2: 37%$; 8 measured groups




Fig. 1 Flow chart illustrating the different phases of the search strategy and study selection process. Key: 3D—Three dimensional, WoS—Web of Science

with 186 participants (93 males, 93 females respectively)) (Fig. 2).

Ankle

In the sagittal plane, no statistical differences were observed between males and females at initial contact (SMD: 0.122, 95% CI -0.380 to 0.623, $p=0.623$, $I^2: 0\%$; 3 measured groups with 62 participants (31 males, 31 females) and peak values during stance (SMD: 0.690, 95% CI -2.388-1.008, $p=0.425$, $I^2: 89\%$; 3 measured groups with 62 participants (31 males, 31 females)) (Fig. 2).

Trunk

Insufficient evidence was synthesised to perform quantitative analysis with only one study [61] reporting outcome measures related to the trunk (forward and lateral inclination).

Quantitative Synthesis: Kinetics

Knee Joint Loads

In the sagittal plane, meta-analysis revealed no statistical differences between males and females for externally applied peak knee flexion moment (SMD: -0.056, 95% CI -0.379 to 0.266, $p=0.731$, $I^2: 0\%$; 5 measured groups with 150 participants (69 males, 81 females)) and internal resultant knee extension moments (SMD: 0.104, 95% CI -0.592 to 0.384, $p=0.677$, $I^2: 0\%$; 3 measured groups with 65 participants (32 males, 33 females)) (Fig. 3).

Regarding frontal plane knee moments, no statistical effect was observed between males and females for externally applied peak knee abduction moments (SMD: 0.078, 95% CI -0.382 to 0.537, $p=0.740$, $I^2: 55\%$; 6 measured groups with 170 participants (79 males, 91 females)) and peak internal resultant knee adduction moments (SMD: -0.270, 95% CI -0.611 to 0.07, $p=0.121$, $I^2: 42\%$;

Table 5 Summary of study characteristics

Study	Participants and classification [67]	COD task	Familiarisation/reliability	Velocity (m.s)	Outcomes	Comments
Beaulieu et al. [85]	15 Female: (21.1 ± 3.6 years, 168.3 ± 5.3 cm, 62.4 ± 4.9 kg) 15 Male: (22.9 ± 3.7 years, 178.2 ± 8.0 cm, 75.1 ± 6.7 kg) Elite soccer players (Tier 4/3)	45° UA cut	Separate practice/familiarisation session	4.0–5.0	3D Kinematics (Sagittal, frontal, transverse) 3D kinetics (Sagittal, frontal, transverse) Peak values and at IC	Only study to control for MC phase: Tested all females in follicular phase of MC
Condello et al. [57]	12 Females: (21.0 ± 2.7 years, 166.9 ± 4.1 cm, 61.9 ± 6.2 kg) 14 Males: (19.6 ± 1.5 years, 178.9 ± 4.7 cm, 73.1 ± 7.2 kg) NCAA Div III soccer players (Tier 3)	60° PP cut (each limb)	Submaximal trials ICCs of outcome variables	Not stated	vGRF, mGRF, contact time, Performance cutting angle, minimum horizontal distance (foot plant width)	Only kinematics and forces measured
Iguchi et al. [58]	12 Females: (21.9 ± 1.2 years, 160.6 ± 5.0 cm, 55.0 ± 5.2 kg) 11 Males (22.9 ± 1.0 years, 170.9 ± 4.5 cm, 64.2 ± 7.6 kg) "Recreationally active" (Tier 1)	60° UA cut	5 PP familiarisation trials	3.0	3D hip and knee kinematics (Sagittal) GRF-impulse (50 ms, 100 ms)	CMI fatigue eliciting protocol Pooled intervention conditions when performing sex comparisons
Ihmels et al. [59]	17 Females: (21.6 ± 2.8 years, 1.7 m ± 0.1, 64.6 ± 7.4 kg) 17 Males: (22.3 ± 3.3 years, 1.8 ± 0.1 m, 78.8 ± 10 kg) "Recreationally active" (Tier 1)	45° PP cut	None	4.0	Peak 3D knee kinematics and kinetics (Sagittal, Frontal and Transverse) Peak 3D ankle kinetics and ROM (Sagittal and Frontal)	Sex differences only observed in women wearing prophylactic braces and not control All conditions were pooled for subsequent sex analysis
Khalid et al. [60]	6 Females (19.33 ± 1.97 years, 1.61 ± 0.04 m, 55.4 ± 9.56 kg) 6 Males (20.17 ± 1.83 years, 1.74 ± 0.02 m, 66.58 ± 6.00 kg) Collegiate soccer players (Tier 2)	45° PP and UA cuts	1 h familiarisation session delivered within 3 days of testing	4.5–5.5	3D Kinetics (Sagittal, frontal, transverse) Peak values	Small sample size of 6 Males and 6 Females Fatigue data pooled when performing sex comparisons
McLean et al. [49]	8 Females: (23.2 ± 3.8 years, 167.3 ± 6.5 cm, 64.1 ± 5.0 kg) 8 Males: (23.2 ± 3.8 years, 177.5 ± 8.3 cm, 73.1 ± 3.8 kg) "Proficient in sidestep manoeuvres"	PP 30–40° cut	Not reported	4.5–5.5	Peak 3D Hip and knee kinematics (Sagittal, frontal and transverse) Rearfoot pronation mGRF	No joint moments examined
McLean et al. [48]	10 Females: 21.1 ± 3.0 years, 176.0 ± 11.1 cm, 76.1 ± 12.4 kg 10 Males: 20.2 ± 1.9 years, 184.7 ± 8.0 cm, 81.9 ± 9.8 kg NCAA DIV I Basketball players (Tier 4)	35–55° path demarcated intended for PP 45° cut	Not reported	4.5–5.5	Peak 3D Hip and Knee Kinematics (Sagittal, frontal, transverse); 3D ankle Kinematics (sagittal and frontal) Peak 3D knee kinetics (Frontal)	Only frontal plane kinetics examined

Table 5 (continued)

Study	Participants and classification [67]	COD task	Familiarisation/reliability	Velocity (m.s)	Outcomes	Comments
McLean et al. [88]	10 Females: 21.1 ± 3.0 years, 176.0 ± 11.1 cm, 76.1 ± 12.4 kg 10 Males: 20.2 ± 1.9 years, 184.7 ± 8.0 cm, 81.9 ± 9.8 kg NCAA DIV I Basketball players (Tier 4)	35–55° path demarcated intended for PP 45° cut and PP 180°	Not reported	4.5–5.5	Peak and IC: 3D Hip and Knee Kinematics (Sagittal, frontal, transverse) 3D ankle Kinematics (sagittal and frontal)	No measurement of kinetics
Nagano et al. 2011 [61]	10 Female (7 Soccer 3 Lacrosse) 20.1 ± 1.4 years, 1.61 ± 0.06 m, 56.8 ± 7.4 kg 10 Male (Soccer) 20.7 ± 1.3 years, 1.75 ± 0.05 m, 66.9 ± 6.2 kg, "Cutting sports"	PP 180° turn	"Several preparation trials"	"Maximal"	3D knee kinematics (sagittal, frontal, transverse) at IC and 75 ms and 150 ms 3D trunk kinematics: (sagittal and frontal) at IC and 75 ms and 150 ms	No knee moments calculated
O'Connor et al. [62]	17 Females: 20.9 ± 1.5 years, 1.68 ± 0.06 m, 62.9 ± 5.9 kg 16 Males: 22.7 ± 2.7 years, 1.81 ± 0.08 m, 86.1 ± 13.5 kg "Recreationally active" (Tier 1)	5 trials of UA 45° cut of dominant leg	Separate familiarisation session delivered one week prior to main testing	4.5–5.0	3D knee kinematics at IC and ROM (sagittal, frontal and transverse planes) 3D kinetics (internal joint moments; sagittal, frontal and transverse)	Multiple dynamic conditions pooled together prior to performing sex comparison
Pollard et al. [89]	12 Females: 19.3 ± 1.1 years, 1.66 m ± 0.05 m, 62.5 ± 6.9 kg 12 Males 19.7 ± 1.5 years NCAA Collegiate soccer players (Tier 3)	7–8 trials of UA 45° right legged cut	Practice session including several Preplanned and unanticipated trials	5.5–6.5	Peak 3D hip and knee Kinematics (Frontal and Transverse) Peak 3D hip and knee kinetics (Frontal and Transverse)	No sagittal angles and kinetics calculated despite their association with KILs
Pollard et al. [86]	15 Females: 19.4 ± 1.5 years, 167.4 ± 8.0 cm, 65.9 ± 7.0 kg 15 Males: 19.6 ± 1.9 years, 179.1 ± 6.0 cm, 74.2 ± 7.0 kg NCAA Collegiate Soccer Players (Tier 3)	4 trials of 5 m right legged PP 45° cut	3–5 practice trials	5.5–7.0	Average 3D hip kinematics (Sagittal Frontal and transverse) Peak hip kinetics (Sagittal Frontal and transverse)	Only examined hip kinematics and kinetics
Schreurs et al. [63]	16 Females: 21.9 ± 2.5 years, 173.6 ± 6.9 cm and 67.1 ± 7.1 kg 13 Males: 22.5 ± 2.3 years, 190.2 ± 7.7 cm and 80.1 ± 10.3 kg "Team sport athletes"	10 trials of PP maximal 45°, 90°, 135°, and 180° CODs	None reported	"Maximal"	3D knee kinematics and ROM (Sagittal) 3D knee kinetics (Sagittal and Frontal)	Pooled COD conditions to perform sex analysis Only knee examined Measures reported at instance of peak valgus

Table 5 (continued)

Study	Participants and classification [67]	COD task	Familiarisation/reliability	Velocity (m.s)	Outcomes	Comments
Sigward and Powers 2006 [52]	15 Females: 19.4 ± 1.5 years, 167.4 ± 8.0 cm, 65.9 ± 7.0 kg 15 Males: 19.6 ± 1.9 years, 179.1 ± 6.0 cm, 74.2 ± 7.0 kg NCAA div I or II Soccer Players (Tier 3 and 4)	4 trials of 5 m right legged PP 45° cut	"Practice trials" Test-retest reliability reported	5.5–7.0	3D knee kinematics (Sagittal, frontal and transverse) 3D knee kinetics (Moment and moment impulse) (Sagittal frontal and transverse)	Only study to report test-retest reliability
Sigward et al. [51]	20 Females: 19.7 ± 0.2 years, 167.0 ± 1.6 cm, 51.8 ± 1.7 kg 20 Males: 19.7 ± 0.2 years, 181.1 ± 1.6 cm, 78.6 ± 1.7 kg NCAA div I or II Soccer Players (Tier 3 and 4)	4 trials of UA 45° cut with 7 m approach	"Practice trials"	4.5–5.5	Peak 3D Kinematics (frontal) Peak 3D Kinetics (Frontal internal) vGRF, pGRF, IGRF	Pooled maturation data to conduct sex comparisons
Sigward et al. [50]	20 Females: 18.25 ± 2.15 years, 1.63 ± 0.07 m, 59.55 ± 7.38 kg 25 Males: 18.76 ± 2.09 years, 1.80 ± 0.07 m, 75.58 ± 8.02 kg NCAA div I or II Soccer Players (Tier 3 and 4)	4 trials of UA 45° and 110° cuts with 7 m approach	"Practice trials"	4.5–5.5	3D Hip Kinematics (Frontal and Transverse) 3D Kinetics (Frontal) vGRF, pGRF, IGRF	Pooled data for angles to conduct sex comparisons
Tanikawa et al. [64]	Pooled male and female: 25.4 ± 3.5 years, 1.67 ± 0.1 m, 60.2 ± 10.1 kg	2 trials of 90° cut at self-selected speed	1 practice trial	Self-selected	3D knee kinematics and kinetics (Sagittal, frontal and transverse)	Trials performed barefoot and exceptionally low approach velocity (2.10 m.s)

°—degrees, 3D—Three-dimensional, cm—centimetre, CMJ—countermovement jump, COD—change of direction, GRF—ground reaction force, IC—initial contact, ICC—intra-class correlation coefficient, kg—kilogram, KJL—knee joint loads, IGRF—lateral ground reaction force, MC—menstrual cycle, mGRF—medial ground reaction force, ms—milliseconds, pGRF—posterior ground reaction force, PP—Preplanned, ROM—range of motion, UA—unanticipated, vGRF—vertical ground reaction force

8 measured groups with 249 participants (129 males, 120 females) (Fig. 3). Meta-analysis of transverse plane kinetics indicated no statistical effect of sex on internal resultant peak knee internal rotation moments (SMD: 0.145, 95% CI -0.264 to 0.553, $p=0.487$, $I^2=39%$; 4 measured groups with 95 participants (47 males, 48 females)) (Fig. 3).

Ground Reaction Force

No statistical differences between males and females were observed for vertical GRF (SMD: 0.121, 95% CI -0.369 to 0.611, $p=0.628$, $I^2=70%$; 7 measured groups with 232 participants (116 males, 116 females)) (Fig. 3).

Subgroup and Moderator Analysis

Post hoc subgroup and moderator analyses were performed on variables that produced significant differences between males and females in the primary meta-analysis as follows: sagittal (peak during stance), frontal and transverse (initial contact) hip kinematics, together with sagittal (peak during stance) and frontal (initial contact) plane knee kinematics.

Hip Flexion During Peak Stance

Subgroup analysis for anticipated vs unanticipated CODs revealed no statistically significant difference between males and females in peak hip flexion during stance;(anticipated SMD: 0.809, 95% CI 0.345–1.274, $I^2=0%$; $Z=3.414$, $p=0.001$, $ES=4$, 8 measured groups, 78 participants (39 males, 39 females); Unanticipated: SMD: 0.214, 95% CI -0.237 to 0.666, $I^2=24%$, $Z=0.930$, $p=0.352$, $ES=3$; 6 measured groups, 77 participants (39 male, 38 female)); Total between group analysis: $Q=3.233$, $p=0.072$. COD angle sub analysis was not possible due to the lack of data on COD angles other than 45°.

Subgroup analysis for sporting code revealed no statistical differences between males and females (soccer and basketball athletes) for peak hip flexion during stance;(soccer SMD: 0.194, 95% CI -0.237 to 0.625, $I^2=0%$, $Z=0.883$, $p=0.377$, $ES=3$; 6 measured groups, 84 participants (42 males, 42 females); basketball SMD: 0.880, 95% CI 0.149–1.612, $I^2=1%$, $Z=2.358$, $p=0.018$, $ES=2$; 4 measured groups, 32 participants (16 males, 16 females)); Total between group analysis: $Q=2.508$, $p=0.113$.

Hip Internal Rotation at Initial Contact

Subgroup analysis for anticipated vs unanticipated CODs revealed no statistically significant difference between males and females in hip internal rotation at initial contact;(anticipated SMD: 0.343, 95% CI -0.197 to 0.882, $I^2=59%$, $Z=1.245$, $p=0.213$, $ES=3$; 6 measured groups,

56 participants (28 males, 28 females); unanticipated SMD: 0.481, 95% CI 0.115–0.848, $I^2=24%$, $Z=2.573$, $p=0.010$, $ES=3$; 6 measured groups, 120 participants (65 males, 55 females); Total Between analysis: $Q=0.174$, $p=0.677$. Insufficient data were present on other COD angles to perform subgroup analysis.

Subgroup analysis for sporting code revealed no statistical differences between males and females in hip internal rotation at initial contact;(soccer SMD: 0.398, 95% CI 0.064–0.731, $I^2=20%$, $Z=2.339$, $p=0.019$, $ES=4$; 8 measured groups, 144 participants (77 males, 67 females); basketball SMD: 0.651, 95% CI 0.129–1.520, $I^2=72%$, $Z=1.685$, $p=0.142$, $ES=2$; 4 measured groups, 32 participants (16 males, 16 females)); Total between analysis: $Q=0.237$, $p=0.626$.

Hip Abduction at Initial Contact

Analysis for the effect of anticipation revealed no statistical differences between sexes in hip abduction at initial contact (anticipated SMD: -0.067, 95% CI -0.600 to 0.446, $I^2=46%$, $Z=-0.245$, $p=0.806$, $ES=3$; 6 measured groups, 56 participants (28 males, 28 females); unanticipated SMD: -0.640, 95% CI -1.010 to -0.271, $I^2=0%$, $Z=-3.401$, $p=0.001$, $ES=3$; 6 measured groups, 120 participants (65 males, 55 females)) Total between group analysis: $Q=3.010$, $P=0.083$. Insufficient data were available to perform COD angle dependent analysis.

Sport subgroup analysis revealed a significant difference between soccer and basketball, with male soccer athletes favouring greater levels of hip abduction at initial contact (soccer SMD: -0.643 95% CI -0.980 to -0.307, $I^2=0%$, $Z=-3.744$, $p<0.001$, $ES=4$; 8 measured groups, 144 participants (77 males, 67 females); basketball SMD: 0.362 (95% CI -0.338 to 1.062, $I^2=0%$, $Z=1.013$, $p=0.311$, $ES=2$; 4 measured groups, 32 participants (16 males, 16 females); Total between group analysis: $Q=6.430$, $p=0.011$.

Knee Flexion During Stance

Subsequent analysis indicated anticipation had no statistical effect on the amount of peak knee flexion observed between males and females during the stance phase of the COD (anticipated SMD: 0.433, 95% CI 0.114–0.753, $I^2=0%$, $Z=2.659$, $p=0.008$, $ES=7$; 156 participants (78 males, 78 females); unanticipated SMD: 0.200, 95% CI -0.343 to 0.744, $I^2=28%$, $Z=0.722$, $p=0.471$, $ES=2$; 53 participants (27 males, 26 females)); Total between group analysis: $Q=0.526$, $p=0.468$. Insufficient data were present for subgroup analysis on COD angle.

Subgroup analysis of sport revealed no statistical effect of sport on peak knee flexion angle between males and females during stance (soccer SMD: 0.311, 95% CI -0.122 to 0.744, $I^2=7%$, $Z=1.406$, $p=0.160$, $ES=3$; 6 measured




Fig. 2 Forest plot illustrating standardised mean differences and 95% confidence intervals for determinants of KJLs between males and females. Key: Dorsi—Dorsiflexion, INT—Internal, IC—Initial contact, ROM—Range of motion, ROT—Rotation




Fig. 3 Forest plot illustrating standardised mean differences and 95% confidence intervals for multiplanar KJLs between males and females. Key: N—Newtons per kilogram, vGRF—Vertical ground reaction force

groups, 84 participants (42 males, 42 females); basketball SMD: 0.803 95% CI 0.081–1.525, $I^2=0\%$, $Z=2.180$, $p=0.029$, $ES=2$; 4 measured groups, 32 participants (16 males, 16 females); Physically active SMD: 0.213, 95% CI –0.236 to 0.663, $I^2=0\%$, $Z=0.929$, $p=0.353$, $ES=3$; 6 measured groups, 77 participants (39 males, 38 females)); Total between analysis: $Q=1.900$, $p=0.387$.

Knee Abduction at Initial Contact

Subgroup analysis indicated no statistical effect of anticipation on knee abduction at initial contact between males and females (anticipated SMD: 0.656, 95% CI 0.068–1.245, $I^2=71\%$, $Z=2.185$, $p=0.029$, $ES=5$; 10 measured groups, 106 participants (53 males and 53 females); unanticipated SMD: 0.726, 95% CI 0.153–1.298, $I^2=12\%$, $Z=2.485$, $p=0.013$, $ES=4$; 8 measured groups, 153 participants (81 males, 72 females)); Total Between group analysis: $Q=0.028$, $P=0.868$.

Analysis of COD angle indicated no significant statistical effect on knee abduction angle between males and females at initial contact: 45° SMD: 0.613, 95% CI 0.073–1.152, $I^2=48\%$, $Z=2.226$, $p=0.026$, $ES=6$; 12 measured groups, 178 participants (91 males, 87 females); 180° SMD: 0.936, 95% CI –0.120–1.993, $I^2=87\%$, $Z=1.737$, $p=0.082$, $ES=2$; 4 measured groups, 36 participants (18 males, 18 females); Total Between group analysis: $Q=0.336$, $P=0.845$.

Sporting code subgroup analysis indicated a significant statistical effect of sport with female basketball players generating greater knee abduction angles at initial contact: soccer SMD: 0.650 95% CI 0.271–1.028, $I^2=37\%$, $Z=3.361$, $p=0.001$, $ES=5$; 12 measured groups, 191 participants (102 males, 89 females); basketball SMD: 1.781, 95% CI 0.889–2.672, $I^2=0\%$, $Z=3.915$, $p<0.001$, $ES=2$; 4 measured groups, 32 participants (16 males, 16 females); total between group analysis: $Q=5.239$, $p=0.022$.

Moderator Analysis

Moderator analysis for hip abduction angle at IC observed a significant statistical effect for age (mean male and female age modelled together: $Q=6.36$, $p=0.0416$), mass (mean male and female mass modelled together: $Q=6.49$, $p=0.0390$) and achieved velocity (mean male and female velocity modelled together: $Q=6.58$, $p=0.0372$). Moderator analysis for knee abduction angle at IC revealed a significant effect for methodological quality score ($B=-0.1565$, $p=0.03$).

Discussion

To the authors' knowledge, this is the first meta-analysis that has examined multiplanar whole body kinematics and knee kinetics between males and females during COD tasks. No effect of sex on knee kinetics was

identified despite extensive literature suggesting apparent sex differences in multiplanar KJLs in COD tasks [48, 50–52, 90], although females were identified as demonstrating greater knee abduction at IC and less peak knee flexion during stance. This is counterintuitive when considering the significant differences observed in sagittal and frontal plane knee motion for females that would increase the moment arm of the GRF vector in the relevant plane, and would amplify the KJLs generated by females [70]. Therefore there may be sex differences in the proportion of the KJL that is determined by the lever arm (lower limb) vs the GRF profile (ground impact profile). It is worth noting that the analysis of resultant internal knee adduction moment fell only marginally outside the 95% confidence interval levels for being greater in females (95% CI –0.611 to 0.07, $p=0.121$), suggesting a potential undetected effect of sex. Considering a lack of differences in KJLs, it is feasible that these high-risk kinematics observed in females could have been offset by differences in the GRF profile (the other component of the KJL) whereby males produce more force resulting in comparable KJLs. This was not the case as meta-analysis revealed no significant differences between males and females for vertical GRF ($p=0.628$). A small number of groups were measured in this analysis (seven), and there were only sufficient data present to analyse vertical GRF. Further research investigating this is required, incorporating analysis of multiplanar forces such as lateral GRFs. This would affect the moment arm of the force in the frontal plane during COD tasks and has been observed in a group of females exhibiting “excessive valgus” [44].

Males were identified as producing greater hip abduction and internal rotation at IC, identified as KJL determinants previously due to the intersegmental relationship between hip and knee positioning further down the kinetic chain [40, 44, 91]. A wider foot-plant would allow greater perpendicular forces to be produced achieving more effective task completion in more mechanically demanding CODs ($>60^\circ$) [92], in line with the faster performance times observed by males in the only study reporting performance time [63]. Interestingly, these hip postures did not translate to greater knee abduction angles in males compared to females despite the determinant relationship with KJLs [45]. This raises the question of other confounding factors such as pelvic width to femoral length ratios [93] being responsible for these knee postures in females. From the above, it is possible there are sex differences in the proportion of the KJL that is generated from lower limb kinematics vs the GRF profile, with females adopting more abducted and extended knee postures. It is worth considering that similar KJLs generated by males and females may not translate to similar injury risk, due to morphological differences in strain

characteristics of the ACL between males and females [94, 95]. Therefore, the evidence concerning differences in KJLs and injury risk between males and females remains contentious and requires further investigation.

This meta-analysis indicates that sex has an effect on lower limb kinematics in COD tasks, namely multiplanar hip and knee kinematics. Females were found to execute COD tasks with greater knee abduction and less knee flexion. Considering the relationship between increased knee abduction and limited knee flexion with ACL strain [28, 30, 96], this observation may partially explain some of the disproportionately greater ACL injury incidence between females and males [19, 97, 98]. This finding also aligns with observational studies identifying limited knee flexion and dynamic knee valgus to be apparent features of non-contact ACL injury [9–11], particularly in females [10, 99–101].

Previous studies have examined differences between males and females in both landing [54], cutting [55] and weight-bearing tasks [53], although these results are somewhat inconclusive. Beaulieu et al. [54] identified females as landing with greater peak knee abduction angles than their male counterparts across a range of unilateral and bilateral drop-landings, although a lack of evidence was observed for knee joint loads and other lower limb kinematics. Benjaminse et al. [55] identified small differences between males and females during cutting, with females generating lower knee flexion angles and greater knee abduction angles and moments. However, they questioned the clinical relevance of these findings due to the lack of statistical power in the majority of included studies and inconsistent effect sizes. A limitation of these studies is the absence of any meta-analysis or regression that may identify differences between groups of males and females. Cronstrom et al. [53] detailed that females exhibited greater knee abduction and excursion across a range of weight bearing activities, including cutting, although this analysis only included frontal plane knee kinematics. Considering non-contact ACL injuries [102], ACL strain [96] and KJLs in CODs are multiplanar by nature and the result of a complex interaction of multiple body segments [45], this justified the current investigation of multiplanar kinematics and kinetics of the lower limb to further understand these potential differences between males and females.

In a number of these studies examining KJLs, data had been pooled across a number of conditions prior to performing sex comparisons [51, 59, 60, 63]. Such practice is questionable due to the confounding effects of task [50, 90], which has been previously identified as biomechanically discrete in relation to multiplanar knee joint loads and braking characteristics [103]. Furthermore, effects of maturation [51, 101], fatigue [60] and externally applied

bracing [59] provide another layer of measurement variability that could skew the interpretation of sex differences in COD mechanics should the data be pooled prior to performing sex comparisons. The current findings of this analysis suggest that more carefully controlled research is required examining sex differences in COD biomechanics. Cronstrom et al. [104] identified knee abduction moments as not being predictive of non-contact ACL injuries in a recent meta-analysis despite contrary prospective findings in the literature [105, 106]. It is worth noting that this was across a range of screening tasks that predominantly consisted of drop landings, which have been identified as generating lower multiplanar KJLs compared to COD tasks [107]. Further prospective research is recommended explicitly in COD to identify the predictive utility of multiplanar knee joint loads in ACL injury. It still remains unexplained as to why there is a gynocentric pattern in non-contact ACL epidemiological literature, although a multifactorial approach that includes anatomical and hormonal contributions must be considered alongside multiplanar knee joint loads.

In addition, embedding a gendered, environmental approach [108] into these recommendations may provide further explanation for the discrepancy between males and females in ACL injury rates in multidirectional sports. This approach would account for sociocultural and socio-economic factors that could affect accumulated motor experience and resistance training history, explaining differences in motor skill and strength [109]. This could confound results when comparing males and females as previously stated [110], and lead to differences being attributed due to sex rather than a modifiable confounding variable. Evidence to support this can be seen in ballet, where there is a substantially lower ACL injury incidence (0.009 per 1000 exposures), and sex is not a risk factor for ACL injury [111]. This can be attributed to females receiving targeted training from an early age in high risk movements such as single legged landing, evidenced through comparable KJLs between males and females during this task [112]. Further investigation is required in multidirectional sports to confirm this notion, through matching participants for skill and resistance training history in COD studies examining KJLs. None of the retrieved studies in this meta-analysis controlled for motor skill levels/skill training history or examined differences in co-ordination, and future research should better account for these factors when examining biomechanical differences between males and females.

It is worth noting that none of the studies included in this analysis included a measurement of lower limb strength to serve as a control for sex comparisons. Hip abductor and external rotator strength have

been identified as predictive of ACL injury in male and female populations [113], with pre-adolescent boys demonstrating greater hip external rotator strength, alongside greater gluteus medius activity in the pre-activation phase of a cutting task [114]. Increased lower limb strength, and eccentric strength in particular, would facilitate greater muscular support of the knee joint and maintain postural integrity through simultaneous co-activation of the hamstrings and quadriceps [115], potentially lowering the KJLs generated during CODs [39, 74]. This would also allow more effective utilisation of the penultimate foot-contact as a braking or preparatory step to also lower KJLs and improve performance times as previously demonstrated [39, 74]. Strength has also been identified as a determinant of COD mechanics [116, 117] and COD performance [118, 119]. Stronger individuals typically adopt a hip dominant strategy with greater levels of knee flexion [116, 117] and reduced internal knee extensor moments [117], thus lowering ACL strain [96]. Considering the above, this raises questions about the strength status of female participants in the retrieved studies, as meta-analysis identified females displaying reduced levels of knee flexion.

Therefore, without directly measuring strength qualities and thus, the absence of a strength control in the synthesised literature, it is postulated that relatively stronger males may have been compared with relatively weaker females, providing some potential explanation for these reported differences in KJLs. Collegiate female athletes participate less frequently in strength based training (1.9–3.0 vs 2.6–3.8 days per week) and for a shorter period of time (26–44 min per week vs 49–70 min per week) [120]. This could be explained from a gendered perspective whereby there are negative societal expectations concerning the appropriateness of muscular strength training in females [108, 121–123]. Therefore, as strength is a modifiable risk factor which can be targeted and trained, the relationship between female participation in strength training, strength, and multiplanar KJLs warrants further investigation to provide further insight about potential sex differences in COD biomechanics and ACL injury rates.

This meta-analysis explored sex differences in multiplanar knee joint loads (KJLs) and whole-body kinematics during change-of-direction (COD) tasks. The study extracted data from 17 retrieved articles involving 451 participants. Recommendations include accounting for resistance training history, motor skill experience and co-ordination, and gender-environmental factors in future research. Considering menstrual cycle phase together with variability and reliability is imperative when examining differences in males and females, due to measures of reliability being reported for only two studies [52, 57] and

only one study controlling for menstrual cycle phase [56]. A range of COD angles were used in the included studies. While most data were extracted from 45° CODs and data from other angles pooled, this may not fully examine biomechanical sex differences at larger angles ($\geq 90^\circ$). However, no significant angle-related effects were observed in the subgroup analysis. All data extracted were the product of discrete point analysis, which captures peak values for injury surrogates such as knee abduction angle or moment. However, this approach overlooks the complexity of COD, which involves multiple phases (approach run, deceleration, redirection, and re-propulsion). By focusing on a single data point, 99% of the remaining cutting cycle is discarded [36]. To gain deeper insights, future research should explore non-contact ACL injury risk using statistical parametric mapping methods, considering the entire cutting cycle. This comprehensive approach would enhance our understanding of potential differences between males and females in COD tasks.

Finally, from the findings of this meta-analysis we cannot discern underlying reasons for observed sex differences in COD biomechanics. Other biomechanical, anatomical and hormonal factors such as quadriceps dominance [124], pelvic width to femoral length ratio [93] and fluctuations in serum sex hormones such as oestrogen in the pre-ovulatory phase [125] all likely interact and may contribute to observed differences in surrogates of non-contact ACL injury risk. Furthermore, there was a failure to examine any differences in co-ordination, resistance or skill training history. It is likely these factors contribute to these sex differences, especially when observing these variables through the lens of a gendered environmental approach [108]. Socioeconomic and gender related factors should be acknowledged regarding their contribution to motor skill and strength levels [110]. Further research incorporating all of these modifiable and non-modifiable risk factors is recommended together with better reporting of female demographics [126] to further understand the interaction and respective contributions these have to COD multiplanar KJLs.

Conclusion

This systematic review and meta-analyses revealed no differences between males and females in multiplanar KJLs despite apparent differences between males and females in COD kinematics. Further research is required to identify if this translates to similar injury risk, considering morphological differences in strain characteristics between males and females. Females exhibit increased knee abduction and limited hip and knee flexion compared to males, whereas males display increased hip abduction and internal rotation. Considering the reported difference in knee abduction

and flexion in CODs, there is a need to focus on this movement pattern during COD technique modification programmes in females. All of the included studies included healthy males and females, predominantly of a collegiate or recreational background, therefore further research is required in ACL deficient and elite populations with considerations for socioeconomic factors, skill training history and strength levels.

Abbreviations

3D	Three dimensional
ACL	Anterior cruciate ligament
COD	Change of direction
GRF	Ground reaction force
HC	Hormonal contraception
I ²	I squared
IC	Initial contact
MSK	Musculoskeletal
KJL	Knee joint load
SD	Standard deviation
SMD	Standardised mean difference

Acknowledgements

Not applicable.

Author Contributions

All authors have made substantial contributions to the conception, drafting and writing of this work as per the adapted regulations of McNutt et al. [127] and approved the submitted version and take accountability for their personal contributions. All authors read and approved the final version.

Funding

No sources of funding were used to assist in the preparation of this article.

Availability of Data and Materials

Not applicable.

Declarations

Ethics Approval and Consent to Participate

Not applicable.

Consent for Publication

Not applicable.

Competing interests

T.D., J.E., M.B., P.J., J.O'D., T.D.S. declare that they have no conflict of interest with the content of this article.

Author details

¹Section of Sport Section of Sport, Exercise and Rehabilitation Sciences, School of Human and Life Sciences, Canterbury Christ Church University, North Holmes Road, Canterbury, Kent CT1 1Q, UK. ²School of Health Sciences, C702 Allerton Building, University of Salford, Salford M6 6PU, UK. ³Department of Sport and Exercise Sciences | Manchester Metropolitan University, 2.01 Institute of Sport, 99 Oxford Road, Manchester M1 7EL, UK.

Received: 26 September 2023 Accepted: 20 March 2024

Published online: 01 April 2024

References

- Martínez-Hernández D, Quinn M, Jones P. Linear advancing actions followed by deceleration and turn are the most common movements preceding goals in male professional soccer. *Sci Med Footb*. 2022;7:25–33.
- Chandler PT, Pinder SJ, Curran JD, Gabbett TJ. Physical demands of training and competition in collegiate netball players. *J Strength Cond Res*. 2014;28(10):2732–7.
- Stølen T, Chamari K, Castagna C, Wisløff U. Physiology of soccer: an update. *Sports Med*. 2005;35:501–36.
- Spencer M, Lawrence S, Rechichi C, Bishop D, Dawson B, Goodman C. Time–motion analysis of elite field hockey, with special reference to repeated-sprint activity. *J Sports Sci*. 2004;22(9):843–50.
- Granero-Gil P, Gómez-Carmona CD, Bastida-Castillo A, Rojas-Valverde D, de la Cruz E, Pino-Ortega J. Influence of playing position and laterality in centripetal force and changes of direction in elite soccer players. *PLoS ONE*. 2020;15(4):e0232123.
- Morgan OJ, Drust B, Ade JD, Robinson MA. Change of direction frequency off the ball: New perspectives in elite youth soccer. *Sci Med Footb*. 2022;6(4):473–82.
- Bloomfield J, Polman R, O'Donoghue P. Physical demands of different positions in FA premier league soccer. *J Sports Sci Med*. 2007;6(1):63.
- Lopes TJA, Simic M, Pappas E. Epidemiology of anterior cruciate ligament reconstruction in Brazil's public health system. *Revista Brasileira de Medicina do Esporte*. 2016;22:297–301.
- Montgomery C, Blackburn J, Withers D, Tierney G, Moran C, Simms C. Mechanisms of ACL injury in professional rugby union: a systematic video analysis of 36 cases. *Br J Sports Med*. 2018;52(15):994–1001.
- Olsen O-E, Myklegust G, Engebretsen L, Bahr R. Injury mechanisms for anterior cruciate ligament injuries in team handball: a systematic video analysis. *Am J Sports Med*. 2004;32(4):1002–12.
- Johnston JT, Mandelbaum BR, Schub D, Rodeo SA, Matava MJ, Silvers-Granelli HJ, et al. Video analysis of anterior cruciate ligament tears in professional american football athletes. *Am J Sports Med*. 2018;46(4):862–8.
- Sanders TL, Maradit Kremers H, Bryan AJ, Larson DR, Dahm DL, Levy BA, et al. Incidence of anterior cruciate ligament tears and reconstruction: a 21-year population-based study. *Am J Sports Med*. 2016;44(6):1502–7.
- Janssen K, Orchard J, Driscoll T, van Mechelen W. High incidence and costs for anterior cruciate ligament reconstructions performed in Australia from 2003–2004 to 2007–2008: Time for an anterior cruciate ligament register by Scandinavian model? *Scand J Med Sci Sports*. 2012;22(4):495–501.
- Zbrojkiewicz D, Vertullo C, Grayson JE. Increasing rates of anterior cruciate ligament reconstruction in young Australians, 2000–2015. *Med J Aust*. 2018;208(8):354–8.
- Herzog MM, Marshall SW, Lund JL, Pate V, Mack CD, Spang JT. Trends in incidence of ACL reconstruction and concomitant procedures among commercially insured individuals in the United States, 2002–2014. *Sports Health*. 2018;10(6):523–31.
- Maniar N, Verhagen E, Bryant AL, Opar DA. Trends in Australian knee injury rates: An epidemiological analysis of 228,344 knee injuries over 20 years. *Lancet Reg Health Western Pac*. 2022;21:100409.
- Boden BP, Dean GS, Feagin JA, Garrett WE. Mechanisms of anterior cruciate ligament injury. *Thorofare, NJ: SLACK Incorporated*; 2000. p. 573–8.
- Gornitzky AL, Lott A, Yellin JL, Fabricant PD, Lawrence JT, Ganley TJ. Sport-specific yearly risk and incidence of anterior cruciate ligament tears in high school athletes: a systematic review and meta-analysis. *Am J Sports Med*. 2016;44(10):2716–23.
- Montalvo AM, Schneider DK, Yut L, Webster KE, Beynonn B, Kocher MS, et al. "What's my risk of sustaining an ACL injury while playing sports?" A systematic review with meta-analysis. *Br J Sports Med*. 2019;53(16):1003–12.
- Gans I, Retzky JS, Jones LC, Tanaka MJ. Epidemiology of recurrent anterior cruciate ligament injuries in National Collegiate Athletic Association sports: the Injury Surveillance Program, 2004–2014. *Orthop J Sports Med*. 2018;6(6):2325967118777823.
- Shah VM, Andrews JR, Fleisig GS, McMichael CS, Lemak LJ. Return to play after anterior cruciate ligament reconstruction in National Football League athletes. *Am J Sports Med*. 2010;38(11):2233–9.
- Shelbourne KD, Nitz P. Accelerated rehabilitation after anterior cruciate ligament reconstruction. *Am J Sports Med*. 1990;18(3):292–9.
- Davies L, Cook J, Leal J, Areia CM, Shirkey B, Jackson W, et al. Comparison of the clinical and cost effectiveness of two management strategies (rehabilitation versus surgical reconstruction) for non-acute anterior

- cruciate ligament (ACL) injury: study protocol for the ACL SNNAP randomised controlled trial. *Trials*. 2020;21:1–16.
24. Morrey MA, Stuart MJ, Smith AM, Wiese-Bjornstal DM. A longitudinal examination of athletes' emotional and cognitive responses to anterior cruciate ligament injury. *Clin J Sport Med*. 1999;9(2):63–9.
 25. Lohmander LS, Ostenberg A, Englund M, Roos H. High prevalence of knee osteoarthritis, pain, and functional limitations in female soccer players twelve years after anterior cruciate ligament injury. *Arthritis Rheum*. 2004;50(10):3145–52.
 26. Ardern CL, Webster KE, Taylor NF, Feller JA. Return to the preinjury level of competitive sport after anterior cruciate ligament reconstruction surgery: two-thirds of patients have not returned by 12 months after surgery. *Am J Sports Med*. 2011;39(3):538–43.
 27. Meeuwisse WH, Tyreman H, Hagel B, Emery C. A dynamic model of etiology in sport injury: the recursive nature of risk and causation. *Clin J Sport Med*. 2007;17(3):215–9.
 28. Markolf KL, Burchfield DM, Shapiro MM, Shepard MF, Finerman GA, Slaughterbeck JL. Combined knee loading states that generate high anterior cruciate ligament forces. *J Orthop Res*. 1995;13(6):930–5.
 29. Shin CS, Chaudhari AM, Andriacchi TP. The influence of deceleration forces on ACL strain during single-leg landing: a simulation study. *J Biomech*. 2007;40(5):1145–52.
 30. Shin CS, Chaudhari AM, Andriacchi TP. The effect of isolated valgus moments on ACL strain during single-leg landing: a simulation study. *J Biomech*. 2009;42(3):280–5.
 31. Shin CS, Chaudhari AM, Andriacchi TP. Valgus plus internal rotation moments increase anterior cruciate ligament strain more than either alone. *Med Sci Sports Exerc*. 2011;43(8):1484–91.
 32. Bates NA, Schilaty ND, Nagelli CV, Krych AJ, Hewett TE. Multiplanar loading of the knee and its influence on anterior cruciate ligament and medial collateral ligament strain during simulated landings and noncontact tears. *Am J Sports Med*. 2019;47(8):1844–53.
 33. Hewett TE, Myer GD, Ford KR, Heidt RS Jr, Colosimo AJ, McLean SG, et al. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: a prospective study. *Am J Sports Med*. 2005;33(4):492–501.
 34. Beaulieu ML, Ashton-Miller JA, Wojtyk EM. Loading mechanisms of the anterior cruciate ligament. *Sports Biomech*. 2023;22(1):1–29.
 35. Dos'Santos T, Thomas C, McBurnie A, Donelon T, Herrington L, Jones PA. The cutting movement assessment score (CMAS) qualitative screening tool: application to mitigate anterior cruciate ligament injury risk during cutting. *Biomechanics*. 2021;1(1):83–101.
 36. Dos'Santos T, Stebbings GK, Morse C, Shashidharan M, Daniels KA, Sanderson A. Effects of the menstrual cycle phase on anterior cruciate ligament neuromuscular and biomechanical injury risk surrogates in eumenorrhic and naturally menstruating women: a systematic review. *Plos ONE*. 2023;18(1):e0280800.
 37. Dempsey AR, Lloyd DG, Elliott BC, Steele JR, Munro BJ. Changing side-step cutting technique reduces knee valgus loading. *Am J Sports Med*. 2009;37(11):2194–200.
 38. Jamison ST, Pan X, Chaudhari AM. Knee moments during run-to-cut maneuvers are associated with lateral trunk positioning. *J Biomech*. 2012;45(11):1881–5.
 39. Jones PA, Herrington L, Graham-Smith P. Braking characteristics during cutting and pivoting in female soccer players. *J Electromyogr Kinesiol*. 2016;30:46–54.
 40. Jones PA, Herrington LC, Graham-Smith P. Technique determinants of knee joint loads during cutting in female soccer players. *Hum Mov Sci*. 2015;42:203–11.
 41. Jones PA, Herrington LC, Graham-Smith P. Technique determinants of knee abduction moments during pivoting in female soccer players. *Clin Biomech*. 2016;31:107–12.
 42. McLean SG, Huang X, van den Bogert AJ. Association between lower extremity posture at contact and peak knee valgus moment during sidestepping: implications for ACL injury. *Clin Biomech*. 2005;20(8):863–70.
 43. Sigward SM, Cesar GM, Havens KL. Predictors of frontal plane knee moments during side-step cutting to 45 and 110 degrees in men and women: implications for anterior cruciate ligament injury. *Clin J Sport Med*. 2015;25(6):529–34.
 44. Sigward SM, Powers CM. Loading characteristics of females exhibiting excessive valgus moments during cutting. *Clin Biomech*. 2007;22(7):827–33.
 45. Donelon TA, Dos'Santos T, Pitchers G, Brown M, Jones PA. Biomechanical determinants of knee joint loads associated with increased anterior cruciate ligament loading during cutting: a systematic review and technical framework. *Sports Med Open*. 2020;6(1):1–21.
 46. Dos'Santos T, Thomas C, Comfort P, Jones PA. Biomechanical effects of a 6-week change of direction speed and technique modification intervention: implications for change of direction side step performance. *J Strength Cond Res*. 2021;36:2780–91.
 47. Dos'Santos T, Thomas C, McBurnie A, Comfort P, Jones PA. Change of direction speed and technique modification training improves 180 degrees turning performance, kinetics, and kinematics. *Sports*. 2021;9(6):73.
 48. McLean SG, Huang X, van den Bogert AJ. Association between lower extremity posture at contact and peak knee valgus moment during sidestepping: implications for ACL injury. *Clin Biomech*. 2005;20(8):863–70.
 49. McLean SG, Lipfert SW, Van Den Bogert AJ. Effect of gender and defensive opponent on the biomechanics of sidestep cutting. *Med Sci Sports Exerc*. 2004;36(6):1008–16.
 50. Sigward SM, Cesar GM, Havens KL. Predictors of frontal plane knee moments during side-step cutting to 45 and 110 degrees in men and women: implications for anterior cruciate ligament injury. *Clin J Sport Med*. 2015;25(6):529–34.
 51. Sigward SM, Pollard CD, Havens KL, Powers CM. Influence of sex and maturation on knee mechanics during side-step cutting. *Med Sci Sports Exerc*. 2012;44(8):1497–503.
 52. Sigward SM, Powers CM. The influence of gender on knee kinematics, kinetics and muscle activation patterns during side-step cutting. *Clin Biomech*. 2006;21(1):41–8.
 53. Cronström A, Creaby MW, Nae J, Ageberg E. Gender differences in knee abduction during weight-bearing activities: a systematic review and meta-analysis. *Gait Posture*. 2016;49:315–28.
 54. Beaulieu ML, McLean SG. Sex-dimorphic landing mechanics and their role within the noncontact ACL injury mechanism: evidence, limitations and directions. *Sports Med Arthrosc Rehabil Ther Technol*. 2012;4:1–13.
 55. Benjaminse A, Gokeler A, Fleisig GS, Sell TC, Otten B. What is the true evidence for gender-related differences during plant and cut maneuvers? A systematic review. *Knee Surg Sports Traumatol Arthrosc*. 2011;19:42–54.
 56. Beaulieu ML, Lamontagne M, Xu L. Lower limb muscle activity and kinematics of an unanticipated cutting manoeuvre: a gender comparison. *Knee Surg Sports Traumatol Arthrosc*. 2009;17(8):968–76.
 57. Condello G, Kernozek TW, Tessitore A, Foster C. Biomechanical analysis of a change-of-direction task in collegiate soccer players. *Int J Sports Physiol Perform*. 2016;11(1):96–101.
 58. Iguchi J, Tateuchi H, Taniguchi M, Ichihashi N. The effect of sex and fatigue on lower limb kinematics, kinetics, and muscle activity during unanticipated side-step cutting. *Knee Surg Sports Traumatol Arthrosc*. 2014;22(1):41–8.
 59. Ihmels WD, Seymore KD, Brown TN. Effect of sex and ankle brace design on knee biomechanics during a single-leg cut. *Am J Sports Med*. 2020;48(6):1496–504.
 60. Khalid AJ, Harris SI, Michael L, Joseph H, Qu X. Effects of neuromuscular fatigue on perceptual-cognitive skills between genders in the contribution to the knee joint loading during side-stepping tasks. *J Sports Sci*. 2015;33(13):1322–31.
 61. Nagano Y, Ida H, Akai M, Fukubayashi T. Relationship between three-dimensional kinematics of knee and trunk motion during shuttle run cutting. *J Sports Sci*. 2011;29(14):1525–34.
 62. O'Connor KM, Monteiro SK, Hoelker IA. Comparison of selected lateral cutting activities used to assess ACL injury risk. *J Appl Biomech*. 2009;25(1):9–21.
 63. Schreurs MJ, Benjaminse A, Lemmink KA. Sharper angle, higher risk? The effect of cutting angle on knee mechanics in invasion sport athletes. *J Biomech*. 2017;63:144–50.
 64. Tanikawa H, Matsumoto H, Komiyama I, Kiriyama Y, Toyama Y, Nagura T. Comparison of knee mechanics among risky athletic motions

- for noncontact anterior cruciate ligament injury. *J Appl Biomech*. 2013;29(6):749–55.
65. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Int J Surg*. 2021;88:105906.
66. Wager E, Wiffen PJ. Ethical issues in preparing and publishing systematic reviews. *J Evid Based Med*. 2011;4(2):130–4.
67. McKay AK, Stellingwerff T, Smith ES, Martin DT, Mujika I, Goosey-Tolfrey VL, et al. Defining training and performance caliber: a participant classification framework. *Int J Sports Physiol Perform*. 2021;17(2):317–31.
68. Dos'Santos T, McBurnie A, Thomas C, Jones PA, Harper D. Attacking agility actions: match play contextual applications with coaching and technique guidelines. *Strength Condit J*. 2022;44(5):102–18.
69. Dos'Santos T, Jones P. 18 Training for change of direction and agility. *Advanced Strength and Conditioning: An Evidence-based Approach*. 2022:328.
70. Donelon TA, Dos'Santos T, Pitchers G, Brown M, Jones PA. Biomechanical determinants of knee joint loads associated with increased anterior cruciate ligament loading during cutting: a systematic review and technical framework. *Sports Med Open*. 2020;6(1):53.
71. McLean SG, Huang X, van den Bogert AJ. Investigating isolated neuromuscular control contributions to non-contact anterior cruciate ligament injury risk via computer simulation methods. *Clin Biomech*. 2008;23(7):926–36.
72. Markolf KL, Burchfield DM, Shapiro MM, Shepard MF, Finerman GA, Slauterbeck JL. Combined knee loading states that generate high anterior cruciate ligament forces. *J Orthop Res*. 1995;13(6):930–5.
73. McBurnie AJ, Dos' Santos T, Jones PA. Biomechanical associates of performance and knee joint loads during A 70–90° cutting maneuver in subelite soccer players. *J Strength Cond Res*. 2019;35:3190–8.
74. Dos'Santos T, Thomas C, McBurnie A, Comfort P, Jones PA. Biomechanical determinants of performance and injury risk during cutting: a performance-injury conflict? *Sports Med*. 2021;51:1983–98.
75. Yu B, Lin C-F, Garrett WE. Lower extremity biomechanics during the landing of a stop-jump task. *Clin Biomech*. 2006;21(3):297–305.
76. Fox AS. Change-of-direction biomechanics: Is what's best for anterior cruciate ligament injury prevention also best for performance? *Sports Med*. 2018;48(8):1799–807.
77. Khalid AJ, Ian Harris S, Michael L, Joseph H, Qu X. Effects of neuromuscular fatigue on perceptual-cognitive skills between genders in the contribution to the knee joint loading during side-stepping tasks. *J Sports Sci*. 2015;33(13):1322–31.
78. Brown SR, Brughelli M, Hume PA. Knee mechanics during planned and unplanned sidestepping: a systematic review and meta-analysis. *Sports Med*. 2014;44:1573–88.
79. Dos'Santos T, Bishop C, Thomas C, Comfort P, Jones PA. The effect of limb dominance on change of direction biomechanics: A systematic review of its importance for injury risk. *Phys Ther Sport*. 2019;37:179–89.
80. Brown SR, Brughelli M, Hume PA. Knee mechanics during planned and unplanned sidestepping: a systematic review and meta-analysis. *Sports Med*. 2014;44(11):1573–88.
81. Brown S, Brughelli M, Hume P. Knee mechanics during planned and unplanned sidestepping: a systematic review and meta-analysis. *Sports Med*. 2014;44(11):1573–88.
82. Giesche F, Stief F, Groneberg DA, Wilke J. Effect of unplanned athletic movement on knee mechanics: a systematic review with multilevel meta-analysis. *Br J Sports Med*. 2021;55:1366–78.
83. Cohen J. *Statistical power analysis for the behavioral sciences*. New York: Academic Press; 2013.
84. Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ*. 1997;315(7109):629–34.
85. Beaulieu ML, Lamontagne M, Xu L, Beaulieu ML, Lamontagne M, Xu L. Lower limb muscle activity and kinematics of an unanticipated cutting manoeuvre: a gender comparison. *Knee Surg Sports Traumatol Arthrosc*. 2009;17(8):968–76.
86. Pollard CD, Sigward SM, Powers CM. Gender differences in hip joint kinematics and kinetics during side-step cutting maneuver. *Clin J Sport Med*. 2007;17(1):38–42.
87. Brooks A, Meade CS, Potter JS, Lohnygina Y, Calsyn DA, Greenfield SF. Gender differences in the rates and correlates of HIV risk behaviors among drug abusers. *Subst Use Misuse*. 2010;45(14):2444–69.
88. McLean S, Walker K, van den Bogert AJ. Effect of gender on lower extremity kinematics during rapid direction changes: an integrated analysis of three sports movements. *J Sci Med Sport*. 2005;8(4):411–22.
89. Pollard CD, Davis IM, Hamill J. Influence of gender on hip and knee mechanics during a randomly cued cutting maneuver. *Clin Biomech*. 2004;19(10):1022–31.
90. Almansouri S, Barclay W, Dighe A, Gong C, Chow M, Sigward S. Can increased gait speed improve knee loading mechanics following anterior cruciate ligament reconstruction? *Med Sci Sports Exerc*. 2020;52(17):248.
91. Kristianslund E, Faul O, Bahr R, Myklebust G, Krosshaug T. Sidestep cutting technique and knee abduction loading: implications for ACL prevention exercises. *Br J Sports Med*. 2014;48(9):779–83.
92. Dos'Santos T, Thomas C, Comfort P, Jones PA. The effect of angle and velocity on change of direction biomechanics: an angle-velocity trade-off. *Sports Med*. 2018;48(10):2235–53.
93. Pantano KJ, White SC, Gilchrist LA, Leddy J. Differences in peak knee valgus angles between individuals with high and low Q-angles during a single limb squat. *Clin Biomech*. 2005;20(9):966–72.
94. Lipps DB, Oh YK, Ashton-Miller JA, Wojtys EM. Morphologic characteristics help explain the gender difference in peak anterior cruciate ligament strain during a simulated pivot landing. *Am J Sports Med*. 2012;40(1):32–40.
95. Schilaty ND, Bates NA, Nagelli C, Krych AJ, Hewett TE. Sex-based differences in knee kinetics with anterior cruciate ligament strain on cadaveric impact simulations. *Orthop J Sports Med*. 2018;6(3):2325967118761037.
96. Bates NA, Schilaty ND, Nagelli CV, Krych AJ, Hewett TE. Multiplanar loading of the knee and its influence on anterior cruciate ligament and medial collateral ligament strain during simulated landings and noncontact tears. *Am J Sports Med*. 2019;47(8):1844–53.
97. Ireland ML. The female ACL: Why is it more prone to injury? *Orthopedic Clinics*. 2002;33(4):637–51.
98. Griffin LY, Agel J, Albohm MJ, Arendt EA, Dick RW, Garrett WE, et al. Non-contact anterior cruciate ligament injuries: risk factors and prevention strategies. *J Am Acad Orthop Surg*. 2000;8(3):141–50.
99. Brophy RH, Stepan JG, Silvers HJ, Mandelbaum BR. Defending puts the anterior cruciate ligament at risk during soccer: a gender-based analysis. *Sports Health Multidiscip Approach*. 2015;7(3):244–9.
100. Stuelcken MC, Mellifont DB, Gorman AD, Sayers MG. Mechanisms of anterior cruciate ligament injuries in elite women's netball: a systematic video analysis. *J Sports Sci*. 2016;34(16):1516–22.
101. Ghasemi M, Sigurdsson HB, Sveinsson P, Briem K. Boys demonstrate greater knee frontal moments than girls during the impact phase of cutting maneuvers, despite age-related increases in girls. *Knee Surg Sports Traumatol Arthrosc*. 2023;31(5):1833–9.
102. McLean SG, Huang X, Su A, Van Den Bogert AJ. Sagittal plane biomechanics cannot injure the ACL during sidestep cutting. *Clin Biomech*. 2004;19(8):828–38.
103. Dos'Santos T, Thomas C, Jones PA. The effect of angle on change of direction biomechanics: comparison and inter-task relationships. *J Sports Sci*. 2021;39(22):2618–31.
104. Cronström A, Creaby MW, Ageberg E. Do knee abduction kinematics and kinetics predict future anterior cruciate ligament injury risk? A systematic review and meta-analysis of prospective studies. *BMC Musculoskelet Disord*. 2020;21(1):1–11.
105. Hewett TE, Myer GD, Ford KR, Heidt RS Jr, Colosimo AJ, McLean SG, et al. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: a prospective study. *Am J Sports Med*. 2005;33(4):492–501.
106. Zazulak BT, Hewett TE, Reeves NP, Goldberg B, Cholweicki J. Deficits in neuromuscular control of the trunk predict knee injury risk: a prospective biomechanical-epidemiologic study. *Am J Sports Med*. 2007;35(7):1123–30.
107. Jones PA, Herrington LC, Munro AG, Graham-Smith P. Is there a relationship between landing, cutting, and pivoting tasks in terms of the characteristics of dynamic valgus? *Am J Sports Med*. 2014;42(9):2095–102.
108. Parsons JL, Coen SE, Bekker S. Anterior cruciate ligament injury: towards a gendered environmental approach. *Br J Sports Med*. 2021;55(17):984–90.

109. Bruton MR, O'Dwyer N, Adams R. Sex differences in the kinematics and neuromuscular control of landing: Biological, environmental and sociocultural factors. *J Electromyogr Kinesiol.* 2013;23(4):747–58.
110. Nimphius S. Exercise and sport science failing by design in understanding female athletes. *Int J Sports Physiol Perform.* 2019;14(9):1157–8.
111. Liederbach M, Dilgen FE, Rose DJ. Incidence of anterior cruciate ligament injuries among elite ballet and modern dancers: a 5-year prospective study. *Am J Sports Med.* 2008;36(9):1779–88.
112. Orishimo KF, Kremenic IJ, Pappas E, Hagins M, Liederbach M. Comparison of landing biomechanics between male and female professional dancers. *Am J Sports Med.* 2009;37(11):2187–93.
113. Khayambashi K, Ghoddosi N, Straub RK, Powers CM. Hip muscle strength predicts noncontact anterior cruciate ligament injury in male and female athletes. *Am J Sports Med.* 2016;44(2):355–61.
114. Jonasson G, Ghasemi M, Jonsdottir U, Briem K, Sveinsson P. Effects of gender and fatigue on strength and activity of gluteus medius muscle during a controlled cutting maneuver in preadolescent athletes. *J Electromyogr Kinesiol.* 2023;70:102779.
115. Hughes G. A review of recent perspectives on biomechanical risk factors associated with anterior cruciate ligament injury. *Res Sports Med.* 2014;22(2):193–212.
116. Spiteri T, Cochrane JL, Hart NH, Haff GG, Nimphius S. Effect of strength on plant foot kinetics and kinematics during a change of direction task. *Eur J Sport Sci.* 2013;13(6):646–52.
117. Davies WT, Ryu JH, Graham-Smith P, Goodwin JE, Cleather DJ. Stronger subjects select a movement pattern that may reduce anterior cruciate ligament loading during cutting. *J Strength Cond Res.* 2022;36(7):1853–9.
118. Jones PA, Dos'Santos T, McMahon JJ, Graham-Smith P. Contribution of eccentric strength to cutting performance in female soccer players. *J Strength Condit Res.* 2022;36(2):525–3.
119. Jones PA, Thomas C, Dos'Santos T, McMahon JJ, Graham-Smith P. The role of eccentric strength in 180 turns in female soccer players. *Sports.* 2017;5(2):42.
120. Nuzzo JL. Narrative review of sex differences in muscle strength, endurance, activation, size, fiber type, and strength training participation rates, preferences, motivations, injuries, and neuromuscular adaptations. *J Strength Cond Res.* 2023;37(2):494–536.
121. Dworkin SL. "Holding back": negotiating a glass ceiling on women's muscular strength. *Sociol Perspect.* 2001;44(3):333–50.
122. Brace-Govan J. Weighty matters: Control of women's access to physical strength. *The Sociological Review.* 2004;52(4):503–31.
123. Salvatore J, Marecek J. Gender in the gym: evaluation concerns as barriers to women's weight lifting. *Sex Roles.* 2010;63:556–67.
124. Pappas E, Shiyko MP, Ford KR, Myer GD, Hewett TE. Biomechanical deficit profiles associated with ACL injury risk in female athletes. *Med Sci Sports Exerc.* 2016;48(1):107.
125. Herzberg SD, Motu'apuaka ML, Lambert W, Fu R, Brady J, Guise J-M. The effect of menstrual cycle and contraceptives on ACL injuries and laxity: a systematic review and meta-analysis. *Orthop J Sports Med.* 2017;5(7):2325967117718781.
126. Elliott-Sale KJ, Minahan CL, de Jonge XAJ, Ackerman KE, Sipilä S, Constantini NW, et al. Methodological considerations for studies in sport and exercise science with women as participants: a working guide for standards of practice for research on women. *Sports Med.* 2021;51(5):843–61.
127. McNutt MK, Bradford M, Drazen JM, Hanson B, Howard B, Jamieson KH, et al. Transparency in authors' contributions and responsibilities to promote integrity in scientific publication. *Proc Natl Acad Sci.* 2018;115(11):2557–60.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.