




Please cite the Published Version

Burden, Richard J , Altini, Marco, Ferrer, Eva, Flood, Tessa R , Lewin, Gary, Nimphius, Sophia, Phillips, Stuart M and Elliott-Sale, Kirsty J  (2024) Editorial: Measure do not guess: a call to action to end assumed and estimated menstrual cycle phases in research. *BMJ Open Sport & Exercise Medicine*, 10 (2). e002095

DOI: <https://doi.org/10.1136/bmjsem-2024-002095>

Publisher: BMJ

Version: Published Version

Downloaded from: <https://e-space.mmu.ac.uk/635174/>



Usage rights:  [Creative Commons: Attribution-Noncommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/)

Additional Information: This is an open access editorial published in *BMJ Open Sport & Exercise Medicine*.

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

Measure do not guess: a call to action to end assumed and estimated menstrual cycle phases in research

Richard J Burden ^{1,2}, Marco Altini,³ Eva Ferrer,^{4,5} Tessa R Flood,² Gary Lewin,⁶ Sophia Nimphius,⁷ Stuart M Phillips,⁸ Kirsty J Elliott-Sale ²

To cite: Burden RJ, Altini M, Ferrer E, *et al.* Measure do not guess: a call to action to end assumed and estimated menstrual cycle phases in research. *BMJ Open Sport & Exercise Medicine* 2024;**10**:e002095. doi:10.1136/bmjsem-2024-002095

Accepted 5 June 2024



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹UK Sports Institute, Manchester, UK

²Department of Sport and Exercise Sciences, Manchester Metropolitan University Institute of Sport, Manchester, UK

³Department of Human Movement Sciences, Vrije Universiteit Amsterdam, Amsterdam, Netherlands

⁴Barça Innovation Hub of Football Club Barcelona, Barcelona, Spain

⁵Sports and Exercise Medicine Unit, Hospital Clinic and Sant Joan de Déu, Barcelona, Spain

⁶Arsenal Women's Football Club, London, UK

⁷School of Medical and Health Sciences, Edith Cowan University, Joondalup, Western Australia, Australia

⁸Department of Kinesiology, McMaster University, Hamilton, Ontario, Canada

Correspondence to

Professor Kirsty J Elliott-Sale; k.elliott-sale@mmu.ac.uk

INTRODUCTION

Several studies have been published in the last year using assumed or estimated menstrual cycle phases. These phases are 'generated' solely on regular menstruation. As more of these studies are published, founded on data that was not actually measured, it becomes imperative to grasp the difference between what is measured versus what is assumed or estimated. We need to measure, not guess, to draw valid and meaningful conclusions in a research topic that is riddled with considerable debate over contradicting outcomes. Hence, this editorial is a call to action directed at editors and reviewers. It is intended to raise the quality, practical implications, application and integrity of future studies investigating the effect of menstrual cycle phases on a given outcome. We do not intend to focus on individual papers but want to ensure that female populations can use their reproductive data accurately and effectively.

THE PROBLEM

Sometimes, a parameter of interest (eg, ovulation) can be measured in various ways (eg, urinary luteinising hormone kits or transvaginal ultrasound), which are open to debate, but nevertheless, these are still measurements; nothing is guessed. When you assume something, you accept it is true or certain to happen without proof. When you estimate something, you take a guess (ie, a rough calculation based on something else). In both cases, you have not used a technique that allows you actually to measure the parameter of interest. You exploit the relationship between a somewhat associated variable (ie, regular menstruation) and a parameter (ie, the remaining phases of the menstrual cycle). As such, it is assumed that an event has occurred (eg, ovulation), and then the timing of that event is estimated (eg, it occurs at the halfway point of the cycle). Assumptions and

estimations are made regardless of what is happening (eg, assuming ovulation has definitely occurred when, in reality, it may not have). These assumptions and estimates are not always harmless; many ovarian hormone profiles exist, and regular menstruation does not preclude menstrual dysfunctions, which go undetected with assumed and estimated phases. When adopting an assumed or estimated approach transparent reporting, including the quantification of confidence in these assumptions and estimations, is of utmost importance. While some limitations are often described, and rarely—if ever—are all of the implications of these assumptions and estimations clearly explained to the reader or user. Studies using an assumed and estimated menstrual cycle phase approach frequently do us a disservice; for example, they use the same terminology when assuming or estimating as other authors use when measuring (eg, using specific phase names associated with measurements—such as 'early luteal'—when they have not measured but assumed that phase and should in fact use the broader term 'non-bleeding day'). When the prevailing narrative is that female populations have been underserved by research, accepting assumptions and estimates is not the progress it is purported to be.

THE SOLUTION

The way to establish menstrual cycle phases in research (ie, laboratory and field-based studies) is clear: in addition to noting regular bleeding, investigators need to measure ovulation and progesterone, noting that the ideal frequency and modality of these measurements are still to be determined. Studies using this approach date back to the early 2000s¹ and guidance on this approach has been produced several times within the last 5 years.^{2 3} Although this approach may



be challenging in field-based studies, it is certainly not impossible.

CALL TO ACTION

Given the problem and solution provided here, we strongly urge reviewers and editors to insist on measurements rather than guesses. If measurements were not made, it is imperative that authors provide (1) the rationale why measurements were not made; (2) a completely transparent description of the limitations associated with their study design and (3) a full disclosure of the implications of their assumptions and estimations on research quality, confidence in the findings, ability—or rather inability—to make scientific inferences, and the specific clinical dangers associated with assumptions and estimations.

X Richard J Burden @rich_burden and Kirsty J Elliott-Sale @ElliottSale

Acknowledgements We acknowledge the work of de Zambotti et al (2024), whose publication helped to shape the narrative used within the present letter; de Zambotti M, Goldstein C, Cook J, Menghini L, Altini M, Cheng P, Robillard R. State of the science and recommendations for using wearable technology in sleep and circadian research. *Sleep*. 2024;47(4):zsad325.

Contributors All authors wrote, reviewed and approved the final manuscript. KE-S is the guarantor and accepts full responsibility for the finished work and controlled the decision to publish.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests The contents of this Editorial are presented clearly, honestly and without fabrication, falsification or inappropriate data manipulation. KE-S, TRF, GL, MA, SN, EF and SP have no professional relationships with companies or manufacturers who will benefit from the contents of the present Editorial. RJB also has no current professional relationships with companies or manufacturers who will benefit from the contents of the present Editorial. In the interest of transparency, he was formerly a paid consultant to Mint Diagnostics, a diagnostics technology company that is no longer trading.

Patient consent for publication Not applicable.

Provenance and peer review Commissioned; internally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Richard J Burden <http://orcid.org/0000-0002-2811-5454>

Kirsty J Elliott-Sale <http://orcid.org/0000-0003-1122-5099>

REFERENCES

- 1 Elliott KJ, Cable NT, Reilly T, *et al*. Effect of menstrual cycle phase on the concentration of bioavailable 17-beta oestradiol and testosterone and muscle strength. *Clin Sci* 2003;105:663–9.
- 2 Janse DE Jonge X, Thompson B, Han A. Methodological recommendations for Menstrual cycle research in sports and exercise. *Med Sci Sports Exerc* 2019;51:2610–7.
- 3 Elliott-Sale KJ, Minahan CL, de Jonge XAKJ, *et al*. Methodological considerations for studies in sport and exercise science with women as participants: a working guide for standards of practice for research on women. *Sports Med* 2021;51:843–61.