



Please cite the Published Version

Hardy, Edward J, Bass, Joseph J, Inns, Thomas B, Piasecki, Mathew, Piasecki, Jessica, Sale, Craig , Morris, Robert H, Lund, Jonathan N, Smith, Ken, Wilkinson, Daniel J, Atherton, Philip J and Phillips, Bethan E  (2024) Exploring the utility of ultrasound to assess disuse atrophy in different muscles of the lower leg. *Journal of Cachexia, Sarcopenia and Muscle*. ISSN 2190-5991

DOI: <https://doi.org/10.1002/jcsm.13583>

Publisher: Wiley

Version: Published Version

Downloaded from: <https://e-space.mmu.ac.uk/635375/>

Usage rights:  [Creative Commons: Attribution 4.0](https://creativecommons.org/licenses/by/4.0/)

Additional Information: This is an open access article which first appeared in *Journal of Cachexia, Sarcopenia and Muscle*

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

Exploring the utility of ultrasound to assess disuse atrophy in different muscles of the lower leg

Edward J. Hardy^{1,2}, Joseph J. Bass¹, Thomas B. Inns¹, Mathew Piasecki¹, Jessica Piasecki³, Craig Sale^{3,4}, Robert H. Morris³, Jonathan N. Lund^{1,2}, Ken Smith¹, Daniel J. Wilkinson¹, Philip J. Atherton¹ & Bethan E. Phillips^{1*} 

¹Centre of Metabolism, Ageing & Physiology (COMAP), MRC-Versus Arthritis Centre for Musculoskeletal Ageing Research (CMAR), and Nottingham NIHR Biomedical Research Centre, University of Nottingham, School of Medicine, Derby, UK; ²Department of Surgery, Royal Derby Hospital, Derby, UK; ³School of Science and Technology, Nottingham Trent University, Nottingham, UK; ⁴Institute of Sport, Manchester Metropolitan University, Manchester, UK

Abstract

Background Skeletal muscle is a highly plastic tissue crucial for many functions associated with whole-body health across the life course. Magnetic resonance imaging (MRI) is the current gold standard for measuring skeletal muscle size. However, MRI is expensive, and access to facilities is often limited. B-mode ultrasonography (U/S) has been proposed as a potential alternative to MRI for the assessment of muscle size. However, to date, no work has explored the utility of U/S to assess disuse muscle atrophy (DMA) across muscles with different atrophy susceptibility profiles, an omission which may limit the clinical application of previous work.

Methods To address this significant knowledge gap, 10 young men ($22 \pm$ years, 24.1 ± 2.3 kg/m²) underwent 15-day unilateral leg immobilization using a knee-brace and air boot. Cross-sectional area (CSA) and muscle thickness (MT) of the tibialis anterior (TA) and medial gastrocnemius (MG) were assessed via U/S before and after immobilization, with CSA and muscle volume assessed via MRI.

Results With both muscles combined, there were good correlations between each U/S and MRI measure, both before (e.g., CSA_{MRI} vs. MT_{U/S} and CSA_{U/S}: $r = 0.88$ and 0.94 , respectively, both $P < 0.0001$) and after (e.g., VOL_{MRI} vs. MT_{U/S} and CSA_{U/S}: $r = 0.90$ and 0.96 , respectively, both $P < 0.0001$) immobilization. The relationship between the methods was notably stronger for MG than TA at each time-point (e.g., CSA_{MRI} vs. MT_{U/S}: MG, $r = 0.70$, $P = 0.0006$; TA, $r = 0.37$, $P = 0.10$). There was no relationship between the degree of DMA determined by the two methods in either muscle (e.g., TA pre- vs. post-immobilization, VOL_{MRI}: 136 ± 6 vs. 133 ± 5 , $P = 0.08$; CSA_{U/S}: 6.05 ± 0.3 vs. 5.92 ± 0.4 , $P = 0.70$; relationship between methods: $r = 0.12$, $P = 0.75$).

Conclusions Both MT_{U/S} and CSA_{U/S} provide comparable static measures of lower leg muscle size compared with MRI, albeit with weaker agreement in TA compared to MG. Although both MT_{U/S} and CSA_{U/S} can discern differences in DMA susceptibility between muscles, neither can reliably assess degree of DMA. Based on the growing recognition of heterogeneous atrophy profiles between muscles, and the topical importance of less commonly studied muscles (i.e., TA for falls prevention in older adults), future research should aim to optimize accessible methods to determine muscle losses across the body.

Keywords Disuse; Imaging; MRI; Muscle; Ultrasound

Received: 14 January 2024; Revised: 18 July 2024; Accepted: 23 July 2024

*Correspondence to: Bethan E. Phillips, School of Medicine, Centre of Metabolism, Ageing & Physiology (COMAP), University of Nottingham Medical School at Derby, Royal Derby Hospital, Derby DE22 3DT, UK. Email: beth.phillips@nottingham.ac.uk
Edward J Hardy and Joseph J Bass contributed equally.

Introduction

Skeletal muscle is the largest organ in the human body, comprising ~40% of whole-body mass in healthy adults.¹ Crucial for many functions associated with whole-body health beyond its most recognized role in locomotion, skeletal muscle also plays a fundamental role in energy homeostasis, oxygen consumption, energy metabolism, and substrate turnover and storage.² A highly plastic tissue, skeletal muscle mass is maintained in health via a dynamic equilibrium between muscle protein synthesis (MPS) and muscle protein breakdown (MPB), with amino acid (AA) nutrition and contractile activity accepted as the most potent anabolic drivers.³ Pairing these drivers leads to increased MPS in response to nutrition⁴ and ultimately skeletal muscle hypertrophy [i.e., as is commonly seen with resistance exercise training (RET)]. Conversely, disuse or inactivity leads to reductions in MPS,⁵ ultimately leading to skeletal muscle atrophy. With skeletal muscle atrophy also a resultant impact of disease (i.e., cancer cachexia⁶), ageing (e.g., in sarcopenia⁷), and traumatic events (e.g., burns⁸ and sepsis⁹), when occurring as a result of decreased or absent contractile activity, it is often referred to as disuse muscle atrophy (DMA). As is seen in response to RET,¹⁰ a reproducible observation in relation to DMA is substantial inter-individual heterogeneity⁵, highlighting the need for accessible methods to determine DMA across different cohorts and individuals.

Over the last five decades, there has been a step-change in the methods available to quantify skeletal muscle size, including the introduction of magnetic resonance imaging (MRI¹¹), computed tomography (CT¹²), and dual-energy X-ray absorptiometry (DXA¹³). Despite each of these methods being widely used, not only for assessment of muscle size, but also in clinical practice for a variety of diagnostic/prognostic endpoints (e.g., the assessment of brain lesions, tumour growth, and osteoporotic progression, respectively), they are each associated with expensive equipment, the need for highly trained operators/interpreters, and in the case of CT and DXA, ionizing radiation exposure.

In more recent years, ultrasound (U/S) has emerged as a potential additional tool for the assessment of muscle size in both young and older healthy cohorts,^{14,15} and more recently in specific clinical cohorts including those with chronic obstructive pulmonary disease (COPD)¹⁶ or intensive care unit patients.¹⁷ Previous studies have reported a positive relationship between U/S-derived measures of muscle thickness (MT_{U/S}) and CT-derived muscle size,¹² DXA-derived lean mass,¹³ and MRI (the gold standard for skeletal muscle size assessment)-derived cross-sectional area (CSA_{MRI}) and muscle volume (VOL_{MRI}).¹⁸

In addition to MT_{U/S}, recent work has shown that muscle size measured as CSA by U/S (CSA_{U/S}) also shows good agreement with MRI.¹⁹ This includes work by Stokes *et al.*, who concluded, after a study of 10-weeks RET to elicit hypertro-

phy and 2-weeks immobilization of the contralateral limb, that CSA_{U/S} was a suitable alternative for measuring vastus lateralis (VL) changes in response to both increased and decreased muscle loading in young men.²⁰ Similarly, Franchi and colleagues reported that RET-induced hypertrophic changes in VL MT_{U/S} correlated with changes in VL CSA_{MRI}, but not VOL_{MRI}.¹⁴ Beyond measures in the VL, Kositsky *et al.* showed that CSA_{U/S} can be used to reliably measure hamstring muscle and tendon size.²¹ Further, Sponbeck *et al.* showed a significant relationship between CSA_{U/S} and CSA_{MRI} across different posterior muscles of the lower leg.¹⁹

Despite this existent body of work, most previous studies that have assessed the utility of U/S (CSA_{U/S} and/or MT_{U/S}) to measure muscle mass or size have reported on one muscle/muscle group only (e.g., VL⁵), with the upper portion of the leg (i.e., quadriceps and hamstrings) most common,²² likely given its functional importance in both athletic performance (e.g., jumping²³) and activities of daily living (e.g., rising from a chair²⁴). However, it has previously been shown that rates of DMA are not uniform across different muscles, even within a similar anatomical region (i.e., the lower leg¹⁸). In addition, the lower leg muscles have been shown to have significant functional importance in relation to gait and balance²⁵ and as such, from a clinical perspective, falls prevention.²⁶

Therefore, the aim of this study was to determine if U/S-derived measures of MT_{U/S} or CSA_{U/S} could be used to accurately estimate changes in muscle size, as assessed by MRI, across different muscles of the lower leg known to have different profiles of atrophy susceptibility (i.e., tibialis anterior (TA) and medial gastrocnemius (MG)¹⁸).

Methods

Ethics approval and participants

This study was reviewed and approved by the University of Nottingham Faculty of Medicine and Health Sciences Research Ethics Committee (FMHS-103-1809) and registered online at ClinicalTrials.gov (NCT04199923). All procedures were conducted in accordance with the Declaration of Helsinki, and all participants provided written informed consent. All ethical guidelines for authorship and publishing in the *Journal of Cachexia, Sarcopenia and Muscle* were also followed.²⁷

Ten recreationally active, young, healthy males (22 ± 4 years, 24.1 ± 2.3 kg/m²) participated in this study. Participants were screened by medical questionnaire, physical assessment, and resting electrocardiogram, with exclusions for cardiovascular, metabolic, and respiratory disorders, or other symptoms of ill-health. Participants had clinically normal blood chemistry and pressure, were not prescribed any

medication, undertook regular activities of daily living and recreation, but had not participated in any exercise training regime in the last 12 months.

Experimental protocol

Each participant underwent 15 days of unilateral limb immobilization (ULI) using a hinged leg brace (Knee Post-Op Cool) and air-boot (Rebound Air Walker, both Ossur, Iceland), and ambulated on crutches (after training) throughout this period. The leg brace was fitted on the dominant leg over a compression sock around the thigh and lower leg, and fixed at 75° knee flexion to ensure no weight bearing could occur and allow sufficient ground clearance of the air-boot (Figure 1). This leg was then also placed into an air-boot with the ankle fixed in a neutral position to ensure no plantar or dorsi-flexion (Figure 1). Signed 'tamper tags' were fitted to indicate if the brace or boot had been modified, which would have resulted in participant exclusion. No participants were excluded.

Prior to and after 15 days of immobilization, each participant visited the research unit for ultrasound and MRI analysis as described below. No adverse events were reported during this study.

Magnetic resonance imaging

Participants were placed into the MRI scanner feet first, supine and instructed to relax for a minimum of 10-min prior

to scanning to normalize fluid shifts in the body. A 1.5T MRI system (Avanto, Siemens, Munich, Germany) was used to collect images of the leg from above the patella, facilitating collection of CSA_{MRI} and VOL_{MRI} measures from the TA and MG. An imaging matrix of 512 x 235 with a resolution of 835 x 835 μm was acquired with a slice thickness of 5 mm using a turbo spin echo sequence with an echo time set to the minimum value of 12 ms and a repetition time of 568 ms to optimize the trade-off between imaging time and contrast for a proton density weighted image. A Siemens peripheral angiography coil was used to maximize the signal to noise ratio of resulting images. Scans were analysed by the same individual using Slicer (v4.10) software, with TA and MG individually segmented by pixel count every third slice before semi-automatic filling between slices and confirming muscle boundaries to generate 3D muscle volumes (Figure 2A), with muscle cross-sectional area measured at 50% of the length of the muscle, determined through the number of slices in each muscle (Figure 2B).

Ultrasound imaging

After the MRI scan, ultrasound images were obtained with the participants leg extended and their ankle relaxed ($\sim 90^\circ$) as per the positioning for the MRI scans. All U/S scans were made with the probe resting on a gel layer without depressing the underlying skin.²⁸ As previously described, TA and MG were scanned at 30% of their length on the mid-sagittal line.¹⁸ TA was measured from the mid-point of the patella on the anterior side of the leg to the fibula end, and MG from the inner knee crease to the fibula end. These anatomical landmarks were chosen to standardize scanning locations and consider variation in leg length between participants. For $MT_{U/S}$ measures along with fibre length (Lf), images were captured using B-mode ultrasonography (Mylab 70, Esaote Biomedica, Italy), with the transducer aligned in the fascicle plane (Figure 2C). Ultrasound Sarcopenia Index (USI)²⁹ was calculated as the ratio between fibre length and muscle thickness (Lf/MT). $CSA_{U/S}$ was measured using panoramic image acquisition²⁸ in the axial-plane at 30% of the muscle length (Figure 2D). Quantification of MT and CSA was then performed using ImageJ (Version 1.53) software, with $MT_{U/S}$ determined as distance between the superficial and deep aponeuroses and an average across three images per muscle used for quantification. All ultrasound scans were performed and analysed in a blinded manner by the same individual.

Statistical analysis

All analyses were performed using GraphPad Prism (v10.1.1). Correlative analysis was undertaken via Pearson's correlation, with r values stated. Columns depict mean \pm SEM, with anal-



Figure 1 Representative unilateral lower leg immobilization using a leg brace and air-boot, with supportive crutches for ambulation.




Figure 2 (A) MRI image of tibialis anterior (TA; green) and medial gastrocnemius (MG; orange) and representative 3D segmentation volume analysis. (B) MRI image with representative cross-sectional area (CSA). Representative ultrasound images of TA (C) and MG (D) with muscle thickness and CSA analysis shown.

ysis via paired *t*-tests within methodology. Bland–Altman analysis is reported as bias (SD of bias) and 95% limits of agreement. Significance was accepted as $P < 0.05$.

Results

Magnetic resonance imaging versus ultrasound for the assessment of muscle volume

Based on the entire data set of both muscles at baseline, there was a significant relationship between CSA_{MRI} and both U/S-derived measures of $MT_{U/S}$ ($r = 0.88$) and $CSA_{U/S}$ ($r = 0.94$) (Figure 3A). There was also a significant relationship between VOL_{MRI} and both $MT_{U/S}$ ($r = 0.91$) and $CSA_{U/S}$ ($r = 0.97$) (Figure 3B), and between the two U/S measures ($r = 0.93$) and the two MRI measures ($r = 0.96$). The *P*-value for each of these correlations was < 0.0001 .

After immobilization these relationships were maintained, with both CSA_{MRI} (Figure 3C) and VOL_{MRI} (Figure 3D) having a significant relationship with both $MT_{U/S}$ (CSA_{MRI} : $r = 0.83$; VOL_{MRI} : $r = 0.90$) and $CSA_{U/S}$ (CSA_{MRI} : $r = 0.90$; VOL_{MRI} : $r = 0.96$). The intra-system relationships were also maintained (U/S: $r = 0.86$; MRI: $r = 0.96$). Unsurprisingly, when both timepoints were combined to offer an enhanced number of data sets for comparison, there remained a significant relationship between both CSA_{MRI} ($MT_{U/S}$: $r = 0.86$; $CSA_{U/S}$: $r = 0.92$) and VOL_{MRI} ($MT_{U/S}$: $r = 0.90$; $CSA_{U/S}$: $r = 0.96$) and each U/S measure. The *P*-value for each of these correlations was < 0.0001 .

When both timepoints (pre- and post-immobilization) were combined but the two muscles (TA and MG) were analysed separately, there was a significant relationship between CSA_{MRI} and both $MT_{U/S}$ ($r = 0.70$, $P = 0.0006$) and $CSA_{U/S}$ ($r = 0.70$, $P = 0.0006$) for MG, and stronger significant relationships between VOL_{MRI} and both $MT_{U/S}$ ($r = 0.90$, $P < 0.0001$) and $CSA_{U/S}$ ($r = 0.90$, $P < 0.0001$) (Figure 3E). For TA, the relationship between CSA_{MRI} and $MT_{U/S}$ was non-significant ($r = 0.37$, $P = 0.10$), and although the relationship with $CSA_{U/S}$ was statistically significant, the relationship was weaker than for MG ($r = 0.57$, $P = 0.009$). Similarly, VOL_{MRI} measures of TA displayed significant but weaker correlations with both $MT_{U/S}$ ($r = 0.48$, $P = 0.03$) and $CSA_{U/S}$ ($r = 0.53$, $P = 0.02$) (Figure 3F).

Absolute values for both muscles via all methods pre- and post-immobilization can be seen in Table S1.

MRI versus ultrasound for the determination of atrophy susceptibility

Using VOL_{MRI} as the previously reported gold standard measure of muscle volume, there were clear differences in the rates of loss between muscles. The overall percentage loss across TA and MG combined was $-5.21 \pm 1.2\%$, with losses in each of these muscles $-2.04 \pm 1.31\%$ and $-8.40 \pm 1.57\%$, respectively. Highlighting inter-individual differences in DMA, overall changes ranged from -16.91 to 7.45% when both muscles were combined, with TA changes of -7.07 to 7.45% and MG changes of -16.91 to -2.37% .





Figure 3 Relationships between muscle size (medial gastrocnemius (MG) and tibialis anterior (TA) combined) at baseline (A and B) and after 15-day unilateral limb immobilization (C and D) in 10 individuals using magnetic resonance imaging (MRI) compared to ultrasound (U/S). MRI measures include cross-sectional area (CSA) and muscle volume (VOL). U/S measures include CSA and muscle thickness (MT). Panels (E) and (F) show the relationship for VOL via MRI compared with both U/S measures for MG and TA, respectively. Analysis via Pearson’s correlation. Significance accepted as $P < 0.05$.

Both U/S and MRI measures each detected significant DMA in the MG (CSA_{MRI}, $P = 0.02$; VOL_{MRI}, $P = 0.002$; MT_{U/S}, $P = 0.008$; CSA_{U/S}, $P = 0.0005$) (Figure 4A) but not the TA (CSA_{MRI}, $P = 0.74$; VOL_{MRI}, $P = 0.08$; MT_{U/S}, $P = 0.60$; CSA_{U/S}, $P = 0.70$) (Figure 4B).

Despite a significant relationship between both MRI measures and each U/S measure at baseline and after immobilization, and that each measure indicated atrophy resistance in the TA (compared to susceptibility in the MG), there was no significant relationship between either MRI measure with ei-



Data shown is mean (SEM)

Figure 4 Skeletal muscle size of (A) medial gastrocnemius and (B) tibialis anterior at baseline (light) and after 15-day unilateral limb immobilization (dark) in 10 individuals using magnetic resonance imaging (MRI) and ultrasound (U/S) methods. MRI measures include muscle volume (VOL) and cross-sectional area (CSA). U/S measures include CSA and muscle thickness (MT). Values are mean \pm SEM. Analysis via paired *t*-tests within methodology. Significance accepted as $P < 0.05$. ** $P < 0.01$; *** $P < 0.001$; ns = non-significant.

ther U/S-derived parameter for degree of DMA over the 15 days of immobilization. Initially analysed with both muscles combined and based on absolute values, this remained true when the two muscles were analysed separately, and when percentage change was used (Table 1).

Recognizing the recent development of the USI as a tool to determine low muscle mass (albeit using the VL), we sought to determine if this method also had utility in determining the degree of DA in the TA and MG as muscles with different atrophy susceptibility profiles. The USI (where a higher value is associated with lower muscle mass) was able to identify the TA as atrophy resistant (pre-immobilization: 5.10 ± 0.35 vs. post-immobilization: 5.74 ± 0.36 , $P = 0.08$) compared with the atrophy susceptible MG (1.90 ± 0.05 vs. 2.30 ± 0.18 post, $P = 0.03$), yet there was no relationship between the degree of DA in either muscle assessed by VOL_{MRI} (the gold standard) compared to the USI ($r = 0.08$, $P = 0.75$).

When the two muscles were grouped together Bland–Altman analysis suggests that compared to VOL_{MRI}, both MT_{U/S} and CSA_{U/S} each appear to underestimate the degree of DMA. This was also true when comparing MT_{U/S} and CSA_{U/S} to CSA_{MRI} when analyzing the two muscles separately (Table 1).

Baseline muscle size versus degree of disuse atrophy

Using the entire data set there was a significant relationship between baseline size and degree of DMA (absolute values)

using both MRI methods (CSA_{MRI}: $r = -0.66$, $P = 0.002$; VOL_{MRI}: $r = -0.78$, $P < 0.0001$) and both U/S methods (CSA_{U/S}: $r = -0.61$, $P = 0.004$; MT_{U/S}: $r = -0.48$, $P = 0.031$), highlighting greater losses in those with larger baseline size. This remained true when DMA was considered as percentage change (data not shown).

However, when the muscles were analysed separately, there was no significant relationship between baseline muscle size and degree of absolute (TA, CSA_{U/S}: $r = -0.15$, $P = 0.68$; MT_{U/S}: $r = -0.52$, $P = 0.13$; MG, CSA_{U/S}: $r = -0.21$, $P = 0.57$; MT_{U/S}: $r = -0.29$, $P = 0.42$) or percentage change in either muscle via U/S. MRI methods did identify a significant relationship between baseline size and absolute DMA, although only via CSA_{MRI} for TA [$r = -0.65$, $P = 0.041$ (VOL_{MRI}: $r = -0.48$, $P = 0.16$)] and VOL_{MRI} for MG [$r = -0.58$, $P = 0.08$ (CSA_{MRI}: $r = -0.37$, $P = 0.29$)]. When DMA was presented as percentage change neither MRI method showed a significant relationship with baseline muscle size for either muscle.

Intra-system assessment of disuse muscle atrophy

Finally, when assessing DMA via the two different U/S measures (i.e., CSA_{U/S} vs. MT_{U/S}) and the two different MRI measures (i.e., CSA_{MRI} vs. VOL_{MRI}), there was a significant relationship for each only when the entire data set was used. There was no relationship between either intra-system measures for TA when the muscles were analysed separately (Table 2), with a significant relationship only for the MRI mea-

Table 1 Relationships between disuse muscle atrophy measured using different imaging techniques after 15-day unilateral limb immobilization in 10 individuals

Absolute change	MT _{U/S}	CSA _{U/S}
CSA _{MRI} - Both muscles	$r = 0.37, P = 0.11$	$r = 0.36, P = 0.12$
CSA _{MRI} - TA	$r = 0.56, P = 0.09$	$r = 0.04, P = 0.92$
CSA _{MRI} - MG	$r = -0.16, P = 0.66$	$r = -0.06, P = 0.88$
VOL _{MRI} - Both muscles	$r = 0.24, P = 0.31$	$r = 0.40, P = 0.08$
VOL _{MRI} - TA	$r = 0.22, P = 0.55$	$r = 0.22, P = 0.53$
VOL _{MRI} - MG	$r = 0.02, P = 0.97$	$r = -0.08, P = 0.82$
Percentage change		
CSA _{MRI} - Both muscles	$r = 0.44, P = 0.05$	$r = 0.09, P = 0.71$
CSA _{MRI} - TA	$r = 0.54, P = 0.11$	$r = -0.009, P = 0.98$
CSA _{MRI} - MG	$r = -0.39, P = 0.27$	$r = -0.35, P = 0.32$
VOL _{MRI} - Both muscles	$r = 0.21, P = 0.39$	$r = 0.21, P = 0.35$
VOL _{MRI} - TA	$r = 0.17, P = 0.65$	$r = 0.12, P = 0.75$
VOL _{MRI} - MG	$r = -0.06, P = 0.87$	$r = -0.26, P = 0.47$
Bland–Altman analysis		
CSA _{MRI} - Both muscles	-0.41 (0.7), -1.8 to 0.96	0.39 (1.2), -1.9 to 2.7
CSA _{MRI} - TA	-0.034 (0.50), -1.0 to 0.95	-0.06 (1.1), -2.2 to 2.3
CSA _{MRI} - MG	-0.8 (0.68), -2.1 to 0.53	0.72 (1.2), -1.6 to 3.1
VOL _{MRI} - Both muscles	-14.18 (17.52), -48.52 to 20.15	-13.38 (17.1), -46.89 to 20.13
VOL _{MRI} - TA	-3.1 (4.9), -13 to 6.5	-3 (4.8), -12 to 6.4
VOL _{MRI} - MG	-25 (19), -62 to 11	-24 (19), -61 to 13

Abbreviations: CSA, cross-sectional area; TA, tibialis anterior; MG, medial gastrocnemius; MRI, magnetic resonance imaging; VOL, volume; MT, muscle thickness; U/S, ultrasound. Analysis via Pearson’s correlation and Bland–Altman analysis. Bland–Altman analysis shows bias (SD of bias) and 95% limits of agreement. Significance accepted as $P < 0.05$.

Table 2 Relationships between muscle atrophy measured using different imaging techniques on the same equipment after 15-day unilateral limb immobilization in 10 individuals

Equipment	Correlation	Bland–Altman analysis
MRI	VOL _{MRI} vs. CSA _{MRI}	Bias (SD), 95% limits of agreement
Both muscles	$r = 0.68, P = 0.0009$	-13.77 (17.06), -47.2 to 19.66
TA	$r = 0.11, P = 0.77$	-3.06 (4.9), -12.66 to 6.54
MG	$r = 0.65, P = 0.04$	-24.48 (18.3), -60.36 to 11.39
Ultrasound	MT _{U/S} vs. CSA _{U/S}	
Both muscles	$r = 0.48, P = 0.03$	-0.81 (1.17), -3.11 to 1.5
TA	$r = 0.48, P = 0.17$	-0.09 (0.94), -1.94 to 1.76
MG	$r = 0.28, P = 0.43$	-1.52 (0.95), -3.37 to 0.33

Analysis via Pearson’s correlation and Bland–Altman analysis. Bland–Altman analysis shows bias (SD of bias) and 95% limits of agreement. Significance accepted as $P < 0.05$.

CSA, cross-sectional area; MG, medial gastrocnemius; MRI, magnetic resonance imaging; MT, muscle thickness; TA, tibialis anterior; U/S, ultrasound; VOL, volume.

ures for MG. Bland–Altman analysis suggests that compared to VOL_{MRI}, CSA_{MRI} appears to underestimate the degree of DMA. Similarly, compared to MT_{U/S}, CSA_{U/S} also appears to underestimate muscle loss.

Discussion

Assessment of changes in skeletal muscle size is an essential aspect of research related to DMA and clinical practice. Although MRI-based measures are considered the gold standard, in recent years U/S has been promoted as a viable alternative for measuring muscle size, including in both hypertrophic¹⁴ and atrophic³⁰ situations, due to its non-invasive and generally accessible nature. However, the ability of U/S to assess DMA in muscles with differing degrees

of atrophy susceptibility has yet to be determined. Here, we demonstrate that static measures of MT_{U/S} and CSA_{U/S} each strongly correlate with MRI-derived measures of both CSA and VOL, before and after a period of immobilization. Moreover, when assessing individual muscles, these correlations were observed in muscles with both atrophy resistant (i.e., TA) and atrophy susceptible (i.e., MG) profiles. However, despite this, neither MT_{U/S} or CSA_{U/S} could resolve the degree of DMA in either muscle, or indeed when both muscles were combined.

Relatively short periods of disuse are known to rapidly induce DMA,³¹ with heterogenous rates of inter-muscle atrophy observed as a result of, for example, prolonged bed-rest.³² Moreover, we have previously characterized this apparent atrophy resistant versus atrophy susceptible (aRaS) paradigm in TA and MG muscles (respectively) in response to ULI,¹⁸ illustrating marked heterogeneity in inter-muscle DMA

even within a single anatomic region (i.e., the lower leg). Herein, we have demonstrated that two complementary U/S measures (i.e., $CSA_{U/S}$ and $MT_{U/S}$) are each viable methods to assess muscle size in concordance to the gold standard measure of VOL_{MRI} , both before and after immobilization. This was true for both atrophy susceptible (MG) and atrophy resistant (TA) muscle, albeit with a stronger correlation in MG compared to TA.

Although not focussed on inter-muscle DMA, previous studies have shown $MT_{U/S}$ be a reliable indicator of muscle size, including both before and after hypertrophic stimuli.¹⁴ However, similar to our findings for degree of atrophy, Franchi and colleagues reported that $MT_{U/S}$ was not able to assess degree of hypertrophy in response to RET when compared to VOL_{MRI} , postulating that this is likely due to regional changes across the muscle which are not accounted for by $MT_{U/S}$ based on its assessment at a single region of the muscle only (i.e., mid-point).¹⁴ Further, although Brook *et al.* reported DMA after 4 days of ULI using $MT_{U/S}$, in this study $MT_{U/S}$ was not compared to MRI, or indeed to any other imaging method. It did however report that both $MT_{U/S}$ and DXA (lean mass) detected changes in the immobilized leg only, and that reductions in $MT_{U/S}$ correlated with declines in MPS.⁵

Considering $CSA_{U/S}$, this method has previously been reported to accurately assess reductions in the size of atrophy susceptible muscles (i.e., MG and quadriceps) in response to sustained (i.e., 70 days) head-down tilt bedrest,²⁸ albeit with less favourable data (vs. CSA_{MRI}) for MG compared to quadriceps. Conversely, although $CSA_{U/S}$ measures of MG did show good agreement with both CSA_{MRI} and VOL_{MRI} at both time-points (i.e., before and after ULI) in this study, it was not able to determine the magnitude of reductions in the size of MG (or TA) when compared to either MRI-derived measure. This discrepancy may be due to sample size (with Scott and colleagues comparing ~700 images from 27 individuals) and/or the shorter duration of immobilization employed in our study.

When exploring intra-system agreement, it is notable that only when both muscles were combined, were $MT_{U/S}$ and $CSA_{U/S}$ correlated. While this relationship is lost in individual muscles, this is likely due to statistical powering as post hoc analysis determines that correlations in MG atrophy between these two U/S methods would likely be observed with a minimum of 18 participants (α error probability = 0.05, power (1- β) = 0.8, $r = 0.28$), a frequently reported sample size in clinical (e.g., critical care cohorts) and healthy volunteer cohorts.³³ Further investigations utilizing U/S methods will help elucidate to determine the utility of ultrasound in clinical scenarios of muscle wasting.

Importantly, while both MRI and U/S methods were able to detect DMA in MG only, there is suggestion of a reduction in TA size via VOL_{MRI} only ($P = 0.08$). This is perhaps unsurprising given that geometric changes in different regions of a muscle are reflected only in VOL_{MRI} through successive

measurement of CSA_{MRI} along the entire muscle length.³⁴ Indeed, although measurement of $CSA_{U/S}$ at 30% of the muscle length (i.e., mid-belly) has been shown to provide the greatest utility to detect changes in muscle size,¹⁹ it is likely that no measure at a single spatial location will truly reflect measures across the whole muscle.³⁵ Specific to TA, it has been reported that only ~60% of variance in TA VOL may be attributable to MT.³⁶ Despite this, although no significant loss in VOL_{MRI} was observed in TA in response to 15 days ULI, previous investigations have observed significant decreases in TA VOL_{MRI} , but only following much longer periods of immobilization (i.e., 56 days bed rest in healthy individuals³²). In addition, recent meta-analysis shows dorsiflexor DMA of only -1.8% after 14 days immobilization³³ across a range of bed-rest studies each employing MRI_{VOL} . As such, if the period of immobilization has been sufficient to elicit significant DMA in the TA, it is likely that this may only have been detected by MRI_{VOL} .

The requirement for accessible methods to accurately determine muscle volume in different clinical contexts is essential to identify patients with low muscle volume, a major contributing factor in both mortality and morbidity,^{37,38} and importantly aid in determining the clinical features of, for example cancer cachexia or sarcopenia.³⁹ In addition, rates of DMA are also predictive of clinical outcomes, with these rates varying greatly between different clinical populations (e.g., ankle fracture vs. intensive care unit patients)³³ and unsurprisingly being associated with severity of illness (i.e., single vs. multi organ failure).³⁰ Effective and easy-to-access measurements of DMA may allow appropriate evaluation and intervention to preserve or reduce muscle loss in sub/clinical populations. For example, the recent development of an USI for the diagnosis of low muscle mass (i.e., sarcopenia) provides an inexpensive and clinically accessible tool centred on changes in muscle geometric proportions (i.e., MT and fibre length).²⁹ Although in this study USI did not reflect varying degrees of DMA between different muscles, the full utility of this marker will clearly develop as the USI is validated in different (e.g.) clinical and ethnic populations, and perhaps different muscles. At present it only pertains to VL, likely due to both its fundamental role in locomotion and activities of daily living (e.g., rising from a chair), and ease of measurement. As significant heterogeneity in atrophy susceptibility between muscles is now clear, and lower leg muscles have been shown essential for gait and balance and therefore falls prevention,²⁶ development of accessible tools such as the USI for application in other muscles may aid our understanding and mitigation of DMA. Further, in scenarios where panoramic image acquisition is not available (i.e., lack of appropriate software/hardware), the USI may be able to discern atrophy susceptibility through the acquisition of traditional static B-mode ultrasound images.

In conclusion, this study demonstrates the utility of $MT_{U/S}$ and $CSA_{U/S}$ to assess size across muscles with divergent atro-

phy susceptibility profiles. Both U/S-derived measures of muscle size were strongly correlated with those determined via MRI; however, neither could determine degree of DMA. Importantly U/S is already utilized within many clinical environments, as such, providing a relatively easy and quick assessment of muscle size in comparison to expensive and more restricted MRI. It is important to note that effective implementation of U/S requires consistent methodological employment (e.g., probe application) to ensure accurate assessment of muscle size. Future research should continue to investigate divergent responses between atrophy resistant/susceptible muscles and the patho/physiological importance of this paradigm, whilst also optimizing clinically accessible methods to assess muscle size in understudied muscles/muscle groups.

Acknowledgements

We are grateful for the technical support of Amanda Gates and Abigail Spicer, and to all our participants for their involvement in this study. The authors certify that they comply with the ethical guidelines for authorship and publishing in the *Journal of Cachexia, Sarcopenia and Muscle*.

References

1. Frontera WR, Ochala J. Skeletal muscle: a brief review of structure and function. *Behav Genet* 2015;**45**:183–195.
2. Stump CS, Henriksen EJ, Wei Y, Sowers JR. The metabolic syndrome: role of skeletal muscle metabolism. *Ann Med*. Taylor & Francis 2006;**38**:389–402.
3. Phillips SM. A brief review of critical processes in exercise-induced muscular hypertrophy. *Sports Med* 2014;**44**:71.
4. Moore DR, Robinson MJ, Fry JL, Tang JE, Glover EI, Wilkinson SB, et al. Ingested protein dose response of muscle and albumin protein synthesis after resistance exercise in young men. *Am J Clin Nutr* 2009;**89**:161–168.
5. Brook MS, Stokes T, Gorissen SHM, Bass JJ, McGlory C, Cegielski J, et al. Declines in muscle protein synthesis account for short-term muscle disuse atrophy in humans in the absence of increased muscle protein breakdown. *J Cachexia Sarcopenia Muscle* 2022;**13**:2005–2016.
6. Wyart E, Bindels LB, Mina E, Menga A, Stanga S, Porporato PE. Cachexia, a systemic disease beyond muscle atrophy. *Int J Mol Sci* 2020;**21**:8592.
7. Cegielski J, Bass JJ, Willott R, Gordon AL, Wilkinson DJ, Smith K, et al. Exploring the variability of sarcopenia prevalence in a research population using different disease definitions. *Aging Clin Exp Res* 2023;**35**:2271–2275.
8. Song J, Clark A, Wade CE, Wolf SE. Skeletal muscle wasting after a severe burn is a consequence of cachexia and sarcopenia. *J Parenter Enteral Nutr* 2021;**45**:1627–1633.
9. Callahan LA, Supinski GS. Sepsis-induced myopathy. *Crit Care Med* 2009;**37**:S354.
10. Franchi MV, Wilkinson DJ, Quinlan JJ, Mitchell WK, Lund JN, Williams JP, et al. Early structural remodeling and deuterium oxide-derived protein metabolic responses to eccentric and concentric loading in human skeletal muscle. *Physiol Rep* 2015;**3**:e12593.
11. Pons C, Borotikar B, Garetier M, Burdin V, Ben SD, Lempereur M, et al. Quantifying skeletal muscle volume and shape in humans using MRI: a systematic review of validity and reliability. *PLoS ONE* 2018;**13**:e0207847.
12. Engelke K, Museyko O, Wang L, Laredo JD. Quantitative analysis of skeletal muscle by computed tomography imaging—state of the art. *J Orthop Transl* 2018;**1**:91–103.
13. Maden-Wilkinson TM, Degens H, Jones DA, McPhee JS. Comparison of MRI and DXA to measure muscle size and age-related atrophy in thigh muscles. *J Musculoskelet Neuronal Interact* 2013;**13**:320–328.
14. Franchi MV, Longo S, Mallinson J, Quinlan JJ, Taylor T, Greenhaff PL, et al. Muscle thickness correlates to muscle cross-sectional area in the assessment of strength training-induced hypertrophy. *Scand J Med Sci Sports* 2018;**28**:846–853.
15. Nijholt W, Scafoglieri A, Jager-Wittenaar H, Hobbelen JSM, van der Schans CP. The reliability and validity of ultrasound to quantify muscles in older adults: a systematic review. *J Cachexia Sarcopenia Muscle* 2017;**8**:702–712.
16. Seymour JM, Ward K, Sidhu PS, Puthucherry Z, Steier J, Jolley CJ, et al. Ultrasound measurement of rectus femoris cross-sectional area and the relationship with quadriceps strength in COPD. *Thorax* 2009;**64**:418–423.
17. Mueller N, Murthy S, Tainter CR, Lee J, Riddell K, Fintelmann FJ, et al. Can sarcopenia quantified by ultrasound of the rectus femoris muscle predict adverse outcome of surgical intensive care unit patients as well as frailty? A prospective, observational cohort study. *Ann Surg* 2016;**264**:1116–1124.
18. Bass JJ, Hardy EJO, Inns TB, Wilkinson DJ, Piasecki M, Morris RH, et al. Atrophy resistant vs. atrophy susceptible skeletal muscles: “aRaS” as a novel experimental paradigm to study the mechanisms of

Funding

This work was funded through a BBSRC grant (BB/R010358/1) and through the Medical Research Council (MRC), United Kingdom (grant no. MR/P021220/1) as part of the MRC-Versus Arthritis Centre for Musculoskeletal Ageing Research awarded to the Universities of Nottingham and Birmingham. This work was also supported by the National Institute for Health Research, United Kingdom, Nottingham Biomedical Research Centre, and an MRC grant award (MR/X005240/1).

Conflicts of Interest

All authors declare that this research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Online supplementary material

Additional supporting information may be found online in the Supporting Information section at the end of the article.

- human disuse atrophy. *Front Physiol* 2021; **12**:1–11.
19. Sponbeck JK, Frandsen CR, Ridge ST, Swanson DA, Swanson DC, Johnson AW. Leg muscle cross-sectional area measured by ultrasound is highly correlated with MRI. *J Foot Ankle Res* 2021; **14**:1–7.
 20. Stokes T, Tripp TR, Murphy K, Morton RW, Oikawa SY, Lam Choi H, et al. Methodological considerations for and validation of the ultrasonographic determination of human skeletal muscle hypertrophy and atrophy. *Physiol Rep* 2021; **9**:e14683.
 21. Kositsky A, Gonçalves BAM, Stenroth L, Barrett RS, Diamond LE, Saxby DJ. Reliability and validity of ultrasonography for measurement of hamstring muscle and tendon cross-sectional area. *Ultrasound Med Biol* 2020; **46**:55–63.
 22. Morse CI, Degens H, Jones DA. The validity of estimating quadriceps volume from single MRI cross-sections in young men. *Eur J Appl Physiol* 2007; **100**:267–274.
 23. Harley YXR, Gibson ASC, Harley EH, Lambert MI, Vaughan CL, Noakes TD. Quadriceps strength and jumping efficiency in dancers. *J Dance Med Sci* 2002; **6**:87–94.
 24. Wearing J, Stokes M, De Bruin ED. Quadriceps muscle strength is a discriminant predictor of dependence in daily activities in nursing home residents. *PLoS ONE* 2019; **14**:e0223016.
 25. Simon SR, Mann RA, Hagy JL, Larsen LJ. Role of the posterior calf muscles in normal gait. *J Bone Joint Surg Am* 1978; **60**:465–472.
 26. Maritz CA, Silbernagel KG. A prospective cohort study on the effect of a balance training program, including calf muscle strengthening, in community-dwelling older adults. *J Geriatr Phys Ther* 2016; **39**:125–131.
 27. von Haehling S, Coats AJS, Anker SD. Ethical guidelines for publishing in the *Journal of Cachexia, Sarcopenia and Muscle*: update 2021. *J Cachexia Sarcopenia Muscle* 2021; **12**:2259–2261.
 28. Scott JM, Martin DS, Ploutz-Snyder R, Matz T, Caine T, Downs M, et al. Panoramic ultrasound: a novel and valid tool for monitoring change in muscle mass. *J Cachexia Sarcopenia Muscle* 2017; **8**:475–481.
 29. Narici M, McPhee J, Conte M, Franchi MV, Mitchell K, Tagliaferri S, et al. Age-related alterations in muscle architecture are a signature of sarcopenia: the ultrasound sarcopenia index. *J Cachexia Sarcopenia Muscle* 2021; **12**:973–982.
 30. Puthuchery ZA, Rawal J, McPhail M, Connolly B, Ratnayake G, Chan P, et al. Acute skeletal muscle wasting in critical illness. *JAMA - J Am Med Assoc* 2013; **310**:1591–1600.
 31. Dirks ML, Wall BT, van de Valk B, Holloway TM, Holloway GP, Chabowski A, et al. One week of bed rest leads to substantial muscle atrophy and induces whole-body insulin resistance in the absence of skeletal muscle lipid accumulation. *Diabetes* 2016; **65**:2862–2875.
 32. Belavý DL, Miokovic T, Armbrecht G, Richardson CA, Rittweger J, Felsenberg D. Differential atrophy of the lower-limb musculature during prolonged bed-rest. *Eur J Appl Physiol* 2009; **107**:489–499.
 33. Hardy EJO, Inns TB, Hatt J, Doleman B, Bass JJ, Atherton PJ, et al. The time course of disuse muscle atrophy of the lower limb in health and disease. *J Cachexia Sarcopenia Muscle* 2022; **13**:2616–2629.
 34. Fukunaga T, Roy RR, Shellock FG, Hodgson JA, Day MK, Lee PL, et al. Physiological cross-sectional area of human leg muscles based on magnetic resonance imaging. *J Orthop Res* 1992; **10**:928–934.
 35. Zabaleta-Korta A, Fernández-Peña E, Torres-Unda J, Francés M, Zubillaga A, Santos-Concejero J. Regional hypertrophy: the effect of exercises at long and short muscle lengths in recreationally trained women. *J Hum Kinet* 2023; **87**:259.
 36. Miyatani M, Kanehisa H, Ito M, Kawakami Y, Fukunaga T. The accuracy of volume estimates using ultrasound muscle thickness measurements in different muscle groups. *Eur J Appl Physiol* 2004; **91**:264–272.
 37. Mourtzakis M, Parry S, Connolly B, Puthuchery Z. Skeletal muscle ultrasound in critical care: a tool in need of translation. *Ann Am Thorac Soc* 2017; **14**:1495–1503.
 38. Wang Y, Luo D, Liu J, Song Y, Jiang B, Jiang H. Low skeletal muscle mass index and all-cause mortality risk in adults: a systematic review and meta-analysis of prospective cohort studies. *PLoS ONE* 2023; **18**:e0286745.
 39. Bozzetti F. Age-related and cancer-related sarcopenia: is there a difference? *Curr Opin Clin Nutr Metab Care* 2024. Available from: <https://pubmed.ncbi.nlm.nih.gov/38488242/>