


Please cite the Published Version

Stanbrook, Roisin , Norrey, John, Kisingo, Alex Wilbard and Jones, Martin  (2021) Dung Beetle Diversity and Community Composition Along a Land Use Gradient in a Savannah Ecosystem of North Western Tanzania. *Tropical Conservation Science*, 14. ISSN 1940-0829

DOI: <https://doi.org/10.1177/19400829211008756>

Publisher: SAGE Publications

Version: Published Version

Downloaded from: <https://e-space.mmu.ac.uk/635585/>

Usage rights:  [Creative Commons: Attribution-Noncommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/)


Additional Information: This is an open access article which first appeared in *Tropical Conservation Science*

Data Access Statement: The data that supports the findings of this study are openly available in the Figshare repository at <https://figshare.com/s/0cbea665c53f76a18086>

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

Dung Beetle Diversity and Community Composition Along a Land Use Gradient in a Savannah Ecosystem of North Western Tanzania

Tropical Conservation Science
Volume 0: 1–15
© The Author(s) 2021
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/19400829211008756
journals.sagepub.com/home/trc


Roisin Stanbrook¹ , John Norrey², Alex Wilbard Kisingo³, and Martin Jones²

Abstract

Habitat loss and degradation are the most widely cited drivers of changes in species abundance and diversity. We explored changes in dung beetle species diversity and composition across different land uses in the north west Tanzanian savannah. We expected a negative response gradient in the diversity and composition of the dung beetle community, from land uses that preserve vegetation and hold native mammal diversity to livestock intensive and heavily grazed areas. Dung beetles were sampled in a protected area and two anthropogenically influenced land use types. Species richness and composition of each land use type, including differences in diversity and functional groups were analyzed and indicator species for each land use gradient were identified. As expected, diversity and community composition varied between areas with less environmental change compared to those impacted anthropogenically. We conclude that conservation of protected areas within African savannahs can provide a functionally rich dung beetle community and subsequently rich ecological functions. The dung beetle species identified by this study as eco-indicators can be used as a benchmark for future studies that use rapid monitoring to assess disturbance in African savannas.

Keywords

dung beetle diversity, bioindicator, functional diversity, Savannah, East Africa

Information on the distribution of biodiversity is the basis of conservation planning and priority setting (Margules & Pressey, 2000; Wilson, 2000; Wu et al., 2014) and the importance of documenting arthropod species occurrences are never as warranted given the recently documented severe and ongoing declines in insect abundance and diversity (Hallmann et al., 2017; Janzen & Hallwachs, 2019; Sánchez-Bayo & Wyckhuys, 2019). Habitat loss and degradation are the most widely cited drivers of these losses, yet to date most surveys have concentrated on historically industrialised nations and not in the emergent economies of Africa and South East Asia where land use conversion from natural habitats to both subsistence and industrial agriculture is intensifying (Cotula et al., 2009).

Limited resources are available to assess the impacts of land use change on ecosystems, thus financial and

time efficient methods such as the use of ecological indicators or ‘eco-indicators’ have gained recognition in recent years (Gerlach et al., 2013; Lawes et al., 2017). Dung beetles are known to be good indicators of disturbance as they are ubiquitous, diverse, easy to sample, and ecologically important (Davis et al., 2001; Estrada

¹Department of Biology, University of Central Florida, Orlando, Florida, United States

²Faculty of Science and Engineering, Manchester Metropolitan University, Manchester, UK

³College of African Wildlife Management, Mweka, Moshi, Tanzania

Received 1 March 2021; Accepted 19 March 2021

Corresponding Author:

Roisin Stanbrook, Department of Biology, University of Central Florida, Libra Drive, Orlando, FL 32816, United States.

Email: roisin.stanbrook@ucf.edu



& Coates-Estrada, 2002; Spector, 2006). In particular, dung beetles have been proposed as effective indicators to analyse the effects of forest fragmentation (Nichols et al., 2007) and reduced impact logging (França et al., 2017) as well as indicators of tropical forest restoration success (Derhé et al., 2016) and forest type (Stanbrook, Wheeler, et al., 2021). However, little research to date has been conducted on the use of dung beetles as indicators of habitat quality in other ecosystems (Carvalho et al., 2020), notably east African grasslands, where increasing agricultural encroachment (Davis, 2002) and commercial livestock grazing (Skarpe, 1991) are driving desertification (Peters et al., 2006).

Alpha diversity and species richness are the most commonly used measurements to evaluate the effects of human disturbance on ecosystems but these metrics assume that all species exert equal influence on ecosystem functioning (Barragán et al., 2011). The relationship between these metrics and ecosystem functioning is typically positive as it greater species diversity leads to functionally richer assemblages which can provide multiple ecosystem services (Manning et al., 2016). The evaluation of functional diversity considers the effects that individual species traits have on ecosystems and the impact their removal may have on ecosystem stability. It has been proposed that anthropogenic disturbance may result in the extirpation of species with certain beneficial functional traits which may in turn affect ecosystem functioning (Díaz et al., 2013; Heilpern et al., 2018; Piccini et al., 2018).

Through dung relocation behaviour, dung beetles provide numerous ecological functions including biological pest control, soil fertilization and seed dispersal (Nichols et al., 2007) which contribute to ecosystem service provision. The execution of these functions is positively related to biomass, body size of the organisms (Nervo et al., 2014; Stanbrook, Harris, et al., 2021), and the complementarity of species and functional guilds (Slade et al., 2007). In particular Slade et al. found that tunnellers, especially nocturnal tunnellers, played a larger role in dung removal compared to rollers; tunnellers alone removed approximately 50% of dung, while rollers alone removed approximately 30% of the dung mass. Tunnelers were also responsible for removing twice the amount of dung compared to rollers in forested sites in Brazil (Batilani-Filho & Hernandez, 2017). Multiple lines of evidence indicate that local and regional scale changes in land use can severely alter patterns of dung beetle species diversity and abundance (Bogoni et al., 2016; Culot et al., 2013) and the decline of dung beetle fauna will impart long term implications on the maintenance of ecosystem integrity (Manning et al., 2016; Nichols et al., 2008).

To investigate the effects of land management intensity on dung beetle communities, we assessed the

diversity of dung beetles along a disturbance gradient: a Game Controlled Area (GCA), a pastoral area, and a community area where a mixture of pastoralism, settlements and small holder farming are the main land-uses. This study describes changes in dung beetle community composition across a land use gradient and we believe is one of the first to document the impact of anthropic pressures on East African dung beetle communities. We predicted that dung beetle richness and diversity would be higher inside the protected and pastoral areas and lowest in the community area. This study aims to describe the dung beetle diversity and functional diversity that are present within each land use type and analyse their potential ecological role. Additionally, we identified which species could be considered ecological indicators for each land use gradient.

Methods

Description of the Study Area

All three study sites were located in the Kwakuchinja wildlife corridor between Lake Manyara National Parks and Tarangire National Parks in Northern central Tanzania (Figure 1). This corridor is part of Kwakuchinja Open Area which covers about 600 km², bounded by parallels 03° 35' 38" and 03° 48' 02" S and meridians 35,048' 21" and 35° 59' 25" E (Martin et al., 2019) and forms part of the Tarangire - Manyara ecosystem (TME) which has high mammal species diversity (Pittiglio et al., 2012). The corridor was classified as 'critical' by Caro et al. (2009) who noted that the high density of human settlements and expansion of croplands were threats affecting the corridors future existence. The vegetation type is primarily microphyllous savannah dominated by *Acacia tortillis* and interspersed with *Commiphora* woodland. Rainfall is characteristically bimodal with short rains from November to January and long rains from February to May. The average rainfall is 600–650 mm *per annum* (Galanti et al., 2006)

Kwakuchinja is used by pastoralists as a transit corridor for goats and cattle between water sources and is greatly impacted by livestock overgrazing and daily fire wood collection. It is located adjacent to Mdori village (3°47'30" S, 35,056' 55" E) within an area owned by the College of African Wildlife Management. Vegetation is heterogeneous, with *Acacia* savannah occurring alongside broadleaf deciduous vegetation around recently established water holes. Human population density is estimated to be 100 – 200 people per km² (Caro et al., 2009). The study site suffers from a paucity in wild mammal species and is dominated by domestic goats and cattle.

Minjingu village is located adjacent to Tarangire National Park (3°46'19" S, 35°51'29" E). The area




Figure 1. Map of Study Locations.

suffers from overgrazing by livestock and there are many agricultural small holdings. The vegetation is dominated by *Acacia tortilis* and *Balanites aegyptiaca*, and by invasive and pioneer species *Solanum incanum* and *Alternanthera pungens*. A variety of mammals including elephant (*Loxodonta africana*), wildebeest (*Connochaetes taurinus*) and zebra (*Equus burchelli*) are present.

The Lokisale Game Controlled Area (GCA) ($3^{\circ} 46' 10''$ S, $35^{\circ} 57' 30''$ E) is a protected area covering 15 km^2 and is also adjacent to Tarangire National Park (TNP). It was gazetted 1974 and categorised as a protected area with sustainable use of natural resources including hunting (Despot Belmonte & Bieberstein, 2016) by the IUCN. Game Controlled Areas (GCA's) are type of protected area provided for in Tanzania's Wildlife Conservation Act of 1974. In accordance with the Act, human settlement and the grazing of livestock are unrestricted, and hunting of wildlife is only permitted under licence (Weldemichel, 2020). Twenty-five large mammal species have been recorded within GCA (Martin et al., 2019) including high abundance of game species including zebra (*Hippotigris quagga*), Grants gazelle (*Nanger*

granti), impala, (*Aepyceros melampus*), Hartebeest (*Alcelaphus buselaphus*) and Buffalo (*Syncerus caffer*). These mammals provide rich and varied dung resources that are expected within savannah regions and required for supporting robust and diverse dung beetle assemblages. Hanski and Cambefort (1991) found that beetle (species) size is strongly influenced by mammal composition, as well as spatial proximity, and other studies have found there is a greater proportion of small-bodied beetles in situations where large mammals are absent (Nichols et al., 2013). There is no fence between the TNP and the GCA allowing the free movement of animals between the two protected areas. Vegetation is primarily *Commiphora-Combretum* deciduous woodlands.

Trapping Methods

Sampling was carried out between 18 June - 10 August 2013. Twelve pitfall traps *per* study site were placed 50 m apart on a linear transect in accordance with (Larsen et al., 2005). Each site contained two transects of six

traps (Figure 1) and trapping was conducted for four consecutive days per transect to provide a trapping effort of eight days *per* site. Each pitfall trap comprised a 1 litre plastic tub sunk into the substrate up to its rim and filled with a saltwater and soap solution. Traps were baited with 40 g of fresh pig dung wrapped in fine gauge mesh netting which allowed dung volatiles to be emitted while excluding dung beetles. The bait was supported above each trap by a tripod of stakes and secured by string. Traps were emptied and re-baited every 24 hours to provide a total sample of 96 samples per site and total of 288 collections for study. Once collected, the dung beetles were transferred into a 70% ethanol for preservation and identification. Preserved specimens form part of a vouchered reference collection housed at the Oxford University Museum of Natural History (OUMNH).

Dung Beetle Identification

We identified the species of Scarabaeidae using taxonomic keys (D'Orbigny, 1913; Ferreira, 1969) unpublished species lists, and collections of the Oxford University Museum of Natural History Museum (OUMNH), with assistance of Darren Mann from OUMNH. Where specific identification was not possible, specimens were identified to genus and then assigned to a morphospecies. The community structure was categorised using the functional classification of African Scarabaeine dung beetle groups proposed by Doube (1990) (see Supplementary Table 1). Functional group classification *sensu* Doube (1990) provides a simple guideline for determining the structure of diverse assemblages of African dung beetles in a way that reflects a species functional role within a habitat. The classification is primarily based on the ways in which dung beetles compete for, use, and disrupt dung.

Environmental Data

Tree canopy cover above each of the 72 traps was measured using a spherical densiometer and expressed as a percentage of total leaf cover. Ground cover (%) was measured using one 1 m² quadrat placed at random within a 20 m radius of each pit fall trap. The percentage of bare ground was calculated by recording the presence or absence of bare ground at 50 cm intervals along a 10 m transect perpendicular to the pitfall trap. Data on classified landcover type was extracted from 30 m resolution landcover maps from 2010 (<https://rcmr.africa.geoportal.com>) for each study location. The distance to the nearest water body was ascertained using nearest neighbour analysis in QGIS 3.10.11 for each pitfall trap. Water body data included lake and river data for Tanzania and was obtained from worldbank.org. Additionally, digital elevation data at 30 m resolution

was extracted for each pitfall trap location and was obtained from SRTM (Shuttle Radar Topography Mission) using data available from usgs.gov.

Data Analysis

Rank abundance curves (RAC) were constructed using the R package 'BiodiversityR' (Kindt & Coe, 2005) to assess abundance and species dominance in each study site. A Shapiro-Wilks test was used to ascertain normality, all variables were found to be normally distributed. We evaluated the changes in mean species richness, diversity, and abundance of dung beetle between locations using generalized linear mixed models (GLMMs). To control for spatial autocorrelation in these models, Management was considered as a fixed factor and transect as a random factor. For species richness and abundance data, a Poisson distribution was assumed (count data) and for species diversity a gamma distribution was used. GLMMs were conducted with the nlme package (Pinheiro et al., 2017). Tukeys post-hoc pairwise comparisons were then used to describe compare differences between species richness, abundance and management using the *glht* function found in the package 'multcomp' (Hothorn et al., 2008) Alpha diversity measurements were calculated for each site using Shannon–Weiner, Simpsons and Berger-Parker indices. The Shannon–Weiner index places greater weight on the addition of new species regardless of rarity while Simpson's index is biased towards the more abundant species within a sample (Magurran, 2004). Thus, both indices were used for comparative purposes to describe the alpha diversity of each site.

The Berger-Parker dominance index was calculated using the package 'BiodiversityR' version 2.10.0. The index is calculated by dividing the number of individuals in the most abundant species by the total number of specimens present in the sample (Berger & Parker, 1970). An increase in the test statistic *d* equates to a decrease in diversity and an increase in species dominance in the sample. It is therefore a useful measure to deduce which sites contain species with a large impact on community structure (Morris et al., 2014) and the resultant diversity of the habitat. Chao1 (Chao, 1984) was used as a species richness estimator to calculate the expected species richness for each site. Richness estimations were then used to compare sampling efficiency.

Kruskal-Wallis with Dunns post-hoc tests were used to assess differences in elevation, distance to the nearest permanent water body, amount of bare ground, and ground and canopy cover for each study location. As sites nearby each other may naturally have more closely related biological communities than those further apart (Soininen et al., 2007), we checked for spatial autocorrelation by performing Moran's I tests (Moran, 1950)

using both dung beetle species richness and abundance data, at trap and transect levels allowing us to examine whether spatial auto-correlation existed in both sets of analysis.

A Non-Metric Multidimensional Scaling (NMDS) analysis using species level abundance data with a Bray-Curtis dissimilarity index was calculated to look at the community composition of dung beetle species between land uses using the R package 'vegan' (Oksanen et al., 2013). We then examined the effects of percent canopy cover, percent ground cover, amount of bare ground elevation and distance to permanent water source on community composition with PERMANOVA on Bray-Curtis distances calculated from species abundances with 999 permutations, using the *adonis* function in the *vegan* package. Because PERMANOVA and NMDS cannot readily account for spatial autocorrelation, spatial effects were not included in these analyses of cumulative patterns. A Chi squared test of association was used to look for a relationship between dung beetle functional groups and land use type. All analyses were carried out in the software programme R v3.5 (R Development Core Team, 2019).

IndVal

The IndVal method (Dufrene & Legendre, 1997) was used to determine characteristic species for each land use type. Most ecological and environmental bio indicators are identified by establishing a strong relationship with some characteristic of their environment. The IndVal method reflects frequency of occurrence (abundance) of species between habitats and is calculated by comparing a species' frequency of occurrence between habitat types (McGeoch, 2007). Species with high IndVals (>70%) thus make reliable indicator species not only because they are specific to a locality, but also because they have a high probability of being sampled in that locality during monitoring and assessment (McGeoch et al., 2002). Species were categorised as an indicator of a particular habitat if the IndVal measurement for habitat fidelity was >70%. Those species with an IndVal measurement between 20–50% were classified as generalists.

Results

Environmental Parameters

Elevation (m asl) was found to be statistically different between study locations (Kruskal–Wallis, $X^2 = 34.00$, $P = <0.05$, $df = 2$). Median elevation was found to be greater in Minjingu with a median elevation 1005 m compared to 986 m in Kwakuchinja and 998 m in Lokisale GCA. Distance to the nearest permanent

water body was also found to be different between locations ($X^2 = 63.12$, $P = <0.05$, $df = 2$) with the median distance varying from 11.76 Km in Kwakuchinja to 7.08 km in Minjingu, and 2.79 km in Lokisale. Vegetation also differed between the three study locations. Canopy cover was significantly different between sites (Kruskal–Wallis, $X^2 = 18.76$, $P = <0.05$, $df = 2$) with Minjingu having significantly less coverage compared to Kwakuchinja and Lokisale GCA (Figure 2). There was no significant difference in the amount of ground cover between study locations.

Variation in Dung Beetle Assemblages Between Land Use Type

Species Richness and Diversity. A total of 1371 individual of 45 species and morphospecies belonging to 8 tribes (*Onthophagini*, *Canthonini*, *Copriini*, *Sisyphini*, *Scarabaeini*, *Gymnopleurini*, *Oniti*, *Oniticellini*) were recorded over the study period of 24 trapping days. The number of individuals collected in the three environments differed significantly (one-way ANOVA, $F = 4.7$, $d.f = 69$, $p < 0.01$). The Kwakuchinja site had the greatest abundance of beetles with 608 individuals representing 44.4% of the total followed by; Lokisale GCA with 509 individuals (37.1%) and the Minjingu site 254 individuals (18.5%).

Species richness also differed by land use type, with Lokisale GCA the most speciose with 21 species (44.6%) followed by Kwakuchinja (17; 36.17%) and Minjingu village (12; 25.53%) (Table 1). The most abundant species overall were, *Onthophagus* sp. 14, *Onthophagus* sp. 2, *Digitonthophagus gazella*, *Sisyphus* sp 1, and *Caccobius* sp 1 with 283 (20.6%), 214 (15.6%), 133 (9.7%) and the two latter ones with 118 (8.6%) and 57 (4.15%) individuals each. These five species accounted for 63% of all individuals collected, demonstrating that all sites were dominated by a few abundant species and fit a geometric series distribution. The protected area at Lokisale GCA was the only site to reach an asymptote of species richness (Figure 3). Using Chao 1 as a population estimator it was calculated that 85.7% of the true species richness was collected across all three sites (Table 1).

Shannon-Weiner, Simpsons and Inverse Simpsons indices were calculated for each site (Table 1). The results of the Shannon-Weiner diversity index indicate that of the three sites sampled Lokisale was the most diverse ($H' = 2.579$) followed by Minjingu ($H' = 1.827$) and Kwachuchinja ($H' = 0.757$). The Simpsons Diversity index resulted in identical evenness and species richness between the Kwachuchinja and Lokisale (0.979). The results of the inverse Simpsons index indicate little difference in species dominance between Kwachuchinja and the Lokisale Game Controlled

Table 1. Species Richness and Diversity Indices for each land use type, Lokisale Game Controlled Area (GCA), Kwakuchinja (KWA) and Minjingu (MIN).

Site	Species richness (s)	Abundance (n)	Shannon ('H)	Simpson ('D)	Berger-Parker	Chao1
Kwakuchinja	17	608	1.8	0.73	0.47	17
Minjingu village	12	254	0.76	0.29	0.84	14
Lokisale GCA	21	509	2.6	0.89	0.23	21




Figure 2. Percent Cover of Bare Ground, Canopy Cover and Ground Cover Between Land Use Type, Lokisale Game Controlled Area (GCA), Kwakuchinja (KWA) and Minjingu (MIN) (Median, Interquartile Range and Whiskers and Outliers). Different letters indicate significant differences with $p < 0.05$. Landcover map generated from 30m resolution Landsat 7 data.

Area (GCA) but due to the greater dominance of *Onthophagus* sp2 at the Minjingu site, the distribution had a steeper slope (Rank abundance curves *per* site shown in Figure 3). The species abundance distribution was calculated using the Berger-Parker index (Berger & Parker, 1970). The results of the index demonstrated decreased diversity in the Minjingu site because of the greater dominance of *Onthophagus* sp2 (84.2%) when compared to that of the dominant species found in the other study sites (Table 1): *Onthophagus* 14 (20.6%) and *Sisysphus crispatus* (23.2%).

NMDS ordination of sampling units based on similarity of species abundance resulted two consistent and segregated groups as follows: (a) one formed by the degraded Kwakuchinja site and (b) two overlapping groups formed by the moderately disturbed Minjingu sites and another overlapping group formed by sites in the GCA (Figure 4). This clear segregation, particularly in terms of degraded vs. less disturbed were supported by ANOSIM tests, which detected a strong effect of landuse type ($R = 0.830$, $p < 0.001$). PERMANOVA indicated that canopy cover, amount of bare ground, study location and elevation all




Figure 3. Rank abundance and species accumulation curves for each land use type in order of land use degradation: Kwakuchinja (KWA) and Minjingu (MIN) and Lokisale Game Controlled Area (GCA)..

had significant effects on community composition (Table 2). The clustering and overlapping of the Minjingu and Lokisale sites is reflective of how strongly habitat characteristics may shape dung beetle community compositions over an acute spatial scale. Species composition in traps located in Kwakachinja sites had little compositional overlap with any of the Minjingu or Lokisale sites, and half of the traps located at the moderately disturbed

Minjingu site comprised a mixture of species compositions from both the protected Lokisale GCA study site and the domestic livestock abundant Kwakachinja site. Three trapping locations in Minjingu were ordinated outside of the main cluster. This is likely due to a lower abundance of individuals per trap with (2,7, and 5 individuals respectively) and were traps dominated by a single species of *Onthophaginae*.




Figure 4. NMDS Plot Based on Dung Beetle Community Composition Between Three Land Use Types Along a Land Use Gradient in a Savannah Ecosystem of North Western Tanzania. In order of land use degradation: Kwakuchinja (KWA), Minjingu (MIN), Lokisale Game Controlled Area. NMDS based on abundance data using Bray Curtis dissimilarity index.

Table 2. Environmental Variables Analysed by PERMANOVA using Adonis. Bolded text denotes statistical significance at $p < 0.05$.

Predictors	F (df)	R ²	P value
Canopy cover	6.87 (1,63)	0.07	0.001
Ground cover	1.82 (1,63)	0.01	0.048
Bare ground	1.97 (1,63)	0.02	0.033
Management	12.76 (2,63)	0.26	0.001
Elevation	1.76 (1,63)	0.01	0.048
Distance to H ₂ O	1.3 (1,63)	0.01	0.187

Analysis of Functional Groups

All individuals were assigned to functional groups based on their dung relocation strategy and body size. The most abundant functional group was Group V containing the small-medium paracoprid, coprophagous species (Doube, 1990). The least abundant functional group was Functional Group VII small, diurnal, coprophagous, endocoprids. In all the sites, Functional Group V (hereafter F.G V) was most dominant comprising 41.17%, 49.98% and 23.80% of the total number of individuals

respectively (Figure 5). Lokisale Game Controlled Area (GCA) was the only site that had all seven functional groups represented and also contained the greatest overall species diversity (Table 1). All individuals identified as F.G VII were trapped in Lokisale Game Controlled Area (GCA) demonstrating that smaller dweller species preferred conditions with increased vegetative cover and varied dung resources. Beetles belonging to FGII, the small telecoprids were entirely absent from the disturbed Minjingu site.

Dung Beetle Species as Indicators of Habitat Quality

The indicator value (IndVal) method (Dufrene & Legendre, 1997) combines measures of habitat fidelity and specificity was used to ascertain which species could be used as indicators of biological change within each study site. A total of 17 species trapped in all study sites could be considered 'indicator species' (McGeoch et al., 2002) (Table 3). *Gymnopleurus sericeifrons* trapped in Lokisale Game Controlled Area (GCA) had the highest %IndVal of all species collected (97%) and can be




Figure 5. Percentage of Each Functional Group Found in Each Land Use Type. Landuse degradation gradient: Kwakuchinja (KWA) -> Minjingu (MIN) -> Lokisale Game Controlled Area.

considered a species characteristic of the site. Lokisale Game Controlled Area (GCA) had a larger complement of more characteristic dung beetle species than both Kwachchinja and the Minjingu site. *Onthophagus* sp13 returned the highest IndVal for the Minjingu site (94%) but could not be considered a reliable indicator of the habitat condition of the Kwachuchinja site as it represented a <70% IndVal. Both species in the genus *Sisyphus*; a small, diurnal, telecoprid trapped in the Lokisale Game Controlled Area (GCA) had >70% indicator values signifying that this habitat is favourable to species that require increased vegetative cover.

Discussion

Abundance and Diversity

Overall we found; (i) a difference in the abundance of dung beetles between non-protected and protected sites (ii) the livestock rich kwakachinja contained the highest abundance, and the moderately disturbed site at Minjingu which contained a mixture of livestock and wildlife contained the least (iii) dung beetle diversity was higher in the protected land use site compared with the unprotected sites and, (iv) dung beetle functional composition changed concomitantly with land use

type. Our hypothesis that a reduction in the vegetative heterogeneity in the disturbed sites would also reduce the species diversity and abundance of dung beetles, causing important changes in species assemblage structure was supported. It is known that dung beetle communities display graded responses to a variety of ecological factors and anthropogenic disturbances including ivermectin use (Krüger & Scholtz, 1998), habitat modification (França et al., 2017), habitat fragmentation (Da Silva & Hernández, 2014), vegetation type (Escobar & Davis, 2010) and dung availability (Raine & Slade, 2019) and this may explain the changes in dung beetle diversity we observed.

The Kwachuchinja site possessed the highest percentage of bare ground and the lowest per cent of ground cover due to excessive grazing but also contained the greatest abundance of dung beetles (608 individuals). According to Jankielsohn et al. (2001), high concentrations of domestic livestock within a concentrated area alter vegetation and soil characteristics by overgrazing and causing soil induration. However, a consequence of the high concentration of livestock is increased dung resource availability which results in closer spatial aggregations of dung and reducing the temporal variation of food availability. The findings of the current study are consistent with those of Verdu et al. (2007), Davis et al.

Table 3. The IndVal Values of the Significant Characteristic Species for Each Land Use Type, Lokisale Game Controlled Area (GCA), Kwakuchinja (KWA) and Minjingu (MIN).

Kwakuchinja			Minjingu Village			Lokisale GCA		
Species	IndVal	P	Species	IndVal	P	Species	IndVal	P
<i>Onthophagus</i> sp14	0.87	0.001	<i>Onthophagus</i> sp13	0.94	0.001	<i>Gymnopleurus sericeifrons</i>	0.97	0.001
<i>Digitonthophagus gazella</i>	0.66	0.001	<i>Sisyphus crispatus</i>	0.79	0.001	<i>Sisyphus</i> (aff. <i>costatus</i>)	0.79	0.001
<i>Onthophagus</i> sp8	0.54	0.001	<i>Cleptocaccobius convexus</i>	0.43	0.001	<i>Sisyphus crispatus</i>	0.73	0.001
<i>Tiniocellus spinipes</i>	0.41	0.001	<i>Onthophagus</i> sp11	0.41	0.001	<i>Onthophagus</i> sp2	0.62	0.001
<i>Onthophagus</i> sp13	0.29	0.001	<i>Caccobius</i> sp1	0.41	0.001			
<i>Anachalcos convexus</i>	0.29	0.001	<i>Onthophagus</i> sp12	0.37	0.001			
<i>Onthophagus</i> sp9	0.25	0.002	<i>Khepher aegyptiorum</i>	0.31	0.002			
<i>Sisyphus</i> (aff. <i>costatus</i>)	0.23	0.031	<i>Onthophagus</i> sp10	0.29	0.002			
<i>Onthophagus vinctus</i>	0.20	0.01	<i>Onthophagus</i> sp7	0.29	0.001			
<i>Onthophagus nigriventris</i>	0.16	0.039	<i>Onthophagus</i> sp9	0.20	0.006			
			<i>Onitis alexis</i>	0.20	0.008			

(2002) and Simelane (2010) who also found that cattle grazing increased dung beetle abundance.

However, both Estrada et al. (1998) and Raine and Slade (2019) note that a large and diverse mammal fauna is also crucial for the maintenance of dung beetle diversity. Lokisale Game Controlled Area (GCA), contains the highest mammalian diversity and contained mega-fauna including elephant (*Loxodonta africana*), along with a wide variety of other herbivores such as zebra (*Equus burchelli*), and bushpig (*Potamochoerus larvatus*). This diversity may promote the co-existence of different species of dung beetles, and both generalist and specialist feeders. The data from both the Kwachuchinja and the Minjingu sites followed a geometric series distribution with both sites dominated by many individuals of a single species. The geometric series is typical of disturbed habitats and describes uneven species abundance, low diversity, and communities dominated by a few species (He & Tang, 2008).

Species dominance patterns were similar in both unprotected habitats, with two or three very dominant species in each site, although different species were dominant in the Kwachachinja compared with the Minjingu village sites. *Onthophagus* sp14 was the dominant species representing 20.6% of the total capture in disturbed, livestock dominated Kwachachinja site, *Onthophagus* 2 represented 84.3% of the total capture in the Minjingu site while *Sisyphus* aff. *costatus* was the most dominant species in the protected Lokisale site.

Jankielsohn et al. (2001) suggests that the abundance, size and ecological role of the dominant species and not just species richness are important factors to consider when assessing the level of disturbance within habitats as not all dung beetles exert the same functionality in the habitats in which they live. Both *Onthophagus* species

found in the disturbed sites are small paracoprids which are often comparatively slow at removing dung from the environment (Simmons & Ridsdill-Smith, 2011). As a result, rates of dung degradation can be reduced, and the important ecosystem services associated with dung beetle fauna can be less obviously effective in habitats dominated by smaller weaker species than those dominated by more effective dung degraders. *Sisyphus* 1, the most dominant species in the protected area is a diurnal, medium telecoprid. Sisyphini are highly effective competitors for dung and are able to remove dung quickly from pats and then bury it in short tunnels beneath the substrate (Tshikae et al., 2008).

Community Structure

Functional groups were not evenly distributed across all habitats as some groups were entirely absent from the disturbed Kwakuchinja and Minjingu study sites and others showed strong preference towards the Lokisale site. Functional group 5 (FGV), the smaller slow-burying paracoprids showed the greatest equitability across all three study sites in terms of abundance. This result is in accordance with those of Davis (1996a) who reported a similar abundance of FGV in habitats dominated by Acacia-Commiphora vegetation on sand substrate. The small telecopied species (*Sisyphus*, *Gymnopleurus*) belonging to FGII were entirely absent from the anthropogenically affected Minjingu site. Jankielsohn et al. (2001) noted that the decreases in grass cover and increases in woody shrub cover found in overgrazed habitats may impede the rolling activity of small telecoprids and decrease their ability to detect dung due to odour attenuation. This may explain the absence of the functional group from the study site.

The low percentage of abundance in FGI: the large telecoprids (*Anachalcos*, *Khepher*, *Scarabaeus*) from the Kwakachinja site is perhaps indicative of the paucity of large herbivorous mammals, other than cattle in the area, and reflects the subsequent lack of appropriate dung resources.

A large and growing body of literature has investigated the effects of declining mammal populations and the replacement of native fauna with livestock on dung beetle communities (Andresen, 1999; Culot et al., 2013; Nichols et al., 2009). Slade et al. (2007) note that the largest beetles belonging to FGI remove a disproportionate amount of dung and their decline and eventual extirpation from disturbed habitats could have profound implications for the maintenance of key ecosystem functions such as dung removal. Slade et al. (2016) describe how when this group is absent from tropical forests a 75% reduction in dung removal from surfaces is observed. The FGVII were found exclusively within the protected Lolkisale Game Controlled Area (GCA) and represented just 7% of the total of all individuals caught. Endocoprids breed within dung pats and are the least effective competitors for dung compared other functional groups (Doube, 1990) and therefore offer little towards the maintenance of ecosystem services (Jankielsohn et al., 2001).

Dung Beetles as Indicators of Ecological Disturbance

This study identified 17 species of dung beetle which can be used to ascertain the direction of ecological change with habitats and be used as an indicator of the ecological integrity of each site (Table 3). The majority of indicator species belonged to two functional groups. FGI (large telecoprids), *Scarabaeus* 1, *Scarabaeus* 2, *Scarabaeus* 3, *Kheper aegyptorium*, *Anachalcos convexus* and FGIII (large paracoprids) *Copris* 1, *Copris* 2, *Copris elphenor*. Our results are consistent with those of Jankielsohn et al. (2001) who suggested that modifications in ground cover caused by overgrazing and trampling has a greater effect on the larger bodied species in dung beetle assemblages, while smaller bodied and those species who are less competitive for dung resources not affected to the same degree. Other studies have also found large species of dung beetle have been shown to be more sensitive to ecosystem disturbance (Nichols et al., 2009) and have shown greater susceptibility to abundance decline in anthropogenically modified habitats (Gardner et al., 2007).

In the Lolkisale Game Controlled Area (GCA) three of the species defined as being indicators (*Sisyphus aff. costatus*, *Sisyphus crispatus*, *Gymnopleurus serceifrons*) belonged to FGII: small, telecoprids whose dung exploitation behaviour requires both sexes to construct and roll a portion of dung away from a dung source and

bury it in a tunnel located some distance from the dropping. This exploitation behaviour requires vegetation cover to be patchy so as not to obstruct rolling behaviour and a solid substrate texture. In his study of differences in African dung beetle communities between habitats (Davis, 1996a) proposes that FGII association is most likely to be found in areas comprising of thicketed vegetation and clay substrate due to the hardness of clay limiting breeding space under dung pats. *Digitonthophagus gazella* returned the highest indicator value for the Kwakachinja site. *Digitonthophagus gazella* is the most widespread dung beetle in tropical and subtropical cattle pastures in the world (Génier & Davis, 2017) and has a natural geographical range of approximately 15million Km² (Scholtz et al., 2009). One of the more significant findings to emerge from this study is the discovery of a single *Euoniticellus parvus* during sampling at the Lokisale study site. The species was thought to be endemic to western Africa: Cote d'Ivoire, Gambia, Sierra Leone (Ferreira, 1972) but was first discovered in eastern Africa in Kenya in 2008 (Gordon & Barbero, 2008). The species was not sampled during the previous and only known comprehensive inventory of northern Tanzanian dung beetles undertaken by in 1975 (Davis & Dewhurst, 1993).

African dung beetle community dynamics and activity is most likely determined by regional rainfall patterns and fluctuations in temperature with maximum diversity observed after rainfall and decreasing as surface conditions become warmer and drier (Davis, 1996b). This effect of seasonality is widespread across the tropics and in areas that have bimodal distribution in rainfall. Although average annual rainfall distribution in Tanzania is unimodal and is influenced by the Intertropical Convergence Zone, northern Tanzania where this study takes place typically experiences both a dry season, which spans from late June to early October and a wet season, with most of the rainfall occurring during the 'long rains' which fall between December and May (Borhara et al., 2020). Environmental monitoring is typically a balance between maintaining adequate sampling effort to detect a statistical signal, while keeping within financial and/or time constraints (Bicknell et al., 2014). While the data presented here is limited to the dry season this research clearly demonstrates the suitability and use of dung beetles as a focal taxonomic group to investigate the effects of disturbance within a restricted timeframe.

Implications for Conservation

The results presented here identify which dung beetle species may provide future effective indicators of habitat modification and provide a cost effective and efficient method of ascertaining the direction of ecological

change within African savannah habitats. A study by Engelbrecht (2010) surveyed the value of invertebrate inventories for protected areas in South Africa and found that 97% of park managers and ecologists wished to have a more inventories of invertebrate groups and described the lack of such information as an impediment to effective conservation planning. While only one of the three sites studied here form part of a legally protected area, all of the study sites are located with the Kwakuchinja wildlife corridor; an area under going rapid change with land converted from mixed savannah to agriculture increasing by 35% and woodlands decreasing by 67% in the years 2002 to 2017 (Martin et al., 2019). Our findings suggest that conservation planning for areas undergoing rapid land use change, such as the locations studied here, should concentrate on setting conservation priorities to conserve the functional integrity of taxa such as dung beetles that contribute greatly to ecosystem services provision as well as protecting biodiversity in general.

Acknowledgments

We are grateful to the College of African Wildlife Management at Mweka and the residents of the surrounding villages who allowed us to use their lands to conduct pitfalling trapping. We also thank Darren Mann of OUMNH for his assistance with species identifications, and Edwin Harris for advice on analyses on a earlier version of this manuscript.

Data Availability Statement

The data that supports the findings of this study are openly available in the Figshare repository at <https://figshare.com/s/0cbea665c53f76a18086>


Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iD

Roisin Stanbrook  <https://orcid.org/0000-0002-6753-9751>

Supplemental material

Supplementary material for this article is available online.

References

Andresen, E. (1999). Seed dispersal by monkeys and the fate of dispersed seeds in a Peruvian rain Forest. *Biotropica*, 31(1), 145. <https://doi.org/10.2307/2663968>

- Barragán, F., Moreno, C. E., Escobar, F., Halffter, G., & Navarrete, D. (2011). Negative impacts of human land use on dung beetle functional diversity. *PLoS One*, 6(3), e17976. <https://doi.org/10.1371/journal.pone.0017976>
- Batilani-Filho, M., & Hernandez, M. I. M. (2017). Decline of ecological functions performed by dung beetles in areas of Atlantic Forest and contribution of rollers and tunnellers in organic matter removal. *Environmental Entomology*, 46(4), 784–793. <https://doi.org/10.1093/ee/nvx091>
- Berger, W. H., & Parker, F. L. (1970). Diversity of planktonic foraminifera in deep-sea sediments. *Science (New York, N. Y.)*, 168(3937), 1345–1347. <https://doi.org/10.1126/science.168.3937.1345>
- Bicknell, J. E., Phelps, S. P., Davies, R. G., Mann, D. J., Struebig, M. J., & Davies, Z. G. (2014). Dung beetles as indicators for rapid impact assessments: Evaluating best practice forestry in the neotropics. *Ecological Indicators*, 43, 154–161. <https://doi.org/10.1016/j.ecolind.2014.02.030>
- Bogoni, J. A., Graipel, M. E., de Castilho, P. V., Fantacini, F. M., Kuhnen, V. V., Luiz, M. R., Maccarini, T. B., Marcon, C. B., de Souza Pimentel Teixeira, C., Tortato, M. A., Vaz-de-Mello, F. Z., & Hernández, M. I. M. (2016). Contributions of the mammal community, habitat structure, and spatial distance to dung beetle community structure. *Biodiversity and Conservation*, 25(9), 1661–1675. <https://doi.org/10.1007/s10531-016-1147-1>
- Borhara, K., Pokharel, B., Bean, B., Deng, L., & Wang, S. Y. S. (2020). On Tanzania's precipitation climatology, variability, and future projection. *Climate*, 8(2), 34–39. <https://doi.org/10.3390/cli8020034>
- Bretz, F., Westfall, P., Heiberger, R. M., Schuetzenmeister, A., & Scheibe, S. (2021). *Package 'multcomp.'*
- Caro, T., Jones, T., & Davenport, T. R. B. (2009). Realities of documenting wildlife corridors in tropical countries. *Biological Conservation*, 142(11), 2807–2811. <https://doi.org/10.1016/j.biocon.2009.06.011>
- Carvalho, R. L., Andersen, A. N., Anjos, D. V., Pacheco, R., Chagas, L., & Vasconcelos, H. L. (2020). Understanding what bioindicators are actually indicating: Linking disturbance responses to ecological traits of dung beetles and ants. *Ecological Indicators*, 108(September 2019), 105764. <https://doi.org/10.1016/j.ecolind.2019.105764>
- Chao, A. (1984). On the estimation of the number of classes in a population. *Board of the Foundation of the Scandinavian Journal of Statistics*, 11(4), 265–270. <https://doi.org/10.1214/aoms/1177729949>
- Cotula, L., Vermeulen, S., Leonard, R. and Keeley, J. (2009). Land Grab Or Development Opportunity? Agricultural Investment And International Land Deals In Africa, IIED/FAO/IFAD, London/Rome. ISBN: 978-1-84369-741-1
- Culot, L., Bovy, E., Zagury Vaz-de-Mello, F., Guevara, R., & Galetti, M. (2013). Selective defaunation affects dung beetle communities in continuous Atlantic rainforest. *Biological Conservation*, 163, 79–89. <https://doi.org/10.1016/j.biocon.2013.04.004>
- D'Orbigny, H. (1913). Synopsis des onthophagides d'Afrique [Synopsis of African onthophagids]. *Annales de La Société Entomologique de France*, 82, 1–743.

- Da Silva, P. G., & Hernández, M. I. M. (2014). Local and regional effects on community structure of dung beetles in a mainland-island scenario. *PLoS One*, 9(10), e111883. <https://doi.org/10.1371/journal.pone.0111883>
- Davis, A. J., Holloway, J. D., Huijbregts, H., Krikken, J., Kirk-Spriggs, A. H., & Sutton, S. L. (2001). Dung beetles as indicators of change in the forests of Northern Borneo. *Journal of Applied Ecology*, 38(3), 593–616. <https://doi.org/10.1046/j.1365-2664.2001.00619.x>
- Davis, A. L. V. (1996a). Community organization of dung beetles (coleoptera: Scarabaeidae): Differences in body size and functional group structure between habitats. *African Journal of Ecology*, 34(3), 258–275. <https://doi.org/10.1111/j.1365-2028.1996.tb00621.x>
- Davis, A. L. V. (1996b). Seasonal dung beetle activity and dung dispersal in selected South African habitats: Implications for pasture improvement in Australia. *Agriculture, Ecosystems and Environment*, 58(2–3), 157–169. [https://doi.org/10.1016/0167-8809\(96\)01030-4](https://doi.org/10.1016/0167-8809(96)01030-4)
- Davis, A. L. V. (2002). Dung beetle diversity in South Africa: Influential factors, conservation status, data inadequacies and survey design. *African Entomology*, 10(1), 53–65.
- Davis, A.L.V., & Dewhurst, C. F. (1993). Climatic and biogeographical associations of Kenyan and northern Tanzanian dung beetles (Coleoptera: Scarabaeidae). *African Journal of Ecology*, 31(4), 290–305. <https://doi.org/10.1111/j.1365-2028.1993.tb00543.x>
- Davis, Adrian L.V., Van Aarde, R. J., Scholtz, C. H., & Delpont, J. H. (2002). Increasing representation of localized dung beetles across achronosequence of regenerating vegetation and natural dune forest in South Africa. *Global Ecology and Biogeography*, 11(3), 191–209. <https://doi.org/10.1046/j.1466-822X.2002.00283.x>
- Derhé, M. A., Murphy, H., Monteith, G., & Menéndez, R. (2016). Measuring the success of reforestation for restoring biodiversity and ecosystem functioning. *Journal of Applied Ecology*, 53(6), 1714–1724. <https://doi.org/10.1111/1365-2664.12728>
- Despot Belmonte, K., & Bieberstein, K. (2016). *How protected areas contribute to achieving global targets for biodiversity*. UNEP-WCMC and IUCN. <https://doi.org/10.1017/S0954102007000077>
- Díaz, S., Purvis, A., Cornelissen, J. H. C., Mace, G. M., Donoghue, M. J., Ewers, R. M., Jordano, P., & Pearse, W. D. (2013). Functional traits, the phylogeny of function, and ecosystem service vulnerability. *Ecology and Evolution*, 3(9), 2958–2975. <https://doi.org/10.1002/ece3.601>
- Doube, B. M. (1990). A functional classification for analysis of the structure of dung beetle assemblages. *Ecological Entomology*, 15(4), 371–383. <https://doi.org/10.1111/j.1365-2311.1990.tb00820.x>
- Dufrene, M., & Legendre, P. (1997). Species assemblages and indicator species: The need for a flexible asymmetrical approach. *Ecological Monographs*, 67(3), 345–366.
- Engelbrecht, I. (2010). Invertebrate species inventories in protected area management: Are they useful? *African Entomology*, 18(2), 235–245. <https://doi.org/10.4001/003.018.0203>
- Escobar, F., & Davis, A. L. V. (2010). How might intensification of farming influence dung beetle diversity (coleoptera: Scarabaeidae) in Maputo special reserve (Mozambique)? *Journal of Insect Conservation*, 14(4), 389–399. <https://doi.org/10.1007/s10841-010-9270-x>
- Estrada, A., & Coates-Estrada, R. (2002). Dung beetles in continuous forest, forest fragments and in an agricultural mosaic habitat island at Los Tuxtlas, Mexico. *Biodiversity and Conservation*, 11(11), 1903–1918. <https://doi.org/10.1023/A:1020896928578>
- Estrada A., Coates-Estrada R., Dada A. & Cammarano P. (1998) Dung and carrion beetles in tropical rainforest fragments and agricultural habitats at Los Tuxtlas, Mexico. *Journal of Tropical Ecology*, 14, 577–559. <http://www.jstor.org/stable/2559949>
- Ferreira, M. C. (1969). Os escarabídeos De africa (sul do saara) [The African beetles (southern Sahara)]. *Revista de Entomologia de Mocambique*, 11, 5–1088.
- Ferreira M. C. (1972) Os escarabídeos de África (Sul do Sáara). *Revista. Ent. Moçamb* 11, 1–1088.
- França, F. M., Frazão, F. S., Korasaki, V., Louzada, J., & Barlow, J. (2017). Identifying thresholds of logging intensity on dung beetle communities to improve the sustainable management of Amazonian tropical forests. *Biological Conservation*, 216, 115–122. <https://doi.org/10.1016/j.biocon.2017.10.014>
- Galanti, V., Preatoni, D., Martinoli, A., Wauters, L. A., & Tosi, G. (2006). Space and habitat use of the African elephant in the Tarangire-Manyara ecosystem, Tanzania: Implications for conservation. *Mammalian Biology*, 71(2), 99–114. <https://doi.org/10.1016/j.mambio.2005.10.001>
- Gardner, T. A., Hernández, M. I. M., Barlow, J., & Peres, C. A. (2007). Understanding the biodiversity consequences of habitat change: The value of secondary and plantation forests for neotropical dung beetles. *Journal of Applied Ecology*, 45(3), 883–893. <https://doi.org/10.1111/j.1365-2664.2008.01454.x>
- Génier, F., & Davis, A. L. V. (2017). Digitonthophagus gazella auctorum: An unfortunate case of mistaken identity for a widely introduced species (coleoptera: Scarabaeidae: Scarabaeinae: Onthophagini). *Zootaxa*, 4221(4), zootaxa.4221.4.8–500. <https://doi.org/10.11646/zootaxa.4221.4.8>
- Gerlach, J., Samways, M., & Pryke, J. (2013). Terrestrial invertebrates as bioindicators: An overview of available taxonomic groups. *Journal of Insect Conservation*, 17(4), 831–850. <https://doi.org/10.1007/s10841-013-9565-9>
- Gordon, R. D., & Barbero, E. (2008). Dung Beetles (Coleoptera: Scarabaeidae: Scarabaeinae) of the Mpala Research Centre and Environs, Laikipia District, Kenya. *Journal of East African Natural History*, 97(2), 135–164. <https://doi.org/10.2982/0012-8317-97.2.135>
- Hallmann, C. A., Sorg, M., Jongejans, E., Siepel, H., Hofland, N., Schwan, H., Stenmans, W., Müller, A., Sumser, H., Hörren, T., Goulson, D., & De Kroon, H. (2017). More than 75 percent decline over 27 years in total flying insect biomass in protected areas. *PLoS One*, 12(10), e0185809. <https://doi.org/10.1371/journal.pone.0185809>

- Hanski, I., & Cambefort, Y. (1991). *Beetle Ecology*. Princeton University Press.
- He, F., & Tang, D. (2008). Estimating the niche preemption parameter of the geometric series. *Acta Oecologica*, 33(1), 105–107. <https://doi.org/10.1016/j.actao.2007.10.001>
- Heilpern, S. A., Weeks, B. C., & Naeem, S. (2018). Predicting ecosystem vulnerability to biodiversity loss from community composition. *Ecology*, 99(5), 1099–1107. <https://doi.org/10.1002/ecy.2219>
- Hothorn, T., Bretz, F., & Westfall, P. (2008). Simultaneous inference in general parametric models. *Biometrical Journal*, 50(3), 346–363. <https://doi.org/10.1002/bimj.200810425>
- Jankielsohn, A., Scholtz, C. H., & Louw, S. V. (2001). Effect of habitat transformation on dung beetle assemblages: A comparison between a south african nature reserve and neighboring farms. *Environmental Entomology*, 30(3), 474–483. <https://doi.org/10.1603/0046-225X-30.3.474>
- Janzen, D. H., & Hallwachs, W. (2019). Perspective: Where might be many tropical insects? *Biological Conservation*, 233, 102–108. <https://doi.org/10.1016/j.BIOCON.2019.02.030>
- Kindt R. & Coe R. (2005). *Tree diversity analysis. A manual and software for common statistical methods for ecological and biodiversity studies*. World Agroforestry Centre (ICRAF), Nairobi (Kenya). ISBN 92-9059-179-X. <http://www.worldagroforestry.org/output/tree-diversity-analysis>.
- Krüger, K., & Scholtz, C. H. (1998). Changes in the structure of dung insect communities after ivermectin usage in a grassland ecosystem. I. Impact of ivermectin under drought conditions. *Acta Oecologica*, 19(5), 425–438. [https://doi.org/10.1016/S1146-609X\(98\)80048-9](https://doi.org/10.1016/S1146-609X(98)80048-9)
- Larsen, T. H., Williams, N. M., & Kremen, C. (2005). Extinction order and altered community structure rapidly disrupt ecosystem functioning. *Ecology Letters*, 8(5), 538–547. <https://doi.org/10.1111/j.1461-0248.2005.00749.x>
- Lawes, M. J., Moore, A. M., Andersen, A. N., Preece, N. D., & Franklin, D. C. (2017). Ants as ecological indicators of rainforest restoration: Community convergence and the development of an ant Forest indicator index in the Australian wet tropics. *Ecology and Evolution*, 7(20), 8442–8455. <https://doi.org/10.1002/ece3.2992>
- Magurran, A. E. (2004). *Measuring biological diversity*. John Wiley & Sons.
- Manning, P., Slade, E. M., Beynon, S. A., & Lewis, O. T. (2016). Functionally rich dung beetle assemblages are required to provide multiple ecosystem services. *Agriculture, Ecosystems and Environment*, 218(February), 87–94. <https://doi.org/10.1016/j.agee.2015.11.007>
- Margules, C. R., & Pressey, R. L. (2000). Systematic conservation planning. *Nature*, 405(6783), 243–253. <https://doi.org/10.1038/35012251>
- Martin, E. H., Jensen, R. R., Hardin, P. J., Kisingo, A. W., Shoo, R. A., & Eustace, A. (2019). Assessing changes in Tanzania's Kwakuchinja wildlife corridor using multitemporal satellite imagery and open source tools. *Applied Geography*, 110, 102051. <https://doi.org/10.1016/j.apgeog.2019.102051>
- McGeoch, M. A. (2007). Insects and bioindication: Theory and progress. In A. J. Stewart, T. R. New, & O. T. Lewis (Eds.), *Insect conservation biology* (pp. 144–173). CABI. <https://doi.org/10.1079/9781845932541.0383>
- McGeoch, M. A., Van Rensburg, B. J., & Botes, A. (2002). The verification and application of bioindicators: A case study of dung beetles in a savanna ecosystem. *Journal of Applied Ecology*, 39(4), 661–672. <https://doi.org/10.1046/j.1365-2664.2002.00743.x>
- Moran, P. (1950). Notes on continuous stochastic phenomena. *Biometrika*, 37(1-2), 17–23. <https://doi.org/doi:10.1093/biomet/37.1-2.17>
- Morris, E. K., Caruso, T., Buscot, F., Fischer, M., Hancock, C., Maier, T. S., Meiners, T., Müller, C., Obermaier, E., Prati, D., Socher, S. A., Sonnemann, I., Wäschke, N., Wubet, T., Wurst, S., & Rillig, M. C. (2014). Choosing and using diversity indices: Insights for ecological applications from the German biodiversity exploratories. *Ecology and Evolution*, 4(18), 3514–3524. <https://doi.org/10.1002/ece3.1155>
- Nervo, B., Tocco, C., Caprio, E., Palestini, C., & Rolando, A. (2014). The Effects of Body Mass on Dung Removal Efficiency in Dung Beetles. *PLoS ONE*, 9(9), e107699. <https://doi.org/10.1371/journal.pone.0107699>
- Nichols, E., Gardner, T. A., Peres, C. A., Spector, S., & The Scarabaeinae Research Network. (2009). Co-declining mammals and dung beetles: An impending ecological cascade. *Oikos*, 118(4), 481–487. <https://doi.org/10.1111/j.1600-0706.2008.17268.x>
- Nichols, E., Larsen, T., Spector, S., Davis, A. L., Escobar, F., Favila, M., & Vulinec, K. (2007). Global dung beetle response to tropical forest modification and fragmentation: A quantitative literature review and meta-analysis. *Biological Conservation*, 137(1), 1–19. <https://doi.org/10.1016/j.biocon.2007.01.023>
- Nichols, E., Spector, S., Louzada, J., Larsen, T., Amezcuita, S., & Favila, M. E. (2008). Ecological functions and ecosystem services provided by scarabaeinae dung beetles. *Biological Conservation*, 141(6), 1461–1474. <https://doi.org/10.1016/j.biocon.2008.04.011>
- Nichols, E., Uriarte, M., Bunker, D. E., Favila, M. E., Slade, E. M., Vulinec, K., Larsen, T., Vaz-De-Mello, F. Z., Louzada, J., Naeem, S., & Spector, S. H. (2013). Trait-dependent response of dung beetle populations to tropical forest conversion at local and regional scales. *Ecology*, 94(1), 180–189. <https://doi.org/10.1890/12-0251.1>
- Oksanen, F.J., et al. (2013) *Vegan: Community Ecology Package*. R package Version 2.4-3. <https://CRAN.R-project.org/package=vegan>
- Peters, D. P. C., Bestelmeyer, B. T., Herrick, J. E., Fredrickson, E. L., Monger, H. C., & Havstad, K. M. (2006). Disentangling complex landscapes: New insights into arid and semiarid system dynamics. *BioScience*, 56(6), 491–501. [https://doi.org/10.1641/0006-3568\(2006\)56\[491:DCLNII\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2006)56[491:DCLNII]2.0.CO;2)
- Piccini, I., Nervo, B., Forshage, M., Celi, L., Palestini, C., Rolando, A., & Roslin, T. (2018). Dung beetles as drivers of ecosystem multifunctionality: Are response and effect traits interwoven? *The Science of the Total Environment*,

- 616-617, 1440–1448. <https://doi.org/10.1016/j.scitotenv.2017.10.171>
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., Heisterkamp, S., & Van Willigen, B. (2017). *Nlme: Linear and nonlinear mixed effects models* (3rd ed., pp. 1–336). R Package.
- Pittiglio, C., Skidmore, A. K., van Gils, H. A. M. J., & Prins, H. H. T. (2012). Identifying transit corridors for elephant using a long time-series. *International Journal of Applied Earth Observation and Geoinformation*, *14*(1), 61–72. <https://doi.org/10.1016/j.jag.2011.08.006>
- R Development Core Team. (2019). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. <https://doi.org/10.1007/978-3-540-74686-7>
- Raine, E. H., & Slade, E. M. (2019). Dung beetle-mammal associations: Methods, research trends and future directions. *Proceedings of the Royal Society B: Biological Sciences*, *286*(1897), 20182002. <https://doi.org/10.1098/rspb.2018.2002>
- Sánchez-Bayo, F., & Wyckhuys, K. A. G. (2019). Worldwide decline of the entomofauna: A review of its drivers. *Biological Conservation*, *232*(January), 8–27. <https://doi.org/10.1016/j.biocon.2019.01.020>
- Scholtz, C. H., Davis, A. L. V., & Kryger, U. (2009). *Evolutionary biology and conservation of dung beetles* (1st ed.). Pensoft.
- Simelane, T. S. (2010). Impacts of traditional land uses on biodiversity outside conservation areas: Effects on dung beetle communities of Vaalbos National Park. *African Journal of Ecology*, *48*(2), 490–501. <https://doi.org/10.1111/j.1365-2028.2009.01138.x>
- Simmons, L. W., & Ridsdill-Smith, T. J. (2011). *Ecology and evolution of dung beetles* (L. W. Simmons & T. J. Ridsdill-Smith, Eds., 1st ed.). Blackwell Publishing Ltd.
- Skarpe, C. (1991). Impact of grazing in savanna ecosystems. *Ambio*, *20*(8), 351–356. <https://doi.org/10.2307/4313864>
- Slade, E. M., Mann, D. J., Villanueva, J. F., & Lewis, O. T. (2007). Experimental evidence for the effects of dung beetle functional group richness and composition on ecosystem function in a tropical forest. *The Journal of Animal Ecology*, *76*(6), 1094–1104. <https://doi.org/10.1111/j.1365-2656.2007.01296.x>
- Slade, E. M., Roslin, T., Santalahti, M., & Bell, T. (2016). Disentangling the “brown world” faecal-detritus interaction web: Dung beetle effects on soil microbial properties. *Oikos*, *125*(5), 629–635. <https://doi.org/10.1111/oik.02640>
- Soininen, J., McDonald R., & Hillebrand H. (2007). The distance decay of similarity in ecological communities. *Ecography*, *30*, 3–12. <https://doi.org/10.1111/j.0906-7590.2007.04817.x>
- Spector, S. H. (2006). Scarabaeine dung beetles (coleoptera: Scarabaeidae: Scarabaeinae): An invertebrate focal taxon for biodiversity research and conservation. *The Coleopterists Bulletin*, *5*, 71–83.2.0.CO;2]
- Stanbrook, R., Wheeler, C. P., Jones, M., & Harris W.E (2021). Habitat Type and Altitude Work in Tandem to Drive the Community Structure of Dung Beetles in Afrotropical Forest. *Journal of Insect Conservation*, *25*, 159–173. <https://doi.org/10.1007/s10841-020-00289-1>
- Stanbrook, R., Harris W.E., Jones, M., & Wheeler C.P. (2021) The Effect of Dung Beetle Size on Soil Nutrient Mobilization in an Afrotropical Forest. *Insects*. *12*(2):141. <https://doi.org/10.3390/insects12020141>
- Tshikae, B. P., Davis, A. L. V., & Scholtz, C. H. (2008). *Trophic associations of a dung beetle assemblage (Scarabaeidae: S carabaeinae) in a woodland Savanna of Botswana*. *Environmental Entomology*, *37*(2), 431–441. <https://doi.org/10.1093/ee/37.2.431>
- Verdú, J. R., Moreno, C. E., Sánchez-Rojas, G., Numa, C., Galante, E., & Halffter, G. (2007). Grazing promotes dung beetle diversity in the xeric landscape of a Mexican Biosphere Reserve. *Biological Conservation*, *140*(3–4), 308–317. <https://doi.org/10.1016/j.biocon.2007.08.015>
- Weldemichel, T. G. (2020). Othering pastoralists, state violence, and the remaking of boundaries in Tanzania’s militarised wildlife conservation sector. *Antipode*, *52*(5), 1496–1518. <https://doi.org/10.1111/anti.12638>
- Wilson, E. O. (2000). A global biodiversity map. *Science (New York, N. Y.)*, *289*(5488), 2279–2279. <https://doi.org/10.1126/science.289.5488.2279>
- Wu, R., Long, Y., Malanson, G. P., Garber, P. A., Zhang, S., Li, D., Zhao, P., Wang, L., & Duo, H. (2014). Optimized spatial priorities for biodiversity conservation in China: A systematic conservation planning perspective. *PLoS One*, *9*(7), e103783. <https://doi.org/10.1371/journal.pone.0103783>