



Please cite the Published Version

Young, Kenneth J. , Leboeuf-Yde, Charlotte, Gorrell, Lindsay, Bergström, Cecilia, Evans, David W., Axén, Iben, Chance-Larsen, Kenneth, Gagey, Olivier, Georgopoulos, Vasileios, Goncalves, Guillaume, Harris, Catherine, Harsted, Steen, Kerry, Roger, Lee, Edward, McCarthy, Christopher , Nim, Casper, Nyirö, Luana, Schweinhardt, Petra and Vogel, Steven (2024) Mechanisms of manipulation: a systematic review of the literature on immediate anatomical structural or positional changes in response to manually delivered high-velocity, low-amplitude spinal manipulation. *Chiropractic & Manual Therapies*, 32 (1). 28 ISSN 2045-709X

DOI: <https://doi.org/10.1186/s12998-024-00549-w>

Publisher: BMC

Version: Published Version

Downloaded from: <https://e-space.mmu.ac.uk/635629/>

Usage rights:  [Creative Commons: Attribution 4.0](https://creativecommons.org/licenses/by/4.0/)

Additional Information: This is an open access article published in *Chiropractic & Manual Therapies*, by BMC.

Data Access Statement: The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Enquiries:


If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

REVIEW

Open Access



Mechanisms of manipulation: a systematic review of the literature on immediate anatomical structural or positional changes in response to manually delivered high-velocity, low-amplitude spinal manipulation

Kenneth J. Young^{1,20*} , Charlotte Leboeuf-Yde^{2,3}, Lindsay Gorrell⁴, Cecilia Bergström⁵, David W. Evans⁶, Iben Axén^{7,8}, Kenneth Chance-Larsen⁹, Olivier Gagey¹⁰, Vasileios Georgopoulos¹¹, Guillaume Goncalves¹, Catherine Harris^{12,13}, Steen Harsted^{14,15}, Roger Kerry¹⁶, Edward Lee¹⁷, Christopher McCarthy¹⁸, Casper Nim^{2,14,15}, Luana Nyirö⁴, Petra Schweinhardt⁴ and Steven Vogel¹⁹

Abstract

Background Spinal manipulation (SM) has been claimed to change anatomy, either in structure or position, and that these changes may be the cause of clinical improvements. The aim of this systematic review was to evaluate and synthesise the peer-reviewed literature on the current evidence of anatomical changes in response to SM.

Methods The review was registered with PROSPERO (CRD42022304971) and reporting was guided by the standards of the PRISMA Statement. We searched Medline, Embase, CINAHL, AMED, Cochrane Library all databases, PEDro, and the Index to Chiropractic Literature from inception to 11 March 2022 and updated on 06 June 2023. Search terms included manipulation, adjustment, chiropractic, osteopathy, spine and spine-related structures. We included primary research studies that compared outcomes with and without SM regardless of study design. Manipulation was defined as high-velocity, low-amplitude thrust delivered by hand to the spine or directly related joints. Included studies objectively measured a potential change in an anatomical structure or in position. We developed a novel list of methodological quality items in addition to a short, customized list of risk of bias (RoB) items. We used quality and RoB items together to determine whether an article was credible or not credible. We sought differences in outcomes between SM and control groups for randomised controlled trials and crossover studies, and between pre- and post-SM outcomes for other study designs. We reported, in narrative form, whether there was a change or not.

Results The search retrieved 19,572 articles and 20 of those were included for review. Study topics included vertebral position (n=3) facet joint space (n=5), spinal stiffness (n=3), resting muscle thickness (n=6), intervertebral disc pressure (n=1), myofascial hysteresis (n=1), and further damage to already damaged arteries (n=1). Eight articles were considered credible. The credible articles indicated that lumbar facet joint space increased and spinal stiffness decreased but that the resting muscle thickness did not change.

*Correspondence:

Kenneth J. Young

kjyoung1@uclan.ac.uk

Full list of author information is available at the end of the article



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Conclusion We found few studies on this topic. However, there are two promising areas for future study: facet joint space and spinal stiffness. A research strategy should be developed with funding for high quality research centres.

Keywords Spinal manipulation, Chiropractic, Osteopathy, Physiotherapy, Systematic review, Mechanism

Background

Spinal manipulation (SM) is an intervention that is commonly sought by people with back and neck pain. Manual therapists, chiropractors and osteopaths, in particular, commonly utilise SM as a therapeutic intervention [1–3]. SM is associated with improved clinical outcomes for certain musculoskeletal disorders [4–7]. As a result, SM is recommended in several treatment guidelines and reviews [8–11]. However, the underlying mechanism(s) of action need to be understood to determine appropriate indications for the application of SM as well as to maximize its therapeutic efficacy. That is, it is important to determine what is inside the “black box” of mechanism(s) of action of SM [12].

There are many theories and assertions on this topic [13], but there is no general consensus on the mechanism(s) of action of SM. It has been claimed that SM can change anatomy, such as repositioning vertebrae [14] or altering the thickness of muscles at rest [15]. It is proposed that these changes may be long-lasting [16]. Other claims include physiological changes, ranging from liberating Innate Intelligence [17], to modification of muscle strength [18] or reducing inflammation [19]. Historically, the nervous system has had a particular interest among chiropractors and osteopaths, as SM has been thought to affect spinal nerves [20, 21], the autonomic nervous system [22], and even the brain [23].

These anatomical and/or physiological changes are then purported to explain any associated clinical improvements, such as increased function, reduced pain, relief from specific diseases, and better health in general [24]. If any of these proposed mechanisms can be supported by evidence, manual therapists will be able to offer to patients a coherent rationale for applying SM.

Any mechanism of manipulation is comprised of two aspects. First, the manipulation must have an effect in the body lasting beyond the application of SM, and this effect must lead to a change in clinical outcome. Both aspects must be investigated in turn to determine potential mechanisms of SM. Although it is possible that there is a cumulative effect from SM or that a minimum dosage is needed to create an effect, this has not been well documented. Therefore, to proceed in a stepwise fashion, it seems reasonable to first determine what the immediate effects may be of one single SM.

There is evidence on what happens within the spine, as a response to various forces applied *during* a

high-velocity low-amplitude (HVLA) SM [25–28], such as the distribution of forces within tissues receiving the manipulation [29], and the amount and direction of displacement of vertebrae during SM [30], but a clear picture of what happens directly afterwards appears to be lacking.

Although there have been systematic reviews on some physiological effects of SM [31–33], to our knowledge, there are no systematic reviews that have attempted to synthesise evidence of the underlying anatomical mechanisms of SM. Therefore, we assessed the state of evidence of a measurable change anatomical structures that occurs following the application of SM.

The overall aim of this systematic review was to evaluate and synthesise the peer-reviewed literature on the immediate changes in or to anatomical structures in response to SM.

Our research objectives were as follows:

1. Identify, evaluate the quality of, and narratively synthesise the evidence that has been published in peer-reviewed research literature regarding immediate anatomical change after a spinal manipulation.
2. Identify gaps in understanding the anatomical effects of spinal manipulation and provide recommendations for future research.

Methods

Advisory board

A research project advisory board was convened for support and guidance, consisting of a chair (chiropractor KJY), an information specialist (CH), 2 chiropractors (CLY, IA), a physiotherapist (RK), an osteopath (SV), a medical doctor/chiropractor (PS), and an anatomist/orthopaedic surgeon (OG). Several had experience with systematic reviews.

Team and roles

In all, 14 people (6 chiropractors, 6 physiotherapists, and 2 osteopaths) were recruited to perform the screening of articles. Several team members practice clinically. One reviewer dropped out before screening was completed and was replaced by KJY. Another reviewer dropped out after the screening process and was replaced by LG. One researcher with a chiropractic background, experienced in systematic reviews (CLY), acted as referee and

supervisor only. The screening of articles was divided between 7 teams of 2 people each.

Protocol registration and reporting

The review was registered with PROSPERO (CRD42022304971) and the reporting was guided by the standards of the Preferred Reporting Items for Systematic Review and Meta-Analysis (PRISMA) Statement [34].

Search strategy

We performed a broad search to capture as many relevant articles as possible and developed our search strategies with an experienced information specialist (CH). The search strategy included relevant subject headings and search terms relating to manipulation and the spine and was adapted for use in each database. We had no resources for a translation service, so we limited the search to the English language. We searched the following databases: Medline (Ovid), Embase (Ovid), CINAHL (EBSCOhost), AMED (EBSCOhost), Cochrane Library all databases (via Wiley), PEDro (<https://pedro.org.au/>), and the Index to Chiropractic Literature (<https://www.chiroindex.org/>). All databases were searched from inception to 11 March 2022; the searches were updated on 06 June 2023. The full search strategy can be found in Additional File 1.

In cases where full-text articles were not available through library services, we emailed the first author, if an email address was published in the article. If there was no email address listed, or if we received no response to an email query, we searched for the first author on ResearchGate and, if found, we sent a full-text article request. If there was no response or the author could not provide us with an article, it was excluded.

Terminology

We used the term “outcome variable” to represent what the researchers measured in each study, referring to the mechanism of manipulation being studied. For instance, if SM is hypothesised to improve a clinical outcome by restoring the position of a vertebra, the mechanism by which SM achieves this end outcome is by changing the position of a vertebra. The outcome variable measured in the experiment would, therefore, be the difference in vertebral position from pre- to post-manipulation. In relation to research findings, we used the term “positive” not as a value judgment, but rather as shorthand to denote when a post-SM change in measurement was reported, and the term “negative”, when no such change was reported.

Eligibility criteria

We included only peer-reviewed articles if they fulfilled certain criteria:

1. We included primary research studies that compared non-treated with treated anatomical structures, regardless of the study design. The articles had to define SM as an HVLA thrust delivered by hand to the spine or directly related joints (i.e., including the sacroiliac or costo-vertebral joints). The measurement of effect must have occurred after a single manipulation session, that is, not after a course of care. If articles did not state a specific time interval between SM and post-SM measurement, but their research designs, or the way the text was written gave the distinct impression that there was little delay between the manipulation and the post-SM measurement, they were included.
2. The SM could not have been combined with any other therapeutic interventions.
3. Studies must have objectively measured a potential change in anatomical structure (the physical attributes of one or more structures in the human body) or a change in position (the relationship of two or more structures to each other). Studies measuring range of motion were considered subjective and were not included, since participants or assessors could consciously or subconsciously influence the position during the measurement.
4. Anatomical change was considered to be distinct from change in physiological state. Therefore, we included articles that reported on resting muscle thickness, as opposed to contracted muscle thickness, because we considered muscle contraction to be a matter better considered under physiological effects of SM. It is possible that a change in resting muscle thickness may be due to a physiological process such as contraction/relaxation. However, there may also be a purely physical mechanism such as stretching. Therefore, we included it.
5. Animal studies were included, because objectively measured anatomical effects of SM are not subject to contextual effects as clinical outcomes may sometimes be.

If studies measured more than one outcome, only the relevant outcome(s) were considered for this review. For the full list of exclusion criteria, please see Additional File 2.

Article selection

All articles retrieved through the literature searches were exported into EndNote X9.3.1 (Clarivate, Philadelphia, 2013). After duplicates were removed, the remaining articles were imported into the web-based Rayyan systematic review management application [35] for reference management and tracking of the screening process. The total number of articles was divided into 7 separate reviews on Rayyan, each given to a pair of reviewers to independently screen titles/abstracts. Full-text versions of the potentially included articles were obtained for screening by the same teams. Detailed written instructions were distributed to the reviewers prior to the screening processes and meetings were held with each pair prior to title/abstract screening to facilitate congruence in approach. In cases of disagreement between reviewers during phase one (title/abstract) or phase two (full text) screening, a third independent reviewer (KJY or LG) was consulted to achieve consensus. Finally, KJY and CB conducted a backward search, manually searching the reference lists from all articles included at the full-text screening stage for any missing articles.

Data extraction

The articles were grouped by topic, and reviewers self-assigned, as much as possible, to 1 or 2 topics, with 2 reviewers independently reviewing each topic. The reviewers were LN, DE, KJY (2 topics), RK, CB, CM, SH, VG, LG, EL GG, KCL, and CN. Each topic pair was overseen by a third reviewer, either KJY, LG or DE, who had knowledge of the topic and was designated as “leader” of the group. Calibration sessions were held by KJY with each team prior to data extraction to help ensure congruency of approach. Study descriptions, methodological quality, and risk of bias (RoB) data were extracted. The 7 pairs of reviewers independently extracted data, with conflicts resolved by discussion between them or with the leader. Findings for each team were reviewed by the leaders of each team. KJY, CLY, and DE reviewed all findings.

Extracted data

Article descriptions

Descriptive information about each article was entered into a table. This included first author/year of publication, mechanism of spinal manipulation investigated, study design, study setting, study cohort, sample sizes of intervention and control groups, control group description, spinal region studied, outcome variable

used, instrument used for measurement, and the time interval between SM and measurement.

Quality assessment and risk of bias

Because the articles we included had used objective measurements of anatomical/positional outcome variables, there were potential areas of technical error introduced during the experiments. For this reason, and informed by a previous publication [22], we considered the techniques used to study the various outcome variables, and developed a novel list of methodological quality items. These items related to technical aspects of the experiments and transparency in methods.

A standard RoB reporting tool was not applicable due to the heterogeneity of study designs. Further, RoB tools are suitable mainly for clinical studies, in which the influence of the study participant is important to account for. However, in the investigations included in our review, study participant influence would be absent, as they would not likely be able to influence technical readings relating to anatomical structures, either consciously or subconsciously. Therefore, study participants did not need to be blinded to treatment or control group allocation. They also did not need to be naïve to the treatment. Thus, we included RoB items only relating to the blinding of assessors and statisticians. We selected only RoB items that we considered appropriate for the relevant study designs, i.e., depending on if they had a control group or not. We also included random allocation when two interventions were compared because it was important that inherent differences between groups was eliminated. We then considered those quality and RoB items together to determine, whether an article was credible or not credible, in a process described below.

Results

Results of each study were extracted, after the quality and RoB items were determined, to avoid reviewer bias of the quality/RoB assessment. The results of each of the studies were extracted from the articles by 3 members of the team. CM reviewed half the articles and SV reviewed the other half, each working in conjunction with KJY, who reviewed all articles. Conflicts and queries were resolved by discussion or consultation with CLY. Results were entered into separate tables for each outcome variable, including the ultimate finding on whether the anatomical structure was affected by the manipulation or not. All tables were consolidated and edited for readability, and each team reviewed and approved their consolidated tables.

Data synthesis

Assessment and weighting of quality and risk of bias

Through consensus discussion, we defined the quality and RoB items by the consequences we assumed that they would have on the credibility of the data. For this purpose, we developed a dichotomous system of weighting. Items were determined to be either “critical” or “important”. “Critical” items were essential to the credibility of the results, whereas “important” items were those that were considered good practice but were not essential, in and of themselves, to a judgement on whether results could be considered credible.

Quality items

We considered 3 quality items to be “critical”. First was “evaluation tool(s) appropriate to measure outcome variable(s)”. The second critical item was “reported the reliability of outcome variable(s)”. The final critical quality item was “measurement tool calibrated” (if appropriate). The remaining quality items on our checklist were assessed as important, but not critical.

Risk of bias items

For non-RCT studies, we included only one RoB item, “assessor blinding to pre-post manipulation status”, and we considered it to be critical. For RCTs and crossover study designs, 2 RoB items were considered critical. First was “random allocation of participants”. The second was “assessor blinding to intervention group”. The remaining RoB item was “statistician blind to intervention/control group”, which is not commonly reported in articles. For this reason, we decided to classify it as important rather than critical.

Assessment of the certainty of evidence

We used the quality and RoB tables to establish “credibility” for each article and outcome variable. If a quality or RoB item was appropriately reported, it was left as white in the table. However, items that had not been reported or were poorly reported, were marked as yellow for important items and red for critical items.

We then made an overall assessment of credibility for each article, based on both the quality and RoB, after which each article was defined as “credible” or “not credible”. Articles were defined as credible if they had 0 red and 0–2 yellow items. Articles were defined as not credible if they had 1 or more red items or 3 or more yellow. Please see Table 1 for the key to interpreting the quality and RoB items as well as explanations of each.

Reporting of results

It was not suitable to pool the results for meta-analysis due to heterogeneity in outcome variables, study design, and participant characteristics. It was also not relevant, because our research question did not require a measurement (e.g., amount of facet joint space increase or cross-sectional area increase in muscle size) but rather just the presence or absence of change post-SM. These results were reported both in detail and summarized in tables as “positive” (i.e., there was a change post-SM) or “negative” (i.e., there was no change).

To excerpt maximum information from this research area, we reported results from all studies, but separately for the “credible” and “not credible” articles. We also included a summary of methodological issues in the Discussion to aid future researchers to improve this research area. Our rules on reporting data were as follows:

- (i) We reported differences between SM and control groups for RCTs and crossover studies and between pre- and post-SM groups for other study designs. When there were several results reported in one article, we used the “best” estimates (i.e., best case scenario of a “positive” outcome or difference pre-post SM). For example, one article reported a positive result when the participant was re-measured while remaining in side-posture position, but there was a “negative” result when the participant was returned to neutral (supine) position for post-SM measurement. In this case, we reported the positive result.
- (ii) If there were no statistically significant differences or statistical significance tests were not reported, we provided, again, the “best” estimate (i.e., selecting the best-case scenario).
- (iii) If no estimates (direct measurements) were reported, but rather only significance values, then we reported those.

Writing and editing the manuscript

To keep the workload achievable and to improve attention to detail in manuscript development, we used an iterative process. Each section of the paper (introduction, methods, results, and discussion) and all tables and figures were disseminated to the research team for comments at different stages. In cases of disagreement, an appropriate member or members of the advisory board were contacted, and discussions were held amongst team members until consensus was reached.

Table 1 Key to interpreting the quality/risk of bias tables in a systematic review of anatomical mechanisms of spinal manipulation

Items included in the methodological quality assessment	Questions asked	Possible answers	
		Yes or N/A (white)	No Important item (yellow) Critical item (red)
Description adequate to reproduce experiment set-up	Did the report provide sufficient information on the investigation, including study setting, population and sample description, human and equipment infrastructure, to allow overall reproduction of study?	Yes	No
Evaluation tool(s) appropriate to measure outcome variable(s)	Was the measurement tool able to measure what the authors stated it measured? Use of a tool capable of accurately measuring the intended outcome is essential to being able to trust the results reported in an article.	Yes	No
Reported reliability of outcome variable(s) reported	Did the authors report that they tested the investigators for intra- and/or inter-examiner reliability and/or the test/measuring tools for reproducibility? Alternatively, at a minimum, was a reliability study in the peer-reviewed literature cited? Consistently accurate interpretation of measurements was considered necessary to the credibility of the results. Note that this is not an assessment of validity on our part, on which we made no judgment. We trusted the authors.	Yes	No
Description of SM adequate to be reproduced by a registered practitioner	Was the spinal manipulation explained either as a written description and/or shown through pictures/diagrams?	Yes	No
Measurement tool calibrated (N/A was included under 'Yes')	Was the measurement tool reported to have been calibrated to ensure accuracy? An uncalibrated tool, or failing to report calibration, meant that results could not be trusted.	Y or N/A	No
Adequately explained how outcome measurements were taken	Was the measurement method described in sufficient detail to understand/reproduce, particularly as regards before-after measurements to ensure they were taken under similar circumstances?	Yes	No
Clinician competence reported	Was the experience/qualification of the person delivering the spinal manipulation reported? Chiropractor/osteopath/physiotherapist was accepted. Physician/doctor/clinician/therapist was not.	Yes	No
Assessor competence reported	Was the experience of the person assessing the outcome variable reported? This could involve previous experience or specific training for the study. N/A is an option because an automated process would have no interpretation to require competence of a human assessor.	Y or N/A	No
Statistics transparent	Was the statistical approach at least relatively transparent, i.e., did it appear to be justified (evident or with a rationale) and reported in detail as to how it related to the results. In studies with small participant numbers, was the analysis of findings well explained?	Y or N/A	No
Items included in the risk of bias assessment			
Non-RCT studies only: Assessor blind to before-after readings (N/A was included under 'Yes')	Was it reported that the observer who recorded/evaluated the findings did so without knowing whether they were pre- or post-intervention? If an assessor knew whether the participant had been manipulated or not, it seemed quite possible that consciously or not, the post SM measurement could have been affected by that knowledge.	Y or N/A	No
RCT/crossover studies only: Random allocation	Was the randomisation process adequate to spread the risks of clustering of characteristics of the sample equally between both groups? We considered it critical that participants be randomly assigned to groups to ensure that the characteristics of the intervention and control groups did not introduce bias.	Yes	No
RCT/crossover studies only: Assessor blind to intervention group	Was it reported that the observer who recorded/evaluated the findings did so without knowing whether they were from the intervention or control group? If an assessor knew the intervention/control status of a participant, the measurement could potentially be affected by that knowledge.	Yes	No
RCT/crossover studies only: Statistician blind to control/intervention findings	Was it reported that the statistician was blind to whether readings were obtained with or without spinal manipulation in RCTs or crossover studies?	Yes	No

Table 1 (continued)

N/A: not applicable

RCT: randomised controlled trial




Fig. 1 PRISMA 2020 flow diagram for new systematic reviews which included searches of databases, registers and other sources. * Records were excluded by humans; no automation tools were used. ** Eleven articles were not available for full-text retrieval after having unsuccessfully attempted to contact the first author of each in two ways, as detailed in the Methods section. *** Three articles were retrieved which had their titles/abstracts in English but the manuscripts in a foreign language and thus were excluded at the full-text screening stage

Results

General

Of the original 19,572 articles (37,902 including duplicates), 20 articles that reported on 20 studies were ultimately included in this review (Fig. 1).

Study descriptions

As shown in Table 2, the reviewed articles included the following outcome variables: vertebral position (n=3) [14, 36, 37], facet joint space (n=5) [38–42], spinal stiffness (n=3) [43–45], resting muscle thickness (n=6) [15, 46–50], intervertebral disc pressure (n=1) [51], myofascial hysteresis (n=1) [52], and further damage to damaged arteries (n=1) [53].

Study designs included both controlled (n=10) and uncontrolled (n=10) studies. The number of study participants ranged from 1 to 250. The most commonly studied participants were healthy adults (n=9). Two studies included animals.

Most studies (n=12) took place in the United States of America, with a private chiropractic or osteopathic school as the most common setting (n=6), while 3 studies explicitly stated that they used a lab setting. SM was most commonly performed on the lumbar spine or lumbopelvic area (n=12).

Methodological quality

As can be seen in Tables 3, 4, 5, 6, and 7, issues relating to the methodological quality varied. Regarding the two critical quality items, the first, “using an appropriate evaluation tool” was consistently lacking in the 3 articles on vertebral position, and the second, “reporting of the reliability of an outcome variable” was not present in 6 of the 11 articles.

Risk of bias

We did not find that critical RoB was a problem in studies using an RCT or crossover design (Tables 3, 4, 5, 6,

Table 2 Descriptions of 20 studies included in a systematic review of anatomical mechanisms of spinal manipulation

First author year of publication	Mechanism(s) of spinal manipulation investigated	Study design	Study setting	Study cohort	Sample size: - N cases - N controls (or control interventions)	Control group description	Spinal region(s)	Outcome variable(s)	Instrument(s) used for measurement	Time interval between spinal manipulation and measurement
Jirout 1972	Change in vertebral position	Uncontrolled intervention	-	Humans with palpatory finding of a "blockade" on lateral flexion of neck	250: - 250 - 0	N/A	Cervical	Vertebral segment position change	Plain radiography	-
Cramer 2000	Increase in facet joint space	RCT	Chiropractic school, USA	Healthy adults, 22-29 yrs	16: - 8 (2 groups of 4) - 8 (2 groups of 4)	(i) Neutral position (ii) Side posture position	Lumbar	Facet joint space change	MRI	"Immediately"
Cramer 2002	Increase in facet joint space	RCT	Chiropractic school, USA	Healthy adults, 22-30 yrs	64: - 32 (2 groups of 16) - 32 (2 groups of 16)	(i) Neutral position (ii) Side posture position	Lumbar	Facet joint space change	MRI	"Immediately"
Cascioli 2003	(i) Increase in facet joint space (ii) Formation of intra-articular gas bubbles	Uncontrolled intervention	Medical teaching hospital, South Africa	Chiropractic students, 23-24 yrs, without cervical complaints	2: - 2 - 0	N/A	Cervical	(i) Facet joint space change (ii) Intra-articular gas bubble formation	Plain radiography	"Immediately"
Lisi 2006	Change in intervertebral disc pressure	Feasibility	-	Healthy adults, 41-42 yrs	1: - 1 - 0	N/A	Lumbar	Intervertebral disc pressure	Pressure sensor probe	Immediately and for 15 s after spinal manipulation
Brenner 2007	Change in multifidus muscle thickness	Case report	Army facility, USA	Male, 33 yrs, chronic LBP	1: - 1 - 0	N/A	Lumbo-pelvic	Thickness of multifidus at rest	Diagnostic ultrasound	"Immediately"
Hausler 2007	Change in spinal stiffness	Crossover	Equine research park at a university, USA	Healthy adult horses with 2 pins in spinous processes to simulate acute back pain	10: - 10 - 10	7-day washout before crossover, Control was no intervention	Thoracic Lumbar	Spinal stiffness	Cable extensometer Pressure sensor mat	-
Raney 2007	Change in lateral abdominal muscle thickness	Prospective case series	Army facility, USA	Adults, 18-53 yrs, acute/sub-acute LBP, SMT responders	9: - 9 - 0	N/A	Lumbo-pelvic	Thickness of lateral abdominal muscles at rest	Diagnostic ultrasound	"Immediately"

Table 2 (continued)

First author year of publication	Mechanism(s) of spinal manipulation investigated	Study design	Study setting	Study cohort	Sample size: - N cases - N controls (or control interventions)	Control group description	Spinal region(s)	Outcome variable(s)	Instrument(s) used for measurement	Time interval between spinal manipulation and measurement
Wynd 2008	Further damage to already damaged arteries	Uncontrolled intervention	University lab, Canada	Dogs	10: -0 -0	N/A	Cervical	Increase in dimensions of vascular lesion	Intravenous diagnostic ultrasound Fluoroscopy	-
Palmer 2009	Change in vertebral position	Retrospective case series	Private chiropractic clinic, USA	Chiropractic patients, 18-65 yrs, non-migraine headache	35: -35 -0	N/A	Cervical	Vertebral segment position change ("atlas laterality")	Plain radiography	"Immediately"
Cramer 2011	Increase in lumbar facet joint space	Feasibility	Chiropractic school, USA	Healthy adults, 25-27 yrs	5: -5 -0	None, but cavitated joints were compared with non-cavitated	Lumbar	Facet joint space change	MRI for gap-ping; Accelerometer for cavitation	"Immediately"
Fritz 2011	Change in spinal stiffness	Prospective case series	-	Physiotherapy patients, 19-60 yrs, LBP with or without leg symptoms	51: -50 -0	N/A	Lumbo-pelvic	Spinal stiffness	Mechanical indentation device	"Immediate"
Konitzer 2011	Change in thickness of transverse and inter-oblisque abdominal muscles	Prospective case series	Army facility, USA	11F, 8M, 21-46 yrs, chronic LBP who met a clinical prediction rule for lumbar stabilization	19: -19 -0	N/A	Lumbo-pelvic	Thickness of transverse and inter-oblisque abdominal muscles at rest	Diagnostic ultrasound	"Immediately"
Puentedura 2011	Change in thickness of transverse abdominal muscles	Crossover, randomised	University, USA	19F, 16M, 21-34 yrs, asymptomatic university staff and students	35: -35 -35	19F, 16M, 21-34 yrs, asymptomatic university staff and students	Lumbar	Thickness of transverse abdominal muscle at rest	Diagnostic ultrasound	"Immediately"
Cramer 2012	Increase in facet joint space	RCT	Chiropractic school, USA	Healthy adults, 18-30 yrs	40: -30 -10	No SMT; Also cavitated joints were compared with non-cavitated	Lumbar	Facet joint space change	MRI for measuring joint space; accelerometer for cavitation	"Immediately"
Barnes 2013	Change in myofascial hysteresis	RCT	Private osteopathic college lab, USA	NR	80: -40 -40	NR	Cervical	Change in tissue texture characteristics	Durometer	"Approx 10 min"

Table 2 (continued)

First author year of publication	Mechanism(s) of spinal manipulation investigated	Study design	Study setting	Study cohort	Sample size: - N cases - N controls (or control interventions)	Control group description	Spinal region(s)	Outcome variable(s)	Instrument(s) used for measurement	Time interval between spinal manipulation and measurement
Wong 2015	Change in spinal stiffness	Non-randomised controlled study	-	18-60 yrs, LBP, SMT responders	107: - 32 (15 previously identified SMT responders, 17 non-responders) - 75	(i) 59 asymptomatic (to session 1); 57 for sessions 2 and 3 who did not receive SMT - 16 w/ LBP that did not receive SMT	Lumbo-pelvic	Spinal stiffness	Mechanical indentation device	"Immediately"
Haavik 2016	Change in pelvic floor muscle thickness	Crossover, unrandomised	Chiropractic school, New Zealand	Healthy pregnant, 18-38 yrs	26: (i) 11 pregnant (ii) 15 non-pregnant (i) 15 non-pregnant (ii) 11 pregnant	Non-pregnant chiro-practic students, 19-32 yrs	Unspecified spine and/or pelvis	Thickness of levator hiatus muscles at rest	Transperineal diagnostic ultrasound	(i) Same day for pregnant participants (ii) Not reported for nonpregnant
Flaum 2017	Change in vertebral position	RCT	Medical school, USA	Asymptomatic medical students, 22-40 yrs	51: - 25 - 26	Asymptomatic medical students, 22-40 yrs	Cervical	Tissue depth to articular pillar compared bilaterally (proxy for vertebral rotation)	Diagnostic ultrasound	"Immediately"
Fosberg 2019	Change in transverse abdominal muscle thickness	RCT	University and private clinic, USA	18-70 yrs, LBP, SMT responders	67: - 33 - 34	18-70 yrs, LBP, SMT responders	Lumbar	Thickness of transverse abdominal muscles at rest	Diagnostic ultrasound	"Immediately"

LBP: Low back pain, MRI: Magnetic resonance imaging, N/A: Not applicable, RCT: Randomised controlled trial, SMT: Spinal manipulative therapy

Table 3 Quality, risk of bias, and credibility assessments for 3 articles on vertebral position in a systematic review of anatomical mechanisms of spinal manipulation

Study description	Jirout 1972	Palmer 2009	Flaum 2017
Proposed mechanism of spinal manipulation	Vertebral position	Vertebral position	Vertebral position
Study Design	Uncontrolled trial	Case series	RCT
Quality			
Description adequate to reproduce experiment set-up	No	No	No
Evaluation tool(s) appropriate to measure outcome variable(s)*	No	No	No
Reliability of outcome variable(s) reported*	No	Yes	No
Description of SM adequate to be reproduced by a registered practitioner	No	No	Yes
Measurement tool calibrated* (N/A was included under 'Yes')	N/A	N/A	N/A
Adequately explained how measurements were taken	No	No	Yes
Clinician competence reported	No	Yes	Yes
Assessor competence reported	No	Yes	Yes
Statistics transparent	Yes	No	No
Risk of bias			
Non-RCT/crossover studies only: Assessor blind to before-after readings* (N/A was included under 'Yes')	No	No	N/A
RCT/crossover studies only: Random allocation*	N/A	N/A	Yes
RCT/crossover studies only: Assessor blind to intervention group*	N/A	N/A	Yes
RCT/crossover studies only: Statistician blind to control/ intervention readings	N/A	N/A	No
Results	3 red 5 yellow	2 red 4 yellow	2 red 3 yellow
Unacceptable quality and risk of bias items			
Overall assessment (Not credible: 1 or more red items and/or 3 or more yellow items)	Not credible	Not credible	Not credible

* Critical item (no asterisk: important item)

White: requirement fulfilled (Yes) or not applicable (N/A)

Red: Critical quality or risk of bias item not fulfilled (No)/Not credible article

Yellow: Important quality or risk of bias item not fulfilled (No)

and 7), as 7 of 9 such articles had no critical deficiencies in this domain. However, in other study designs, 7 of 11 articles did not report that outcome measurements had been taken by assessors who were blinded to the previous assessment.

Credibility

The RoB/quality tables (Tables 3, 4, 5, 6 and 7) show that 8 of the 20 studies were considered credible, whereas the other 12 were considered not credible. Notably, two of the studies, reported by Konitzer [46] and Fritz [43], met all our quality and RoB criteria.

The numbers of articles reporting on credible studies were as follows: facet joint space (n = 4/5), spinal stiffness (n = 1/3), and resting muscle thickness (n = 3/6). Therefore, 1 of the 5 studies on facet joint space was not credible; 2 of the 3 studies on spinal stiffness were not credible and 3 of the 6 studies on resting muscle thickness were not credible. In addition, all the articles reporting on studies on vertebral position (n = 3), intervertebral disc (IVD) pressure (n = 1), further damage to damaged arteries (n = 1), and myofascial hysteresis (n = 1) were found to be not credible.

Results of credible studies by outcome variable

Results from the 8 credible studies are shown below, reported by outcome variable.

Facet joint space (n = 4/5) Four of 5 studies on changes to facet joint space were considered credible. They all reported an increase in lumbar spine facet joint space post-side-posture manipulation for the “up” side facet joints, but only if the participant was re-scanned using magnetic resonance imaging (MRI) while maintaining side posture position. When returned to neutral position the increased joint space disappeared.

Spinal stiffness (n = 1/3) Only 1 of the 3 studies on spinal stiffness was found credible and reported immediate reduced spinal stiffness post-SM.

Resting muscle thickness (n = 3/6) Three of six studies on changes to resting muscle thickness were considered credible. These studies, using diagnostic ultrasound, reported no statistically significant differences in either the transverse or internal oblique abdominal muscles post-manipulation.

Table 4 Quality, risk of bias, and credibility assessments for 5 articles on facet joint space in a systematic review of anatomical mechanisms of spinal manipulation

Study description	Cramer 2000	Cramer 2002	Cascioli 2003	Cramer 2011	Cramer 2012
Proposed mechanism of spinal manipulation	Increase in facet joint space	Increase in facet joint space	Increase in facet joint space and formation of intra-articular gas bubbles	Increase in lumbar facet joint space	Increase in facet joint space
Study design	RCT	RCT	Uncontrolled intervention	Feasibility	RCT
Quality					
Description adequate to reproduce experiment set-up	Yes	Yes	Yes	Yes	Yes
Evaluation tool(s) appropriate to measure outcome variable(s)*	Yes	Yes	Yes	Yes	Yes
Reliability of outcome variable(s) reported*	Yes	Yes	No	Yes	Yes
Description of SM adequate to be reproduced by a registered practitioner	Yes	Yes	No	Yes	Yes
Measurement tool calibrated* (N/A was included under 'Yes')	N/A	N/A	N/A	N/A	N/A
Adequately explained how measurements were taken	Yes	Yes	Yes	Yes	Yes
Clinician competence reported	No	Yes	No	No	No
Assessor competence reported	Yes	Yes	No	Yes	Yes
Statistics transparent	Yes	Yes	N/A	N/A	Yes
Risk of Bias					
Non-RCT/crossover studies only: Assessor blind to before-after readings* (N/A was included under 'Yes')	N/A	N/A	No	Yes	N/A
RCT/crossover studies only: Random allocation*	Yes	Yes	N/A	N/A	Yes
RCT/crossover studies only: Assessor blind to intervention group*	Yes	Yes	N/A	N/A	Yes
RCT/crossover studies only: Statistician blind to control/intervention readings	No	No	N/A	N/A	No
Results					
Unacceptable quality and risk of bias items	0 red 2 yellow	0 red 1 yellow	2 red 3 yellow	0 red 1 yellow	0 red 2 yellow
Overall assessment (Not credible: 1 or more red items and/or 3 or more yellow items)					
	Credible	Credible	Not credible	Credible	Credible

* Critical item (no asterisk: important item)

White: requirement fulfilled (Yes) or not applicable (N/A)

Red: Critical quality or risk of bias item not fulfilled (No)/Not credible article

Yellow: Important quality or risk of bias item not fulfilled (No)

Green: Credible article

Results of not credible studies by outcome variable

Below are shown the results from the 12 not credible articles, reported by outcome variable.

Vertebral position (n=3/3) The results of these 3 not credible studies were conflicting; 2 articles reported post-SM changes in vertebral position, using plain radiography, whereas one reported no change post-SM using ultrasound.

Facet joint space (n=1/5) The 1 not credible study had only 2 participants measured for the relevant outcome variable as part of a larger study, in which all other participants also received traction before and after SM, and so were not considered controls. No change in facet joint space and no presence of pneumarthrosis (discrete bubble

of intra-articular gas) was reported post-SM using computed tomography (CT).

Spinal stiffness (n=2/3) The 2 not credible studies found a reduction in spinal stiffness post-SM. One study used a mechanical indentometer; the other studied horses using a cable extensometer with a pressure sensor mat.

Resting muscle thickness (n=3/6) Results in the 3 not credible studies were conflicting. One reported a pre-post-SM difference in resting transverse abdominal muscle thickness. One reported a difference in the resting thickness of pelvic floor muscles in pregnant women but not in non-pregnant women. The third reported no difference in thickness for multifidus muscles. All studies made measurements using ultrasound.

Table 5 Quality, risk of bias, and credibility assessment for 3 articles on spinal stiffness in a systematic review of anatomical mechanisms of spinal manipulation

Study description	Hausler 2007	Fritz 2011	Wong 2015
Proposed mechanism of spinal manipulation	Spinal stiffness	Spinal stiffness	Spinal Stiffness
Study Design	Crossover	Prospective case series	Non-randomised controlled study
Quality			
Description adequate to reproduce experiment set-up	Yes	Yes	Yes
Evaluation tool(s) appropriate to measure outcome variable(s)*	Yes	Yes	Yes
Reliability of outcome variable(s) reported*	No	Yes	Yes
Description of SM adequate to be reproduced by a registered practitioner	Yes	Yes	Yes
Measurement tool calibrated* (N/A was included under 'Yes')	Yes	Yes	No
Adequately explained how measurements were taken	Yes	Yes	Yes
Clinician competence reported	No	Yes	No
Assessor competence reported	No	N/A (mechanical indentometer)	N/A (mechanical indentometer)
Statistics transparent	Yes	Yes	Yes
Risk of Bias			
Non-RCT/crossover studies only: Assessor blind to before-after readings* (N/A was included under 'Yes')	N/A	N/A (mechanical indentometer)	N/A (mechanical indentometer)
RCT/crossover studies only: Random allocation*	Yes	N/A	N/A
RCT/crossover studies only: Assessor blind to intervention group*	No	N/A	N/A
RCT/crossover studies only: Statistician blind to control/intervention readings	No	N/A	N/A
Results			
Unacceptable quality and risk of bias items	2 red 3 yellow	0 red 0 yellow	1 red 1 yellow
Overall assessment (Not credible: 1 or more red items and/or 3 or more yellow items)	Not credible	Credible	Not credible

* Critical item (no asterisk: important item)

White: requirement fulfilled (Yes) or not applicable (N/A)

Red: Critical quality or risk of bias item not fulfilled (No)/Not credible article

Yellow: Important quality or risk of bias item not fulfilled (No)

Green: Credible article

Intervertebral disc pressure (n = 1/1) There was 1 not credible study which found increased disc pressure post-SM for at least 15 s using a pressure probe inserted into the disc.

Myofascial hysteresis (n = 1/1) The 1 not credible RCT used a durometer to measure hysteresis with mixed results that we found difficult to interpret.

Further damage to already damaged arteries (n = 1/1) The 1 not credible study on this topic was an uncontrolled intervention. The article reported no further damage post-SM to vertebral arteries of dogs that were damaged prior to manipulation with an angioplasty cutting balloon. This study was stopped early for ethical reasons once no effect was detected.

Please see Table 8 for the results and credibility assessments of all studies.

Post-hoc analysis

Cross-referencing credibility with positive (change after SM) or negative (no change after SM) results of SM resulted in a mixed picture. Thus, there was no relationship between the credibility of studies with positive or negative results of these studies. For a visual summary of the findings by outcome variable, credibility, and whether an article reported positive or negative results, see Table 9.

Discussion

Summary of main findings

This systematic review included 8 articles that we considered to be credible and 12 that we regarded as not credible. They dealt with 7 proposed SM mechanisms of action: change in vertebral position, facet joint space, spinal stiffness, resting muscle thickness, IVD pressure, myofascial hysteresis, and further damage to damaged

Table 6 Quality, risk of bias, and credibility assessment for 6 articles on muscle thickness in a systematic review of anatomical mechanisms of spinal manipulation

Study description	Brenner 2007	Raney 2007	Konitzer 2011	Puentedura 2011	Haavik 2016	Fosberg 2019
Proposed mechanism of spinal manipulation	Change in resting thickness of multifidous muscles	Change in resting thickness of lateral abdominal muscles	Change in resting thickness of transverse and internal oblique abdominal muscles	Change in resting thickness of transverse abdominal muscles	Change in resting thickness of pelvic floor muscles	Change in resting thickness of transverse abdominal muscles
Study Design	Case report	Prospective case series	Prospective case series	Crossover	Crossover	RCT
Quality						
Description adequate to reproduce experiment set-up	Yes	Yes	Yes	Yes	No	Yes
Evaluation tool(s) appropriate to measure outcome variable(s)*	Yes	Yes	Yes	Yes	Yes	Yes
Reliability of outcome variable(s) reported*	Yes	Yes	Yes	Yes	Yes	Yes
Description of SM adequate to be reproduced by a registered practitioner	Yes	Yes	Yes	Yes	No	Yes
Measurement tool calibrated* (N/A was included under 'Yes')	N/A	N/A	N/A	N/A	N/A	N/A
Adequately explained how measurements were taken	Yes	Yes	Yes	Yes	No	Yes
Clinician competence reported	No	Yes	Yes	No	Yes	Yes
Assessor competence reported	Yes	Yes	Yes	Yes	Yes	Yes
Statistics transparent	Yes	Yes	Yes	Yes	Yes	Yes
Risk of Bias						
Non-RCT/crossover studies only: Assessor blind to before-after readings* (N/A was included under 'Yes')	No	No	Yes	N/A	N/A	N/A
RCT/crossover studies only: Random allocation*	N/A	N/A	N/A	Yes	No	Yes
RCT/crossover studies only: Assessor blind to intervention group*	N/A	N/A	N/A	Yes	No	Yes
RCT/crossover studies only: Statistician blind to control/ intervention readings	N/A	N/A	N/A	No	No	No
Results						
Unacceptable quality and risk of bias items	1 red 1 yellow	1 red 0 yellow	0 red 0 yellow	0 red 2 yellow	2 red 4 yellow	0 red 1 yellow
Overall assessment (Not credible: 1 or more red items and/or 3 or more yellow items)	Not credible	Not credible	Credible	Credible	Not credible	Credible

* Critical item (no asterisk: important item)

White: requirement fulfilled (Yes) or not applicable (N/A)

Red: Critical quality or risk of bias item not fulfilled (No)/Not credible article

Yellow: Important quality or risk of bias item not fulfilled (No)

Green: Credible article

arteries. We include results from articles that we considered credible as well as ones we considered not credible, in order to report as fully and fairly as possible, any information published to this point on the immediate anatomical/positional changes in response to SM.

The credible articles reported that, post-SM, there were: (i) changes in facet joint space, (ii) changes in spinal stiffness, but (iii) no changes in resting muscle thickness after SM.

A comparison between the results of the credible and not credible articles revealed that the latter:

(i) disagreed with the credible articles on facet joint space/pneumarthrosis (discrete bubble of intra-articular gas), (ii) agreed on spinal stiffness, but (iii) had conflicting results for resting muscle thickness.

In addition, the not credible articles reported (i) conflicting results between them on vertebral position, (ii) change in IVD pressure and (iii) change in myofascial hysteresis, but (iv) no further damage to already damaged arteries after SM.

Table 7 Quality, risk of bias, and credibility assessment for 1 article on intervertebral disc pressure, 1 article on cervical myofascial hysteresis, and 1 article on further damage to already damaged arteries in a systematic review of anatomical mechanisms of spinal manipulation

Study description	Lisi 2006	Barnes 2013	Wynd 2008
Proposed mechanism of spinal manipulation	Change in lumbar intervertebral disc pressure	Cervical myofascial hysteresis	Further damage to already damaged arteries
Study Design	Feasibility	RCT	Uncontrolled intervention
Quality			
Description adequate to reproduce experiment set-up	Yes	No	Yes
Evaluation tool(s) appropriate to measure outcome variable(s)*	Yes	Yes	Yes
Reliability of outcome variable(s) reported*	No	Yes	No
Description of SM adequate to be reproduced by a registered practitioner	Yes	No	Yes
Measurement tool calibrated* (N/A was included under 'Yes')	Yes	No	N/A
Adequately explained how measurements were taken	Yes	No	Yes
Clinician competence reported	Yes	Yes	Yes
Assessor competence reported	Yes	Yes	No
Statistics transparent	Yes	No	Yes
Risk of Bias			
Non-RCT/crossover studies only: Assessor blind to before-after readings* (N/A was included under 'Yes')	No	N/A	No
RCT/crossover studies only: Random allocation*	N/A	Yes	N/A
RCT/crossover studies only: Assessor blind to intervention group*	N/A	Yes	N/A
RCT/crossover studies only: Statistician blind to control/ intervention readings	N/A	No	N/A
Results Unacceptable quality and risk of bias items	2 red 0 yellow	1 red 5 yellow	2 red 1 yellow
Overall assessment (Not credible: 1 or more red items and/or 3 or more yellow items)	Not credible	Not credible	Not credible

* Critical item (no asterisk: important item)

White: requirement fulfilled (Yes) or not applicable (N/A)

Red: Critical quality or risk of bias item not fulfilled (No)/Not credible article

Yellow: Important quality or risk of bias item not fulfilled (No)

Comparisons to the literature

To our knowledge, this is the first systematic review dealing exclusively with the immediate anatomical/positional changes in response to SM. However, a systematic review on spinal mobilisation (i.e., not HVLA manipulation) reported on articles that evaluated mainly clinical outcomes but also on some anatomical mechanisms [33]. Specifically, they included 4 articles that evaluated spinal stiffness, and 3 of the 4 reported reduced spinal stiffness after mobilisation (not HVLA SM). These findings thus aligned with the results in our 3 credible articles that reported reduced spinal stiffness post-SM.

Methodological considerations of the present systematic review

Literature search Important positive aspects of our literature search were that an experienced information specialist (CH) ensured that we consulted all relevant databases for this topic and that all databases were interrogated using the different input parameters necessary to retrieve the relevant articles. We also used broad search

parameters with no date limit, to capture all relevant articles on the topic. The search was updated to ensure we captured any more recent articles prior to submission of our review. We limited the search to articles in English only. Although, therefore, we may have failed to include every relevant published article, we believe this to be a minor limitation, as most articles dealing with SM are typically published in the English language. Also, we did not follow the PRESS guideline recommendation [54] to perform a review of the search strategy, which might have affected the quality and comprehensiveness of our search.

Inclusion/exclusion criteria To ensure that SM was the most likely reason for any potential change in an outcome variable, we included only articles on studies that performed pre- and post-SM measurements and reported this for a single SM session. If some effects of SM require multiple sessions to manifest, we would have missed out on such information. We also excluded articles that used SM in combination with any other therapeutic intervention. Therefore, if SM requires facilitation by another

Table 8 Results reported in 20 articles included in systematic review of anatomical mechanisms of spinal manipulation. Table shows results grouped by proposed mechanism of spinal manipulation (column 2), with detailed results (column 4), overall credibility (column 5), and whether results were positive or negative (column 6)

Change in vertebral position

First author year of publication	Proposed mechanism(s) of spinal manipulation investigated/outcome variable	Study design Total number (N) of participants	Detailed results relating to SMT v. control (best case scenario - for the definition of best-case scenario please see text in the methods section)	Overall credibility assessment (RoB+quality) Credible (green)/ Not credible (red)	Positive (green)* or negative (red)**
Jirout 1972	Change in cervical vertebral position	Uncontrolled intervention N=250	Conflicting changes in vertebral segmental tilt from pre- to post-manipulation. In 65 (26%) anterior tilt increased in all measured segments using plain radiography. In 95 (38%) anterior tilt decreased in lower segments and increased in upper segments. In 37 (15%) anterior tilt increased in lower segments and decreased in upper segments. In 26 (10%) anterior tilt increased or decreased in only 2-3 segments. In 6 (2%) anterior tilt decreased or posterior tilt increased in all measured segments. In 21 (8%) there was no change. Statistical significance not tested.	Not credible	Positive
Palmer 2009	Change in cervical vertebral position	Retrospective case series N=47	Lateral position of atlas reduced from 1.79 degrees ± 1.10 degrees to 0.28 degrees ± 0.40 degrees pre- to post-manipulation (p<0.001).	Not credible	Positive
Flaum 2017	Change in cervical vertebral position	RCT N=25	No statistically significant differences in ultrasound-measured articular process rotation for the post-manipulation group.	Not credible	Negative

Increase in facet joint space

Cramer 2000	Increase in lumbar facet joint space	RCT N=16	0.7mm average increase in joint space for intervention group (pre- v. post-lumbar manipulation) versus 0.0mm change in control group (pre- v. post-side-posture positioning with post-intervention MRI scans performed in side-posture position. Not stated whether statistically significant.	Credible	Positive
Cramer 2002	Increase in lumbar facet joint space	RCT N=64	1.89 mm average difference in joint space for lumbar manipulation group v. no manipulation group (p<0.000) from pre- to post-MRI scan with post-intervention scans performed in side-posture position. Statistically significant.	Credible	Positive
Cascioli 2003	i) Increase in cervical facet joint space ii) Formation of intra-articular gas bubbles	Uncontrolled intervention N=2	i) No increase in joint space ii) No formation of gas bubbles	Not credible	Negative
Cramer 2011	Increase in lumbar facet joint space	Feasibility N=5	Average 0.5mm ± 0.6mm pre- to post-intervention increase in joint space for manipulated joints v. non-manipulated joints (-0.2 ± 0.6mm), measured in side posture position. Statistical significance not tested.	Credible	Positive
Cramer 2012	Increase in facet joint space	RCT N=40	Average increase in joint space of (0.75 mm) in ‘upside’ manipulated joints compared to upside non-manipulated (side-posture positioning only) joints (0.52 mm), when measured in side-posture position. Statistically significant (p=.03). Joints that cavitated (determined using accelerometry) gapped more than those that did not (mean 0.56 v. 0.22 mm, p=.01).	Credible	Positive

Change in spinal stiffness

Hausler 2007	Change in spinal stiffness	Crossover N=10 horses	Mean statistically significant increase in displacement (decreased stiffness) of 15% (range, 7% to 25%) post-manipulation, compared with 0% (range, -4% to 7%) after the control intervention.	Not credible	Positive
Fritz 2011	Change in spinal stiffness	Prospective case series N=48	Significant immediate changes in global stiffness (p=0.034) and terminal stiffness (p=0.027) were found pre- to post-manipulation.	Credible	Positive
Wong 2015	Change in spinal stiffness	Non-randomised controlled study N=107	The spinal stiffness of SMT responders was significantly reduced after each SMT. No change for SMT non-responders or asymptomatic participants who did not receive SMT.	Not credible	Positive

Table 8 (continued)

Change in muscle thickness					
Brenner 2007	Change in resting multifidous muscle thickness	Case report N=1	No change greater than the calculated minimal detectable change in the multifidus thickness at rest at both the L4-5 and L5-S1 levels either immediately post-manipulation or 24 hours later.	Not credible	Negative
Raney 2007	Change in lateral abdominal muscle resting thickness	Prospective case series N=9	Average decrease of transversus abdominis muscle thickness of 0.48mm (13.9%) pre- to post-manipulation. Average decrease of internal oblique muscle thickness of 0.57mm (5.7%) pre- to post-manipulation. Not stated if statistically significant but was greater than the minimum detectable change of 0.4mm.	Not credible	Positive
Konitzer 2011	Change in resting thickness of transverse and internal oblique abdominal muscles	Prospective case series N=19	No statistically significant differences were found in either the transverse or internal oblique abdominal muscles post-manipulation.	Credible	Negative
Puentedura 2011	Change in resting thickness of transverse abdominal muscles	Crossover, randomized N=35	No statistically significant differences for transverse abdominal muscle thickness at rest pre- to post-manipulation.	Credible	Negative
Haavik 2016	Change in pelvic floor resting muscle thickness	Comparison of 2 study groups both included in a crossover non-randomised experiment i) N=11 pregnant ii) N=15 nonpregnant	Levator hiatal area at rest increased (interpreted as relaxation) from 12.2cm ² ± 2.1cm ² to 14.0cm ² ± 2.0cm ² in pregnant group post-spinal manipulation compared to pre-manipulation. No significant change was reported pre- to post-manipulation in the nonpregnant group. No significant change after passive movements.	Not credible	Positive
Fosberg 2019	Change in transverse abdominal resting muscle thickness	RCT N=67	No significant differences in transverse abdominal muscle thickness between participants with low back pain who received manipulation compared with the participants with LBP who received sham manipulation. Experiment stopped prior to reaching power calculation requirement because no difference was found between groups.	Credible	Negative

Change in intervertebral disc pressure

Lisi 2006	Change in lumbar intervertebral disc (IVD) pressure	Feasibility N=1	Increased IVD pressure post-manipulation in prone position (110kPa to 150kPa) and side lying position (150kPa to 165kPa) for 15 seconds, when recording stopped.	Not credible	Positive
------------------	---	--------------------	--	--------------	----------

Change in myofascial hysteresis

Barnes 2013	Change in myofascial hysteresis (as measured by 4 unitless components)	RCT N=40	Statistically significant change reported (but not specified) for motoricity, frequency, and fixation. "Suggestive trend" reported for mobility.	Not credible	Positive
--------------------	--	-------------	--	--------------	----------

Damage to arteries

Wynd 2008	Further damage to already damaged arteries	Uncontrolled intervention N=10 dogs	No statistically significant differences between the overall pre- and post-manipulation lesion dimensions.	Not credible	Negative
------------------	--	--	--	--------------	----------

* Positive result: spinal manipulation led to a change

** Negative result: spinal manipulation did not lead to a change

Table 9 Synthesis of findings by outcome variable and author in a systematic review of 20 articles of anatomical mechanisms of spinal manipulation

Post-manipulation outcome variable (number of included articles)	Positive (+) and negative (-) results for each credible article			Positive (+) and negative (-) results for each not credible articles		
	Author year (+ or - results)	N of reds	N of yellows	Author	N of reds	N of yellows
Vertebral position (n = 3)				Jirout (+)	3	5
				Palmer (+)	2	4
				Flaum (-)	2	3
Facet joint space (n = 5)	Cramer 2000 (+)	0	2	Cascioli (-)	2	3
	Cramer 2002 (+)	0	1			
	Cramer 2011 (+)	0	1			
	Cramer 2012 (+)	0	2			
Spinal stiffness (n = 3)	Fritz (+)	0	0	Hausler (+)	3	3
				Wong (+)	1	2
Resting muscle thickness (n = 6)	Konitzer (-)	0	0	Brenner (-)	1	1
	Puentedura (-)	0	2	Raney (-)	1	0
	Fosberg (-)	0	1	Haavik (+)	3	4
Intervertebral disc pressure (n = 1)				Lisi (+)	1	0
Myofascial hysteresis (n = 1)				Barnes (+)	0	5
Further damage to damaged arteries (n = 1)				Wynd (-)	2	1

Positive (+): article reported a change after spinal manipulation
 Negative (-): article reported no change after spinal manipulation
 Red: critical quality or risk of bias item(s)
 Yellow: important quality or risk of bias item(s)

intervention for the effects to manifest, we would have missed those changes. There may have been differences among team members in interpretation of inclusion/exclusion criteria which could have led to articles being missed out, although we mitigated this possibility with calibration sessions and written instructions, so this is unlikely.

Full text screening Eleven articles included at the title/abstract screening stage were not available for the full-text screening process. Nine of those were published in chiropractic subluxation-focused journals, thus not available through mainstream library systems, with authors either not found on ResearchGate or not responding to requests for articles. This missingness of data may have affected our conclusions, but our experience is that the “grey” chiropractic journals do not attract high quality articles, so we do not believe that our conclusions were impacted.

Quality/risk of bias It is well known that clinical studies will more easily produce positive findings if the human factor is allowed to play an essential role by voluntarily or subconsciously introducing bias. This is the reason why systematic reviews assess the RoB before drawing conclusions on the validity of results. This phenomenon was clearly shown in a previous systematic review on the “effect” of spinal manipulative therapy (SMT) on non-musculoskeletal conditions [55]. All studies that were considered to have failed in preventing the “human factor” reported positive results, whereas none of the high-quality studies found there to be an “effect” of SMT [55].

We emphasized RoB only in relation to aspects that clearly could be influenced by the beliefs and wishes of the researchers. Experimental/basic science (e.g. laboratory) studies, are susceptible to fewer RoB items. Instead, commercial and university laboratories are often subjected to accreditation procedures [56, 57],

which are specific to the requirements of their area of activities and relate more to technicalities than to systematic human errors. Thus, the use of appropriate and calibrated tools that are operated by competent personnel would be paramount in preventing random errors in studies that rely on technical assessments, beyond that of human subjective observations and reactions.

Results in studies that deal with anatomical/positional changes after SM are, thus, unlikely to be influenced by the beliefs and wishes of study participants and clinicians, whereas the beliefs and wishes of the assessor and statistician could have an impact. Failing to use a reliable outcome variable, measuring changes with a non-calibrated machine, or allowing amateurs to conduct the study, could possibly induce bias but more likely result in random errors.

Therefore, in addition to the RoB, we accounted for several technical aspects, defined as “quality”. Quality issues are likely to cause non-systematic errors, as opposed to bias, which may cause systematic errors. Examples of quality issues that must be assured to prevent non-systematic errors are skills of assessment personnel and the calibration of measurement tools. To accurately judge the technical quality of studies, we ensured that each review team had at least one member with expertise in the area.

Nonetheless, the evaluation and weighting of methods was a subjective matter, and other investigators may have judged differently. There may also have been differences among team members in the interpretation of quality and RoB criteria, which could have led to differences in results. However, we mitigated that possibility with multiple calibration sessions and repeated consultations on iterations of the tables among the team members.

Results of studies The results were extracted from each article only after the quality and RoB items were determined, to avoid reviewer bias of the quality/RoB assessment. To prevent biased results interpretation, extraction of the study outcomes and interpretation of findings were performed by other teams than the experts in the area, although the experts were invited to critically review the extracted findings and interpretations.

Synthesis Part of the standard synthesis in systematic reviews is to identify ‘gaps in the literature’. However, these ‘gaps’ may not be areas that have not been studied, just areas that have not been studied *well*. By listing all the studies that have been conducted in this area and detailing all the methodological errors that we identified, we show which areas *can* be studied (or which may be too difficult to study with current technology) and also indicate how they may be studied *better* in the future.

Methodological considerations for reviewed articles and potential future studies

General comments In the hope of being helpful to future researchers, we provide some methodological comments regarding the techniques used to study the potential anatomical/positional effects of SM.

Only 8 of the 20 studies were assessed as credible. Consequently, this indicates that technical experimental studies have not been prioritised in environments that were created for such purposes, i.e., taking advantage of relevant equipment and skilled researchers working in dedicated research laboratories. Assessors for these procedures should be highly experienced or adequately trained. In addition, when there is a human element, it is important to establish inter-/intra-rater reliability of the assessors, before undertaking the study. Further, researchers must strive towards the use of measurement tools that are validated, and it is important to remember that some need to be calibrated. The use of frameworks like COSMIN (Consensus-based Standards for the selection of health Measurement Instruments) [58] may be helpful in this process. Also, when measuring positions or spaces, it would be of utmost importance to place study subjects in identical positions before and after the SM, to prevent ‘normal’ aberrations and measurement errors due to distortion.

Items relating to relevant RoB should be observed, in particular, blinding of before-after readings in studies without control groups. Statisticians should be blinded to control/intervention readings. Statistical methods, cut points, etc. should be determined a priori, rather than after any results have been returned.

Comments relating to each outcome variable Vertebral position

Two of 3 studies on vertebral position used radiographs to try to capture very small post-SM changes (<2 degrees of rotation [14] or <4mm of displacement). However, we note that the use of plain radiography to detect very small changes in vertebral position is debatable, at best, and in our opinion, the use of radiography for these measurements was not appropriate [59]. Instead, we suggest that future investigations could use computed tomography, which offers much greater resolution. The precision and likely error tolerance of the measurement instrument should be stated in future studies.

The third study used ultrasound to measure paraspinous tissue thickness as a proxy for vertebral rotation. However, the pressure on the ultrasound head was not measured. This is important, as increased pressure could compress tissues and distort readings. Nevertheless, this method could hold promise for developing a method of measuring vertebral rotation without the use of ionising radiation (Tables 2 and 3).

Facet joint space

All 4 credible articles found an increase in facet joint space post-SM. However, as they were conducted by the same team of researchers, these results should be confirmed by at least 1 independent team.

MRI is likely optimal for viewing facet joint space increase that is retained after the SM event, as bone and capsule detail can be captured.

Theories exist on the potential role of SM on intra-articular meniscoids/discoids, synovial folds, and adhesions within the facet joints [60–65]. We found no relevant articles that investigated any of these anatomical variations/pathologies. We suggest that if someone wanted to study them, they may be visualised using high resolution imaging such as MRI (Tables 2 and 4).

Spinal stiffness

The use of the mechanical indentometer in 2 of the 3 studies on spinal stiffness was useful because it removed the human element from the interpretation of the measurements. Calibration of the tool is important in this area (Tables 2 and 5).

Resting muscle thickness

Since 3 credible studies all showed absence of changes to resting muscle thickness, we believe that this area no longer warrants further study. We found the study design in one of the not credible articles [50] overly complex and difficult to understand; it seemed to report on 2 crossover studies, with important differences between the intervention and control groups (Tables 2 and 6).

Intervertebral disc pressure

The 1 study conducted on IVD pressure showed that it is difficult to study. The use of a pressure-measuring probe to physically penetrate the disc is not attractive to study participants and seems ethically challenging, as it damages the disc tissue to an unknown extent. Perhaps an indirect method of measuring IVD pressure could be developed in the future, for instance using fluid diffusion into or out of the disc, as measured by MRI, as an indicator of pressure. Alternatively, perhaps candidates for IVD replacement could be included as participants (Tables 2 and 7).

Further damage to already damaged arteries

In vivo studies of arterial walls are also difficult to perform. In addition, it seems unlikely that SM can cause arterial damage de novo [66–72]. The assumption tested in this article is interesting, if one thinks that it is mainly arteries with pre-existing damage or pathology

that are susceptible to further damage by SM. Hence, the authors devised a method to “pre-damage” vertebral arteries in dogs, as a proxy for “naturally occurring” damage or pathology. The types of lesions created were not predictable, for which reason the usefulness of this method is unclear.

The assessment method in this study was complex, using a fluoroscopically guided ultrasound probe as the measurement tool. However, magnetic resonance angiography would offer better resolution, or a micro-video-camera could allow direct visualisation of arterial damage.

Since vertebral artery dissection is so rare and only temporally linked to SM [73, 74], the justification of sacrificing animals to study this should be considered (Tables 2 and 7).

Myofascial hysteresis

There were several unclear elements in this article, making it difficult for us to interpret how well the outcome variables in the article related to the concept of hysteresis (Tables 2 and 7).

Conclusions

Clinical perspectives

Although this review is primarily valuable to researchers, clinicians should also benefit from our findings. It is a common clinical observation that patients can experience sudden relief immediately after SM. In our experience, when this happens, they may ask: “What exactly happened when you cracked my back?” As this review describes, there is no easy answer because of the many theories and few facts. Nevertheless, we suggest the following, which clinicians can modify to suit their practice and patients. Regarding anatomical/positional changes, it would be possible to say: “There is no simple answer, because the spine is a difficult area to study. It seems likely that the manipulation/adjustment causes some physical changes, but it is not known exactly how. Presently, though, we are fairly confident that the facets, i.e., the small joints at the back of the spine, open up a little bit. There also seems to be a measurable change in the stiffness of the spine immediately after manipulation. We assume that these changes are part of what helps you feel better.”

Research perspectives

- There has been little research on anatomical mechanisms of SM, and most of the articles we found were not credible according to our assessment methods. The few studies that have been published are on a wide variety of topics, performed by a

small number of researchers, and were often small studies (only 7 studies recruited more than 50 participants) that were not followed-up by other similar studies. It seems that there has been no coherent research planning strategy undertaken by any of the manual therapy professions to investigate the anatomical/positional mechanisms of SM. Therefore, there is an opportunity to develop research centres with areas of expertise that can lead high-quality studies in these areas concentrating on anatomically feasible outcome variables.

- Cut points for meaningful changes should be established and should incorporate information such as normal variations, repeatability, and inter- and intra-examiner reliability. The time between the application of SM and the measurement of the potential effect is also important to establish, to infer mechanism(s).
- The results of our review indicate that the 2 most promising areas for further study are changes to facet joint space and spinal stiffness after SM.
- However, after having established what actually happens anatomically in response to SM, it would be important to continue by investigating whether these mechanisms also have a lagged effect and/or result in physiological reactions. Then, this potential chain of events must be linked to the clinical picture, that is, reduction of pain or improvement in function.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12998-024-00549-w>.

Additional file 1.

Additional file 2.

Acknowledgements

Thanks to Amber Beynon for assistance with article screening.

Author contributions

Concept and research design: KJY, CLY, CH. Coordination and quality assurance of article screening, critical appraisal and data extraction: KJY, CLY, PS, IA, LG, DE, OG, SV. Article screening, data extraction, and critical appraisal: KJY, CN, CB, CM, DE, EL, GG, KCL, LG, LN, RK, SH, VG, SV. Writing and final editing of the manuscript: KJY, CLY. All co-authors edited and approved the final manuscript. Humans wrote this manuscript with no assistance from artificial intelligence except for Microsoft Word autocorrect for help with spelling and grammar.

Funding

No specific funding was received for this project. Authors have been funded by their respective sites of employment (see affiliations) and none of the authors is dependent on present or future funding relating to mechanisms of spinal manipulation. CH is part-funded by the National Institute for Health and Care Research (NIHR) Applied Research Collaboration (ARC) North West Coast (NWC). The views expressed are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethical approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

IA is Co-Editor-in-Chief of *Chiropractic and Manual Therapies*. The editorial management system automatically blinded her from the submitted manuscript, and she had no part in this manuscript's editorial or peer-review process.

Author details

¹Allied Health Research Unit, University of Central Lancashire, Preston, UK. ²Department of Regional Health Research, University of Southern Denmark, Odense, Denmark. ³University of Central Lancashire, Preston, UK. ⁴Department of Chiropractic Medicine, Integrative Spinal Research Group, Balgrist University Hospital, University of Zurich, Zurich, Switzerland. ⁵Department of Clinical Sciences, Obstetrics and Gynecology, Umeå University, Umeå, Sweden. ⁶School of Sport, Exercise and Rehabilitation Sciences, University of Birmingham, Edgbaston, Birmingham, UK. ⁷Karolinska Institutet, Institute of Environmental Medicine, Nobels V. 13, 177 77 Stockholm, Sweden. ⁸The Norwegian Chiropractors' Research Foundation «Et Liv I Bevegelse», Lilleakerveien 31, 0283 Oslo, Norway. ⁹Department of Health and Functioning, Faculty of Health and Social Sciences, Western Norway University of Applied Sciences, Bergen, Norway. ¹⁰Université de Paris-Saclay, Gif-sur-Yvette, France. ¹¹Advanced Physiotherapist Practitioner, University of Nottingham, A25 Academic Rheumatology, Clinical Sciences Building, City Hospital, Nottingham, UK. ¹²Synthesis, Economic Evaluation and Decision Science (SEEDS) Group, Health Technology Assessment Unit | Applied Health Research Hub, University of Central Lancashire, Preston, UK. ¹³Methodological Innovation, Development, Adaptation and Support (MIDAS) Group, NIHR Applied Research Collaboration North West Coast (ARC NWC), Liverpool, UK. ¹⁴Center for Muscle and Joint Health, Department of Sports Science and Clinical Biomechanics, University of Southern Denmark, Odense, Denmark. ¹⁵Medical Research Unit, Spine Centre of Southern Denmark, Lillebaelt Hospital, University Hospital of Southern Denmark, Middelfart, Denmark. ¹⁶School of Health Sciences, Medical School, Queen's Medical Centre, University of Nottingham, Room B228a, Nottingham, UK. ¹⁷Division of Physiotherapy and Rehabilitation Sciences, University of Nottingham, Nottingham, England. ¹⁸Department of Health Professions, Faculty of Health and Education, Brooks Building, Manchester Metropolitan University, Manchester, UK. ¹⁹University College of Osteopathy, 275 Borough High Street, London, UK. ²⁰Health Sciences, Doornfontein Campus, University of Johannesburg, Johannesburg, South Africa.

Received: 26 January 2024 Accepted: 27 August 2024

Published online: 11 September 2024

References

1. Plunkett A, Fawkes C, Carnes D. Osteopathic practice in the United Kingdom: a retrospective analysis of practice data. *Plos One*. 2022;17(7):e0270806.
2. National Board of Chiropractic Examiners. Practice analysis of chiropractic 2020. Greeley: National Board of Chiropractic Examiners; 2020.
3. Beliveau PJH, Wong JJ, Sutton DA, Simon NB, Bussi eres AE, Mior SA, et al. The chiropractic profession: a scoping review of utilization rates, reasons for seeking care, patient profiles, and care provided. *Chiropr Man Ther*. 2017;25(1):35.
4. Murthy VMSMPH, Sibbritt DWP, Adams JP. An integrative review of complementary and alternative medicine use for back pain: a focus on

- prevalence, reasons for use, influential factors, self-perceived effectiveness, and communication. *Spine J.* 2015;15(8):1870–83.
5. Goertz CM, Pohlman KA, Vining RD, Brantingham JW, Long CR. Patient-centered outcomes of high-velocity, low-amplitude spinal manipulation for low back pain: a systematic review. *J Electromyogr Kinesiol.* 2012;22(5):670–91.
 6. Masaracchio M, Kirker K, States R, Hanney WJ, Liu X, Kolber M. Thoracic spine manipulation for the management of mechanical neck pain: a systematic review and meta-analysis. *Plos One.* 2019;14(2):e0211877.
 7. Franke H, Franke J-D, Fryer G. Osteopathic manipulative treatment for chronic nonspecific neck pain: a systematic review and meta-analysis. *Int J Osteopath Med.* 2015;18(4):255–67.
 8. Organization WH. WHO guideline for non-surgical management of chronic primary low back pain in adults in primary and community care settings. Geneva: World Health Organization; 2023.
 9. National Institute for Health and Care Excellence. Low back pain and sciatica in over 16s: assessment and management. London: National Institute for Health and Care Excellence (NICE); 2016.
 10. Beroes JM, Booth MS, Miake-Kye I, Paige NM, Shekelle PG. Association of spinal manipulation therapy with clinical benefit and harm for acute low back pain systematic review and meta-analysis. *JAMA J Am Med Assoc.* 2017;317(14):1451.
 11. Coronado RA, Gay CW, Bialosky JE, Carnaby GD, Bishop MD, George SZ. Changes in pain sensitivity following spinal manipulation: a systematic review and meta-analysis. *J Electromyogr Kinesiol.* 2012;22(5):752–67.
 12. Howick J, Glasziou P, Aronson JK. Evidence-based mechanistic reasoning. *J R Soc Med.* 2010;103(11):433–41.
 13. Evans DW. Mechanisms and effects of spinal high-velocity, low-amplitude thrust manipulation: previous theories. *J Manip Physiol Ther.* 2002;25(4):251–62.
 14. Palmer J, Dickholtz M. Improvement in radiographic measurements, posture, pain and quality of life in non-migraine headache patients undergoing upper cervical chiropractic care: a retrospective practice based study. *Ann Vertebr Subluxation Res.* 2009;2009(1):1–11.
 15. Puentedura EJ, Landers MR, Hurt K, Meissner M, Mills J, Young D. Immediate effects of lumbar spine manipulation on the resting and contraction thickness of transversus abdominis in asymptomatic individuals. *J Orthop Sports Phys Ther.* 2011;41(1):13–21.
 16. Marcon AR, Murdoch B, Caulfield T. The “subluxation” issue: an analysis of chiropractic clinic websites. *Arch Physiother.* 2019;9(1):11.
 17. Palmer DD. The chiropractor’s adjuster: textbook of the science, art and philosophy of chiropractic for students and practitioners. Portland: Portland Printing House Co; 1910.
 18. Keller TS, Colloca CJ. Mechanical force spinal manipulation increases trunk muscle strength assessed by electromyography: a comparative clinical trial. *J Manip Physiol Ther.* 2000;23(9):585–95.
 19. Duarte FCK, Funabashi M, Starmer D, Partata WA, West DWD, Kumbhare DA, et al. Effects of distinct force magnitude of spinal manipulative therapy on blood biomarkers of inflammation: a proof of principle study in healthy young adults. *J Manip Physiol Ther.* 2022;45(1):20–32.
 20. Senzon SA. The chiropractic vertebral subluxation part 10: integrative and critical literature from 1996 and 1997. *J Chiropr Humanit.* 2018;25:146–68.
 21. Young KJ. Evaluation of publicly available documents to trace chiropractic technique systems that advocate radiography for subluxation analysis: a proposed genealogy. *J Chiropr Humanit.* 2014;21(1):1–24.
 22. Picchiottino M, Leboeuf-Yde C, Gagey O, Hallman DM. The acute effects of joint manipulative techniques on markers of autonomic nervous system activity: a systematic review and meta-analysis of randomized sham-controlled trials. *Chiropr Manual Ther.* 2019;27(1):17.
 23. Meyer A-L, Amorim M-A, Schubert M, Schweinhardt P, Leboeuf-Yde C. Unravelling functional neurology: Does spinal manipulation have an effect on the brain? A systematic literature review. *Chiropr Manual Ther.* 2019;27(1):60.
 24. Vernon H. Historical overview and update on subluxation theories. *J Chiropr Hum.* 2010;17(1):22–32.
 25. Gorrell LM, Nyirö L, Pasquier M, Pagé I, Heneghan NR, Schweinhardt P, et al. Spinal manipulation characteristics: a scoping literature review of force-time characteristics. *Chiropr Manual Ther.* 2023;31(1):1–36.
 26. Funabashi M, Son J, Pecora CG, Tran S, Lee J, Howarth SJ, et al. Characterization of thoracic spinal manipulation and mobilization forces in older adults. *Clin Biomech Bristol.* 2021;89:105450.
 27. Funabashi M, Kawchuk GN, Vette AH, Goldsmith P, Prasad N. Tissue loading created during spinal manipulation in comparison to loading created by passive spinal movements. *Sci Rep.* 2016;6(1):38107.
 28. Herzog W, Conway PJ, Kawchuk GN, Zhang Y, Hasler EM. Forces exerted during spinal manipulative therapy. *Spine.* 1993;18(9):1206–12.
 29. Herzog W. The biomechanics of spinal manipulation. *J Bodyw Mov Ther.* 2010;14(3):280–6.
 30. Herzog W, Kats M, Symons B. The effective forces transmitted by high-speed, low-amplitude thoracic manipulation. *Spine.* 2001;26(19):2105–10.
 31. Kovanur-Sampath K, Treffel L, Thomson OP, Rodi JD, Fleischmann M, Tumilty S. Changes in biochemical markers following a spinal manipulation—a systematic review update. *J Man Manip Ther.* 2023; 1–23.
 32. Kovanur-Sampath KMO, Mani RP, Cotter JP, Gisselman ASPT, Tumilty SP. Changes in biochemical markers following spinal manipulation—a systematic review and meta-analysis. *Musculoskelet Sci Pract.* 2017;29:120–31.
 33. Aguirrebeña IL, Newham D, Critchley DJ. Mechanism of Action of Spinal Mobilizations: A Systematic Review. *Spine.* 2016;41(2):159–72.
 34. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ.* 2021;372:n71.
 35. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan—a web and mobile app for systematic reviews. *Syst Rev.* 2016;5(210):1–10.
 36. Jirout J. The effect of mobilisation of the segmental blocade on the sagittal component of the reaction on lateroflexion of the cervical spine. *Neuroradiology.* 1972;3(4):210–5.
 37. Flaum TB, Arfa Mirza A, Rusnack FM-A, Apoznanski TE, Munarova A, Mazzie J, et al. Palpatory and ultrasound assessment of cervical dysfunctions and the effect of cervical high-velocity, low-amplitude (HVLA) technique. *AAO J.* 2017;27:9–16.
 38. Cramer GD, Tuck NR, Knudsen JT, Fonda SD, Schliesser JS, Fournier JT, et al. Effects of side-posture positioning and side-posture adjusting on the lumbar zygapophysial joints as evaluated by magnetic resonance imaging: a before and after study with randomization. *J Manip Physiol Ther.* 2000;23(6):380–94.
 39. Cramer GD, Gregerson DM, Knudsen JT, Hubbard BB, Ustas LM, Cantu JA. The effects of side-posture positioning and spinal adjusting on the lumbar Z joints: a randomized controlled trial with sixty-four subjects. *Spine.* 2002;27(22):2459–66.
 40. Cramer GD, Ross KDCP, Pocius JMS, Cantu JADC, Laptook EDC, Fergus MDC, et al. Evaluating the relationship among cavitation, zygapophysial joint gapping, and spinal manipulation: an exploratory case series. *J Manip Physiol Ther.* 2011;34(1):2–14.
 41. Cramer GD, Ross KDCP, Raju PKP, Cambron JDCMPPH, Cantu JADC, Bora PM, et al. Quantification of cavitation and gapping of lumbar zygapophysial joints during spinal manipulative therapy. *J Manip Physiol Ther.* 2012;35(8):614–21.
 42. Cascioli V, Corr P, Till AG. An investigation into the production of intra-articular gas bubbles and increase in joint space in the zygapophysial joints of the cervical spine in asymptomatic subjects after spinal manipulation. *J Manip Physiol Ther.* 2003;26(6):356–64.
 43. Fritz JM, Koppenhaver SLSL, Kawchuk GNGN, Teyhen DSDS, Hebert JJJJ, Childs JDJD. Preliminary investigation of the mechanisms underlying the effects of manipulation: exploration of a multi-variate model including spinal stiffness, multifidus recruitment, and clinical findings. *Spine.* 2011;36(21):1772–81.
 44. Haussler KK, Hill AE, Puttlitz CM, McLwraith CW. Effects of vertebral mobilization and manipulation on kinematics of the thoracolumbar region. *Am J Vet Res.* 2007;68(5):508–16.
 45. Wong AYL, Parent EC, Dhillon SS, Prasad N, Kawchuk GN. Do participants with low back pain who respond to spinal manipulative therapy differ biomechanically from nonresponders, untreated controls or asymptomatic controls? *Spine.* 2015;40(17):1329–37.
 46. Konitzer LN, Gill NW, Koppenhaver SL. Investigation of abdominal muscle thickness changes after spinal manipulation in patients who meet a clinical prediction rule for lumbar stabilization. *J Orthop Sports Phys Ther.* 2011;41(9):666–74.
 47. Fosberg KK, Puentedura E, Schmitz B, Jain TK, Cleland JA. The effects of thrust joint manipulation on the resting and contraction thickness of transversus abdominis in patients with low back pain: a randomized control trial. *J Manip Physiol Ther.* 2020;43(4):339–55.

48. Brenner AK, Gill NW, Buscema CJ, Kiesel K. Improved activation of lumbar multifidus following spinal manipulation: a case report applying rehabilitative ultrasound imaging. *J Orthop Sports Phys Ther.* 2007;37(10):613–9.
49. Raney NH, Teyhen DS, Childs JD. Observed changes in lateral abdominal muscle thickness after spinal manipulation: a case series using rehabilitative ultrasound imaging. *J Orthop Sports Phys Ther.* 2007;37(8):472–9.
50. Haavik HBPP, Murphy BADCMP, Kruger JBMP. Effect of spinal manipulation on pelvic floor functional changes in pregnant and nonpregnant women: a preliminary study. *J Manip Physiol Ther.* 2016;39(5):339–47.
51. Lisi AJ, Oneill CW, Lindsey DP, Cooperstein R, Cooperstein E, Zucherman JF. Measurement of in vivo lumbar intervertebral disc pressure during spinal manipulation: a feasibility study. *J Appl Biomech.* 2006;22(3):234–9.
52. Barnes PL, Laboy F 3rd, Noto-Bell L, Ferencz V, Nelson J, Kuchera ML. A comparative study of cervical hysteresis characteristics after various osteopathic manipulative treatment (OMT) modalities. *J Bodyw Mov Ther.* 2013;17(1):89–94.
53. Wynd S, Anderson T, Kawchuk G. Effect of cervical spine manipulation on a pre-existing vascular lesion within the canine vertebral artery. *Cerebrovasc Dis.* 2008;26(3):304–9.
54. McGowan J, Sampson M, Salzwedel DM, Cogo E, Foerster V, Lefebvre C. PRESS peer review of electronic search strategies: 2015 guideline statement. *J Clin Epidemiol.* 2016;75:40–6.
55. Côté P, Hartvigsen J, Axén I, Leboeuf-Yde C, Corso M, Shearer H, et al. The global summit on the efficacy and effectiveness of spinal manipulative therapy for the prevention and treatment of non-musculoskeletal disorders: a systematic review of the literature. *Chiropr Man Ther.* 2021;29(1):8.
56. Zima T. Accreditation of medical laboratories—system, process, benefits for labs. *J Med Biochem.* 2017;36(3):231–7.
57. New US. System for lab accreditation starts May 1998. *J Res Nat Inst Stand Technol.* 1997;102(5):605.
58. Mokkink LB, Prinsen CAC, Bouter LM, de Vet HCW, Terwee CB. The CONsensus-based standards for the selection of health measurement INstruments (COSMIN) and how to select an outcome measurement instrument. *Revista brasileira de fisioterapia (São Carlos (São Paulo, Brazil)).* 2016;20(2):105–13.
59. Sigler DC, Howe JW. Inter- and intra-examiner reliability of the upper cervical X-ray marking system. *J Manip Physiol Ther.* 1985;8(2):75–80.
60. Scholz C, Hubbe U, Kogias E, Klingler J-H. Incomplete resection of lumbar synovial cysts—evaluating the risk of recurrence. *Clin Neurol Neurosurg.* 2015;136:29–32.
61. Manchikanti L, Pampati V, Fellows B, Rivera J, Beyer CD, Damron KS. Role of one day epidural adhesiolysis in management of chronic low back pain: a randomized clinical trial. *Pain Physician.* 2001;4(2):153–66.
62. Schulte TL, Filler TJ, Struwe P, Liem D, Bullmann V. Intra-articular meniscoid folds in thoracic zygapophysial joints. *Spine.* 2010;35(6):E191–7.
63. Inami S, Kaneoka K, Hayashi K, Ochiai N. Types of synovial fold in the cervical facet joint. *J Orthop Sci.* 2000;5(5):475–80.
64. Farrell SF, Khan S, Osmotherly PG, Sterling M, Cornwall J, Rivett DA. Lateral atlantoaxial joint meniscoid volume in individuals with whiplash associated disorder: a case-control study. *Musculoskelet Sci Pract.* 2018;33:46–52.
65. Farrell SF, Osmotherly PG, Cornwall J, Sterling M, Rivett DA. Cervical spine meniscoids: an update on their morphological characteristics and potential clinical significance. *Eur Spine J.* 2017;26(4):939–47.
66. Gorrell LM, Sawatsky A, Edwards WB, Herzog W. Vertebral arteries do not experience tensile force during manual cervical spine manipulation applied to human cadavers. *J Man Manip Ther.* 2023;31(4):261–9.
67. Gorrell LM, Kuntze G, Ronsky JL, Carter R, Symons B, Triano JJ, et al. Kinematics of the head and associated vertebral artery length changes during high-velocity, low-amplitude cervical spine manipulation. *Chiropract Man Ther.* 2022;30(1):28.
68. Piper SL, Howarth SJ, Triano J, Herzog W. Quantifying strain in the vertebral artery with simultaneous motion analysis of the head and neck: a preliminary investigation. *Clin Biomech (Bristol, Avon).* 2014;29(10):1099–107.
69. Herzog W, Leonard TR, Symons B, Tang C, Wuest S. Vertebral artery strains during high-speed, low amplitude cervical spinal manipulation. *J Electromyogr Kinesiol.* 2012;22(5):740–6.
70. Symons BP, Leonard T, Herzog W. Internal forces sustained by the vertebral artery during spinal manipulative therapy. *J Manip Physiol Ther.* 2002;25(8):504–10.
71. Wuest S, Symons B, Leonard T, Herzog W. Preliminary report: biomechanics of vertebral artery segments C1–C6 during cervical spinal manipulation. *J Manip Physiol Ther.* 2010;33(4):273–8.
72. Fagundes C, Herzog W. Strain of the vertebral artery during passive neck movements and spinal manipulation of the cervical spine: an observational study. *J Bodyw Mov Ther.* 2024;40:569–74.
73. Chung CLRDC, Côté PDCP, Stern PDC, L'Espérance GMD. The association between cervical spine manipulation and carotid artery dissection: a systematic review of the literature. *J Manip Physiol Ther.* 2015;38(9):672–6.
74. Church EW, Sieg EP, Zalatimo O, Hussain NS, Glantz M, Harbaugh RE. Systematic review and meta-analysis of chiropractic care and cervical artery dissection: no evidence for causation. *Cureus.* 2016;8(2):e498.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.