






Please cite the Published Version

Smith, Kieran , Taylor, Guy S , Peeters, Wouter, Walker, Mark, Perazzolo, Simone, Atabaki-Pasdar, Naeimeh, Bowden Davies, Kelly A , Karpe, Fredrik, Hodson, Leanne , Stevenson, Emma J and West, Daniel J  (2024) Elevations in plasma glucagon are associated with reduced insulin clearance after ingestion of a mixed-macronutrient meal in people with and without type 2 diabetes. *Diabetologia*, 67 (11). pp. 2555-2567. ISSN 0012-186X

DOI: <https://doi.org/10.1007/s00125-024-06249-7>

Publisher: Springer

Version: Published Version

Downloaded from: <https://e-space.mmu.ac.uk/635680/>

Usage rights:  Creative Commons: Attribution 4.0

Additional Information: The version of record of this article, first published in *Diabetologia*, is available online at Publisher's website: <http://dx.doi.org/10.1007/s00125-024-06249-7>

Data Access Statement: The raw data supporting the conclusions of this article will be made available upon reasonable request.

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)



Elevations in plasma glucagon are associated with reduced insulin clearance after ingestion of a mixed-macronutrient meal in people with and without type 2 diabetes

Kieran Smith^{1,2,3} · Guy S. Taylor^{2,3} · Wouter Peeters^{2,3} · Mark Walker⁴ · Simone Perazzolo^{5,6} · Naeimeh Atabaki-Pasdar^{1,7,8} · Kelly A. Bowden Davies^{2,9} · Fredrik Karpe^{1,7} · Leanne Hodson^{1,7} · Emma J. Stevenson^{2,3} · Daniel J. West^{2,3}

Received: 25 March 2024 / Accepted: 1 July 2024
© The Author(s) 2024

Abstract

Aims/hypothesis The temporal suppression of insulin clearance after glucose ingestion is a key determinant of glucose tolerance for people without type 2 diabetes. Whether similar adaptations are observed after the ingestion of a mixed-macronutrient meal is unclear.

Methods In a secondary analysis of data derived from two randomised, controlled trials, we studied the temporal responses of insulin clearance after the ingestion of a standardised breakfast meal consisting of cereal and milk in lean normoglycaemic individuals ($n=12$; Lean-NGT), normoglycaemic individuals with central obesity ($n=11$; Obese-NGT) and in people with type 2 diabetes ($n=19$). Pre-hepatic insulin secretion rates were determined by the deconvolution of C-peptide, and insulin clearance was calculated using a single-pool model. Insulin sensitivity was measured by an oral minimal model.

Results There were divergent time course changes in insulin clearance between groups. In the Lean-NGT group, there was an immediate post-meal increase in insulin clearance compared with pre-meal values ($p<0.05$), whereas insulin clearance remained stable at baseline values in Obese-NGT or declined slightly in the type 2 diabetes group ($p<0.05$). The mean AUC for insulin clearance during the test was ~40% lower in the Obese-NGT (1.3 ± 0.4 l min⁻¹ m⁻²) and type 2 diabetes (1.4 ± 0.7 l min⁻¹ m⁻²) groups compared with Lean-NGT (1.9 ± 0.5 l min⁻¹ m⁻²; $p<0.01$), with no difference between the Obese-NGT and type 2 diabetes groups. HOMA-IR and glucagon AUC emerged as predictors of insulin clearance AUC, independent of BMI, age or insulin sensitivity (adjusted $R^2=0.670$). Individuals with increased glucagon AUC had a 40% reduction in insulin clearance AUC (~ -0.75 l min⁻¹ m⁻²; $p<0.001$).

Conclusions/interpretation The ingestion of a mixed-macronutrient meal augments differing temporal profiles in insulin clearance among individuals without type 2 diabetes, which is associated with HOMA-IR and the secretion of glucagon. Further research investigating the role of hepatic glucagon signalling in postprandial insulin kinetics is warranted.

Trial registration ISRCTN17563146 and ISRCTN95281775

Keywords Glucagon · Insulin clearance · Insulin resistance · Obesity · Postprandial glycaemia · Type 2 diabetes

Abbreviations

Adipo-IR	Adipocyte insulin resistance	ISR	Pre-hepatic insulin secretion rate
AU	Arbitrary units	Lean-NGT	Lean normoglycaemic
CEACAM1	Carcinoembryonic antigen-related cell adhesion molecule 1	NGT	Normal glucose tolerance
GIP	Glucose-dependent insulinotropic polypeptide	Obese-NGT	Normoglycaemic with central obesity
GLP-1	Glucagon-like peptide 1	PPAR- α	Peroxisome proliferator-activated receptor α
ICR	Insulin clearance rate	Rd_{ins}	Insulin extraction rate
		S_i	Measure of insulin sensitivity
		ϕ_{total}	Measure of beta cell function

Extended author information available on the last page of the article

Research in context

What is already known about this subject?

- The suppression of hepatic insulin clearance after ingestion of an oral glucose load is a key determinant of glucose tolerance in people without type 2 diabetes
- Hepatic insulin clearance is affected by gastrointestinal processes that differ between the ingestion of a mixed-macronutrient meal and a glucose solution

What is the key question?

- How does insulin clearance regulate postprandial glycaemia after ingestion of a mixed-macronutrient meal representative of daily living in people with and without type 2 diabetes?

What are the new findings?

- Insulin clearance is reduced in individuals with obesity, both with and without type 2 diabetes, compared with lean people with normoglycaemia. Indices of hepatic insulin resistance and increased glucagon exposure predict variances in insulin clearance
- Average insulin clearance decreased by 40% in individuals with increased plasma glucagon
- There is a temporal increase in postprandial insulin clearance in lean individuals, negating the increase in insulin secretion

How might this impact on clinical practice in the foreseeable future?

- The ingestion of a mixed-macronutrient meal augments differing temporal profiles in insulin clearance in people without type 2 diabetes that possibly relate to differences in glucagon secretion. Studies investigating the role of hepatic glucagon signalling on postprandial insulin kinetics are warranted

Introduction

Plasma insulin concentrations reflect the balance between the secretion and clearance of insulin [1, 2]. In healthy individuals, ~70% of newly secreted insulin is extracted by the liver during its first passage through the portal system, which is the major site for insulin clearance [1, 2]. Because of the liver's capacity to rapidly adjust the insulin clearance rate (ICR), the secretion and hepatic extraction of insulin are tightly related [1]. Hepatic ICR is reduced in individuals with obesity [3–5], central adiposity and hepatic steatosis [6–8]. By enabling a larger fraction of secreted insulin to reach systemic circulation, reduced ICR may serve as an adaptive mechanism to preserve normal glucose tolerance (NGT) in response to insulin resistance [9]. Alterations in ICR may, therefore, precede changes in glucose tolerance [5, 10].

ICR is a dynamic process that changes in response to feeding [9] and macronutrient manipulation [11] on a minute-by-minute basis [1]. Although humans spend most of the day in the fed state [12], few studies have examined the temporal change in ICR in response to meal ingestion. Studies using oral glucose loads have shown there is an immediate and pronounced suppression of ICR in people without type 2 diabetes, which is described as a

key determinant of glucose tolerance [3, 9]. However, the use of oral glucose solutions has inherent limitations and does not reflect the hormonal responses or gastrointestinal processes that occur following the consumption of a mixed-macronutrient meal [13, 14]. This requires consideration as ICR may be regulated by the rate of gastric emptying [15] and the secretion of incretin peptides [16] and glucagon [11, 17], both of which are affected by meal type [18] and composition [13, 19]. Whether the adaptive processes reported after the ingestion of glucose [3, 9] are observed after the consumption of a mixed-macronutrient meal remains unknown.

In the current study, we sought to provide insight into postprandial insulin kinetics following the ingestion of a standardised mixed-macronutrient breakfast meal in people with and without type 2 diabetes. We also aimed to identify potential metabolic or hormonal determinants of ICR.

Methods

Participants and study design This study is a secondary analysis of data derived from two randomised, controlled trials investigating the postprandial metabolic effects

Table 1 Participant anthropometric and metabolic characteristics

Characteristic	Lean-NGT (<i>n</i> =12)	Obese-NGT (<i>n</i> =11)	Type 2 diabetes (<i>n</i> =19)	ANOVA <i>p</i> value	Lean-NGT vs Obese-NGT	Lean-NGT vs type 2 diabetes	Obese-NGT vs type 2 diabetes
Women/men (<i>n/n</i>)	0/12	0/11	6/13	–	–	–	–
Age (years)	36±11	35±7	50±5	<0.001	0.879	<0.001	<0.001
BMI (kg/m ²)	23.7±1.8	33.7±2.4	32.7±5.7	<0.001	<0.001	<0.001	0.767
Waist (cm)	80.3±6.4	110.8±10.3	105.5±14.7	<0.001	<0.001	<0.001	0.460
Hip (cm)	95.4±4.6	115.2±10.7	107±14.0	<0.001	<0.001	0.021	0.140
WHR	0.84±0.04	0.97±0.07	0.99±0.06	<0.001	<0.001	<0.001	0.601
Fasting biochemistry							
HbA _{1c} (mmol/mol)	–	–	56.7±8.8	–	–	–	–
HbA _{1c} (%)	–	–	7.3±0.8	–	–	–	–
Glucose (mmol/l)	4.4±0.4	4.5±0.3	7.1±1.6	<0.001	0.958	<0.001	<0.001
Insulin (pmol/l)	52.5±20.1	107.6±51.5	86.7±54.1	0.017	0.013	0.133	0.370
ISR (pmol min ⁻¹ m ⁻²)	40.8 (37.2, 54.9)	80.6 (66.7, 112.7)	100.8 (85.8, 119.4)	<0.001 ^a	0.009	<0.001	0.938
ICR (l min ⁻¹ m ⁻²)	1.1±1.1	0.8±1.2	1.4±0.4	0.002	0.784	0.017	0.003
Glucagon (pmol/l)	7.0±3.3	12.3±4.6	15.8±6.4	<0.001	0.053	<0.001	0.190
GLP-1 (pmol/l)	17.7±4.9	26.3±12.7	34.1±13.8	<0.001	0.148	<0.001	0.162
GIP (pmol/l)	12.2±7.2	13.4±6.0	15.4±7.8	0.525	–	–	–
Triglycerides (mmol/l)	1.1 (1.0, 1.2)	1.6 (1.2, 2.0)	1.9 (1.0, 3.1)	0.017 ^a	0.056	0.025	1.00
NEFA (μmol/l)	508±199	558±196	731±275	0.030	0.868	0.037	0.142
GGT (U/l)	19.6 (13.6, 27.1)	40.3 (18.8, 55.2)	34.8 (27.4, 48.5)	0.010 ^a	0.046	0.013	1.00
AST (U/l)	25.2 (19.8, 34.5)	22.5 (20.6, 32.1)	24.0 (18.5, 33.4)	0.479	–	–	–
HOMA-IR	1.8 (1.1, 2.3)	3.3 (2.2, 4.5)	3.9 (2.1, 5.9)	0.002 ^a	0.015	0.002	1.00
Adipo-IR	26.2±13.5	61.8±43.4	60.8±38.2	0.010	0.025	0.015	0.986

Continuous data are presented as mean±SD or median (IQR), whereas categorical data are presented as *n*

Dashes represent data that are not available

All *p* values were adjusted for multiple comparisons

^aData were analysed by a Kruskal–Wallis *H* test

AST, aspartate aminotransferase; GGT, γ -glutamyltransferase

of protein supplementation in people with [11, 20] and without [21] type 2 diabetes (clinical trial registration no. ISRCTN17563146 and no. ISRCTN95281775). Exclusion criteria for participation included: age >65 years; history of gastrointestinal disease or medication use known to affect gastrointestinal function; BMI >40 kg/m²; breakfast skippers; or known dietary intolerances. In addition, participants with type 2 diabetes who were treated with exogenous insulin or glucagon-like peptide 1 (GLP-1) receptor agonists, who had an HbA_{1c} of >80 mmol/mol (9.5%) or who had a duration of type 2 diabetes of <1 year were ineligible. Both trials received ethical approval from the local National Health Service Research Ethics Committee (#18/NE/0372) and from the Research Ethics Committee of the Faculty of Medical Sciences, Newcastle University (#1512/4830/2018). All participants provided written informed consent prior to enrolment.

The study population included 19 individuals with type 2 diabetes (13 men and six women), and 12 lean (Lean-NGT)

and 11 centrally obese (Obese-NGT) normoglycaemic adult men (Table 1). In all participants, sex and gender were self-reported. All participants were of white European descent (self-reported) and had stable body mass (± 1 kg for >2 months). NGT was defined as fasting and postprandial blood glucose concentrations of <5.6 mmol/l and >7.8 mmol/l, respectively, and a return of blood glucose concentrations to preprandial concentrations within 120 min [22, 23]. Participants with type 2 diabetes (HbA_{1c} 56.7±8.8 mmol/mol [7.3±0.8%]) had a median (IQR) duration of diabetes of 4 years (3, 8 years) and were treated by diet and lifestyle modifications (*n*=2), sulfonylurea monotherapy (*n*=1), metformin monotherapy (*n*=5), the combination of metformin with a sulfonylurea (*n*=7) or sodium–glucose cotransporter 2 inhibitor (SGLT2i) (*n*=3), or the combination of metformin, sulfonylurea and a thiazolidinediones (*n*=1). All medications used by the type 2 diabetes group were kept consistent and unaltered throughout; this was to avoid any effects of a missed dose on postprandial glycaemic excursions, which

may also affect insulin kinetics [24]. The type 2 diabetes group is representative of the wider English type 2 diabetes population with respect to age, HbA_{1c} and anti-hyperglycaemic medication use [25]. Individuals in the Lean-NGT and Obese-NGT groups were not taking any medication known to affect glucose tolerance or insulin sensitivity.

Experimental protocol Participants arrived via pre-arranged transport to the research facilities after a ~12 h overnight fast. Once rested, participants consumed a breakfast meal consisting of cereal (Cheerios, Nestle, UK) and whole milk, providing 387 kcal from 58% carbohydrates, 27% fat and 15% protein. Venous blood samples were collected from a cannula placed in a forearm vein at baseline ($t=0$ min) and at $t=15, 30, 45, 60, 90, 120, 150, 180$ and 240 min after ingestion of the meal. Meals were consumed within 10 min, and participants remained seated throughout.

Blood sampling For the analysis of aspartate aminotransferase, alanine aminotransferase, γ -glutamyltransferase, C-peptide, insulin, NEFA and triglycerides, blood samples were collected in serum collection tubes. Plasma glucagon, total glucose-dependent insulinotropic polypeptide (GIP) and total GLP-1 were measured from blood samples collected in EDTA tubes containing a protease and dipeptidyl-peptidase IV inhibitor [11]. Collected blood samples were centrifuged at 2500 rev/min at 4°C for 10 min with the corresponding supernatant frozen at -80°C until analysis.

Analytical methods Blood glucose concentrations were measured from venous whole blood using the enzymatic-amprometric method (Biosen C_Line, EKF Diagnostics, UK). C-peptide, insulin, glucagon, GIP and GLP-1 were measured by ELISA, as described [11]. Liver enzymes, NEFA and triglycerides were measured by routine clinical chemistry using a benchtop clinical analyser (Daytona⁺, Randox Laboratories, UK). NEFA were measured for 60 min post meal to coincide with their suppression [12]. Due to several participants in the Lean-NGT group demonstrating alanine aminotransferase levels below the limit of detection (<12.8 U/l), these data are not included in the analyses.

Calculations AUC was calculated using the trapezoidal method and time-averaged for the duration of interest. Adipocyte insulin resistance (Adipo-IR) was estimated from the product of fasting NEFA and fasting insulin concentrations [9]. The suppression of NEFA (%) was taken as the change in NEFA concentrations from baseline to nadir values divided by baseline, and then multiplied by 100. HOMA-IR, an indirect measurement of insulin resistance (primarily hepatic), was determined from fasting insulin and glucose concentrations. Oral minimal models were used to compute measures of insulin sensitivity (S_i) and beta cell function

(Φ_{total}) using MATLAB (version R2022b, MathWorks, USA; www.uk.mathworks.com) and SAAM II software (version 2.3.3, Nanomath, USA; www.nanomath.us) [26].

Insulin secretion and clearance Pre-hepatic insulin secretion rates (ISRs) were calculated from the deconvolution of C-peptide concentrations using a two-compartmental model and population-derived metrics of C-peptide kinetics [27]. Basal ICR was calculated as fasting ISR divided by fasting insulin, whereas during the feeding test, postprandial ICR was calculated using a single-pool model [9]. As the clearance of insulin by peripheral tissues is constant over a wide physiological range of insulin concentrations [2], differences in ICR from this model primarily reflect the removal of insulin by the liver. Prior to modelling, plasma insulin concentrations were smoothed by cubic spline fitting, and the derivative of insulin over time was calculated from smoothed data using 5 min intervals. The rates of insulin extraction (Rd_{ins}) and ICR over a given time interval (t) were calculated using:

$$Rd_{\text{ins}}(t) = \text{ISR}(t) - [dI(t)/dt] \times V$$

$$\text{ICR}(t) = Rd_{\text{ins}}(t)/I(t)$$

Where I is insulin concentrations at time point (t), and V is the distribution volume for insulin (141 ml kg⁻¹ [28]). The average volume of plasma that was cleared of insulin per min (1 min⁻¹ m⁻²) during the feeding test was calculated as:

$$\text{ICR AUC}_{240} = (\text{AUC [ISR}(t)/I(t)]_{240 \text{ min}} - [\log_e(I_{240 \text{ min}}) - \log_e(I_{0 \text{ min}})] \times V)/240,$$

where $I_{0 \text{ min}}$ and $I_{240 \text{ min}}$ are insulin concentrations at time points $t=0$ min and $t=240$ min, respectively [9]. Total insulin extraction (pmol min⁻¹ m⁻²) was calculated as:

$$Rd_{\text{ins}} \text{ AUC}_{240} = (\text{ISR AUC}_{240} - [I_{240 \text{ min}} - I_{0 \text{ min}}] \times V)/240$$

Hepatic insulin delivery (pmol/min) was estimated as the sum of ISR and plasma insulin concentrations delivered to the liver via arterial circulation; the latter was assumed to be a product of hepatic plasma flow (0.576 l min⁻¹ m²) and plasma insulin concentrations [28].

Statistical analysis A posteriori sample size calculation was performed using previously published data in lean and obese individuals [3]. To detect a 25% difference in ICR AUC with a power of 0.8 and an α of 0.05, it was calculated that ten participants in each group were required.

Normality was assessed using the Shapiro–Wilk test, and non-normally distributed data underwent log₁₀ transformation. Differences between means were analysed using a one-way ANOVA for normally distributed data or a Kruskal–Wallis H test when the parametric assumption was not met. Corrections for multiple comparisons were performed by applying the Tukey post hoc comparison test for ANOVA or the Dunn–Bonferroni post hoc procedure for the




Fig. 1 Postprandial glucose (a), insulin (b), ISR (c), ICR (d), Rd_{ins} (e), hepatic insulin delivery (f), glucagon (g), GLP-1 (h) and NEFA (i) responses to the ingestion of a mixed-macronutrient breakfast meal in Lean-NGT (red circles, dashed line [$n=12$]) and Obese-NGT (blue circles, dotted line [$n=11$]) individuals, and in people with type 2 diabetes (grey circles, solid line [$n=19$]). The arrow depicts the time of meal ingestion. In (f), the purple area represents the maximum hepatic insulin uptake capacity (i.e. ~ 2000 pmol/min) [2]. In (g),

glucagon data in the Obese-NGT group are presented as $n=10$. All data are presented as mean \pm SEM and were analysed by a two-way, mixed-model ANOVA with Tukey post hoc comparisons. p values were adjusted for multiple comparisons: * $p<0.05$ Lean-NGT vs Obese-NGT; † $p<0.05$ Obese-NGT vs type 2 diabetes; ‡ $p<0.05$ Lean-NGT vs type 2 diabetes. For clarity, significant main effects for time (i.e. change from baseline) are not presented in the figure

Kruskal–Wallis H test. To detect differences in postprandial responses between populations, a two-way, mixed-model ANOVA with Tukey post hoc comparisons was performed. Univariate analyses were performed using Spearman's ρ test to detect correlations between ICR AUC₂₄₀ and metabolic and hormonal measures. A multivariate linear regression model was used to identify predictors of ICR AUC₂₄₀. Unless stated, all data in the text and in the tables are presented as mean \pm SD or as median (IQR) for excessively skewed data. All modelling was performed using MATLAB (version R2022b, MathWorks, USA; www.uk.mathworks.com) and inferential statistics were performed using SPSS Statistics (version 28; IBM, USA; www.ibm.com/spss). Significance was accepted as $p<0.05$, and adjusted p values for multiple comparisons are presented.

Results

Participant characteristics Participant characteristics are shown in Table 1. Individuals in the type 2 diabetes group were ~ 14 years older than the Lean-NGT and Obese-NGT groups ($p<0.001$), but had similar BMI, waist-to-hip ratio and liver enzymes as the Obese-NGT group. Markers of insulin resistance (Adipo-IR and HOMA-IR) were similar between the Obese-NGT and type 2 diabetes groups but were higher compared with Lean-NGT (Table 1).

Fasting metabolic parameters Fasting blood glucose concentrations were similar between the respective NGT groups but were elevated in those with type 2 diabetes ($p<0.001$). Fasting plasma insulin concentrations were greater in Obese-NGT vs

Table 2 Postprandial metabolic responses to the ingestion of a standardised mixed-macronutrient breakfast meal in Lean-NGT and Obese-NGT volunteers and people with type 2 diabetes

Variable	Lean-NGT (n=12)	Obese-NGT (n=11)	Type 2 diabetes (n=19)	ANOVA p value	Lean-NGT vs Obese-NGT	Lean-NGT vs type 2 diabetes	Obese-NGT vs type 2 diabetes
Glucose							
Peak (mmol/l)	5.8±0.9	6.2±0.7	11.8±3.3	<0.001	0.932	<0.001	<0.001
AUC ₂₄₀ (mmol/l)	4.4±0.4	4.9±0.4	10.0±2.4	<0.001	0.718	<0.001	<0.001
Insulin							
Peak (pmol/l)	230.7±100.1	531.5±271.2	419.9±246.7	0.008	0.007	0.068	0.393
AUC ₆₀ (pmol/l)	100.2±63.5	270.8±158.5	201.2±128	0.003	0.002	0.031	0.339
AUC ₂₄₀ (pmol/l)	69.6±22.8	181.6±97.7	268.2±157.3	<0.001	0.002	<0.001	0.496
ISR							
Peak (pmol min ⁻¹ m ⁻²)	294.8 (179.0, 346.0)	396.3 (328.7, 4749)	339.5 (301.9, 395)	0.029 ^a	0.024	0.359	0.472
AUC ₆₀ (pmol min ⁻¹ m ⁻²)	216.5±60.4	342.0±82.8	305.7±122.5	0.011	0.011	0.048	0.598
AUC ₂₄₀ (pmol min ⁻¹ m ⁻²)	100.7±20.6	170.0±42.5	266.4±98.4	<0.001	0.062	<0.001	0.003
GLP-1							
Peak (pmol/l)	26.0 (24.2, 44.8)	38.1 (35.9, 46.6)	47.7 (38.5, 56.6)	0.005 ^a	0.600	0.024	0.248
AUC ₂₄₀ (pmol/l)	25.3±6.8	33.3±6.8	41.5±14.3	<0.001	0.071	<0.001	0.216
Glucagon^b							
Peak (pmol/l)	7.9 (7.0, 9.8)	15.2 (11.5, 21.7)	26.6 (21.5, 38.2)	<0.001 ^a	0.081	<0.001	0.235
Nadir (pmol/l)	3.6 (1.8, 4.6)	8.2 (4.7, 9.0)	10.0 (4.1, 13.4)	0.005 ^a	0.121	0.003	1.00
AUC ₂₄₀ (pmol/l)	7.1±4.5	11.0±4.2	16.9±7.4	<0.001	0.053	<0.001	0.097
NEFA							
Nadir (µmol/l)	125.2±40.4	254.1±132.1	382.7±160.9	<0.001	0.007	<0.001	0.026
AUC ₆₀ (µmol/l)	245.2±80.9	338.5±141.0	561.3±152.4	<0.001	0.036	<0.001	0.004
Suppression (%)	-75.1 (-77.3, -69.9)	-51.4 (-69.3, -45.0)	-49.1 (-66.4, -42.8)	<0.001 ^a	0.012	<0.001	0.999
Modelling variables							
ICR AUC ₂₄₀ (l min ⁻¹ m ⁻²)	1.9±0.5	1.3±0.4	1.4±0.7	0.014	0.034	0.021	0.989
R _d _{ins} AUC ₂₄₀ (pmol min ⁻¹ m ⁻²)	91.9±18.4	154.8±38.4	242.0±89.0	<0.001	<0.001	0.012	0.004
Hepatic insulin delivery AUC ₂₄₀ min (pmol/min)	220.0±45.8	506.8±141.5	713.2±316.5	<0.001	0.012	<0.001	0.055
log ₁₀ S _i	2.5 (2.1, 2.6)	1.4 (1.2, 1.5)	0.6 (0.4, 0.8)	<0.001	<0.001	<0.001	0.177
log ₁₀ ϕ _{total}	2.0 (1.8, 2.3)	2.0 (1.9, 2.1)	1.4 (1.2, 1.5)	<0.001	0.849	<0.001	<0.001

Data are presented as mean±SD or median (IQR)

All *p* values were adjusted for multiple comparisons

^aData were analysed by a Kruskal–Wallis *H* test

^bPostprandial glucagon data in the Obese-NGT group were available on *n*=10

Lean-NGT (*p*=0.011), which was reflected by the increase in basal ISR (*p*=0.009) since fasting ICR values were similar. Compared with Lean-NGT, basal ISR was greater in the type 2 diabetes group (*p*<0.001), although fasting plasma insulin concentrations were similar, with the latter caused by the increase in fasting ICR in the type 2 diabetes group vs the

Lean-NGT group (*p*=0.017) (Table 1). Fasting glucagon concentrations, which are consistent with the presence of hepatic steatosis [29], were increased in the type 2 diabetes group (*p*<0.001) and tended to be greater in Obese-NGT (*p*=0.053) compared with Lean-NGT, with no difference between Obese-NGT vs the type 2 diabetes group (Table 1).

Postprandial blood glucose and insulin kinetics Postprandial glycaemic excursions (AUC_{240}) were comparable in Lean-NGT and Obese-NGT but were markedly greater in the type 2 diabetes group (Fig. 1a, Table 2). The change in blood glucose from fasting to peak concentrations was similar in Lean-NGT (1.5 ± 0.9 mmol/l) and Obese-NGT groups (1.7 ± 0.7 mmol/l), but greater in the type 2 diabetes group (4.6 ± 2.1 mmol/l; $p < 0.0001$ vs NGT). Both plasma insulin concentrations (Fig. 1b) and ISR (Fig. 1c) were elevated post meal in the type 2 diabetes and Obese-NGT groups, compared with Lean-NGT. While ISR and plasma insulin returned to baseline values within ~ 90 min post meal in the NGT groups, plasma insulin and ISR excursions were prolonged in the type 2 diabetes group. Compared with Lean-NGT, insulin AUC_{240} was approximately threefold greater in the Obese-NGT and type 2 diabetes groups ($p < 0.01$), with no difference between the last two (Table 2). Conversely, overall ISR AUC_{240} was greater in type 2 diabetes compared with both Lean-NGT ($p < 0.001$) and Obese-NGT ($p = 0.003$), but was not statistically different between Obese-NGT vs Lean-NGT ($p = 0.062$).

As shown in Fig. 1d, the postprandial temporal responses in ICR differed between the groups. In Lean-NGT individuals, there was a distinct and immediate increase in ICR from pre-meal values (all $p < 0.05$), whereas postprandial ICR remained at baseline values in Obese-NGT. In contrast, there was a modest suppression in ICR at $t = 30$ min and $t = 45$ min post meal in the type 2 diabetes group ($p < 0.03$). Accordingly, ICR was reduced at several points during the postprandial period in the Obese-NGT and type 2 diabetes groups, compared with Lean-NGT (all $p < 0.018$; Fig. 1d). Overall, the mean ICR (ICR AUC_{240}) was ~ 35 – 45% lower in the Obese-NGT and type 2 diabetes groups compared with the Lean-NGT group ($p = 0.034$ and $p = 0.021$, respectively), with no difference between Obese-NGT and type 2 diabetes groups (Table 2). Compared with Lean-NGT, insulin extraction (Rd_{ins} AUC_{240}), defined as the average amount of secreted insulin removed per min, was $\sim 40\%$ and $\sim 62\%$ greater in the Obese-NGT and type 2 diabetes groups, respectively (Fig. 1e, Table 2). The increase in Rd_{ins} in the Obese-NGT and type 2 diabetes groups was attributable to the greater hepatic delivery of insulin (i.e. during the first and second pass [30]) compared with Lean-NGT (Fig. 1f, Table 2).

Plasma glucagon and incretin responses After the meal, glucagon increased in the type 2 diabetes group ($t = 15$ min to $t = 45$ min; all $p < 0.001$), whereas there was no change from fasting values in the NGT groups (Fig. 1g). Compared with Lean-NGT, glucagon AUC_{240} was elevated in type 2 diabetes ($p < 0.001$) and appeared greater in Obese-NGT ($\sim 55\%$; $p = 0.053$). Glucagon AUC_{240} was similar in the type 2 diabetes and Obese-NGT groups. Similar temporal responses

were seen for GLP-1 (Fig. 1h). Accordingly, GLP-1 AUC_{240} was $\sim 63\%$ greater in type 2 diabetes compared with Lean-NGT ($p < 0.001$) but was similar between the type 2 diabetes and Obese-NGT groups (Table 2). There was a tendency for GLP-1 AUC_{240} to be elevated in the Obese-NGT group compared with Lean-NGT ($\sim 31\%$; $p = 0.071$). GIP was similar between groups (data not shown).

Postprandial NEFA suppression In the Lean-NGT group, there was an immediate and significant postprandial decline in NEFA concentrations from fasting values (all $p < 0.01$), whereas the post-meal decline in NEFA was delayed in Obese-NGT and type 2 diabetes groups, with statistical reductions from baseline values only being observed from $t = 45$ min and $t = 60$ min (all $p < 0.01$; Fig. 1i). Overall, the absolute suppression of NEFA (%) was $\sim 30\%$ greater in Lean-NGT compared with both Obese-NGT and type 2 diabetes groups ($p < 0.012$), with no difference between the Obese-NGT and type 2 diabetes groups (Table 2).

Insulin sensitivity and beta cell function There was a stepwise decline in insulin sensitivity (S_i) from Lean-NGT to Obese-NGT to type 2 diabetes ($p < 0.0001$). Similar patterns were observed when expressing insulin sensitivity using the oral glucose insulin sensitivity index, whereas insulin sensitivity measured by the Matsuda index was similar between Obese-NGT and type 2 diabetes groups (electronic supplementary material [ESM] Table 1). ϕ_{total} was reduced in the type 2 diabetes group compared with the NGT group ($p < 0.001$), consistent with beta cell dysfunction, whereas ϕ_{total} was similar in the Obese-NGT and Lean-NGT groups (Table 2).

Metabolic associations and predictors of ICR ICR AUC_{240} was inversely associated with HOMA-IR ($r = -0.826$; $p < 0.001$), Adipo-IR ($r = -0.681$; $p < 0.001$), glucagon AUC_{240} ($r = -0.670$; $p < 0.001$ [$n = 41$]) and GLP-1 AUC_{240} ($r = -0.623$; $p < 0.001$), as well as waist-to-hip ratio ($r = -0.525$; $p < 0.001$). There was also a positive correlation between ICR and S_i ($r = 0.619$; $p < 0.001$).

Due to one Obese-NGT participant not having available postprandial glucagon data, a multivariate linear regression model was performed on $n = 41$ (Table 3). The model demonstrated that \log_{10} HOMA-IR and glucagon AUC_{240} were the strongest predictors of \log_{10} ICR AUC_{240} in both the unadjusted model (adjusted $R^2 = 0.636$; $p < 0.001$) and the Adipo-IR-, BMI-, age- and S_i -adjusted model (adjusted $R^2 = 0.670$; $p < 0.001$). Confirming this, we separated the whole group into high or low based on the group median for HOMA-IR (2.7 arbitrary units [AU]) or glucagon AUC_{240} (10.7 pmol/l). ICR $AUC_{240 \text{ min}}$ was reduced by 40% (~ -0.75 $l \text{ min}^{-1} \text{ m}^{-2}$) in the ‘high’ groups compared with the ‘low’ groups (all $p < 0.001$; Fig. 2). As elevations in plasma glucagon concentrations are implicated with hepatic steatosis

Table 3 Independent predictors of ICR from a multivariate linear regression model

Variable	log ₁₀ ICR AUC ₂₄₀		SE	β	F	R ²	Adjusted R ²	p value
	B	95% CI B						
		Lower	Upper					
Model 1					24.424	0.562	0.539	<0.001
Constant	0.305**	0.238	0.372	0.033				
log ₁₀ HOMA-IR	-0.375**	-0.492	-0.259	0.057	-0.722			
S _i	<0.000	0.000	0.000	0.000	0.095			
Model 2					18.451	0.672	0.636*	<0.001
Constant	0.393**	0.311	0.474	0.040				
log ₁₀ HOMA-IR	-0.196*	-0.360	-0.033	0.081	-0.377			
S _i	<0.000	0.000	0.000	0.000	-0.083			
Adipo-IR	<0.000	0.000	0.000	0.000	-0.324			
Glucagon AUC ₂₄₀	-0.007*	-0.012	-0.001	0.003	-0.300			
Model 3					14.544	0.720	0.670	<0.001
Constant	0.227	-0.453	0.775	0.302				
log ₁₀ HOMA-IR	-0.255*	-0.429	-0.081	0.086	-0.491			
S _i	<0.000	0.000	0.000	0.000	-0.006			
Adipo-IR	<0.000	0.000	0.000	0.000	-0.243			
Glucagon AUC ₂₄₀	-0.008*	-0.013	-0.003	0.003	-0.362			
Age	0.004*	0.001	0.007	0.002	0.251			
BMI	0.001	-0.008	0.009	0.004	0.028			

Data were analysed by multivariate regression model on $n=41$

* $p<0.05$

** $p<0.01$




Fig. 2 Differences in ICR AUC₂₄₀ in individuals classified as high or low HOMA-IR or glucagon AUC₂₄₀. High (grey bars) and low (white bars) groups were defined as those above or below the group median for HOMA-IR (2.7 AU) or glucagon AUC₂₄₀ (10.7 pmol/l), respectively. Data are presented as median and IQR. *** $p<0.001$ between low and high groups

[29], independent of insulin resistance, these results suggest that reduced ICR is a consequence of intra-hepatic lipid accumulation rather than insulin resistance per se.

Discussion

The temporal suppression of ICR in response to an oral glucose load has been described as a key determinant of glucose tolerance [9]. However, the metabolic handling of a mixed-macronutrient meal differs compared with the consumption of glucose alone [13]. In the present study, we examined postprandial ICR responses after the ingestion of a standardised mixed-macronutrient breakfast meal in Lean-NGT and Obese-NGT individuals, and in people with type 2 diabetes. Our main findings identify a unique temporal variation in ICR that differs between Lean-NGT and Obese-NGT individuals. Specifically, in response to the meal, we observed an immediate increase in ICR in the Lean-NGT group, whereas ICR remained at baseline values in the Obese-NGT group. Although there were differing ICR responses in the NGT groups, overall postprandial glycaemic excursions were similar. Furthermore, we show that overall variances in postprandial ICR are determined by HOMA-IR and elevated plasma glucagon concentrations. Taken together, these findings demonstrate the adaptive responses of ICR to regulate postprandial glycaemia in the presence of insulin resistance.

While our data are in general agreement that ICR is reduced in obese, insulin-resistant individuals [3, 4, 8, 9, 31], this was not reflected by a greater post-meal suppression in ICR in the Obese-NGT group. These findings contrast with what has been reported following an oral glucose load [3, 9]; however, direct comparisons between studies are challenging due to differences in the test meals provided. Indeed, variances in nutrient absorption kinetics [15] and the subsequent islet [17] and incretin [16] responses between meals offer a plausible explanation for the reported variances in temporal ICR. The possible influence of hormonal responses on ICR is suggested by the current data, as shown by the strong inverse associations of GLP-1 AUC₂₄₀ ($r=-0.623$) and glucagon AUC₂₄₀ ($r=-0.670$) with ICR AUC₂₄₀. However, our reported post-meal suppression of ICR in the type 2 diabetes group is akin to what is seen after the ingestion of glucose [3, 9]. As our calculated ISRs were twofold greater than what is reported after glucose ingestion [9], this may reflect the inability of the diabetic liver to extract insulin [32], independent of allostatic load (i.e. nutrient or hormonal stimuli). Nevertheless, as we did not perform an oral glucose feeding test, future studies examining the interactions between nutrient composition and postprandial insulin kinetics are warranted.

In our study, we observed a synchronous increase in ISR and ICR in the Lean-NGT group, such that the clearance of insulin overshadowed its secretion, resulting in reduced overall plasma insulin concentrations. Despite this response, overall postprandial glycaemic excursions were minor in Lean-NGT, implying that a proficient insulinaemic response to maintain NGT was achieved. This is consistent with the notion that the balance between insulin secretion and clearance prevents insulin excess [33] or insufficiency [9] and their adverse metabolic consequences. Previous work has demonstrated that during periods of low blood glucose, a temporal increase in ICR negates the increase in postprandial ISR as a possible means to prevent hypoglycaemia [34]. In this regard, there is evidence to support a sensory function of the portal vein that alters the hepatic arteriovenous insulin gradient and hepatic glucose metabolism in response to nutrients and/or hormones [30, 35, 36]. Such a feedback loop may be present here given our observation of a complete post-meal suppression of glucagon in Lean-NGT individuals, which combined with the increase in ISR is suggestive that hepatic glucose production was suppressed [37]. This also offers an explanation for the observed temporal responses of ICR in the Obese-NGT group, for whom glucagon was less suppressed and overall glucagon excursions greater, yet NGT was well maintained. It is, therefore, reasonable to infer that during periods of altered portal nutrient/hormonal flux, the temporal responses in ICR reflect an adaptive mechanism to protect not only against hyperglycaemia [9], but also hypoglycaemia [33].

In the present study, the Obese-NGT and type 2 diabetes groups showed an impaired suppression of NEFA, which is proposed to impair ICR [38]. Mechanistically, increased delivery of NEFA to the liver blunts hepatic insulin signalling and activates peroxisome proliferator-activated receptor α (PPAR- α) [39], which reduces the transcription of carcinoembryonic antigen-related cell adhesion molecule 1 (CEACAM1) [39–41]. CEACAM1 facilitates the internalisation of receptor-bound insulin for endosomal degradation and its specific liver inactivation impairs ICR [40]. Our data suggest that the increased secretion of glucagon may impair ICR, independent of hepatic NEFA delivery. As elevations in plasma glucagon are a metabolic disturbance of hepatic steatosis [29], independent of insulin resistance or glucose tolerance [42], these observations likely describe reduced ICR as a consequence of hepatic steatosis [8]. In support, the infusion of exogenous glucagon was recently shown to directly impair ICR in healthy adults [17]. Indeed, increased hepatic glucagon signalling augments PPAR- α activity [43], which reduces the expression of CEACAM1 [39] and impairs ICR [40]. Of note, glucagon-stimulated PPAR- α activation is augmented further by the presence of fatty acids [44]. These observations may be of interest for the pathophysiology of hepatic steatosis that is described by increased plasma glucagon concentrations [29] and increased hepatic lipid flux [45]; these ‘dual-hit’ signals may offer an explanation as to why CEACAM1 expression [46] and ICR [8, 31] are reduced in people with increased liver fat.

We observed an inverse association between GLP-1 AUC₂₄₀ and ICR AUC₂₄₀, and the substitution of glucagon AUC₂₄₀ with GLP-1 AUC₂₄₀ in our multivariate analyses did not change our findings (see ESM Results; ESM Table 2). As our measurement of plasma glucagon is specific for pancreatic-derived glucagon, this observation is not due to cross-reactivity with the other proglucagon-derived peptide, GLP-1, or ‘gut-glucagon’. Previously, the blockage of the endogenous GLP-1 receptor with exendin(9–39)NH₂ was also reported to increase ICR [16]. As there are no known GLP-1 receptors in the liver, and exendin-4 shares ~50% amino acid homology with glucagon, this may reflect reduced signalling via the hepatic glucagon receptor.

It has been suggested that the postprandial ICR profile in those with type 2 diabetes is due to the reduced hepatic delivery of insulin (i.e. beta cell dysfunction), rather than an impairment in the liver to adjust the clearance of insulin [3]. In contrast, our data suggest that the impaired suppression of ICR in individuals with type 2 diabetes reflects a maladaptive hepatic response and not a primary beta cell defect. For instance, overall ISR and hepatic insulin delivery were increased in the type 2 diabetes group, but this was not associated with marked deviations in ICR compared with the Obese-NGT group. Thus, although there was a temporal increase in insulin extraction in those with type 2 diabetes, this should not be interpreted that the

hepatic handling of insulin is intact or functional. It must be recognised, however, that type 2 diabetes is a heterogeneous condition with varying degrees of residual beta cell function. While our data suggest that the impaired suppression of ICR is a primary hepatic defect, this may not be the case for people with more advanced type 2 diabetes and reduced insulin secretory capacity. Moreover, although our reported postprandial ICR profile in the type 2 diabetes group is akin to previous findings in people with type 2 diabetes after drug withdrawal [9], as all anti-hyperglycaemic medications were kept unaltered in our study, we cannot exclude the potential effects of concomitant medication use on postprandial insulin kinetics which may cloud our findings.

We acknowledge that there are further limitations associated with our study which include the secondary, cross-sectional nature of the analyses, the small sample size and the age imbalance between the type 2 diabetes and NGT groups. Furthermore, sex distributions of participants between groups were unmatched in our study. Previous analyses have shown that, although there is an overall reduction in ICR in female individuals compared with male individuals [47], the temporal response in ICR following glucose ingestion between insulin-resistant and insulin-sensitive individuals is qualitatively unaffected by biological sex [3], suggesting that our findings may be generalisable across sexes. Nonetheless, as our sample size did not allow for subgroup analyses stratified by biological sex, future studies examining potential differences in postprandial insulin kinetics in phenotypically matched (age, insulin sensitivity and glucose tolerance) female and male individuals are warranted. As all participants were of white European descent, our data also do not reflect the potential ethnic differences in postprandial ICR [28]. In our study, we did not directly quantify hepatic ICR; however, the use of indirect approaches that incorporate insulin and C-peptide kinetic modelling offer an alternative to calculate ICR in vivo during dynamic tests [48]. Moreover, direct measurements of intra-hepatic fat content and insulin sensitivity were not available in our analysis. Although our findings describe a potential link between increased plasma glucagon concentrations and reduced ICR, we were unable to establish causality. As elevations in plasma glucagon are a reported consequence of intra-hepatic fat deposition [29, 42], future prospective studies examining the role of glucagon on ICR in well described cohorts of people with and without hepatic steatosis are warranted.

Previous studies have assessed postprandial differences in ICR in obese and insulin-resistant populations after ingestion of glucose [4, 13] or mixed-macronutrient solutions [13]. Their findings, however, are limited by the reported static indices of ICR (i.e. the ratio of C-peptide/insulin or ISR/insulin AUCs). As demonstrated here, the use of these estimates alone has significant interpretative

limitations for the study of postprandial hyperinsulinaemia and masks the dynamic changes in insulin extraction that occur post meal. As failure to appropriately regulate the liver–insulin gate is associated with declining blood glucose control [9, 47, 49], the appropriate assessment of ICR after feeding has clear clinical significance.

To our knowledge, this is the first study to investigate the temporal clearance of insulin after a standardised mixed-macronutrient meal. By using a time-responsive model of ICR, our data provide new understanding into the hepatic extraction of insulin during physiological, non-steady-state conditions expected to occur in free-living. Altogether, we show that after ingestion of a mixed-macronutrient meal, ISR and ICR coordinate to produce a plasma insulin profile to maintain euglycaemia in normoglycaemic individuals with differing temporal profiles. In Lean-NGT individuals, there is a time course increase in ICR, thus constraining plasma insulin concentrations, and possibly in response to the complete postprandial suppression in glucagon, thereby minimising the risk of hypoglycaemia. In contrast, the adaptive glycaemic processes that involve ICR and ISR are dysregulated in type 2 diabetes. Our data stress the importance of taking into consideration differences in the secretion and clearance of insulin between groups when investigating the aetiology of postprandial hyperinsulinaemia in obese and type 2 diabetes populations.

Supplementary Information The online version of this article (<https://doi.org/10.1007/s00125-024-06249-7>) contains peer-reviewed but unedited supplementary material.

Acknowledgements The authors thank the study participants for their time, effort and commitment, and the research teams at the Newcastle National Institute for Health Research Clinical Research Facility, Newcastle upon Tyne, for their assistance with data collection.

Data availability The raw data supporting the conclusions of this article will be made available upon reasonable request.

Funding This work was supported by a grant awarded to DJW and EJS (grant number: BH172513) and an equipment award to KS from Arla Foods Ingredients group P/S (Viby J, Denmark). Arla Foods Ingredients group P/S had no role in the collection, analysis, or interpretation of data. Funding was supported by a Philanthropic Award to DJW from the Francis James Bell Endowment Fund, County Durham Community Foundation, UK.

Authors' relationships and activities The authors declare that there are no relationships or activities that might bias, or be perceived to bias, their work.

Contribution statement KS designed and conducted the research, analysed the data and wrote the manuscript. SP performed oral minimal modelling. GST, WP, MW, SP, NA-P, KABD, FK, LH, EJS and DJW reviewed and edited the manuscript and contributed to interpretation of the data. All authors provided final approval of the final manuscript. KS, EJS and DJW acquired funding. DJW is the guarantor of this work and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Meier JJ, Veldhuis JD, Butler PC (2005) Pulsatile insulin secretion dictates systemic insulin delivery by regulating hepatic insulin extraction in humans. *Diabetes* 54(6):1649–1656. <https://doi.org/10.2337/diabetes.54.6.1649>
- Ferrannini E, Wahren J, Faber OK, Felig P, Binder C, DeFronzo RA (1983) Splanchnic and renal metabolism of insulin in human subjects: a dose-response study. *Am J Physiol* 244(6):E517–E527. <https://doi.org/10.1152/ajpendo.1983.244.6.E517>
- Mittendorfer B, Patterson BW, Smith GI, Yoshino M, Klein S (2022) β Cell function and plasma insulin clearance in people with obesity and different glycemic status. *J Clin Invest* 132(3):e154068. <https://doi.org/10.1172/JCI154068>
- Fu Z, Wu Q, Guo W et al (2022) Impaired insulin clearance as the initial regulator of obesity-associated hyperinsulinemia: novel insight into the underlying mechanism based on serum bile acid profiles. *Diabetes Care* 45(2):425–435. <https://doi.org/10.2337/dc21-1023>
- Meistas MT, Margolis S, Kowarski AA (1983) Hyperinsulinemia of obesity is due to decreased clearance of insulin. *Am J Physiol* 245(2):E155–E159. <https://doi.org/10.1152/ajpendo.1983.245.2.E155>
- Smith GI, Polidori DC, Yoshino M et al (2020) Influence of adiposity, insulin resistance, and intrahepatic triglyceride content on insulin kinetics. *J Clin Invest* 130(6):3305–3314. <https://doi.org/10.1172/JCI136756>
- Gastaldelli A, Cusi K, Pettiti M et al (2007) Relationship between hepatic/visceral fat and hepatic insulin resistance in nondiabetic and type 2 diabetic subjects. *Gastroenterology* 133(2):496–506. <https://doi.org/10.1053/j.gastro.2007.04.068>
- Zaharia OP, Antoniou S, Bobrov P et al (2023) Reduced insulin clearance differently relates to increased liver lipid content and worse glycemic control in recent-onset type 2 and type 1 diabetes. *Diabetes Care* 46(12):2232–2239. <https://doi.org/10.2337/dc23-1267>
- Gastaldelli A, Abdul Ghani M, DeFronzo RA (2021) Adaptation of insulin clearance to metabolic demand is a key determinant of glucose tolerance. *Diabetes* 70(2):377–385. <https://doi.org/10.2337/db19-1152>
- Galderisi A, Polidori D, Weiss R et al (2019) Lower insulin clearance parallels a reduced insulin sensitivity in obese youths and is associated with a decline in beta-cell function over time. *Diabetes* 68(11):2074–2084. <https://doi.org/10.2337/db19-0120>
- Smith K, Taylor GS, Walker M et al (2023) Pre-meal whey protein alters postprandial insulinemia by enhancing β -cell function and reducing insulin clearance in T2D. *J Clin Endocrinol Metab* 108(8):e603–e612. <https://doi.org/10.1210/clinem/dgaa069>
- Ruge T, Hodson L, Cheeseman J et al (2009) Fasted to fed trafficking of fatty acids in human adipose tissue reveals a novel regulatory step for enhanced fat storage. *J Clin Endocrinol Metab* 94(5):1781–1788. <https://doi.org/10.1210/jc.2008-2090>
- Alsalam W, Tura A, Pacini G et al (2016) Mixed meal ingestion diminishes glucose excursion in comparison with glucose ingestion via several adaptive mechanisms in people with and without type 2 diabetes. *Diabetes Obes Metab* 18(1):24–33. <https://doi.org/10.1111/dom.12570>
- Carr RD, Larsen MO, Jelic K et al (2010) Secretion and dipeptidyl peptidase-4-mediated metabolism of incretin hormones after a mixed meal or glucose ingestion in obese compared to lean, nondiabetic men. *J Clin Endocrinol Metab* 95(2):872–878. <https://doi.org/10.1210/jc.2009-2054>
- Marathe CS, Rayner CK, Jones KL, Horowitz M (2018) Impact of variations in duodenal glucose load on insulin clearance in health and type 2 diabetes. *Acta Diabetol* 55(2):205–207. <https://doi.org/10.1007/s00592-017-1073-z>
- Shah A, Holter MM, Rimawi F et al (2019) Insulin clearance after oral and intravenous glucose following gastric bypass and gastric banding weight loss. *Diabetes Care* 42(2):311–317. <https://doi.org/10.2337/dc18-1036>
- Gray SM, Goonatilake E, Emrick MA et al (2023) High doses of exogenous glucagon stimulate insulin secretion and reduce insulin clearance in healthy humans. *Diabetes* 73(3):412–425. <https://doi.org/10.2337/db23-0201>
- Schusdziarra V, Dangel G, Klier M, Henrichs I, Pfeiffer EF (1981) Effect of solid and liquid carbohydrates upon postprandial pancreatic endocrine function. *J Clin Endocrinol Metab* 53(1):16–20. <https://doi.org/10.1210/jcem-53-1-16>
- Jacobsen SH, Olesen SC, Dirksen C et al (2012) Changes in gastrointestinal hormone responses, insulin sensitivity, and β -cell function within 2 weeks after gastric bypass in non-diabetic subjects. *Obes Surg* 22(7):1084–1096. <https://doi.org/10.1007/s11695-012-0621-4>
- Smith K, Taylor GS, Brunsgaard LH et al (2022) Thrice daily consumption of a novel, premeal shot containing a low dose of whey protein increases time in euglycemia during 7 days of free-living in individuals with type 2 diabetes. *BMJ Open Diabetes Res Care* 10(3):e002820. <https://doi.org/10.1136/bmjdr-2022-002820>
- Smith K, Taylor GS, Allerton DM et al (2021) The postprandial glycaemic and hormonal responses following the ingestion of a novel, ready-to-drink shot containing a low dose of whey protein in centrally obese and lean adult males: a randomised controlled trial. *Front Endocrinol (Lausanne)* 12:696977. <https://doi.org/10.3389/fendo.2021.696977>
- Wolever TM, Chiasson JL, Csima A et al (1998) Variation of postprandial plasma glucose, palatability, and symptoms associated with a standardized mixed test meal versus 75 g oral glucose. *Diabetes Care* 21(3):336–340. <https://doi.org/10.2337/diacare.21.3.336>
- Shankar SS, Vella A, Raymond RH et al (2016) Standardized mixed-meal tolerance and arginine stimulation tests provide reproducible and complementary measures of beta-cell function: results from the foundation for the national institutes of health biomarkers consortium investigative series. *Diabetes Care* 39(9):1602–1613. <https://doi.org/10.2337/dc15-0931>
- Merovci A, Tripathy D, Chen X et al (2021) Effect of mild physiologic hyperglycemia on insulin secretion, insulin clearance, and insulin sensitivity in healthy glucose-tolerant subjects. *Diabetes* 70(1):204–213. <https://doi.org/10.2337/db20-0039>
- Dennis JM, Henley WE, McGovern AP et al (2019) Time trends in prescribing of type 2 diabetes drugs, glycaemic response and risk factors: a retrospective analysis of primary care data, 2010–2017. *Diabetes Obes Metab* 21(7):1576–1584. <https://doi.org/10.1111/dom.13687>
- Breda E, Cavaghan MK, Toffolo G, Polonsky KS, Cobelli C (2001) Oral glucose tolerance test minimal model indexes of

- β -cell function and insulin sensitivity. *Diabetes* 50(1):150–158. <https://doi.org/10.2337/diabetes.50.1.150>
27. Van Cauter E, Mestrez F, Sturis J, Polonsky KS (1992) Estimation of insulin secretion rates from C-peptide levels. Comparison of individual and standard kinetic parameters for C-peptide clearance. *Diabetes* 41(3):368–377. <https://doi.org/10.2337/diab.41.3.368>
 28. Polidori DC, Bergman RN, Chung ST, Sumner AE (2016) Hepatic and extrahepatic insulin clearance are differentially regulated: results from a novel model-based analysis of intravenous glucose tolerance data. *Diabetes* 65(6):1556–1564. <https://doi.org/10.2337/db15-1373>
 29. WewerAlbrechtsen NJ, Faerch K, Jensen TM et al (2018) Evidence of a liver-alpha cell axis in humans: hepatic insulin resistance attenuates relationship between fasting plasma glucagon and glucagonotropic amino acids. *Diabetologia* 61(3):671–680. <https://doi.org/10.1007/s00125-017-4535-5>
 30. Edgerton DS, Scott M, Farmer B et al (2019) Targeting insulin to the liver corrects defects in glucose metabolism caused by peripheral insulin delivery. *JCI Insight* 5(7):e126974. <https://doi.org/10.1172/jci.insight.126974>
 31. Kotronen A, Juurinen L, Hakkarainen A et al (2008) Liver fat is increased in type 2 diabetic patients and underestimated by serum alanine aminotransferase compared with equally obese nondiabetic subjects. *Diabetes Care* 31(1):165–169. <https://doi.org/10.2337/dc07-1463>
 32. Bonora E, Zavaroni I, Coscelli C, Butturini U (1983) Decreased hepatic insulin extraction in subjects with mild glucose intolerance. *Metabolism* 32(5):438–446. [https://doi.org/10.1016/0026-0495\(83\)90004-5](https://doi.org/10.1016/0026-0495(83)90004-5)
 33. Salehi M, Gastaldelli A, D'Alessio DA (2014) Altered islet function and insulin clearance cause hyperinsulinemia in gastric bypass patients with symptoms of postprandial hypoglycemia. *J Clin Endocrinol Metab* 99(6):2008–2017. <https://doi.org/10.1210/jc.2013-2686>
 34. Salehi M, DeFronzo R, Gastaldelli A (2022) Altered insulin clearance after gastric bypass and sleeve gastrectomy in the fasting and prandial conditions. *Int J Mol Sci* 23(14):7667. <https://doi.org/10.3390/ijms23147667>
 35. Moore MC, Flakoll PJ, Hsieh PS et al (1998) Hepatic glucose disposition during concomitant portal glucose and amino acid infusions in the dog. *Am J Physiol* 274(5):E893–902. <https://doi.org/10.1152/ajpendo.1998.274.5.E893>
 36. Lindgren O, Pacini G, Tura A, Holst JJ, Deacon CF, Ahrén B (2015) Incretin effect after oral amino acid ingestion in humans. *J Clin Endocrinol Metab* 100(3):1172–1176. <https://doi.org/10.1210/jc.2014-3865>
 37. Camastra S, Muscelli E, Gastaldelli A et al (2013) Long-term effects of bariatric surgery on meal disposal and β -cell function in diabetic and nondiabetic patients. *Diabetes* 62(11):3709–3717. <https://doi.org/10.2337/db13-0321>
 38. Hennes MM, Dua A, Kissebah AH (1997) Effects of free fatty acids and glucose on splanchnic insulin dynamics. *Diabetes* 46(1):57–62. <https://doi.org/10.2337/diab.46.1.57>
 39. Ramakrishnan SK, Khuder SS, Al-Share QY et al (2016) PPAR α (Peroxisome proliferator-activated receptor α) activation reduces hepatic CEACAM1 protein expression to regulate fatty acid oxidation during fasting-refeeding transition. *J Biol Chem* 291(15):8121–8129. <https://doi.org/10.1074/jbc.M116.714014>
 40. Poy MN, Yang Y, Rezaei K et al (2002) CEACAM1 regulates insulin clearance in liver. *Nat Genet* 30(3):270–276. <https://doi.org/10.1038/ng840>
 41. Ben-HaroushSchyr R, Al-Kurd A, Moalem B et al (2021) Sleeve gastrectomy suppresses hepatic glucose production and increases hepatic insulin clearance independent of weight loss. *Diabetes* 70(10):2289–2298. <https://doi.org/10.2337/db21-0251>
 42. WewerAlbrechtsen NJ, Junker AE, Christensen M et al (2018) Hyperglucagonemia correlates with plasma levels of non-branched-chain amino acids in patients with liver disease independent of type 2 diabetes. *Am J Physiol Gastrointest Liver Physiol* 314(1):G91–G96. <https://doi.org/10.1152/ajpgi.00216.2017>
 43. Longuet C, Sinclair EM, Maida A et al (2008) The glucagon receptor is required for the adaptive metabolic response to fasting. *Cell Metab* 8(5):359–371. <https://doi.org/10.1016/j.cmet.2008.09.008>
 44. Berglund ED, Kang L, Lee-Young RS et al (2010) Glucagon and lipid interactions in the regulation of hepatic AMPK signaling and expression of PPARalpha and FGF21 transcripts in vivo. *Am J Physiol Endocrinol Metab* 299(4):E607–614. <https://doi.org/10.1152/ajpendo.00263.2010>
 45. Smith GI, Shankaran M, Yoshino M et al (2020) Insulin resistance drives hepatic de novo lipogenesis in nonalcoholic fatty liver disease. *J Clin Invest* 130(3):1453–1460. <https://doi.org/10.1172/JCI134165>
 46. Lee W (2011) The CEACAM1 expression is decreased in the liver of severely obese patients with or without diabetes. *Diagn Pathol* 6:40. <https://doi.org/10.1186/1746-1596-6-40>
 47. Bizzotto R, Trico D, Natali A et al (2021) New insights on the interactions between insulin clearance and the main glucose homeostasis mechanisms. *Diabetes Care* 44(9):2115–2123. <https://doi.org/10.2337/dc21-0545>
 48. Piccinini F, Bergman RN (2020) The measurement of insulin clearance. *Diabetes Care* 43(9):2296–2302. <https://doi.org/10.2337/dc20-0750>
 49. Bizzotto R, Jennison C, Jones AG et al (2021) Processes underlying glycemic deterioration in type 2 diabetes: an IMI DIRECT study. *Diabetes Care* 44(2):511–518. <https://doi.org/10.2337/dc20-1567>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Authors and Affiliations

Kieran Smith^{1,2,3}  · **Guy S. Taylor**^{2,3} · **Wouter Peeters**^{2,3} · **Mark Walker**⁴ · **Simone Perazzolo**^{5,6} · **Naeimeh Atabaki-Pasdar**^{1,7,8} · **Kelly A. Bowden Davies**^{2,9} · **Fredrik Karpe**^{1,7} · **Leanne Hodson**^{1,7}  · **Emma J. Stevenson**^{2,3} · **Daniel J. West**^{2,3} 

✉ Kieran Smith
Kieran.Smith@ocdem.ox.ac.uk

✉ Daniel J. West
Daniel.West@newcastle.ac.uk

¹ Oxford Centre for Diabetes, Endocrinology and Metabolism, Churchill Hospital, University of Oxford, Oxford, UK

² Human Nutrition and Exercise Research Centre, Population Health Sciences Institute, Newcastle University, Newcastle upon Tyne, UK

³ School of Biomedical, Nutritional, and Sport Sciences, Newcastle University, Newcastle upon Tyne, UK

⁴ Translational and Clinical Research Institute, Newcastle University, Newcastle upon Tyne, UK

⁵ Nanomath LLC, Spokane, WA, USA

⁶ Department of Pharmaceutics, University of Washington, Seattle, WA, USA

⁷ NIHR Oxford Biomedical Research Centre, Oxford University Hospital Trust, Oxford, UK

⁸ Genetic and Molecular Epidemiology Unit, Lund University Diabetes Centre, Department of Clinical Science, Lund University, Malmö, Sweden

⁹ Sport and Exercise Sciences, Manchester Metropolitan University, Manchester, UK